

HUMAN EXPOSURE STUDIES OF AIRBORNE PARTICLES FROM COMMON SOURCES

Bohgard, Mats; Albin, Maria; Dierschke, Katrin; Gudmundsson, Anders; Hagerman, Inger; Isaxon, Christina; Jakobsson, Jonas; Jönsson, Bo A; Nielsen, Jörn; Nilsson, Patrik; Pagels, Joakim; Xu, Yiyi; Wierzbicka, Aneta

2014

Link to publication

Citation for published version (APA):

Bohgard, M., Albin, M., Dierschke, K., Gudmundsson, A., Hagerman, I., Isaxon, C., Jakobsson, J., Jönsson, B. A., Nielsen, J., Nilsson, P., Pagels, J., Xu, Y., & Wierzbicka, A. (2014). *HUMAN EXPOSURE STUDIES OF AIRBORNE PARTICLES FROM COMMON SOURCES*. Paper presented at 13th International Conference on Indoor Air Quality and Climate, Indoor Air 2014, Hong Kong, Hong Kong.

Total number of authors:

General rights

Unless other specific re-use rights are stated the following general rights apply:
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study
- You may not further distribute the material or use it for any profit-making activity or commercial gain
 You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: https://creativecommons.org/licenses/

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117 221 00 Lund +46 46-222 00 00

Topic C3: Nanoparticles in indoor environment

HUMAN EXPOSURE STUDIES OF AIRBORNE PARTICLES FROM COMMON SOURCES

Mats BOHGARD^{1,*}, Maria ALBIN², Katrin DIERSCHKE², Anders GUDMUNDSSON¹, Inger HAGERMAN³, Christina ISAXON¹, Jonas JAKOBSSON¹, Bo AG JÖNSSON², Jörn NIELSEN², Patrik NILSSON¹, Joakim H PAGELS¹, Yiyi XU², and Aneta WIERZBICKA¹

Keywords: Aerosols, nanoparticles, chamber studies, biomarkers

INTRODUCTION

Several epidemiological studies show correlation between exposure to airborne particles and adverse health effects. Exposure measures used are often based only on mass concentration of particles with sizes below certain limits (PM10, PM2.5, PM1, i.e. particulate matter with diameters less than 10, 2.5 and 1 μ m, respectively) regardless the shape, morphology and chemical composition of the particles. There is a need for information based on precise experimental methods in order to increase our understanding on the mechanisms behind the adverse effects and to determine the exact nature of exposures which can cause the adverse effects.

The aims of the study are i) to develop settings for experimental human exposure studies for commonly abundant indoor air particle types and ii) to experimentally determine physiological effects on human test subjects.

METHODOLOGIES

A stainless steel walk-in chamber of the volume 22 m³ is used (Figure 1). Air can only enter and leave the chamber through a well-controlled ventilation system. Systems for exposure of various commonly abundant indoor particles are being developed. Test subjects have been and are being exposed to particles from candle lights, terpene-ozone reactions, metal inert gas welding, zeolite particles from detergent residues, cooking and diesel engine in a controlled way with respect to concentration, size distribution and particle characteristics. Concentrations and size distributions are in accordance with what can be found in indoor environments where air is contaminated by particles from these sources. Test persons (three or four at the time are sitting in the chamber) are being exposed to one particle type at one occasion and clean (virtually particle-free) air at another occasion, according to a double blind protocol. A number of biomarkers from the test subjects are determined (e.g. markers of inflammation in body fluids, ECG-registration with heart rate variability (HRV) analysis and

¹Ergonomics and Aerosol Technology, Lund University, Sweden

²Division of Occupational and Environmental Medicine, Lund University, Sweden,

³Department of Cardiology, Karolinska Institute, Karolinska University Hospital, Huddinge, Sweden.

^{*}Corresponding email: Mats.Bohgard@design.lth.se

respiratory physiological parameters). Biochemical biomarkers are measured immediately after and 24 hours after exposure.

RESULTS AND DISCUSSION

Exposure studies show that effects of airborne particles from different sources on biomarkers can be studied in a well-controlled experimental model systems. Various test aerosols could be artificially created and added to the chamber air with realistic properties compared to what can be found in real environments.

Aerosols with particles from candle lights, terpene-ozone reactions, welding, diesel engine were achieved with realistic particle concentrations and characterized with respect to mass and number concentrations and chemical composition (Isaxon et al. 2013 I ; Isaxon et al. 2013 II., Pagels et al. 2009; Wierzbicka et al. 2014). Mass concentrations achieved were typically 200 $\mu g/m^3$ for candle smoke, 80 $\mu g/m^3$ for terpene-ozone generated particles, 1 mg/m^3 for welding fume and 300 $\mu g/m^3$ for diesel exhaust particles. In addition to these particle types, zeolite particles from detergent residues in textiles (Gudmundsson et al. 2007) are being generated in mass concentrations of 300 $\mu g/m^3$ and particles from frying hamburgers in concentrations of 300-400 $\mu g/m^3$.

Results on effects on biomarkers have, until today, been observed for candle light smoke: significant increase of high frequency HRV compared to clean air and compared to terpene-ozone-generated particles (Hagerman et al. 2014). Exposure to diesel exhaust caused a significant decrease in the high frequency of heart rate variability spectrum. Heart rate variability is since long a well-recognized, noninvasive, independent method, for cardiovascular risk prediction in high prevalent groups (Malik, 1996). The studies, so far show significant negative impact on HRV of diesel exhaust compared to clean air and significant positive impact of candle light smoke on HRV compared to clean air and compared to terpene-ozone generated particles.

CONCLUSION

Experimental chamber human exposure methodology have been developed in order to examine effects of exposures to common indoor airborne particles. The studies have so far shown significant negative effects on heart rate variability of diesel exhaust compared to particle-free air and significant positive effects on heart rate variability of candle light smoke compared to particle-free air and exposure to terpene-ozone generated particles, for healthy individuals.




Figure 1. Simplified sketch of the experimental set-up for the chamber exposure studies.

ACKNOWLEDGEMENT

We would like to thank all test subjects. Many thanks also to staff members Ulla BK Andersson, Eva Assarsson, Margareta Berglund, and Anna-Therese Gunnskog who participated in the exposure studies.

This study was financed by the Swedish Research Council FORMAS and performed within the framework of Metalund, the Centre for Medicine and Technology for Working Life and Society, a competence centre at Lund University, Sweden, supported by FORTE, the Swedish Research Council for Health, Working Life and Welfare. Additional support was obtained from VINNOVA - Swedish Governmental Agency for Innovation Systems.

REFERENCES

Gudmundsson A, Löndahl J & Bohgard M (2007) Methodology for identifying particle sources in indoor environments. *Journal of Environmental Monitoring*, 9(8) 831-838.

Hagerman I, Isaxon C, Gudmundsson A et al. (2014) Effects on heart rate variability by artificially generated indoor nano-sized particles in a chamber study. *Atmospheric Environment*, 88 165-171.

Isaxon C, Dierschke K, Pagels J et al. (2013) A novel system for source characterization and controlled human exposure to nanoparticle aggregates

- generated during gas-metal arc welding. *Aerosol Science and Technology*, 47(1) 52-59. I.
- Isaxon C, Dierschke K, Pagels J et al. (2013) Realistic indoor nano-aerosols for a human exposure facility. *Journal of Aerosol Science*, 60 55-66. II.
- Malik M. (1996) Heart rate variability. Standards of measurement, physiological interpretation and clinical use. Task group of the Europeans society of cardiology and the North American society of pacing and electrophysiology. *Circulation* 93, 1043-1065.
- Pagels J, Wierzbicka A, Nilsson E et al. (2009) Chemical composition and mass emission factors of candle smoke particles. *Journal of Aerosol Science*, 40(3) 193-208.
- Wierzbicka A, Nilsson P, Rissler J et al. (2014) Detailed diesel exhaust characteristics including particle surface area and lung deposited dose for better understanding of health effects in human chamber exposure studies. *Atmospheric Environment*, 86 212-219.



HUMAN EXPOSURE STUDIES OF AIROBORNE PARTICLES FROM COMMON SOURCES

MATS BOHGARD, M. ALBIN, K. DIERSCHKE, A. GUDMUNDSSON, I. HAGERMAN, C. ISAXON,J. JAKOBSSON, B.A.G JÖNSSON, J. NIELSEN, P. NILSSON, J. H. PAGELS, Y. XU, A. WIERZBICKA







1 billion nanoparticles /min.

Many studies show adverse effects of airborne particles.

Measures used are often based on mass concentrations (PM10, PM2.5, PM1, i.e. mass of particles with diameters less than 10, 2.5 and 1 μ m)

The aim is to experimentally determine physiological effects on human test subjects of particles from common sources.









ECG-recordings Beat-to-beat-variability

RR₁ RR₂ RR₃

Results

SOURCE	CONCENT Mass (μg/ m ⁻³)	Number	MEDIAN DIAMETER (nm)	NUMBER OF TEST PERSONS	CHANGE IN HF OF HRV (%)	INCREASE/ DECREASE	р
Candle	200 ± 30	8 ·10 ⁵	Bimodal 23; 270	18	+22 ± 5	↑	0.01
Terpene. Ozone	80 ± 10	3·10 ⁴	100	17	-7±5	Я	0.4
Diesel	280 ± 30	4 · 10 ⁵	90	18	-15 ± 5	\	0.04