

### Molecular and anatomical evidence for a three-way association between Pinus sylvestris and the ectomycorrhizal fungi Suillus bovinus and Gomphidius roseus

Olsson, Pål Axel; Munzenberger, B; Mahmood, Shahid; Erland, Susanne

Published in: Mycological Research

10.1017/S0953756200002823

2000

### Link to publication

Citation for published version (APA):

Olsson, P. A., Munzenberger, B., Mahmood, S., & Erland, S. (2000). Molecular and anatomical evidence for a three-way association between Pinus sylvestris and the ectomycorrhizal fungi Suillus bovinus and Gomphidius roseus. Mycological Research, 104(11), 1372-1378. https://doi.org/10.1017/S0953756200002823

Total number of authors:

#### General rights

Unless other specific re-use rights are stated the following general rights apply: Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights

- Users may download and print one copy of any publication from the public portal for the purpose of private study
- You may not further distribute the material or use it for any profit-making activity or commercial gain
   You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: https://creativecommons.org/licenses/

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**LUND UNIVERSITY** 

PO Box 117 221 00 Lund +46 46-222 00 00

Download date: 23. Oct. 2025

## Molecular and anatomical evidence for a three-way association between *Pinus sylvestris* and the ectomycorrhizal fungi *Suillus bovinus* and *Gomphidius roseus*

#### Pål Axel OLSSON<sup>1</sup>, Babette MÜNZENBERGER<sup>2</sup>, Shahid MAHMOOD<sup>1</sup> and Susanne ERLAND<sup>1</sup>

- <sup>1</sup> Department of Microbial Ecology, Ecology Building, Lund University, SE-223 62 Lund, Sweden.
- <sup>2</sup> Institute of Primary Production and Microbial Ecology, Dr.-Zinn-Weg 18, D-16225 Eberswalde, Germany. E-mail: pal\_axel.olsson@mbioekol.lu.se

Received 28 October 1999; accepted 14 February 2000.

Many intimate associations between different species of ectomycorrhizal fungi are inferred on the basis of the consistent cooccurrence of their fruit bodies. *Suillus bovinus* and *Gomphidius roseus*, where the latter never occurs without the former, is one
example. This association was examined with PCR identification and light microscopy. *S. bovinus* and *G. roseus* were unambiguously
separated on the basis of RFLPs of the PCR-amplified ITS region of ribosomal DNA. Tuberculate mycorrhizas of *Pinus sylvestris*sampled under fruit bodies of *G. roseus* and *S. bovinus* were investigated and the majority were identified as mixed associations
involving both *G. roseus* and *S. bovinus*. Tuberculate mycorrhizas, which macroscopically resemble the ones of *Suillus* species,
contained typical chlamydospores of *G. roseus* and they had haustoria where *G. roseus* hyphae penetrated the cortical root cells. Pine
seedlings collected near the fruit bodies of the two species were mainly colonised by *S. bovinus*. Mycelial rhizomorphs collected
under the fruit bodies of *G. roseus* were identified as *S. bovinus*, while both fungal species were present at the base of *G. roseus* fruit
bodies. The significance of these observations and the possibility that *G. roseus* acts as a parasite are discussed.

#### **INTRODUCTION**

Some ectomycorrhizal fungi show consistent associations with each other as revealed by co-occurrence of fruit bodies and in mycorrhizal root tips, demonstrated by morphological observations (Brand 1992, Agerer 1992). An association between Suillus species and members of the Gomphidiaceae has for long been suspected because several Gomphidiaceae species only occur where their Suillus 'partner' is present. Recently studied examples involve Gomphidius roseus with Suillus bovinus, Chroogomphus rutilus with S. granulatus, and C. helveticus with different suilloid fungi (Agerer 1991). The 'mixed' mycorrhizas of these fungi show the typical appearance of mycorrhizas formed by Suillus species, but those of Gomphidiaceae may form haustoria in the cortical cells of the root (Brand 1992, Agerer 1990, 1991). The genera Suillus and Gomphidius are now considered to be closely related, both belonging to the group of suilloid fungi (Bruns

Suillus bovinus and G. roseus both have narrow host ranges among ectomycorrhizal plants and typically form mycorrhizas with members of the *Pinaceae* (Molina *et al.* 1992). Agerer (1990, 1991) described the morphology of the *S. bovinus—G. roseus—P. sylvestris* mycorrhiza. PCR-amplification of the internal transcribed spacer (ITS) region of the ribosomal DNA and subsequent digestion with endonucleases is a method

commonly used for the identification of mycobionts in single mycorrhizas (Erland 1995, Gardes & Bruns 1996, Dahlberg et al. 1997, Erland et al. 1999, Jonsson et al. 1999, Mahmood, Finlay & Erland 1999). The ITS-region has proved variable enough to separate most ectomycorrhizal species investigated, but conserved enough not to show much intraspecific variation, at least on a regional level (Kårén et al. 1997).

The aims of this study were to: (1) apply PCR-identification to study the interaction between *S. bovinus* and *G. roseus* in field-collected mycorrhizas and fungal rhizomorphs; (2) establish the frequency of detection of a three-way association between *S. bovinus*, *G. roseus* and *P. sylvestris* in mycorrhizas under fruitbodies of the two fungal species; (3) determine the identity of fungal rhizomorphs connected to fruitbodies and mycorrhizas of *S. bovinus* and *G. roseus*; and (4) investigate the anatomy of the three-way association between *S. bovinus*, *G. roseus* and *P. sylvestris*.

#### MATERIALS AND METHODS

#### Sampling sites and sampling technique

Roots and fungal material were sampled at three sites for molecular identification (site 1–3) and at one site for microscopic studies (site 4). Sampling was performed on 1–4

different points at each site where *G. roseus* fruit bodies appeared within 10 cm of *S. bovinus* fruit bodies.

Site 1: A moraine-filled path in a *Picea abies* forest with few *Pinus sylvestris* trees in southern Sweden (56° 23′ N, 14° 20′ E, September 1997).

Site 2: A P. abies forest with few P. sylvestris in southern Sweden (56° 29′ N, 14° 13′ E, October 1997) with a ground vegetation of Vaccinium myrtillus and Deschampsia flexuosa.

Site 3: A moraine-filled path in a mixed *P. abies* and *P. sylvestris* forest in southern Sweden (56° 15′ N, 14° 04′ E, August 1998).

Site 4: A pure stand of *P. sylvestris* on a dystric cambisol (derived from glacial outwash sediments) in northern Germany with a ground vegetation of *Vaccinium myrtillus* and *Deschampsia flexuosa* (52° 49′ N, 13° 50′ E, October 1993).

#### Sample preparation for PCR identification

Soil cores of  $10 \times 20 \times 7.5$  cm (width  $\times$  length  $\times$  depth), which at the surface contained fruit bodies of G. roseus and S. bovinus, were collected at sites 1-3 (4 cores at site 1; 4 cores at site 2; 1 core at site 3). The cores were transferred to the laboratory in plastic bags for sampling of mycorrhizal roots, fungal rhizomorphs and fruit bodies under dissecting microscope within 24 h. The soil cores from sites 1 and 3 had a poorly developed organic soil layer and therefore mycorrhizal root tips and rhizomorphs occurred mainly in the mineral soil. In the soil cores from site 2, mycorrhizal root tips and rhizomorphs occurred at the interface between the organic and the mineral soil layers. Samples were rinsed in water and stored frozen (-80 °C) until DNA extraction. Tuberculate mycorrhizas consisting of 5-20 root tips (see Treu 1990) were divided into fragments containing 3-5 root tips that were used for PCR identification.

#### DNA extraction

DNA was extracted from thawed mycorrhizas, rhizomorphs and fruitbody tissue following the procedures described by Erland et al. (1994) with slight modifications. Six hundred μl of extraction buffer (200 mм Tris-HCl pH 8.5, 250 mм NaCl, 25~mM EDTA,  $0.5\,\%$  SDS,  $0.2\,\%$  mercaptoethanol) and 10~µlof proteinase K (10 mg ml<sup>-1</sup>) were added to samples and incubated for 15 min at 65° for cellular lysis. The tissues were crushed with pestles and sand in Eppendorf tubes. The tubes were frozen in liquid nitrogen and thawed at 65° three times and then incubated at 65° for one hour. The samples were centrifuged for 10 min at 16000 g to remove cell debris. The aqueous phase was extracted first with 500 µl of phenol/ chloroform/isoamylalcohol (25/24/1, v/v/v), then with 500 µl of chloroform/isoamylalcohol (24/1, v/v) to remove the residual phenol after centrifugation at 16000 g for 10 minfor each extraction. The DNA was precipitated at  $-20^\circ$  for 1 h in cold ethanol and pelleted by centrifugation at 16 000 gfor 30 min. The pellets were washed again with 500 µl of 70% ethanol and centrifuged at 16000 g for 10 min. The pellets were left to air dry at room temperature, dissolved in 50 µl Tris-EDTA buffer (1 mm EDTA, 10 mm Tris-HCl; pH 8.0),

and stored at  $-20^{\circ}$ . The DNA concentration of the extracts was determined spectrophotometrically at 260 nm.

1373

#### PCR identification

The primers used for PCR were: ITS1 and ITS4 (White *et al.* 1990) to amplify the ITS region of the rDNA. The 50  $\mu l$  reaction mixture for PCR contained the following final concentrations or total amounts: 500 pmol of each primer, 20 mm Tris (pH 8.3), 200  $\mu m$  dNTP, 1.5 mm MgCl $_2$ , 50 mm KCl and 2.5 units of Taq polymerase (Perkin–Elmer). To this mixture about 10 ng of DNA from fruitbodies or mycelial rhizomorphs or between 10 and 60 ng of DNA from mycorrhizas was added. PCR-amplification was performed with one cycle of 10 min at 94°, followed by 35 cycles of 30 s at 94°, 30 s at 53°, 1.5 min at 72° and one cycle of 10 min at 72°.

RFLP analysis of the PCR products was carried out with the restriction endonucleases HinfI and TaqI (Boehringer–Mannheim) by digesting 18  $\mu$ l of PCR product with 2  $\mu$ l of enzyme cocktail for 4 h, according to the manufacturers recommendations. The PCR products and their restriction fragments were visualised using 1% agarose + 1% Nusieve agarose gels stained with ethidium bromide.

#### Microscopical investigation


Tuberculate mycorrhizas from site 4 were fixed for microscopy and then divided into smaller pieces, which were embedded acording to Münzenberger, Kottke & Oberwinkler (1992). Semi-thin sections (0.5  $\mu$ m) of the mycorrhizal root tips were cut on an Ultracut Reichert ultramicrotome. The sections were stained with New Fuchsin-Crystal Violet (solution with 0.3 g New Fuchsin, 0.3 Crystal Violet, 40 ml glycerol and 60 ml sodium tetraborate; Merck) for light microscopy.

#### **RESULTS**

# PCR amplification and identification of DNA from S. bovinus and G. roseus

PCR amplification of the ITS-region from fruit bodies of *S. bovinus* and *G. roseus* gave products of 700 bp and 780 bp respectively (Fig. 1). Cleaving with the restriction enzymes *Hinf*I and *Taq*I, both gave distinctive RFLP patterns by which the two species could be distinguished (Fig. 1).

When between 5 and 70 ng total genomic DNA from mycorrhizal roots were used for PCR amplification, mixed *G. roseus* and *S. bovinus* DNA was detected over the whole range. DNVA-extracts from fruit bodies of *S. bovinus* and *G. roseus* were mixed in different ratios to get information about the proportions needed for amplification of the ITS region of both species simultaneously. When 50 ng of mixed DNA was used for PCR amplification, DNA of each species could be detected if constituting 1% of the total DNA. When the total amount of DNA was reduced to 5 ng, DNA of *G. roseus* was usually amplified if constituting at least 7% of the mixture and DNA from *S. bovinus* was amplified if constituting at least 4%. This



**Fig. 1.** Agarose gel electrophoresis of PCR amplified ribosomal DNA (ITS region) from 1. fruitbody of *G. roseus*, 2. mycorrhizal root tip containing both *G. roseus* and *S. bovinus* and 3. fruitbody of *S. bovinus*. Bands either show the uncut PCR product or the RFLP pattern after digestion with *Hinf*I or *Taq*I.

shows that in a mixture of *G. roseus* and *S. bovinus*, less *S. bovinus* total genomic DNA is needed to amplify the ITS-region of that fungus.

#### PCR identification of tuberculate mycorrhizas

Pinus sylvestris tuberculate mycorrhizas, which resembled Suillus mycorrhizas, were found under the fruitbodies of G. roseus. In total, 10 tuberculate mycorrhizas originating from sites 1–3 were used for PCR identification (Table 1). Six of these were identified as three-way associations between P. sylvestris, G. roseus, and S. bovinus (Fig. 1). Three-way associations were found at all three sites investigated. The tubercules consisted of 5–20 Y-shaped mycorrhizal root tips. Three of the tubercules were identified as S. bovinus and from one tubercule no DNA could be amplified. From some of the tuberculate mycorrhizal systems with both S. bovinus and G. roseus, several subsamples from the same tubercule were taken and these consistently showed PCR-products of both fungi.

#### Microscopical investigation of tuberculate mycorrhizas

Tuberculate mycorrhizas are formed by many root tips and enveloped by a common hyphal mantle. Fungal rhizomorphs originated from between the root tips in the tuberculate mycorrhizas (Fig. 2). The rhizomorphs had enlarged vessel hyphae surrounded by smaller hyphae. The individual root tips within the tubercules were also enveloped by hyphal mantles, which connected them (Fig. 3).

The hyphae of *G. roseus* within the tuberculate mycorrhizas penetrated the root cell wall and formed haustoria in cortical cells of the outer cortex (Figs 4–7). Several sections indicate that only a few cells of the cortex are infected at the same time. The presence of *G. roseus* was also evident from the occurrence of its characteristic chlamydospores (Agerer 1990). These were present at various depths in the hyphal mantle (Fig. 8).

#### Pine seedlings

Three one year-old pine seedlings were removed from the soil cores sampled at site 1. The mycorrhizal root tips of these seedlings were subjected to PCR-based identification (Table 1). Two of the seedlings were mainly colonized by *S. bovinus* according to PCR identification. One of these seedlings had small tuberculate mycorrhizas consisting of 2 to 6 root tips. DNA was successfully amplified from 7 out of 12 of these tubercules of which five were *S. bovinus*, one a mixture of *S. bovinus* and *G. roseus* and one was unidentified. Seven root tips with *S. bovinus* were identified in the second seedling out of 15 single mycorrhizas. DNA amplification from the third seedling was only successful for one out of 17 root tips, which showed the RFLP pattern of *S. bovinus*. Rhizomorphs connected to the seedlings were always identified as *S. bovinus* when DNA amplification was successful (Table 1).

#### Mycelial rhizomorphs in soil

White rhizomorphs were commonly found in close proximity to the fruit bodies of *S. bovinus* and *G. roseus*. All rhizomorphs from which DNA was successfully amplified were identified as *S. bovinus* (Table 1). *S. bovinus* rhizomorphs were found under *G. roseus* at all three sites. One rhizomorph also contained DNA from an unidentified fungal species. DNA of *G. roseus* was not detected in any of the rhizomorphs.


Mycelium attached to the base of G. roseus fruit bodies

**Table 1.** PCR-identification of samples of root and mycelium collected under fruit bodies of *S. bovinus* and *G. roseus* at three sites in southern Sweden. The total number of samples analyzed is given and the identity of material where PCR amplification was positive is indicated. Mixed associations refer to samples where both *S. bovinus* and *G. roseus* were identified.


Type of sample	Total number	Mixed association	Suillus bovinus	Unidentified
Mature pine trees				
Tuberculate mycorrhizas	10	6	3	_
Rhizomorphs connected to tuberculate mycorrhizas	2	_	1	<del></del>
One-year old pine seedlings				
Mycorrhizal root tips	44	1	13	1
Rhizomorphs	12	_	3	<del></del>
Rhizomorphs under fruit bodies				
At the base of G. roseus fruit bodies	4	4	_	<del></del>
Under G. roseus	20	_	6	1*
Under S. bovinus	6	_	2	_

<sup>\*</sup> Detected on the same root tip as S. bovinus.


P. A. Olsson and others



Figs 2–3. Section through a *Pinus sylvestris–Suillus bovinus–Gomphidius roseus* tuberculate mycorrhiza. Fig. 2. A differentiated rhizomorph can be seen (Rm, arrows) between two mycorrhizal tips; note the randomly distributed enlarged hyphae. Fig. 3. Two mycorrhizal root tips connected by a thick hyphal mantle (Hm) containing numerous vacuolar bodies. The Hartig net (Hn) is well established showing finger-like structures (arrow). Bar  $= 50 \mu m$ .



Figs 4–7. Section through a tuberculate mycorrhizal system of Pinus sylvestris–Suillus bovinus–Gomphidius roseus. Fig. 4. A penetrating intracellular hypha of Gomphidius roseus in an outer cortical cell (arrow). Bar =  $50 \mu m$ . Fig. 5. Enlarged section of the penetrated cortical cell (represented as a rectangle in Fig. 4); the invading hypha (arrow) surrounded by the tonoplast of the vacuole is visible. Bar =  $20 \mu m$ . Fig. 6. A cortical cell filled with intracellular hyphae (arrow); note the irregularly shaped nucleus. Bar =  $50 \mu m$ . Fig. 7. Another cortical cell showing a haustorium of Gomphidius roseus. Bar =  $50 \mu m$ .



**Fig. 8.** Section through a mycorrhizal root tip of a tuberculate mycorrhizal system of *Pinus sylvestris–Suillus bovinus–Gomphidius roseus* revealing chlamydospores of *Gomphidius roseus* of different age (arrows) embedded in the hyphal mantle. Bar = 50 µm.

produced both *G. roseus* and *S. bovinus* PCR products (Table 1).

#### DISCUSSION

Here we show that the tuberculate mycorrhizal system is an important structure in the interaction between the ectomycorrhizal fungi *S. bovinus* and *G. roseus*. Tuberculate *P. sylvestris* mycorrhizas were analyzed from four different sites and interactions were detected at all sites. Chlamydospores and haustoria were identified as structures of *G. roseus* inside tuberculate mycorrhizas of *S. bovinus* (see also Agerer 1990, 1991), which otherwise show the morphological structure of *Suillus* mycorrhizas. *S. bovinus* colonization was also present on *P. sylvestris* seedlings near the fruit bodies sampled, but here, a mixed association was only observed once.

The distribution of S. bovinus was successfully tracked by PCR identification and the fungus was present in the mycorrhizas, in the rhizomorphs originating from the mycorrhizas, and rhizomorphs under fruit bodies of both fungal species. The presence of G. roseus was only detected in the S. bovinus mycorrhizas and as fruit bodies. Similarly, Agerer (1990) could not identify G. roseus in rhizomorphs connected to mycorrhizas with both species. However, since S. bovinus was present on the stipe of G. roseus fruit bodies, it is likely that G. roseus hyphae are present in, or connected to, S. bovinus rhizomorphs and initiate fruit body formation from these. If present in an amount lower than 7% of the total fungal DNA, G. roseus DNA would not be detected by our method, as indicated from the results where the proportions needed for amplification of both species simultaneously were tested. In an interaction that may be similar to the one described here, Agerer (1991) found rhizomorphs of an

unidentified *Suillus* species attached to the fruit bodies of *Gomphidius glutinosus*. Agerer (1991) also found *Rhizopogon* hyphae in the stipe of *G. glutinosus* and concluded that *Rhizopogon* can grow into the stipe of this fungus. Another explanation could be that the *G. glutinosus* fruit bodies are formed from hyphae associated with *Rhizopogon* rhizomorphs and that these are included in the fruit body by a purely physical interaction.

The test of PCR amplification of in vitro mixed DNA from S. bovinus and G. roseus indicates that the amount of G. roseus DNA in the tuberculate mycorrhizas may have been considerable. According to our results, over 7% of the DNA should have been of *G. roseus* origin for amplification to occur. In the mycorrhizas, plant DNA as well as substances inhibiting the PCR reaction may also have influenced the results, making the detection of both species at low concentration more difficult. In some cases the PCR products of G. roseus were of equal brightness to the PCR products of S. bovinus, whilst in other cases those of S. bovinus were brighter (see Fig. 1). Chlamydospores may make a major contribution to the DNA of G. roseus in the mycorrhizas. The fact that G. roseus was not detected in rhizomorphs except at the base of its fruitbodies in this study, indicates that if present in the rhizomorphs they have a low biomass. Hyphae of G. roseus (which are amyloid in contrast to Suillus hyphae) were not found in the rhizomorphs of S. bovinus in microscopical investigations (Agerer 1990).

Our finding that in soil *G. roseus* never occurred without *S. bovinus*, while *S. bovinus* often occurred alone, suggests that *S. bovinus* is a host for *G. roseus*. Agerer (1991) found that *G. roseus* could not be isolated unless co-cultivated with *S. bovinus*, but once established on the growth medium was it possible to grow pure cultures of the fungus. Mycorrhizas of

*G. roseus* have been observed but they seem to be very infrequent (Agerer 1994). The close association of *G. roseus* with *S. bovinus*, the formation of haustoria in the plant root cells and the observation that *G. roseus* does not seem to form an extensive soil mycelium, suggests that *G. roseus* acts as a parasite on either the fungal host, the plant host, or both. Further studies of these species in experimental systems may reveal their true functional relationships.

In studies where PCR identification of mycorrhizas is applied to study community structure of ectomycorrhizal fungi, it is relatively common that two PCR products are yielded from the same mycorrhiza. In some studies up to 20% of root tips are colonized by two fungi (Erland 1995, Erland et al. 1999). This may be due to a succession of fungal species or to competition between different fungi to colonize the root. It may also be due to close associations between species such as described here for S. bovinus and G. roseus. Suilloid fungi are very rarely detected on mycorrhizal roots in community structure studies even when their fruit bodies are abundantly present (Gardes & Bruns 1996). It is therefore difficult to obtain information from the literature on how often mycorrhizas formed by Suillus species are associated with other fungi. Future studies focusing on mycorrhizas connected to fruit bodies may reveal new types of close interactions between species of ectomycorrhizal fungi.

#### **ACKNOWLEDGEMENTS**

Financial support from the Swedish Council for Forestry and Agricultural Research is gratefully acknowledged. We thank Ingvar Månsson and Ulf Olsson for information about sites for the fungal species.

#### REFERENCES

- Agerer, R. (1990) Studies on ectomycorrhizae XXIV. Ectomycorrhizae of Chroogomphus helveticus and C. rutilus (Gomphidiaceae, Basidiomycetes) and their relationship to those of Suillus and Rhizopogon. Nova Hedwigia 50: 1–63.
- Agerer, R. (1991) Studies on ectomycorrhizae XXXIV. Mycorrhizae of *Gomphidius glutinosus* and of *G. roseus* with some remarks on *Gomphidiaceae* (*Basidiomycetes*). *Nova Hedwigia* **53**: 127–170.
- Agerer, R. (1992) Studies on ectomycorrhizae XLIV. Ectomycorrhizae of Boletopsis leucomelaena (Thelephoraceae, Basidiomycetes) and their relationship to an unidentified ectomycorrhiza. Nova Hedwigia 55: 501–518.
- Agerer, R. (1994) Ectomycorrhizae in the fungal community: with special emphasis on interactions between ectomycorrhizal fungi. In Mycorrhizas in

- Integrated Systems from Genes to Plant Development (C. Azcon-Aguilar & J. M. Barea, eds): 52–57. Fourth European Symposium on Mycorrhizas, Granada
- Brand, F. (1992) Mixed associations of fungi in ectomycorrhizal roots. In Mycorrhizas in Ecosystems (I. J. Alexander, A. H. Fitter, D. H. Lewis & D. J. Read, eds), pp. 142–147. CAB International, Wallingford.
- Bruns, T. D., Szaro, T. M., Gardes, M., Cullings, K. W., Pan, J. J., Taylor, D. L., Horton, T. R., Kretzer, A., Garbeletto, M. & Li, Y. (1998) A sequence database for the identification of ectomycorrhizal basidiomycetes by phylogenetic analysis. *Molecular Ecology* 7: 257–272.
- Dahlberg, A., Jonsson, L. & Nylund, J.-E. (1997) Species diversity and distribution of biomass above- and below-ground among ectomycorrhizal fungi in an old-growth Norway spruce forest in south Sweden. *Canadian Journal of Botany* 75: 1323–1335.
- Erland, S. (1995) Abundance of Tylospora fibrillosa ectomycorrhizas in a South Swedish spruce forest measured by RFLP analysis of the PCR-amplified rDNA ITS-region. Mycological Research 99: 1425–1428.
- Erland, S., Henrion, B., Martin, F., Glover, L. A. & Alexander, I. J. (1994) Identification of the ectomycorrhizal basidiomycete *Tylospora fibrillosa* Donk by RFLP analysis of the PCR-amplified ITS and IGS regions of ribosomal DNA. *New Phytologist* **126**: 525–532.
- Erland, S., Jonsson, T., Mahmood, S. & Finlay, R. D. (1999) Below-ground ectomycorrhizal community structure in two *Picea abies* forests in southern Sweden. *Scandinavian Journal of Forest Research* 14: 193–198.
- Gardes, M. & Bruns, T. D. (1996) Community structure of ectomycorrhizal fungi in a *Pinus muricata* forest: above- and below-ground views. *Canadian Journal of Botany* 74: 1572–1583.
- Jonsson, T., Kokalj, S., Finlay, R. D. & Erland, S. (1999) Ectomycorrhizal community structure in a limed spruce forest. Mycological Research 103: 501–508.
- Kårén, O., Högberg, N., Dahlberg, A., Jonsson, L. & Nylund, J. E. (1997) Interand intraspecific variation in the ITS-region of rDNA of ectomycorrhizal fungi in Fennoscandia as detected by endonuclease analysis. *New Phytologist* 136: 313–326.
- Mahmood, S., Finlay, R. D. & Erland, S. (1999) Effects of repeated harvesting of forest residues on the ectomycorrhizal community in a Swedish spruce forest. New Phytologist 142: 577–585.
- Molina, R., Massicotte, H. B. & Trappe, J. M. (1992) Ecological role of specificity phenomena in ectomycorrhizal plant communities: potentials for interplant linkages and guild development. In *Mycorrhizas in Ecosystems* (I. J. Alexander, A. H. Fitter, D. H. Lewis & D. J. Read, eds): 106–112. CAB International, Wallingford.
- Münzenberger, B., Kottke, I. & Oberwinkler, F. (1992) Ultrastructural investigations of *Arbutus unedo-Laccaria amethystea* mycorrhiza synthesized in vitro. *Trees* 7: 40–47.
- Treu, R. (1990) Suillus plorans. In Colour Atlas of Ectomycorrhizae (R. Agerer, ed.): pl. 46. Einhorn Verlag, Schwäbisch Gmünd.
- White, T. J., Bruns, T., Lee, S. & Taylor, J. (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In *PCR Protocols: a guide to methods and applications* (M. A. Innis, D. H. Gelfand, J. J. Sninsky & T. J. White, eds): 315–322. Academic Press, San Diego.