



# LUND UNIVERSITY

## Time to extinction in relation to mating system and type of density regulation in populations with two sexes

Saether, BE; Engen, S; Lande, R; Moller, AP; Bensch, Staffan; Hasselquist, Dennis; Beier, J; Leisler, B

*Published in:*  
Journal of Animal Ecology

*DOI:*  
[10.1111/j.0021-8790.2004.00869.x](https://doi.org/10.1111/j.0021-8790.2004.00869.x)

2004

[Link to publication](#)

### *Citation for published version (APA):*

Saether, BE., Engen, S., Lande, R., Moller, AP., Bensch, S., Hasselquist, D., Beier, J., & Leisler, B. (2004). Time to extinction in relation to mating system and type of density regulation in populations with two sexes. *Journal of Animal Ecology*, 73(5), 925-934. <https://doi.org/10.1111/j.0021-8790.2004.00869.x>

*Total number of authors:*  
8

### **General rights**

Unless other specific re-use rights are stated the following general rights apply:  
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117  
221 00 Lund  
+46 46-222 00 00

# Time to extinction in relation to mating system and type of density regulation in populations with two sexes

BERNT-ERIK SÆTHER\*, STEINAR ENGEN†, RUSSELL LANDE‡,  
ANDERS PAPE MØLLER§, STAFFAN BENSCH¶, DENNIS HASSELQUIST¶,  
JOSEF BEIER\*\* and BERND LEISLER\*\*

\*Institute for Biology, Norwegian University of Science and Technology, Realfagsbygget, N-7491 Trondheim, Norway; †Department of Mathematical Sciences, Norwegian University of Science and Technology, N-7491 Trondheim, Norway; ‡Division of Biology 0116, University of California at San Diego, La Jolla, CA 92093, USA; §Laboratoire Parasitologie Evolutive, CNRS UMR 7103, Université Pierre et Marie Curie, Bat. A, 177, quai St Bernard, Case 237, F-75252 Paris Cedex 5, France; ¶Department of Animal Ecology, Ecology Building, Lund University, S-223 62 Lund, Sweden; and \*\*Max Planck Research Centre for Ornithology, Vogelwarte Radolfzell, Schlossallee 2, D-78315 Radolfzell, Germany

## Summary

1. Population viability models usually consider only the female segment of the population. However, random variation in sex ratio as well as the mating system may also affect variation in fitness among females. Here we develop population models incorporating demographic stochasticity in both sexes. Furthermore, we consider the effects on the estimated time to extinction and whether density regulation acts only on females or on total population size.

2. We applied these models to two populations of polygynous great reed warblers *Acrocephalus arundinaceus* L. with differences in population trends to investigate the importance of considering sex in population viability models.

3. Demographic stochasticity was larger in a polygynous than in a monogamous mating system.

4. The estimated time to extinction was considerably shorter for a monogamous than for a polygynous mating system, particularly if density regulation acted only on females than rather on the total population.

5. This study demonstrates that structure of mating system must be included when making population viability analysis based on counts of total population sizes. It is especially important to model the specific effects of density regulation on the two sexes.

*Key-words:* demographic stochasticity, density regulation, mating system, sex-specific population dynamics, great reed warbler.

*Journal of Animal Ecology* (2004) **73**, 925–934

## Introduction

Analyses of population dynamics or demography usually consider only the female segment of the population (Caswell 2001). It is assumed implicitly that there are a sufficient number of males available for the females to find an appropriate mate, and that the numbers of males do not affect the dynamics or the demographic characteristics of the females. However, several lines of

evidence do indicate that this assumption may not always be valid. One of the most extreme examples of how processes affecting the numbers of each sex may be strongly interrelated is provided by infanticide that occurs when the death of the male defending or controlling the females in an area leads to severe reduction in the reproductive success, because the new incoming male will kill as many offspring fathered by the previous male as possible (Hausfater & Hrdy 1984; Swenson *et al.* 1997). More indirect effects of variation in the proportion of males also may be present. In the moose *Alces alces* L. differences in the adult sex ratio explained a large proportion of both temporal and spatial variation in several reproductive traits (Solberg

Correspondence: Bernt-Erik Sæther, Institute for Biology, Norwegian University of Science and Technology, Realfagsbygget, N-7491 Trondheim, Norway. Fax: +4773591309; E-mail: Bernt-Erik.Sather@bio.ntnu.no

et al. 1999; Sæther, Solberg & Heim 2003; Sæther et al. 2004). Similar effects have also been recorded in red deer *Cervus elaphus* L. (Clutton-Brock et al. 2002). These examples suggest that in many cases it is important to consider the effects of both sexes on demographic processes in a population (Mysterud, Coulson & Stenseth 2002).

In some populations many individuals fail to obtain a mate (Newton 1998). Dale (2001) showed that in many small isolated bird populations a high proportion of males was unmated, probably because of low return rates of females due to female-biased dispersal in birds (Clarke, Sæther & Røskaft 1997). Similarly, in several bird species a proportion of female floaters may also be present (Sæther & Fonstad 1981; Smith & Arcese 1989; Newton 1998). The effects of a surplus of one sex in the population will depend on the mating system. In a strictly monogamous system the lack of males will immediately reduce the reproductive output of females. In contrast, in a polygynous mating system these effects are likely to be greatly reduced because a single male can fertilize multiple females (Emlen & Oring 1977). However, the intensity of sexual selection measured by the adult sex ratio in the population may also have other less evident consequences including effects on the reproductive rate of females (see reviews in Møller & Legendre 2001; Kokko & Brooks 2003; Møller 2003).

Demographic stochasticity is random variation among individuals in fitness (Lande, Engen & Sæther 2003). Theoretical analyses have demonstrated that demographic stochasticity has a major impact on the dynamics of small populations on the order of 100 individuals or less (Lande 1998; Brook, Burgman & Frankham 2000; Lande et al. 2003). However, some evidence suggests that once sex and mating system are incorporated into the models, significant effects of sex ratios on extinction risk may occur even at much larger populations (Legendre et al. 1999). Demographic stochasticity will therefore strongly affect the mean time to extinction (Lande 1993; Sæther & Engen 2003). Engen, Lande & Sæther (2003) showed in a population model with two sexes that the demographic variance  $\sigma_d^2$  governing fluctuations in population size in a constant environment could be partitioned into three components. The first component arises from variation among individual females in survival and reproduction of both sexes ( $\sigma_{df}^2$ ). We have shown previously that in small passerine birds this component ranges from 0.20 to 0.65 (Sæther et al. 1998, 2000a,b; 2002a, in press; Tufto et al. 2000) when only female offspring are considered. The second component is generated by the binomial distribution of the number of surviving males ( $\sigma_{dm}^2$ ). Finally, the third component  $\sigma_{sex}^2$  is due to stochastic fluctuations in the adult sex ratio. Thus, in a population with two sexes the demographic variance can be decomposed as:

$$\sigma_d^2 = q\sigma_{df}^2 + (1-q)\sigma_{dm}^2 + \sigma_{sex}^2, \quad \text{eqn 1}$$

where  $q$  is the expected sex ratio in the breeding population,  $s$  the adult survival rate and  $\sigma_{dm}^2 = s(1-s)$ . Engen et al. (2003) showed that the sex ratio component is especially large at small population sizes due to large sex ratio fluctuations. Temporal autocorrelations in the sex ratio, which may be substantial with long generation times, also will have a large effect on  $\sigma_{sex}^2$ .

Allee effects that reduce population growth rates at small population sizes (Allee et al. 1949) can strongly affect population dynamics and may cause rapid extinction of small populations (Dennis 1989, 2002). Lande (1998) demonstrated that demographic stochasticity by itself could induce a stochastic Allee effect that will cause most sample paths in small populations to decrease. Furthermore, stochastic fluctuations in the sex ratio, resulting in a lack of mates, may also lead to reduced reproductive rate at low population sizes and have direct negative effects on the mean population growth rate. Because demographic stochasticity operates differently in monogamous and polygynous mating systems (Engen et al. 2003), this may indicate that the risk of extinction at small population sizes may be related to the mating system.

In addition to changing the influence of the stochastic factors, using the total population size may also affect the deterministic component of the population dynamics (e.g. the strength and form of density dependence). In most demographic or population dynamical analyses density dependence is considered only within the female segment of the population. For instance, in many polygynous mammals the males visit the females only during a short period during the rut (Dubois et al. 1996). However, in many cases the number of males will also affect the demographic characteristics of the females. For instance, in many temperate ungulates, changes in age and sex-composition affect the demography and the population dynamics in complicated ways (Coulson et al. 2001; Clutton-Brock & Coulson 2002; Clutton-Brock et al. 2002; Mysterud et al. 2002).

The purpose of the present study is to partition the demographic variance into the three components in two populations of the great reed warbler *Acrocephalus arundinaceus* L. with different dynamical characteristics. We will then examine how assumptions regarding the structure of the mating system affect the stochastic components of the population dynamics, dependent on whether the density regulation operates on total population size or only within the female segment of the population. To obtain a proper understanding of magnitude of the effects of mating system our major comparison involves the two extremes of a strictly monogamous and a polygynous mating system. We explore how this affects the estimated time to extinction of the two populations. This will enable us to examine how mating system and form of density regulation interact to influence the viability of small bird populations.

**Population model****DENSITY-DEPENDENCE ACTING ON TOTAL  
POPULATION SIZE**

Let  $N$  and  $X$  be the total population size and the number of females, respectively. Then, assuming that the density regulation depends on the total population size and acts only on the fecundity so that mean fecundity for a given environment  $\mathbf{Z}$  is a function of the population size,  $f(N, \mathbf{Z})$ . Writing  $\bar{f}(N)$  for the mean of  $f(N, \mathbf{Z})$  over temporal environmental fluctuations in  $\mathbf{Z}$ , the carrying capacity is determined by  $\bar{f}(K) = 0$ . For this type of density regulation Engen *et al.* (2003) showed, approximating the coloured noise process by a process with white noise, that in a polygynous mating system

$$\sigma_{sex}^2 = q(1-q)\bar{f}^2(1+s)/(1-s). \quad \text{eqn 2}$$

In a monogamous mating system the effects of sex ratio fluctuations become more complicated because fluctuations in the sex ratio may also create an Allee effect, that is, an unstable equilibrium at some small population size (Dennis 1989, 2002; Lande 1998). Engen *et al.* (2003) showed in a monogamous mating system that:

$$\sigma_{sex}^2 = \frac{1}{4}\bar{f}^2\left(1 - \frac{2}{\pi}\right)G_\epsilon \approx 0.0908\bar{f}^2G_\epsilon. \quad \text{eqn 3}$$

Here  $G_\epsilon = 1 + 2\sum_{h=1}^{\infty} \rho_\epsilon(h)$ , where  $\rho_\epsilon(h)$  is the correlation function for the noise, required for correction of the white noise approximation.

**DENSITY REGULATION ACTING WITHIN THE  
FEMALE SEGMENT OF THE POPULATION**

To analyse the effects of density regulation operating only in the female segment of the population we consider the model obtained by substituting  $f(X/q, \mathbf{Z})$  for  $f(N, \mathbf{Z})$  where  $q$  denotes the mean proportion of females in the population. The carrying capacity  $K_f$  for the females is then defined by  $\bar{f}(K_f/q) = 0$ , so that  $K_f = qK$ , making the two models of density regulation directly comparable because they will actually generate population fluctuations around the same total population size. We will now consider how fluctuations in the sex ratio will generate additional variance in the change in population size from one year to another,  $\Delta N$ , for two types of mating systems. We show in Appendix I (eqns A4 and A6) that when the density regulation operates within the female segment of the populations,

$$\sigma_{sex}^2 = q(1-q)\left(\bar{f} + N\frac{d\bar{f}}{dN}\right)^2(1+s)/(1-s) \quad \text{eqn 4}$$

for a polygynous mating system, and

$$\sigma_{sex}^2 = \frac{1}{4}\bar{f}^2\left(1 - \frac{2}{\pi}\right)G(s) + \frac{1}{4}\left(N\frac{d\bar{f}}{dN}\right)^2\left(1 - \sqrt{\frac{2}{\pi N}}\right)^2(1+s)/(1-s) \quad \text{eqn 5}$$

for a monogamous mating system. These expressions for  $\sigma_{sex}^2$  differ from those derived by Engen *et al.* (2003) due to  $f(N, \mathbf{Z})$  being replaced by the stochastic term  $f(X/q, \mathbf{Z})$ . In eqn 4 this gives the term  $N df/dN$ , whereas in eqn 5 it generates the last term that is proportional to  $(N df/dN)^2$ . For further details, see Appendix I.

**SPECIES**

The great reed warbler is a medium-sized passerine bird that winters in sub-Saharan tropical Africa and breeds in Europe and Asia. The males return to the Swedish breeding grounds in early May, and the females about 2 weeks later. The great reed warbler has a facultative polygynous mating system (Dyrce 1986, 1988); at Lake Kvismaren, about 40% of the territorial males form pair bonds with two to five females, whereas about 20% of the territorial males remain unpaired (Bensch & Hasselquist 1991; Hasselquist, Bensch & von Schantz 1995; Hasselquist 1998). In the German population, about 11% of the males form pair bonds with two females and 14% of the territorial males remain unpaired (Leisler *et al.* 1995). In general, the proportion of polygynous males as well as the frequency of extra-pair mating was smaller in the German than in the Swedish population (Hasselquist *et al.* 1995; Leisler *et al.* 1995, 2000; Hasselquist, Bensch & von Schantz 1996; Leisler & Wink 2000).

**STUDY AREAS***Swedish population*

A population of great reed warblers was studied at Lake Kvismaren (59°10'N, 15°25'E) in southern Central Sweden in 1986–2002. Each year, fieldwork was carried out daily from late April to early May when the first great reed warblers arrive, to the beginning of August when the last young become fledged. All territorial males and breeding females were mist-netted the first year they settled in our study area, in most cases very soon after arrival (Bensch & Hasselquist 1991; Bensch *et al.* 1998; Hasselquist 1998). A high degree of philopatry was found in the Swedish population (Hansson *et al.* 2002).

*German population*

The German study site is located 290 m above sea level in a fishpond area north-west of Erlangen (49°40'N, 10°51'E) in northern Bavaria (see Leisler *et al.* 1995 for a closer description). The study area was monitored by

regular visits during the whole breeding period from the end of April to the middle of July. Adults were colour-ringed individually, whereas nestlings were ringed with metal rings.

**Methods**

**POPULATION CENSUSES**

An examination of the time-series of the population fluctuations (Fig. 1) shows different patterns in the two study areas. The Swedish population has fluctuated during the last years around a mean value. In contrast, the German population has declined over a period of several years.

**ESTIMATION OF PARAMETERS**


The demographic variance in the female segment of the population was estimated as the weighted mean across years of the fitness contributions of the females (number of female offspring born during the year that survive for at least 1 year plus 1 if the female herself survives) to next generations (Engen, Bakke & Islam 1998; Sæther *et al.* 1998, 2000a,b; Sæther & Engen 2002a).

In order to estimate  $\sigma_{dm}^2$ , including male offspring as well as females (Engen *et al.* 2003), we assume an equal number of adult males and females in the two breeding populations (i.e.  $q = 0.5$ ) and write  $R_{TOT} = M + F +$

$J = O + J$  for the total contribution of the female to the next generation. Here  $M$  and  $F$  are the numbers of male and female offspring that survive to the next generation ( $O = M + F$ ), and  $J$  is an indicator variable for the mother's survival to the next generation. We can then estimate according to Appendix C in Engen *et al.* (2003) the contribution  $\sigma_{dm}^2$  to  $\sigma_d^2$  from the females when only data on female offspring are available. Based on the recapture rate of adults, we assume an adult survival rate  $s = 0.51$  ( $n = 12$  years) and  $s = 0.43$  ( $n = 25$  years) in the Swedish and German population, respectively. Furthermore, assuming no sexual difference in adult survival, the component of the demographic variance generated by binomial distribution of male survival then becomes  $\sigma_{dm}^2 = s(1 - s)(1 - q)$ .

The final component in the demographic variance of the total population (eqn 1) is the contribution due to stochastic temporal variation in the sex ratio  $\sigma_{sex}^2$ . This contribution will depend on the mating system (see eqns 2, 3, A4, A6).

Based on the differences in the pattern of population fluctuations in the two different study areas (Fig. 1), we fitted different population models to the two populations. In the declining German population, we assumed a population model without density regulation. We assume that the change in the logarithm of population size among years ( $\Delta \ln N$ ) is normally distributed. If  $r$  is the deterministic growth rate, i.e.  $\exp(r) = E(N_{t+1}/N_t | N_t)$  and  $\sigma_e^2$  is the environmental stochasticity (Engen, Sæther & Møller 2001), the stochastic growth rate



**Fig. 1.** Fluctuations in total (○) and female (●) population size of the Swedish population of great reed warblers at Lake Kvismare and of the German population in the Fränkischer Weiher region.

$$s = E(\Delta \ln N | N = n) \approx r - \sigma_e^2/2 - \sigma_d^2/2N \quad \text{eqn 5a}$$

and

$$\text{var}(\Delta \ln N | N = n) \approx \sigma_e^2 + \sigma_d^2/N. \quad \text{eqn 5b}$$

If  $X_t = N_t$ , the likelihood function (omitting the constant factors) then becomes:

$$L(r, \sigma_e^2) = \prod_{i=2}^k \frac{1}{\sqrt{(\sigma_e^2 + \sigma_d^2 \exp(-X_{i-1}))}} \exp\left\{-\frac{[X_i - X_{i-1} - r + \sigma_e^2/2 + (\sigma_d^2 \exp(-X_{i-1}))/2]^2}{2(\sigma_e^2 + \sigma_d^2 \exp(-X_{i-1}))}\right\} \quad \text{eqn 6}$$

that is maximized numerically with respect to the two unknown parameters  $r$  and  $\sigma_e^2$ .

For the Swedish population we assumed a logistic type of density regulation (Lande *et al.* 2003), where

$$E(\Delta \ln N | N) = r_1 \left[ 1 - \frac{N - 1}{K - 1} \right]. \quad \text{eqn 7}$$

Here  $r_1$  is the specific growth rate at the smallest possible population size (excluding extinction) at  $N = 1$ . In many passerine species a logistic model describes



the pattern of density regulation quite well (Sæther *et al.* 2000a; Sæther & Engen 2002b). Following Engen *et al.* (1998) and Sæther *et al.* (2000a), the population parameters  $r_1$ ,  $K$  and  $\sigma_e^2$  were then estimated by maximum likelihood. Parametric bootstrapping (Efron & Tibshirani 1993) was used to estimate the bias in the estimate of  $r_1$  (see procedures in Sæther *et al.* 2002a).

#### TIME TO EXTINCTION

We examined the effect of mating system on the extinction process by simulating the process approximating  $\Delta N$  using a normally distributed variable with mean and variance defined by eqns A1 and A5. Under logistic density regulation we have  $\bar{f}(N) = f_0(1 - \beta N)$  so that  $\beta = [1 - (1 - s)f_0]/K$ . The Swedish population showed stationary fluctuations around  $K$  (Fig. 1), making it difficult to obtain estimates of the reproductive rate  $f_0$  at low densities (see Aanes *et al.* 2002). Thus, we assumed  $f_0$  to be the maximum number of female fledglings produced during a year, assuming an even sex ratio at fledging (Bensch *et al.* 1999), multiplied by the maximum annual survival rate, giving  $\hat{f}_0 = 2.13 * 0.82 = 1.75$  (see also Sæther, Engen & Matthysen 2002b for a similar approach). In the declining German population (Fig. 1), we assumed that the reproductive rate currently recorded was representative of  $f_0$ , giving  $\hat{f}_0 = 0.52$ . In this population we assumed  $K$  to be large ( $K = 1000$ ) to produce a small value of  $\beta$  and hence very weak density dependence in  $\bar{f}(N)$ .

Although these approximations may be rather rough at very small population sizes, Engen *et al.* (2003) found the model to be a fair approximation down to population sizes of five to 10 individuals. Because the populations are likely to spend only short periods at very small population sizes, we chose to use the approximations all the way down to extinction at  $N = 1$  individual.

#### Results

All components of the demographic variance were larger in the stationary Swedish population than in the declining German population (Table 1). The demographic variance in the female segment of the population constituted 65% and 73% of the contribution from females to the demographic variance in the total Swedish and German populations, respectively. A large influence of mating system was found on the contribution to demographic stochasticity due to sex ratio fluctuations. In both populations  $\sigma_{sex}^2$  was about five times higher for the polygynous mating system than for the monogamous mating system (Table 1). As a consequence, the total demographic variance assuming a monogamous mating system was about half the value of the demographic variance for the polygynous mating system (Table 1). The environmental stochasticity in the total population size was, however, smaller for a polygynous ( $\hat{\sigma}_e^2 = 0.0056$  in the Swedish and

**Table 1.** Components of demographic stochasticity for monogamous and polygynous mating systems in the Swedish and German populations of the great reed warbler.  $\sigma_f^2$  is the demographic variance in the female segment of the population,  $\sigma_d^2$  the demographic variance for the total population, and  $\sigma_{df}^2$ ,  $\sigma_{dm}^2$  and  $\sigma_{sex}^2$  the contribution to  $\sigma_d^2$  from females, males and sex ratio fluctuations, respectively


Population	Mating system						
	$\sigma_f^2$	$\sigma_{df}^2$	$\sigma_{dm}^2$	$\sigma_{sex}^2$	$\sigma_d^2$	$\sigma_{sex}^2$	$\sigma_d^2$
Germany	0.360	0.491	0.122	0.816	1.184	0.166	0.533
Sweden	0.520	0.798	0.125	0.744	1.268	0.143	0.667

$\hat{\sigma}_e^2 = 0.0323$  in the German population) mating system than when assuming monogamous mating ( $\hat{\sigma}_e^2 = 0.0192$  in the Swedish and  $\hat{\sigma}_e^2 = 0.0423$  in the German population).


The great reed warbler has a facultative polygynous mating system (Dyrce 1986, 1988). In the study population in Kvismare, on average 38% of the males were polygynous (Hasselquist 1998), whereas in Germany the proportion of males with more than one mate varied from 8 to 35% during a 5-year period (Leisler & Wink 2000). For such a polygynous mating system the risk of extinction of the total population was dependent on the type of density regulation. If density regulation acted on the whole population, the risk of extinction in a polygynous mating system was orders of magnitude larger for the Swedish population than when density regulation operated within only the female segment of the population (Fig. 2). For instance, in the latter case the cumulative probability of extinction after 5000 years was 0.0059, compared to 0.2239 if the density regulation acted on the whole population. A similar pattern was also present in the rapidly declining German population (Fig. 1), although the effect of differences in the pattern of density regulation was less pronounced (Fig. 3) than for the Swedish population (Fig. 2).

So far, we have assumed that all females obtain a mate in such a polygynous mating system. However, a surplus of non-breeding females has been recorded in many bird species (Newton 1998). In the Swedish population, the expected time to extinction decreased with a decrease in the proportion of females being mated (Fig. 4). This decline was smaller for density regulation acting on total population size than when it acted within the female segment of the population.

The risk of extinction of the total population is larger if we assume a monogamous mating system than under a polygynous mating system (Figs 2, 3). The magnitude of the difference in extinction risk between the mating system in the stationary Swedish population was much higher for density regulation acting only on the females than when it acted on total population size.



**Fig. 2.** The cumulative probability of extinction of the Swedish population of great reed warblers in relation to mating system (solid line: monogamy, dotted line: polygyny) and type of density regulation.




**Fig. 3.** The cumulative probability of extinction of the German population of great reed warblers in relation to mating system (solid line: monogamy, dotted line: polygyny) and type of density regulation.

## Discussion

Our analyses have shown that sex ratio fluctuations may strongly affect the estimated time to extinction. These effects are influenced strongly by the mating system. In general, the time to extinction becomes shorter in a monogamous than in polygynous mating system (Figs 2, 3). This difference was larger in the Swedish population when density regulation acts only within the female segment than when it operates on the total population size. This demonstrates that sex effects must be included when making population predictions based on counts of total population sizes, and supports previous results obtained by simulation studies (Brook *et al.* 1999; Brook, Burgman & Frankham 2000).

The mathematical results are based on the same simplifying assumptions as in Engen *et al.* (2003), implying several first-order approximations. These are valid for large populations but were shown to give sufficient accuracy even down to quite small population sizes. Furthermore, we also used a first-order approximation (a linearization) for comparing the two models of density regulation. The accuracy of this approximation also depends on the population size, but as the new term involved is always relatively small, the errors involved at small population sizes will also be small. Thus, we believe that our model gives a relatively accurate representation of the fluctuations even for small populations.



**Fig. 4.** The expected time to extinction ( $\log_{10} T$ ) of the Swedish population of great reed warblers in relation to the probability of a female being mated for different types of density regulation. A polygynous mating system is assumed.

Our analyses are based on several simplifying assumptions that may not always be valid (see Møller & Legendre 2001; Kokko & Brooks 2003; Møller 2003). First, we assume an age-independent adult survival rate that also is independent of sex. Especially in polygynous species sex-specific differences in survival have been recorded (Promislow, Montgomerie & Martin 1992; 1994). A second assumption is that the reproductive

success of the females in the polygynous mating system is independent of the proportion of males in the population. Such effects arise if low availability of males results in more females forming pair bonds with polygynous males and thus obtaining less help with parental care, causing reduction in their reproductive success (see Møller & Legendre 2001 for further discussion). For instance, great reed warbler females with secondary status when feeding nestlings have poorer reproductive success than females with primary status (Bensch & Hasselquist 1994; Bensch 1996; Sejberg, Bensch & Hasselquist 2000). Thirdly, we assume that the sex ratio in the population fluctuates only stochastically. Some evidence now suggests that offspring sex ratio may be affected by population structure through largely unknown feedback mechanisms (Sæther *et al.* 2003; 2004); however, no such evidence has been found in the Swedish great reed warbler population (Bensch *et al.* 1999). Finally, the density dependence is assumed to affect only the fecundity rate, which is not necessarily true for sedentary small passerines (Arcese *et al.* 1992). Only further analyses with the use of an explicit simulation model will reveal how these effects will influence the present results.

Nevertheless, even with the simplifying assumptions, our modelling effort was able to demonstrate the influence of variation in mating system on the expected time to extinction of the great reed warbler population (Figs 2, 3). In general, time to extinction for the total population size was shorter for a monogamous than for a polygynous mating system. This is in accordance with the results of Legendre *et al.* (1999), modelling the effects of stochastic variation in sex ratio for establishment success. They showed that when the mating system changed from polygyny to monogamy a large increase in the initial population size was necessary for keeping the extinction risk constant. The large proportion of unmated females present in many monogamous breeding populations (Newton 1998) also suggests that such an extreme monogamous mating system may be a valid description of the mating system in many bird species. Thus, predictions of extinction risk for total population size must include explicit assumptions about the mating system.

These effects of mating system were strongly dependent on how density regulation acted on the total population (Figs 2–4). In polygynous passerine birds such as the pied flycatcher *Ficedula hypoleuca* (Pallas) there is some evidence of intrasexual aggression among females (Slagsvold & Lifjeld 1994), which may indicate the potential for density-dependence within the female segment of the population (Arcese 1989). Also, in the great reed warbler female behaviour seems to be influenced by the presence of other females (Bensch & Hasselquist 1992). Accordingly, removal experiments in passerine bird species have shown the presence of a surplus of non-breeding females during the breeding season (Sæther & Fønstad 1981). Even in species with weaker territorial defence than in many passerines

there is some evidence that density regulation may operate mainly among females. For instance, in some ungulates males and females use different areas during most of the year (e.g. Mouflon *Ovis ammon* L. (Dubois *et al.* 1996). Furthermore, even in species where the sexes are less spatially segregated (e.g. red deer (Conradt, Clutton-Brock & Thomson 1999) and white-tailed deer *Odocoileus virginianus* (Boddaert) (McCullough 1979) density regulation acts mainly within the female segment of the population. In contrast, in some other ungulate populations density dependence seems to operate through total population size (Myerud *et al.* 2002), due probably to greater sexual overlap in resource use.

To summarize, our analyses of two populations of great reed warbler demonstrate that predictions of future population sizes based on total population counts must include explicit assumptions on how density regulation acts and the structure of the mating system (Brook *et al.* 1999, 2000). In particular, analyses of the time to extinction for the total population, which is central for the field of population viability analyses (see Beissinger & McCullough 2002), is likely to be strongly biased if such information is unavailable.

### Acknowledgements

This study was financed by grants from the EU-commission (project 'METABIRD'), the US National Science Foundation and the Research Council of Norway (Strategic University Programme). The studies of the Swedish population has been financed by grants from the Swedish Natural Research Council (VR), the Swedish Research Council for Environment, Agricultural Sciences and Spatial Planning (Formas), Stiftelsen Olle Engkvist Byggmästare, Crafoordska Stiftelsen, Carl Tryggers Stiftelse, Schwartz Stiftelse, Lunds Djurskuddsfond and Kvismare Bird Observatory. During the years 1994–2000 the German study was supported financially by the Deutsche Forschungsgemeinschaft. B.-E. S. acknowledges especially the hospitality of Prof. P.H. Harvey during a stay at the Department of Zoology, University of Oxford, where this paper was written, T. Coulson for discussion and B. W. Brook for comments on the manuscript.

### References

- Aanes, S., Engen, S., Sæther, B.-E., Willebrand, T. & Marcström, V. (2002) Sustainable harvesting strategies of willow grouse in a fluctuating environment. *Ecological Applications*, **12**, 281–290.
- Allee, W.C., Emerson, A.E., Park, O., Park, T. & Schmidt, K.P. (1949) *Principles of Animal Ecology*. Saunders, Philadelphia, PA.
- Arcese, P. (1989) Intrasexual competition, mating system and natal dispersal in song sparrows. *Animal Behaviour*, **38**, 958–979.
- Arcese, P., Smith, J.N.M., Hochachka, W.M., Rogers, C.M. & Ludwig, D. (1992) Stability, regulation, and the determination of abundance in an insular Song Sparrow population. *Ecology*, **73**, 805–822.



- Beissinger, S.R. & McCullough, D.R., eds (2002) *Population Viability Analysis*. University of Chicago Press, Chicago.
- Bensch, S. (1996) Female mating status and reproductive success in the great reed warbler: is there a potential cost of polygyny that requires compensation? *Journal of Animal Ecology*, **65**, 283–296.
- Bensch, S., Hansson, B., Wester Dahl, H. & Hasselquist, D. (1999) Do females adjust the offspring sex ratio to skew in breeding sex ratio? *Journal of Evolutionary Biology*, **12**, 1104–1109.
- Bensch, S. & Hasselquist, D. (1991) Territory infidelity in the polygynous great reed warbler *Acrocephalus arundinaceus*: the effect of variation in territory attractiveness. *Journal of Animal Ecology*, **60**, 857–871.
- Bensch, S. & Hasselquist, D. (1992) Evidence for active female choice in a polygynous warbler. *Animal Behaviour*, **44**, 301–311.
- Bensch, S. & Hasselquist, D. (1994) Higher rate of nest loss among primary than secondary females: infanticide in the great reed warbler? *Behavioral Ecology and Sociobiology*, **35**, 309–317.
- Bensch, S., Hasselquist, D., Nielsen, B. & Hansson, B. (1998) Higher fitness for philopatric than for immigrant males in a semi-isolated population of great reed warblers. *Evolution*, **52**, 877–883.
- Brook, B.W., Burgman, M.A. & Frankham, R. (2000) Differences and congruencies between PVA packages: the importance of sex ratio for predictions on extinction risk. *Conservation Ecology*, **4**, 6 [online].
- Brook, B.W., Cannon, J.R., Lacy, R.C., Mirande, C. & Frankham, R. (1999) A comparison of the population viability analysis packages GAPPs, INMAT, RAMAS and VORTEX for the whooping crane (*Grus americana*). *Animal Conservation*, **2**, 23–31.
- Caswell, H. (2001) *Matrix Population Models*, 2nd edn. Sinauer, Sunderland, MA.
- Clarke, A.L., Sæther, B.-E. & Røskaft, E. (1997) Sex biases in avian dispersal: a reappraisal. *Oikos*, **79**, 429–438.
- Clutton-Brock, T.H. & Coulson, T.N. (2002) Comparative ungulate dynamics: the devil is the detail. *Philosophical Transactions of the Royal Society London, Series B*, **357**, 1285–1298.
- Clutton-Brock, T.H., Coulson, T.N., Milner-Gulland, E.J., Thomson, D. & Armstrong, H.M. (2002) Sex differences in emigration and mortality affect optimal management of deer populations. *Nature*, **415**, 633–637.
- Conradt, L., Clutton-Brock, T.H. & Thomson, D. (1999) Habitat segregation in ungulates: are males forced into sub-optimal foraging habitats through indirect competition by females? *Oecologia*, **119**, 367–377.
- Coulson, T., Catchpole, E.A., Albon, S.D., Morgan, B.J.T., Pemberton, J.M., Clutton-Brock, T.H., Crawley, M.J. & Grenfell, B.T. (2001) Age, sex, density, winter weather and population crashes in Soay sheep. *Science*, **292**, 1528–1531.
- Dale, S. (2001) Female-biased dispersal, low female recruitment, unpaired males, and the extinction of small and isolated bird populations. *Oikos*, **92**, 344–356.
- Dennis, B. (1989) Allee effects: population growth, critical density, and the chance of extinction. *Natural Resource Modeling*, **3**, 481–538.
- Dennis, B. (2002) Allee effects in stochastic populations. *Oikos*, **96**, 389–401.
- Dubois, M., Khazraïe, K., Guilhem, C., Maublanc, M.-L. & Pendu, Y.L. (1996) Philopatry in mouflon rams during the rutting season: Psycho-ethological determinism and functional consequences. *Behavioural Processes*, **35**, 93–100.
- Dyrce, A. (1986) Factors affecting facultative polygyny and breeding results in the great reed warbler (*Acrocephalus arundinaceus*). *Journal für Ornithologie*, **127**, 447–461.
- Dyrce, A. (1988) Mating systems in European marsh-nesting Passeriformes. *Proceedings of the International Ornithological Congress*, **19**, 2613–2623.
- Efron, B. & Tibshirani, R.J. (1993) *An Introduction to the Bootstrap*. Chapman & Hall, New York.
- Emlen, S.T. & Oring, L.W. (1977) Ecology, sexual selection, and the evolution of mating systems. *Science*, **197**, 215–223.
- Engen, S., Bakke, Ø. & Islam, A. (1998) Demographic and environmental stochasticity – concepts and definitions. *Biometrics*, **54**, 830–836.
- Engen, S., Lande, R. & Sæther, B.-E. (2003) Demographic stochasticity and Allee effects in populations with two sexes. *Ecology*, **84**, 2378–2386.
- Engen, S., Sæther, B.-E. & Møller, A.P. (2001) Stochastic population dynamics and time to extinction of a declining population of barn swallows. *Journal of Animal Ecology*, **70**, 789–797.
- Hansson, B., Bensch, S., Hasselquist, D. & Nielsen, B. (2002) Restricted dispersal in a long-distance migrant bird with patchy distribution, the great reed warbler. *Oecologia*, **130**, 536–542.
- Hasselquist, D. (1998) Polygyny in great reed warblers: a long-term study of factors contributing to male fitness. *Ecology*, **79**, 2376–2390.
- Hasselquist, D., Bensch, S. & von Schantz, T. (1995) Low frequency of extra-pair paternity in the polygynous great reed warbler, *Acrocephalus arundinaceus*. *Behavioural Ecology*, **6**, 27–38.
- Hasselquist, D., Bensch, S. & von Schantz, T. (1996) Correlation between male song repertoire, extra-pair paternity and offspring survival in the great reed warbler. *Nature*, **381**, 229–232.
- Hausfater G. & Hrdy, S.B., eds (1984) *Infanticide: Comparative and evolutionary perspectives*. Aldine, NY.
- Kokko, H. & Brooks, R. (2003) Sexy to die for? Sexual selection and the risk of extinction. *Annales Zoologici Fennici*, **40**, 207–219.
- Lande, R. (1993) Risks of population extinction from demographic and environmental stochasticity and random catastrophes. *American Naturalist*, **142**, 911–927.
- Lande, R. (1998) Demographic stochasticity and Allee effect on a scale with isotropic noise. *Oikos*, **83**, 353–358.
- Lande, R., Engen, S. & Sæther, B.-E. (2003) *Stochastic Population Dynamics in Ecology and Conservation*. Oxford University Press, Oxford.
- Legendre, S., Clobert, J., Møller, A.P. & Sorci, G. (1999) Demographic stochasticity and social mating system in the process of extinction of small populations: the case of passerines introduced to New Zealand. *American Naturalist*, **153**, 449–463.
- Leisler, B., Beier, J., Heine, G. & Siebenrock, K.-H. (1995) Age and other factors influencing mating status in German great reed warblers (*Acrocephalus arundinaceus*). *Japanese Journal of Ornithology*, **44**, 169–180.
- Leisler, B., Beier, J., Staudter, H. & Wink, M. (2000) Variation in extra-pair paternity in the polygynous great reed warbler (*Acrocephalus arundinaceus*). *Journal für Ornithologie*, **141**, 77–84.
- Leisler, B. & Wink, M. (2000) Frequencies of multiple paternity in three *Acrocephalus* species (Aves Sylviidae) with different mating systems (*A. palustris*, *A. arundinaceus*, *A. paludicola*). *Ethology Ecology and Evolution*, **12**, 237–249.
- McCullough, D.R. (1979) *The George Reserve Deer Herd: Population Ecology of a K-Selected Species*. University of Michigan Press, Ann Arbor.
- Møller, A.P. (2003) Sexual selection and extinction: why sex matters and why asexual models are insufficient. *Annales Zoologici Fennici*, **40**, 221–230.
- Møller, A.P. & Legendre, S. (2001) Allee-effect, sexual selection and demographic stochasticity. *Oikos*, **92**, 27–34.
- Mysterud, A., Coulson, T. & Stenseth, N.C. (2002) The role of males in the dynamics of ungulate populations. *Journal of Animal Ecology*, **71**, 907–915.

- Newton, I. (1998) *Population Limitation in Birds*. Academic Press, San Diego.
- Promislow, D.E.L., Montgomerie, R. & Martin, T.E. (1992) Mortality costs of sexual dimorphism in birds. *Proceedings of the Royal Society of London, Series B*, **250**, 143–150.
- Promislow, D.E.L., Montgomerie, R. & Martin, T.E. (1994) Sexual selection and survival in North American waterfowl. *Evolution*, **48**, 2045–2050.
- Sæther, B.-E., Engen, S., Lande, R., Arcese, P. & Smith, J.N.M. (2000a) Estimating the time to extinction in an island population of song sparrows. *Proceedings of the Royal Society of London, Series B*, **267**, 621–626.
- Sæther, B.-E. & Engen, S. (2002a) Including uncertainties in population viability analysis using Population Prediction Intervals. *Population Viability Analysis* (eds S.R. Beissinger & D.R. McCullough), pp. 191–212. University of Chicago Press, Chicago.
- Sæther, B.-E. & Engen, S. (2002b) Pattern of variation in avian population growth rates. *Philosophical Transactions of the Royal Society, Series B*, **357**, 1185–1196.
- Sæther, B.-E. & Engen, S. (2003) The routes to extinction. *Macroecology* (eds T. Blackburn & K. Gaston), pp. 218–236. Blackwell Scientific Publishers, Oxford.
- Sæther, B.-E., Engen, S., Islam, A., McCleery, R. & Perrins, C. (1998) Environmental stochasticity and extinction risk in a population of a small songbird, the great tit. *American Naturalist*, **151**, 441–450.
- Sæther, B.-E., Engen, S. & Matthysen, E. (2002b) Demographic characteristics and population dynamical patterns of solitary birds. *Science*, **295**, 2070–2073.
- Sæther, B.-E., Engen, S., Møller, A.P., Weimerskirch, H., Visser, M.E., Fiedler, W., Matthysen, E., Lambrechts, M.M., Badyaev, A., Becker, P.H., Brommer, J.E., Bukacinski, D., Buckinska, M., Christensen, H., Dickinson, J., du Feu, C., Gehlbach, F.R., Heg, D., Hötter, H., Merilä, J., Nielsen, J.T., Rendell, W., Robertson, R.J., Thomson, D.L., Török, J. & Van Hecke, P. (in press) Life history variation predicts the effects of demographic stochasticity on avian population dynamics. *American Naturalist*.
- Sæther, B.-E. & Fonstad, T. (1981) A removal experiment showing unmated females in a breeding population of chaffinches. *Animal Behaviour*, **29**, 147–149.
- Sæther, B.-E., Lande, R., Engen, S., Both, C. & Visser, M. (2002a) Density-dependence and stochastic variation in the dynamics of a newly established small songbird population. *Oikos*, **99**, 331–337.
- Sæther, B.-E., Solberg, E.J. & Heim, M. (2003) Effects of altering sex ratio structure on the demography of an isolated moose population. *Journal of Wildlife Management*, **67**, 455–466.
- Sæther, B.-E., Solberg, E.J., Heim, M., Stacy, J., Jakobsen, K.S. & Olstad, R. (2004) Offspring sex ratio in moose *Alces alces* in relation to paternal age: an experiment. *Wildlife Biology*, **10**, 51–57.
- Sæther, B.-E., Tufto, J., Engen, S., Jerstad, K., Røstad, O.W. & Skåtan, J.E. (2000b) Population dynamical consequences of climate change for a small temperate songbird. *Science*, **287**, 854–856.
- Sejberg, D., Bensch, S. & Hasselquist, D. (2000) Nestling provisioning in polygynous great reed warblers: do males bring larger prey to compensate for fewer nest visits? *Behavioral Ecology and Sociobiology*, **47**, 213–219.
- Slagsvold, T. & Lifjeld, J.T. (1994) Polygyny in birds: the role of competition between females for male parental care. *American Naturalist*, **143**, 59–94.
- Smith, J.N.M. & Arcese, P. (1989) How fit are floaters? Consequences of alternative territorial behaviours in a non-migratory sparrow. *American Naturalist*, **133**, 830–845.
- Solberg, E.J., Sæther, B.-E., Strand, O. & Loison, A. (1999) Dynamics of a harvested moose population in a variable environment. *Journal of Animal Ecology*, **68**, 186–204.
- Swenson, J.E., Sandegren, F., Söderberg, A., Bjärvall, A., Franzén, R. & Wabakken, P. (1997) Infanticide caused by hunting of male bears. *Nature*, **386**, 450–451.
- Tufto, J., Sæther, B.-E., Engen, S., Arcese, P., Jerstad, K., Røstad, O.W. & Smith, J.N.M. (2000) Bayesian meta-analysis of demographic parameters in three small, temperate passerines. *Oikos*, **88**, 273–281.

Received 24 June 2003; accepted 12 February 2004

## Appendix I

Here we assume that density regulation operates only within the female segment of the population so that mean fecundity for a given environment  $\mathbf{Z}$  is a function of the female population size,  $f(X/q, \mathbf{Z})$ . Here  $N$  is the total population size,  $X$  the number of females in the population and  $q$  is the mean proportion of females.

### POLYGYNOUS MATING SYSTEM

For a polygynous mating system and density regulation depending on total population size, Engen *et al.* (2003) showed that the yearly change in population size could be written approximately as:

$$\Delta N = N[f(N)q - (1-s)] + \sqrt{N}\sigma_d\delta_d + N\sigma_e\delta_e, \quad \text{eqn A1}$$

where  $\sigma_e^2 = \text{var}[f(N, \mathbf{Z})q + s(\mathbf{Z})|N]$ . Here  $s(\mathbf{Z})$  is the annual survival rate in environment  $\mathbf{Z}$  and  $s$  is the mean of  $s(\mathbf{Z})$  over the temporal environmental fluctuations in  $\mathbf{Z}$ . The variables  $\delta_d$  and  $\delta_e$  are inde-

pendent standard normally distributed so that  $\text{var}(\Delta N|N) = \sigma_d^2 N + \sigma_e^2 N^2$ .

The demographic variance component  $\sigma_d^2$  is given by eqn 1. The stochastic fluctuation in the sex ratio generates a variance-component given by:

$$\sigma_{sex}^2 = q(1-q)f^2(1+s)/(1-s),$$

which increases with the adult survival due to autocorrelations in the sex ratio generated by this survival. This temporal autocorrelation has been corrected for in eqn A1 that is a model with white noise (Engen *et al.* 2003).

The demographic term in eqn A1 may alternatively be written in the form:

$$\sigma_d\delta_d = \sigma_{dm}\sqrt{1-q}\delta_{dm} + \sigma_{df}\sqrt{q}\delta_{df} + \sigma_{sex}\delta_{sex} \quad \text{eqn A2}$$

where  $\delta_{dm}$ ,  $\delta_{df}$  and  $\delta_{sex}$  are independent variables with zero mean and unit variance.

Conditioned on  $N$  the number of females is approximately binomially distributed (see Engen *et al.* 2003)

and the standardized variable in eqn 3 describing the sex ratio fluctuations then becomes:

$$\delta_{sex} = (X/N - q) / \sqrt{q(1 - q)/N}.$$

We then have:

$$X/q = N + \delta_{sex} \sqrt{N(1 - q)/q}.$$

When the density regulation depends on the number of females only, we find the first order approximation

$$\bar{f}(X/q) \approx \bar{f}(N) + \frac{d\bar{f}}{dN} \delta_{sex} \sqrt{N(1 - q)/q}. \quad \text{eqn A3}$$

Replacing  $\bar{f}(N)$  by this expression in eqn A1, ignoring higher-order terms, we find the only difference is that the factor  $\bar{f}$  in the coefficient of  $\delta_{sex}$  is replaced by  $\bar{f} + (d\bar{f}/dN)N$ . Hence, the demographic variance component due to sex ratio fluctuations turns out to be:

$$\sigma_{sex}^2 = q(1 - q) \left( \bar{f} + N \frac{d\bar{f}}{dN} \right)^2 (1 + s)/(1 - s). \quad \text{eqn A4}$$

Because density regulation implies that the derivative of  $\bar{f}(N)$  is negative, we see that  $\sigma_{sex}^2$  is likely to be somewhat smaller for density regulation operating only within the female segment of the population than when it acts on total population size.

MONOGAMOUS MATING SYSTEM

Following Engen *et al.* (2003) we assume that newborn individuals are females with probability  $q = 0.5$  and a strictly monogamous mating system where the number of females mated is the minimum of the number of males and females in the population. Then, when density regulation acts through  $N$ , the reproductive rate has

mean  $\bar{f}(N)X$  if  $X < N - X$  and otherwise  $\bar{f}(N)(N - X)$ , which may be rewritten as  $\bar{f}(N)(N/2)(1 - |\delta_{sex}|/\sqrt{N})$ . As was the case for the polygynous mating system, our aim is to examine the effects of density regulating operating only within the female segment of the population. This is obtained by replacing  $\bar{f}(N)$  with  $\bar{f}(X/q)$ . Using the same approach as for the polygynous mating system, and observing that  $\delta_{sex}$  and  $|\delta_{sex}|$  are independent, we find that the mean of  $\Delta N$  is unaffected, with an Allele effect as described by Engen *et al.* (2003), that is:

$$E(\Delta N | N) = N \left[ \frac{1}{2} \bar{f}(N) \left( 1 - \sqrt{\frac{2}{\pi N}} \right) - (1 - s) \right]. \quad \text{eqn A5}$$

However, the demographic variance component  $\sigma_{sex}^2$  now includes an additional term

$$\frac{1}{4} \left( N \frac{d\bar{f}}{dN} \right)^2 \left( 1 - \sqrt{\frac{2}{\pi N}} \right)^2 (1 + s)(1 - s), \text{ giving:}$$

$$\sigma_{sex}^2 = \frac{1}{4} \bar{f}^2 \left( 1 - \frac{2}{\pi} \right) G(s) + \frac{1}{4} \left( N \frac{d\bar{f}}{dN} \right)^2 \left( 1 - \sqrt{\frac{2}{\pi N}} \right)^2 (1 + s)(1 - s). \quad \text{eqn A6}$$

The factor  $G(s)$  corrects for the temporal autocorrelations in the sex ratio. More precisely,  $G(s) = 1 + \sum_{h=1}^{\infty} \rho(h, s)$ , where

$$\rho(h, s) \approx \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \left| \left( xs^h + y\sqrt{1 - s^{2h}} \right) \right| \frac{1}{2\pi} e^{-(x^2 + y^2)/2} dx dy.$$

By inspection of eqn A6 we see that the value of  $\sigma_{sex}^2$  will now be larger when the density regulation operates only within the female segment of the population. This is in contrast to the patterns found for the polygynous mating system.