



# LUND UNIVERSITY

## Some classical mixing formulas in the time domain

Kristensson, Gerhard; Rikte, Sten; Andersson, Michael; Sihvola, Ari

1998

[Link to publication](#)

*Citation for published version (APA):*

Kristensson, G., Rikte, S., Andersson, M., & Sihvola, A. (1998). *Some classical mixing formulas in the time domain*. Paper presented at International Symposium on Electromagnetic Theory, 1998, Thessaloniki, Greece.

*Total number of authors:*

4

### General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117  
221 00 Lund  
+46 46-222 00 00

## SOME CLASSICAL MIXING FORMULAS IN THE TIME DOMAIN

Gerhard Kristensson, Sten Rikte, and Michael Andersson

*Department of Electromagnetic Theory, Lund University, P.O. Box 118, S-221 00 Lund, Sweden*

Ari H. Sihvola

*Electromagnetics Laboratory, Helsinki University of Technology*

*P.O. Box 3000, FIN-02015 HUT, Finland*

**Abstract:** This paper discusses a time domain approach to the effective-medium modeling of dielectrically heterogeneous materials. Special emphasis is put on the dispersive behavior of the mixture. Specifically, the Debye model is treated. In terms of the geometry of the mixture, the present analysis is carried out for inclusions of ellipsoidal or spherical form.

### MIXTURE FORMULAS IN THE TIME DOMAIN

The classical mixing theories use a static or quasistatic treatment of the inclusions embedded in a host medium and the polarizabilities of the inclusions. In the time domain, the evaluation of the effective permittivity operator requires evaluation of inverse Fourier transforms or calculation of convolutions and operator inverses.

In the present paper we consider linear, isotropic, temporally dispersive dielectric materials. In a time domain setting, the constitutive relations have the general form, see [2]:

$$\begin{cases} \mathbf{D}(\mathbf{r}, t) = \varepsilon_0 [\epsilon \mathbf{E}](\mathbf{r}, t) = \varepsilon_0 \{ \varepsilon_\infty(\mathbf{r}) \mathbf{E}(\mathbf{r}, t) + (\chi * \mathbf{E})(\mathbf{r}, t) \} \\ \mathbf{B}(\mathbf{r}, t) = \mu_0 \mathbf{H}(\mathbf{r}, t) \end{cases}$$

where the vacuum permittivity is  $\varepsilon_0$  and the vacuum permeability  $\mu_0$ , and the operation denoted by  $*$  stands for the temporal convolution. The function  $\varepsilon_\infty(\mathbf{r}) \geq 1$  describes the instantaneous (optical) response of the material. The dispersive part is described by the susceptibility kernel  $\chi(\mathbf{r}, t)$  as a function of time. Due to causality, the susceptibility kernel  $\chi$  vanishes for  $t < 0$ , but it can have a discontinuity in the origin, i.e., it can happen that  $\chi(t = 0^-) \neq 0$ . Although this option of discontinuity across the origin is sometimes doubted in the literature (see, for example [1, p. 310], [4]), it does not violate causality. In fact, the Debye model, which models the dispersive behavior of polar liquids in the microwave regime excellently, is an example of a case where  $\chi(t = 0^-) \neq 0$ .

### NON-DISPERSIVE BACKGROUND—DEBYE MODEL FOR ELLIPSOIDAL INCLUSIONS

Consider a mixture of dispersive ellipsoidal inclusions in a dispersionless background. The ellipsoidal inclusions are located in random positions with their axes aligned in the host medium. The time domain Maxwell Garnett formula, written for a non-dispersive background medium,  $\varepsilon_b$ , and dispersive inclusions with optical response  $\varepsilon_\infty$  and susceptibility kernel  $\chi(t)$  implies that the effective permittivity dyadic of the mixture is

$$\epsilon_{\text{eff}} = \epsilon_{\text{eff}}^x \hat{x}\hat{x} + \epsilon_{\text{eff}}^y \hat{y}\hat{y} + \epsilon_{\text{eff}}^z \hat{z}\hat{z}, \quad i = x, y, z$$

where the effective permittivity in the  $i$ th direction is

$$\epsilon_{\text{eff}}^i = \epsilon_{\infty, \text{eff}}^i + \chi_{\text{eff}}^{i*} = \varepsilon_b + \varepsilon_b f (\varepsilon_\infty - \varepsilon_b + \chi^*) [\varepsilon_b + N_i (1 - f) (\varepsilon_\infty - \varepsilon_b + \chi^*)]^{-1}$$

This expression is well defined unless  $\varepsilon_b + N_i (1 - f) (\varepsilon_\infty - \varepsilon_b) = 0$ . The permittivity operators (functions) contain the depolarization factors  $N_i$  of the ellipsoids [3]. Specifically, for a Debye-mixture,  $\chi(t) = \beta \exp[-t/\tau]$ ,  $t > 0$ , the optical response and the susceptibility functions are

$$\begin{cases} \varepsilon_{\infty, \text{eff}}^i = \varepsilon_b + \varepsilon_b f \frac{\varepsilon_\infty - \varepsilon_b}{\varepsilon_b + N_i (1 - f) (\varepsilon_\infty - \varepsilon_b)} \\ \chi_{\text{eff}}^i(t) = \frac{f \varepsilon_b^2 \beta H(t)}{[\varepsilon_b + N_i (1 - f) (\varepsilon_\infty - \varepsilon_b)]^2} \exp \left[ - \left( \frac{1}{\tau} + \frac{N_i (1 - f) \beta}{\varepsilon_b + N_i (1 - f) (\varepsilon_\infty - \varepsilon_b)} \right) t \right] \end{cases}$$

### A MIXTURE OF TWO DISPERSIVE MATERIALS

If the background medium and the inclusions are both dispersive, the time domain Maxwell Garnett formula can still be written in the standard form. For spherical inclusions we have

$$\epsilon_{\text{eff}} = \epsilon_{\infty, \text{eff}} + \chi_{\text{eff}}(t) *$$

where the optical response of the mixture is

$$\epsilon_{\infty, \text{eff}} = \epsilon_{\infty, b} + 3\epsilon_{\infty, b} f \frac{\epsilon_{\infty, i} - \epsilon_{\infty, b}}{\epsilon_{\infty, i} + 2\epsilon_{\infty, b} - f(\epsilon_{\infty, i} - \epsilon_{\infty, b})}$$

We now focus on the special case of spherical Debye inclusions in a Debye background material, which susceptibility kernels we denote by

$$\begin{cases} \chi_b(t) = \frac{\epsilon_{s, b} - \epsilon_{\infty, b}}{\tau_b} \exp\left(-\frac{t}{\tau_b}\right) H(t) \equiv \alpha_b \exp(-t\beta_b) H(t) \\ \chi_i(t) = \frac{\epsilon_{s, i} - \epsilon_{\infty, i}}{\tau_i} \exp\left(-\frac{t}{\tau_i}\right) H(t) \equiv \alpha_i \exp(-t\beta_i) H(t) \end{cases}$$

In the case  $\tau_b \neq \tau_i$ , we get the following explicit expression for the effective susceptibility kernel:

$$\chi_{\text{eff}}(t) = c_1 \chi_b(t) + c_2 \chi_i(t) + c_3 \chi_+(t) + c_4 \chi_-(t) + c_5 \chi_b(t) \frac{t}{\tau_b}$$

where the coefficients are


$$\begin{cases} c_1 = 1 + a_1 \left(1 - \frac{\alpha_+}{\beta_+ - \beta_b} - \frac{\alpha_-}{\beta_- - \beta_b}\right) + a_3 \left(\frac{\alpha_+ \beta_b}{(\beta_+ - \beta_b)^2} + \frac{\alpha_- \beta_b}{(\beta_- - \beta_b)^2}\right) \\ c_2 = a_2 \left(1 - \frac{\alpha_+}{\beta_+ - \beta_i} - \frac{\alpha_-}{\beta_- - \beta_i}\right) \\ c_3 = -\epsilon_{\infty, \text{eff}} + \epsilon_{\infty, b} - a_1 \frac{\alpha_b}{\beta_b - \beta_+} - a_2 \frac{\alpha_i}{\beta_i - \beta_+} - a_3 \frac{\alpha_b \beta_b}{(\beta_+ - \beta_b)^2} \\ c_4 = -\epsilon_{\infty, \text{eff}} + \epsilon_{\infty, b} - a_1 \frac{\alpha_b}{\beta_b - \beta_-} - a_2 \frac{\alpha_i}{\beta_i - \beta_-} - a_3 \frac{\alpha_b \beta_b}{(\beta_- - \beta_b)^2} \\ c_5 = a_3 \left(1 - \frac{\alpha_+}{\beta_+ - \beta_b} - \frac{\alpha_-}{\beta_- - \beta_b}\right) \end{cases}$$

where

$$\begin{cases} a_1 = \frac{\epsilon_{\infty, \text{eff}} - \epsilon_{\infty, b}}{\epsilon_{\infty, b}} - \frac{\epsilon_{\infty, \text{eff}} - \epsilon_{\infty, b}}{\epsilon_{\infty, i} - \epsilon_{\infty, b}} \left(1 + \frac{\epsilon_{s, i} - \epsilon_{\infty, i}}{\epsilon_{\infty, b}} \frac{\tau_b}{\tau_i - \tau_b}\right) \\ a_2 = \frac{\epsilon_{\infty, \text{eff}} - \epsilon_{\infty, b}}{\epsilon_{\infty, i} - \epsilon_{\infty, b}} \left(1 + \frac{\epsilon_{s, b} - \epsilon_{\infty, b}}{\epsilon_{\infty, b}} \frac{\tau_i}{\tau_i - \tau_b}\right) \\ a_3 = -\frac{(\epsilon_{\infty, \text{eff}} - \epsilon_{\infty, b})(\epsilon_{s, b} - \epsilon_{\infty, b})}{\epsilon_{\infty, b}(\epsilon_{\infty, i} - \epsilon_{\infty, b})} \\ 2\beta_{\pm} = f_i \alpha_i + \beta_i + f_b \alpha_b + \beta_b \pm \sqrt{(f_i \alpha_i - \beta_i + f_b \alpha_b + \beta_b)^2 + 4f_i \alpha_i (\beta_i - \beta_b)} \\ \alpha_{\pm} = \frac{\beta_i - \beta_b}{\frac{(\beta_+ - \beta_b)}{(\beta_{\pm} - \beta_b)} - \frac{(\beta_+ - \beta_i)}{(\beta_{\pm} - \beta_i)}} \end{cases}$$

and

$$\chi_{\pm}(t) = \alpha_{\pm} \exp(-\beta_{\pm} t) H(t)$$



**Figure 1:** The susceptibility kernels for water, ethanol, ethanol-in-water mixture (20% ethanol), and water-in-ethanol mixture (80% ethanol).

Consequently,  $\chi_{\text{eff}}(t)$  is a linear combination of four Debye kernels and one modified Debye kernel. We also have that

$$\min(\beta_i, \beta_b) < \beta_- < \max(\beta_i, \beta_b) < \beta_+, \quad \alpha_{\pm} > 0$$

This implies that the relaxation time  $\tau_+ = 1/\beta_+$  is less than both the relaxation times  $\tau_i$  and  $\tau_b$ , whereas  $\tau_- = 1/\beta_-$  assumes a value between  $\tau_i$  and  $\tau_b$ .

Numerical results for a Debye-in-Debye medium (ethanol and water, 20-80% and 80-20%) are displayed in Figure 1. The figure shows that although the mixture curves resemble in form the Debye-kernels of water and ethanol, they are not exactly exponentially decaying. Another interesting detail of Figure 1 is the fact that the water-in-ethanol mixture has a slightly stronger memory for times after 40 ps compared to both water and ethanol, although the difference is very small.

#### REFERENCES

- [1] J. D. Jackson. *Classical Electrodynamics*. John Wiley & Sons. New York. second edition. 1975.
- [2] A. Karlsson and G. Kristensson. Constitutive relations, dissipation and reciprocity for the Maxwell equations in the time domain. *J. Electro. Waves Applic.*, 6(5/6), 537-551, 1992.
- [3] L. D. Landau, E. M. Lifshitz, and L. P. Pitaevskii. *Electrodynamics of Continuous Media*. Pergamon, Oxford, 1984.
- [4] W. S. Weiglhofer and A. Lakhtakia. On causality requirements for material media. *AEÜ Int J. Electron. Commun.*, 50(6), 389-391, 1996.