



LUND UNIVERSITY

Glacial geology and palaeo-ice dynamics of two ice-sheet margins, Taymyr Peninsula, Siberia and Jameson Land, East Greenland

Alexanderson, Helena

2002

[Link to publication](#)

Citation for published version (APA):

Alexanderson, H. (2002). *Glacial geology and palaeo-ice dynamics of two ice-sheet margins, Taymyr Peninsula, Siberia and Jameson Land, East Greenland*. [Doctoral Thesis (compilation), Quaternary Sciences]. Quaternary Sciences, Department of Geology, Lund University.

Total number of authors:

1

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

LUNDQUA Thesis 48

Glacial geology and palaeo-ice dynamics of two ice-sheet margins, Taymyr Peninsula, Siberia and Jameson Land, East Greenland

Helena Alexanderson

Avhandling

att med tillstånd från Naturvetenskapliga Fakulteten vid Lunds Universitet för avläggande av filosofie doktorexamen, offentligen försvaras i Geografiska institutionens föreläsningssal, Sölvegatan 13, Lund, fredagen den 25 oktober 2002 kl. 13.15

Lund 2002

Lund University, Department of Geology, Quaternary Geology

Organization LUND UNIVERSITY Department of Geology, Quaternary Geology	Document name DOCTORAL DISSERTATION	
	Date of issue 25 September 2002	
	Sponsoring organization	
Author(s) Helena Alexanderson		
Title and subtitle Glacial geology and palaeo-ice dynamics of two ice-sheet margins, Taymyr Peninsula, Siberia and Jameson Land, East Greenland		
Abstract <p>The North Taymyr ice-marginal zone (NTZ) on the Taymyr Peninsula, Arctic Siberia and the Ugleelv Valley on Jameson Land, East Greenland, have been investigated with the aim of reconstructing the glacial history, including depositional processes and environments. Geomorphological, sedimentological, stratigraphical and remote sensing methods have been combined to give a comprehensive view of developments in the two areas. Optically stimulated luminescence (OSL) and radiocarbon (^{14}C) dates provide the chronological control.</p> <p>The Kara Sea shelf was glaciated three times during the Weichselian, each time with a smaller ice cover than before. The ice sheets caused a reversal of the fluvial drainage towards the south on the Taymyr Peninsula and, during the Early-Middle Weichselian, also the damming of proglacial lakes. The youngest ice-advance, but probably also the older ones, was warm-based and 'surge-like'. After it had reached its maximum position, the margin froze to its base and compressional flow took place there. The North Taymyr ice-marginal zone (NTZ) was initiated during an Early Weichselian retreat stage (c. 80 ka BP) and added to during the Middle (c. 65 ka BP) and Late Weichselian (<20 ka BP) ice advances, thus revealing a complex history. It comprises ice-marginal and supraglacial landsystems dominated by 2-3 km wide thrust-block moraines. Large areas are still underlain by remnant glacier ice and a supraglacial landscape with numerous ice-walled lakes and kames is forming even today. The proglacial landsystem is characterised by subaqueous or terrestrial environments, depending on altitude and time of formation.</p> <p>The sedimentary succession in the Ugleelv area comprises three tills, glaciolacustrine, glaciofluvial and aeolian sediments. The depositional history started in the early Saalian with a prograding delta in a lake dammed by an outlet glacier in Scoresby Sund. Soon thereafter, and also once during the Weichselian, Jameson Land was inundated by glaciers emanating from Liverpool Land in the east. These glaciers were warm-based and deposited glaciofluvial sediments, local tills and small end moraines. Conditions were less dynamic later in the glacial cycles. During most of the Weichselian the Ugleelv area was ice free and aeolian activity took place, but in the late Saalian the Greenland ice sheet expanded eastwards over the area. This ice was mainly cold-based but in places temporarily warm-based, where a lodgement till was deposited. During deglaciation there was substantial glaciofluvial erosion.</p> <p>The Kara Sea ice sheet and the Greenland ice sheet behave differently through a glacial cycle. The shelf-centred Kara Sea ice experiences large shifts in areal extent and disappears completely during interglacials. 'Individual' ice advances seem to be governed by internal ice dynamics rather than by climatic changes. Contrary to this, the Greenland ice sheet is relatively stable and the climatically driven expansion or retreat of its outlet glaciers through the fjords represents the major changes.</p>		
Key words: sedimentology, geomorphology, glacial history, ice-marginal processes, ice dynamics, Pleistocene, Arctic Siberia, East Greenland		
Classification system and/ or index terms (if any):		
Supplementary bibliographical information: 250 copies		Language English
ISSN and key title: 0281-3033 LUNDQUA THESIS		ISBN 91-86746-48-0
Recipient's notes	Number of pages 17 + 4 app.	Price 120 SEK
	Security classification	

Distribution by Quaternary Geology, Lund University, Tornavägen 13, SE-223 63 Lund, Sweden
 I, the undersigned, being the copyright owner of the abstract of the above-mentioned dissertation, hereby grant to all reference sources permission to publish and disseminate the abstract of the above-mentioned dissertation.

Signature 

Date 2 September 2002

Contents

Introduction	2
Study areas	4
Methods	4
Results - summaries of papers	4
Fieldwork and authorship contributions	4
Paper I	4
Paper II	5
Paper III	6
Paper IV	7
Discussion	8
Glaciations and landsystems	8
Ice-sheet limits	9
The nature of glacial cycles	10
Conclusions	11
Implications and ideas for the future	12
Acknowledgements	12
Svensk sammanfattning	13
References	14

Appendices

- I: **Alexanderson, J.H.** 2000: Landsat mapping of ice-marginal features on the Taymyr Peninsula, Siberia – image interpretation versus geological reality. *Geological Quarterly* 44(1), 15-25.
- II: **Alexanderson, H., Hjort, C., Möller, P., Antonov, O. & Pavlov, M.** 2001: The North Taymyr ice-marginal zone, Arctic Siberia – a preliminary overview and dating. *Global and Planetary Change* 31(1-4), 427-445.
- III: **Alexanderson, H., Adrielsson, L., Hjort, C., Möller, P., Antonov, O., Eriksson, S. & Pavlov, M.** 2002: Depositional history of the North Taymyr ice-marginal zone, Siberia – a landsystem approach. *Journal of Quaternary Science* 17(4), 361-382.
- IV: **Adrielsson, L. & Alexanderson, H.:** Two cycles of ice-sheet and coastal mountain glaciation in central East Greenland. Manuscript submitted to *Boreas*.

Glacial geology and palaeo-ice dynamics of two ice-sheet margins, Taymyr Peninsula, Siberia and Jameson Land, East Greenland

by

Helena Alexanderson

Quaternary Geology, Department of Geology, Lund University

Sölvegatan 13, SE-223 62 Lund, Sweden

'To see a world in a grain of sand'

William Blake

This thesis is based on four papers listed below as appendix I-IV and referred to in the text by their Roman numerals. The papers are reprinted with the permission of the Polish Geological Institute (paper I), Elsevier (paper II) and John Wiley & Sons, Ltd. (paper III). Paper IV has been submitted to *Boreas*.

App. I: **Alexanderson, J.H.** 2000: Landsat mapping of ice-marginal features on the Taymyr Peninsula, Siberia – image interpretation versus geological reality. *Geological Quarterly* 44(1), 15-25.

App. II: **Alexanderson, H., Hjort, C., Möller, P., Antonov, O. & Pavlov, M.** 2001: The North Taymyr ice-marginal zone, Arctic

Siberia – a preliminary overview and dating. *Global and Planetary Change* 31(1-4), 427-445.

App. III: **Alexanderson, H., Adrielsson, L., Hjort, C., Möller, P., Antonov, O., Eriksson, S. & Pavlov, M.** 2002: Depositional history of the North Taymyr ice-marginal zone, Siberia – a landsystem approach. *Journal of Quaternary Science* 17(4), 361-382.

App. IV: **Adrielsson, L. & Alexanderson, H.**: Two cycles of ice-sheet and coastal mountain glaciation in central East Greenland. Manuscript submitted to *Boreas*.



Folded glacier ice below a thin layer of melted-out diamicton, exposed in a slump headwall at White Lake, the Taymyr Peninsula, Siberia.

Introduction

Landscapes in high northern latitudes are largely shaped by ice sheets, which have repeatedly grown and decayed during the Quaternary. Sediments and landforms hold information about these events and can be used to reconstruct past environments and climate. This reconstruction is an ever ongoing work and our views are constantly changing as new data are brought forward.

Ice sheets grow and decay due to external and internal mechanisms. The ultimate driving force behind extensive glaciations is believed to be the Milankovitch astronomic orbital cycles (Berger *et al.* 1984). Repeated ice-lobe advances alternating with periods of ice-front stagnation or retreat during a single ice age are also generally considered to be driven by climatic events. However, this is only partly true, as other factors, such as topographic setting and subglacial conditions (substrate, temperature, hydrology) may hamper or facilitate individual glacier advances (e.g. Clark 1994) and even affect ice-sheet behaviour (Oerlemans 1982; Clark *et al.* 1999). Thus, to interpret the geological record correctly it is necessary to know whether an ice-lobe advance was caused exclusively by climatic changes or was mainly triggered by other mechanisms.

Two distinct modes of ice-sheet behaviour through a glacial cycle are indicated by the striking differences between the glacial records from northern Eurasia and Greenland. The Greenland ice sheet has long remained stable with only relatively small oscillations (Funder *et al.* 1998), while during the same time a build-up of ice sheets and large-scale rapid advances of low-gradient ice-sheet lobes seem to have occurred along the coast of arctic Eurasia (e.g. Tveranger *et al.* 1999).

Sedimentological data from southern Scandinavia (Adriellsson 2001), North America (Clark 1994) and the northern Siberian coast (this study), indicate that special physical and hydrological conditions, occurring at the base of the ice sheet, caused instabilities that resulted in the rapid spread of ice masses. This suggests a complex connection between climatic factors and ice-sheet dynamics, and highlights the importance of comparative glacial-geological studies of areas with contrasting ice-dynamic behaviour.

The spatial distribution of ice sheets influences the oceanic and atmospheric circulation, as well as the hydrology of surrounding continental areas. For

example, depending on the ice-sheet distribution in arctic Eurasia, several large, north-flowing rivers may become blocked, leading to the creation of large proglacial lakes and to reversed (southward) drainage (Arkhipov *et al.* 1995; Mangerud *et al.* 2001). Ice-sheet limits are also important input parameters in computer models of climate, isostasy and rheology (e.g. Lambeck *et al.* 1998; Siebert & Marsiat 2001).

Several contradicting hypotheses have been put forward regarding the spatial and temporal extents of Eurasian ice sheets, and also about their geographical origins, especially for the Late Weichselian – the Last Glacial Maximum (Rutter 1995; Svendsen *et al.* 1999). The propositions have ranged from a marine-based panarctic ice sheet (Grosswald 1980; Grosswald & Hughes 2002) to limited local highland glaciation (Velichko *et al.* 1997; Pavlidis *et al.* 1997). Astakhov (1992, 1997, 1998) incorporated a temporal aspect and suggested that the large ice sheets occurred in the Early Weichselian or Saalian and that the Late Weichselian glaciation in Siberia was restricted.

Work within the QUEEN (Quaternary Environment of the Eurasian North) and 'Eurasian Ice Sheets' programmes, of which this study is a part, has led to a revision of the view on glaciations in the Eurasian Arctic (Larsen *et al.* 1999; Thiede *et al.* 2001). We now believe that the greatest ice extent was during the Saalian, and during the Weichselian the maximum glaciation was successively displaced towards the west, occurring in the east (Kara Sea region) during the Early Weichselian and culminating in the west (Scandinavia) during the Late Weichselian.

The glacial history of East Greenland is better known, in large part due to the results of the PONAM (Late Cenozoic Evolution of the Polar North Atlantic Margins) programme in the early 1990's (Funder *et al.* 1998). Most work has, however, been concentrated on outlet glaciers in fjords and less has been done in the highland areas in between.

The aim of my work has been to reconstruct the glacial history, including processes and environments, at former ice-sheet margins, in this case at the eastern flanks of the Kara Sea ice sheet and the Greenland ice sheet (Fig. 1). The more specific objectives have been to describe sediments and landforms, map ice-sheet limits, recognise the depositional processes and, from all this, deduce the nature of the ice sheets and glacial cycles.




Fig. 1. Map of the Arctic with insets of my study areas: the Taymyr Peninsula, Siberia and Jameson Land, East Greenland. On Taymyr the fieldwork areas are, from left to right, the Astronomical Lakes, Barometric Lake and White Lake. EW is Early Weichselian, LW is Late Weichselian and R is river. Positions of the ice-marginal zones south of the Byrranga Mountains are mainly from Kind & Leonov (1982).

Study areas

My main fieldwork area has been the Taymyr Peninsula in north-central Siberia. The major target there was the North Taymyr ice-marginal zone (NTZ), which is the northernmost of several ice-marginal zones on Taymyr (Fig. 1). During two summers, in 1998 and 1999, the 750-km-long NTZ was extensively studied in three areas: the Astronomical Lakes (Ozera Astronomicheskoye; 76°17'N, 96°19'E), Barometric Lake (Ozero Barometricheskoye; 75°35'N, 98°24'E) and White Lake (Ozero Beloye; 75°57'N, 101°54'E) areas. These sites are located 80-100 km apart on a SW-NE transect (Fig. 1). In the summer of 2000 I worked in the inner-most parts of the Ugleelv Valley on Jameson Land in East Greenland (Fig. 1), where more than 100 m of Quaternary sediments are exposed.

Methods

To get an overview of the northwestern Taymyr Peninsula and the NTZ, I studied a suite of satellite images (Landsat, Corona, KATE; see Table 1 in paper III) and analysed them both visually and digitally (paper I). I did the latter analyses by using the GIS programmes IDRISI (Eastman 1997) and ArcView (Environmental Systems Research Institute). Aerial photographs were available for the fieldwork areas on both Taymyr and Jameson Land and have been used to map sediments and landforms.

Digital elevation models (DEM's) of the Taymyr Peninsula and of Jameson Land have been extracted from the Global Land One-km Base Elevation (GLOBE) data set (GLOBE Task Team *et al.* 1999). I have also created more detailed DEM's of the fieldwork areas from digitised topographic maps (scale 1:100,000; 1:250,000). The DEM's have, for example, been used to estimate the extent and drainage-ways of ice-dammed lakes.

I have mapped landforms both in the field and by remote sensing. Detailed logging of sediments, including bed contacts, lithology, grain size and structures in sections along lake shores, rivers and valley sides formed a basis for interpreting depositional processes and environments. Palaeocurrent measurements and fabric analyses were also made and samples were taken for dating, laboratory analyses and documentation. The radiocarbon (¹⁴C-) dating was done at the

Radiocarbon Dating Laboratory in Lund, Sweden and the optically stimulated luminescence (OSL) dating at the Nordic Laboratory for Luminescence Dating at Risø, Denmark.

Latitude/longitude coordinates were measured with a Magellan GPS Pioneer. Altitudes are either taken directly from topographical maps or were measured by a Thommen HM30 digital altimeter, calibrated at fix-points on the topographical maps. I have investigated many more sites than are presented here, so the original site numbers have been kept to aid correlation with the public PANGAEA database (<http://www.pangaea.de>), from which all sedimentary logs from Taymyr can be retrieved.

Results – summaries of papers

Fieldwork and authorship contributions

The work behind the co-authored papers (II-IV) has been shared as follows: For papers II and III, the greater part of the fieldwork along the NTZ was done by Christian Hjort and me, with contributions from Oleg Antonov, Saskia Eriksson, Per Möller and Maksim Pavlov. I did most of the sedimentological logging and interpretation, a large part of the geomorphological mapping and also the bulk of the remote sensing. I was responsible for writing and received input from Lena Adrielsson, Oleg Antonov, Christian Hjort and Per Möller. For paper IV, Lena Adrielsson and I contributed equally to both fieldwork and writing.

Paper I

Alexanderson, J.H. 2000: Landsat mapping of ice-marginal features on the Taymyr Peninsula, Siberia – image interpretation versus geological reality. *Geological Quarterly* 44(1), 15-25.

The aim of this paper was to survey fieldwork localities and to acquire a regional geological overview of the North Taymyr ice-marginal zone on the Taymyr Peninsula, Siberia (Fig. 1). I did this by visually and digitally studying a Landsat 5 Multispectral Scanner (MSS) image (153/006; 8 Aug. 1992). The computer-based analyses were mainly made in IDRISI (Eastman 1997).

The ice-marginal zone, which is >2 km wide and many kilometres long, is clearly visible on the

satellite image and I could also distinguish different ground types by classifying spectral signatures. Two areas were ground-truthed during fieldwork and this helped in interpreting the ground types geologically. A supervised classification method ('Maximum likelihood'), based on Bayesian probability theory, proved to give the result which best corresponded to reality. Seven ground types (spectral classes) were distinguished, described and interpreted:

- *Dark-coloured bedrock and peat.* The bedrock includes dolerites and greenschists and is found on topographic highs. The peat areas are dominated by dark moss with sparse grasses and herbs and are located mainly within topographic depressions.

- *Light-coloured bedrock.* This class is made up of limestones, shales, sandstones, granites and, locally, also unconsolidated Cretaceous sand.

- *Unvegetated cobbles, gravel and sand.* This ground type has no or very sparse (<20%) vegetation cover. It is a diverse class that encompasses, for example, fluvial bars, beaches, boulder fields, windblown hilltops, sediment cliff sections and even some frost-shattered bedrock outcrops.

- *Vegetation-covered fine diamict and coarse sorted material.* This class is the most common on the satellite image and predominates in high, relatively dry terrain. The ground is >90% covered by vegetation. It consists mainly of thin silty till and, concentrated in some areas, of glaciofluvial and glaciolacustrine sands and gravels.

- *Tussocky and dry silt.* This ground type corresponds to a hummocky tundra landscape with frost-boiled silt and gully formation. In lowland areas, it consists of glaciolacustrine silts and in higher terrain of silty till.

- *Well-vegetated silt.* This is a very patchy ground type where moss-tussocks dominate and it forms large undulating areas. The class has been interpreted as silty glaciolacustrine sediments.

- *Waterlogged mud.* The class occurs on damp to wet grasslands on floodplains, recent deltas and around lakes. It consists of muddy alluvial sediments, mainly Holocene in age.

Satellite image analysis is advantageous in that it is relatively low-cost, covers large areas and can be done quickly and systematically. However, there are some disadvantages, such as poor spatial and spectral resolution. In this case the results were largely satisfactory, but to correctly interpret the identified spectral classes geologically, it was necessary to consider individual areas and their topographic position, relative size and association with other classes.

Paper II

Alexanderson, H., Hjort, C., Möller, P., Antonov, O. & Pavlov, M. 2001: The North Taymyr ice-marginal zone, Arctic Siberia – a preliminary overview and dating. *Global and Planetary Change* 31(1-4), 427-445.

This paper gives an overview of the geographical extent and the general geomorphology and sedimentology of the North Taymyr ice-marginal zone (the NTZ), a c. 750 km long ice-marginal zone built up in front of an ice sheet emanating from the Kara Sea shelf area (Fig. 1). It also presents a preliminary chronology. The NTZ is most prominent and continuous in its central parts, on which this paper concentrates. For the rest of its length, west of the Astronomical Lakes and north of the Leningradskaya River, the zone is less distinct.

The NTZ is a complex physiographic feature, composed of several glacial elements of differing age. It marks a distinct change in the landscape, from a distal weathered terrain to a proximal, fresh landscape littered with crystalline erratics. Morphologically, the base of the NTZ is made up of large, arcuate ice-marginal moraines with a silty till cover. These moraines are ice-cored and pitted with lakes. Superimposed are smaller ice-marginal hills and ridges, which consist of sand, gravel or diamict sediments and occur mainly on the southern (distal) side. Distal to the moraines are sandur plains and valley trains, and also deltas related to two different glacial-lake levels, at 120-140 m a.s.l. and at c. 80 m a.s.l. Other shoreline indicators, such as beach ridges, corresponding to these levels have also been found. In the low-lying areas just outside the NTZ are large expanses of laminated, fine-grained glaciolacustrine sediments.

Ice-dammed lakes were formed when ice sheets, advancing from the north, blocked the normal, northward drainage and also supplied increased volumes of meltwater. The lakes occupied the Lower Taymyr, Shrenk and Trautfetter river valleys and were linked to the Taymyr Lake basin south of the Byrranga Mountains. From Taymyr Lake, the drainage was either westward, through the Pyasina River valley out onto the southern Kara Sea shelf, or eastward to Khatanga Bay and the Laptev Sea. South-directed glaciofluvial and glaciolacustrine sediments at, for example, Lake Engelgardt support the reversed, southward drainage through the Byrranga Mountains.

The depositional history of the NTZ can be divided into three major events (Fig. 2). The




Fig. 2. Glaciation curve for the eastern flank of the Kara Sea ice sheet. The vertical, light grey shading represents the NTZ area. For locations see Fig. 1.

chronology is based on optically stimulated luminescence (OSL) and radiocarbon (^{14}C) dating.

1. *Early Weichselian, c. 80 ka BP.* A retreating ice-sheet front made a temporary halt at the NTZ and the large moraines and some superimposed ridges were formed together with deltas and glaciolacustrine sediments in the 120-140-m lake. During the early stage of the ice-dammed lake, a brief marine incursion may have occurred. After deglaciation, a detached debris-rich ice-margin body was left and it was gradually covered by its own melt-out till and incorporated into the permafrost. This event is most likely rather closely preceded by an earlier retreat stage, the 'Ledyanaya event' (Möller *et al.* 1999a, b), with glaciomarine delta sedimentation south of the Byrranga Mountains.

2. *Middle Weichselian, c. 65 ka BP.* The ice front once again stood at the NTZ and most of the superimposed ridges belong to this phase. Deltas and glaciolacustrine sediments were deposited in the 80-m lake, which was fed both by rivers and directly from the ice sheet. The lower lake level may be attributed to a lowered threshold, a new opened-up passage or simply to reduced isostatic downloading. There is also at least a 10 ka time difference between Events 1 and 2.

3. *Late Weichselian, <20 ka, >9.5 ka BP.* A rather thin ice sheet from the Kara Sea shelf inundated some coastal lowlands and large river valleys and dislocated and deformed older sediments. The central parts of the ice sheet melted away earlier than the frontal areas, where remnant glacier ice is still left under a thin (<1 m) till cover. No glacial lake of any consequence was dammed by this ice.

Paper III

Alexanderson, H., Adrielsson, L., Hjort, C., Möller, P., Antonov, O., Eriksson, S. & Pavlov, M. 2002: Depositional history of the North Taymyr ice-marginal zone, Siberia – a landsystem approach. *Journal of Quaternary Science* 17(4), 361-382.

In this paper the North Taymyr ice-marginal zone (NTZ) is described in more detail and the processes responsible for its formation and the nature of the ice sheets involved are discussed. Information from satellite imagery, geomorphological mapping and sedimentological logging has been combined in a landsystem approach. The NTZ landsystem (Fig. 3), consisting of the sediment-landform associations mentioned below, has many similarities to subpolar ice-marginal landsystems, but there are some differences, such as the lack of subglacial landforms. It is concluded that:

- *The ice-marginal association* consists of large thrust-block moraines, two types of small push moraines, kames, subaqueous fans and large-scale deformation of sediment and ice. It formed where debris was concentrated and moved to the ice surface due to compressive flow at a margin frozen to its base. The push moraines, kames and subaqueous fans occur on the distal side of the large moraines, and the latter two are in places arranged in concentric or reticulate patterns.

- *The supraglacial association* is underlain by remnant glacier ice and is characterised by an ablation terrain with numerous glaciokarst lakes and kames. There is a zonation reflecting the relative abundance of sediment and ice, from a sediment-dominated distal zone to an ice-rich proximal zone. The surface cover is a supraglacial melt-out till, mostly consisting of redeposited shell-bearing marine sediments. The landscape is still under development by back and downwasting processes, which eventually should lead to topographic inversion.

- *The proglacial association* is either subaqueous or terrestrial, depending on altitude for different areas and times of deposition. It consists of ice-




Fig. 3. A schematic representation of the landsystem of the North Taymyr ice-marginal zone.

contact deltas at two levels (c. 130 and 80 m a.s.l.), fine-grained glaciolacustrine sediments, glacial lake shorelines, sandar and valley fills.

- *The subglacial sediment-landform association* has not been studied in any detail. It seems, however, as if the subglacial drainage system was a distributed one, but that a switch to a channelised system may have taken place at the margin.

- *The paraglacial association*, consisting of alluvial and aeolian sediments, thaw lakes, patterned ground and palsas, has formed since the active ice retreated from the NTZ.

A schematic model, divided into four phases, for the dynamics of the Late Weichselian ice advance onto the Taymyr Peninsula has been constructed. In phase 1, cold ice was formed on the Kara Sea shelf. As it advanced over the permafrozen shallow shelf, much debris was incorporated and the subglacial drainage also became blocked (phase 2). Then the porewater pressure under the ice increased until the ice was lifted and started to slide over subglacial water and a deforming bed (phase 3). The ice moved on to present land areas until the driving force at the margin was reduced and the ice slowed down. The margin then froze to its bed (phase 4) and a compressional-flow regime was developed since ice from behind still pushed on.

The formation of the NTZ derives from three successively smaller glacial events during the Weichselian (Fig. 2). The oldest parts of the NTZ were formed c. 80 ka BP and represent a stagnation or readvance phase during the deglaciation of an Early Weichselian ice sheet that at its maximum

reached far south of the Byrranga Mountains (Hjort *et al.* 2002). New deposits and landform segments were added to the NTZ during the Middle Weichselian, around 65 ka BP. Proglacial lakes were dammed by the ice during both these phases, at c. 130 and 80 m a.s.l., respectively. During the Late Weichselian, after 20 ka BP, a comparatively small, thin ice sheet advanced on to the northwesternmost Taymyr Peninsula. Like its predecessors it caused a reversal of the drainage towards the south, in this case, however, only leading to an elevated level and increased sedimentation rates in the Taymyr Lake basin (Möller *et al.* 1999a; Hjort *et al.* 2002). This last event was of short duration and the ice had collapsed by c. 12 ¹⁴C ka.

Paper IV

Adriellsson, L. & Alexanderson, H.: Two cycles of ice-sheet and coastal mountain glaciation in central East Greenland. Manuscript submitted to *Boreas*.

This paper presents the glacial history of the Ugleelv Valley on central Jameson Land, East Greenland, as inferred from sedimentological and geomorphological data. Stratigraphy gives a relative chronology and optically stimulated luminescence dates provide some absolute age control. Correlations with the Jameson Land succession (e.g. Funder *et al.* 1998) and with regional ice-core and IRD-records (Johnsen *et al.* 1992; Stein *et al.* 1996) are discussed.

Three tills have been identified, differentiated by their lithological content and stratigraphic position, but the thickest deposits are




Fig. 4. Glaciation curve for the Scoresby Sund area. The hatched areas represent the Greenland ice sheet and its outlet glaciers, in contrast to local ice caps (grey). The vertical, light grey shading shows the location of the Ugleelv area. The $\delta^{18}\text{O}$ -record is from the Renland ice core (Johnsen *et al.* 1992) and the ice-rafted debris (IRD) record from core PS1726 on the continental slope off Scoresby Sund (Stein *et al.* 1996; stretched to fit a linear timescale). Additional data also from Funder *et al.* (1998) and Hansen *et al.* (1999; Scoresby Sund).

glaciolacustrine deltaic and ice-marginal glaciofluvial sediments. Ice-free intervals are primarily represented by aeolian deposits. The depositional history in the Ugleelv Valley is as follows (Fig. 4):

- *A glacial lake dammed by an outlet glacier in Scoresby Sund, early Scoresby Sund glacial (Saalian).* A delta prograded from southwest into an ice-dammed lake on central Jameson Land, implying a debris-rich, warm-based glacier in Scoresby Sund.

- *Liverpool Land mountain glaciation 1, early Scoresby Sund glacial (Saalian).* Glaciers expanded from the Liverpool Land ice caps in the east and reached the Ugleelv area. A local lodgement till was deposited, showing that the glacier was warm-based. During deglaciation there was significant deposition of thick glaciofluvial sediments in subglacial and ice-marginal positions.

- *Periglacial conditions, Scoresby Sund glacial (Saalian).* At the end of deglaciation and during a following ice-free period, a cold and dry climate prevailed. Aeolian sediments were deposited and composite wedges and cryoturbation structures were

formed.

- *Main ice-sheet advance, late Scoresby Sund glacial (Saalian).* The Greenland ice sheet overrode the area from the west, depositing lodgement till and crystalline erratics. Although mainly cold-based, spatial and temporal changes in basal temperature occurred. Meltwater eroded most of the present Ugleelv Valley, c. 135 ka BP.

- *Liverpool Land mountain glaciation 2, early? Weichselian, age <109 ka BP.* A glacier from the east (Liverpool Land) moved up the Ugleelv Valley while eroding its substrate and forming a U-shaped valley. It tectonised older glaciolacustrine sediments, as well as depositing a deformation till and small end moraines.

- *Aeolian activity, Weichselian, around c. 109, 27 and 10 ka BP.* On at least three occasions during the Weichselian, aeolian sediments have been deposited in the Ugleelv Valley. Both wind directions from the south (27 ka) and the north (10 ka) are recorded. Lee-side dunes within the valley are associated with the winds from the south and indicate arid conditions and no ice cover in the area at that time. The northerly winds are related to the formation of permanent snowpatches.

Two glacial cycles have thus been identified and they seem to follow a similar climatic pattern. Both glacial cycles began with the expansion of outlet glaciers from the main ice sheet and the growth of coastal mountain glaciers, caused mainly by relatively high precipitation. The glaciers were active, warm-based and contained much debris. Later, the climate got colder and drier, and the coastal glaciers probably became cold-based and retreated. Glaciers were less dynamic during the later parts of the glacial cycles. During the Saalian the main ice sheet advanced over Jameson Land. A drift limit southeast of the Ugleelv area, assumed to be erosional (Möller *et al.* 1994), may be explained by variations in basal thermal regime, either by spatial variation during the advance or by temporal variation at the ice-sheet margin during climatic warming. During the middle-late Weichselian the Ugleelv Valley was ice free, although a long-lasting glacier existed in Scoresby Sund (Hansen *et al.* 1999).

Discussion

Glaciations and landsystems

Three types of ice-dynamic behaviour, characterised mainly by basal thermal regime, have been

recognised for the glaciations in my study areas. They are distinguished by different sediment-landform associations (papers III, IV). Climate and pre-existing topography initially control the type of glaciation, and thus the type of resulting landsystem, but as the ice sheet grows, subglacial bed conditions (substrate, hydrology) become increasingly important. For a single ice sheet, ice-dynamic behaviour may, and usually does, change with time and from place to place.

A warm-based 'surge-like' ice advance, with a margin that froze to its bed after reaching the maximum extent, has been inferred for the Late Weichselian Kara Sea glaciation on Taymyr, based mainly on the sediment-landform association of the NTZ (Fig. 3; paper III). An entirely cold-based ice would not have been able to reach that far inland without being very thick in its central parts on the shelf, and such a thick ice sheet is rejected by the lack of postglacial uplift (Lambeck 1995, 1996; Hjort *et al.* 2002). The older parts of the NTZ, older deformation tills elsewhere on Taymyr (unpublished data from this study; Funder *et al.* 1999; Möller *et al.* 2000) and the appearance of older moraines south of the Byrranga Mountains (Fig. 1; Kind & Leonov 1982) suggest that previous glacial events may have been similar. The southwestern sector of the Kara Sea ice sheet also seems to have been dynamic, at least partly warm-based and possessing a low-gradient profile during its maximum (Tveranger *et al.* 1999; Henriksen *et al.* 2001).

The present morphology of the NTZ is controlled by its buried glacier ice, melted-out supraglacial material and the glaciotectionic activity at its formation (Fig. 3). However, there are very few signs of glacial overriding north (proximal) of the NTZ (paper III). Landsystems similar to the NTZ are found in other parts of the Arctic (e.g. Sharpe 1988) and also along the former southern margins of the Laurentide (Mooers 1990; Ham & Attig 1996) and Scandinavian ice sheets (e.g. Wysota 1999), although in the latter areas the buried ice has melted long ago, leaving a hummocky moraine landscape. The interpretations from these other areas vary, but generally involve a warm-based ice-sheet lobe with compressive flow at its margin, located at the periphery of a large ice sheet in a relatively low-relief area and, at least partly, lying on top of permafrozen sedimentary bedrock or unconsolidated sediments. The conclusions drawn for the NTZ support the conception that transverse zones of hummocky moraines may not always

represent passive marginal stagnation but could rather be the remnants of actively formed zones of ice-cored moraine (Dyke & Savelle 2000).

The Greenland ice sheet expanded onto Jameson Land during the Saalian, leaving scattered erratic boulders and lodgement till (paper IV; Möller *et al.* 1994). Most likely the ice sheet was cold-based on the Jameson Land plateau, as on other slightly higher areas between faster-moving outlet glaciers in the fjords (cf. England *et al.* 2000; Wilch & Hughes 2000). However, the deposition of lodgement till requires a warm-based ice and a thawed bed must have developed temporarily.

The main sediments and landforms resulting from this glaciation are characterised by glacial meltwater. On Jameson Land valley erosion, e.g. of the Ugleelv Valley, replaced the lateral meltwater channels most commonly associated with the retreat of cold ice (e.g. Kleman *et al.* 1992; Dyke 1993; Ó Cofaigh *et al.* 1999). This could be due to the flat topography of the area, with a broad ice-sheet front that mainly allowed frontal, not lateral, drainage.

The valley glaciers that extended to Jameson Land from the Liverpool Land mountains were active, erosive and carried much debris. These glaciers left small end moraines in the Ugleelv Valley and also glacially eroded their substrate (U-shaped valleys, local till). The pre-Saalian topography is largely unknown, but at least during the youngest advance, the glaciers were confined to valleys (paper IV). They do, however, differ from ordinary valley glaciers in that they moved upvalley on Jameson Land.

Ice-sheet limits

The maximum extensions of the Weichselian Kara Sea ice sheets are fairly well known on the Taymyr Peninsula (this study; Hjort *et al.* 2002), although two different moraines have been suggested as alternatives for terminal moraines of the most widespread glaciation, the one in the Early Weichselian (Kind & Leonov 1982; Astakhov 1998; C. Siegert *et al.* 1999). Evidence for both one and two Early-Middle Weichselian glacial advances from the Kara Sea shelf have been found on the mainland between the Taymyr Peninsula and the Pechora River (this study; Manley *et al.* 2001; Henriksen *et al.* 2001), on Severnaya Zemlya (Bolshiyonov & Makeyev 1995) and on the adjacent shelf (Knies *et al.* 2000, 2001; Kleiber *et al.* 2001). These Kara Sea ice sheets are generally assumed to have been confluent with the Barents Sea ice sheets (e.g. Svendsen *et al.* 1999) and the geographical limits

are relatively well agreed upon.

The most questioned terminal ice margins are those for the Late Weichselian ice sheet and, especially, whether or not the ice was then confluent with the Barents Sea ice sheet. No evidence of inundation by a shelf-based ice has so far been recognised on Severnaya Zemlya (Bolshiyarov & Makeyev 1995; Raab *et al.* 2001; Berger *et al.* 2002; Per Möller pers.comm. 2002) or on the West Siberian and Russian mainland (Svendsen *et al.* 1999). Apart for Taymyr, the Late Weichselian ice-sheet limits are therefore believed to have been situated offshore (Fig. 1), and support for an ice sheet on the central Kara Sea shelf at that time is also provided by marine records (Polyak *et al.* 1997, 2002a, b). Recent marine-geological investigations seem to locate the southern limit of this ice sheet, based on till and moraine-ridge distribution, on the central shelf (Polyak *et al.* 2002a). So far the data are, however, inconclusive about whether the last Kara Sea ice sheet was separated from or confluent with the Barents Sea ice sheet to the west. Indications of glacial lakes and dammed north-flowing rivers are found in the southern Kara Sea and, thus, suggest a connection between the two ice sheets (Niessen *et al.* 2002; Polyak *et al.* 2002b). However, present-day land south of the Kara Sea does not seem to have been flooded (Mangerud *et al.* 1999; Svendsen 2001) and likely, the confluence was, if it existed, only brief, breaking up prior to 13 ka BP (Polyak *et al.* 2002b). The available data may favour a local ice mass on the central Kara Sea shelf, but are really too scarce yet to give a definite answer to this question.

As evidenced by western erratics and lodgement till, the Greenland ice sheet has, at some time, crossed both Jameson Land and Liverpool Land (Funder 1972). This glacial event is older than the Weichselian and has been placed in the Saalian (Scoresby Sund glaciation; Möller *et al.* 1994; Funder *et al.* 1998) and it has been proposed that the ice sheet terminated on the present shelf at this stage.

The boundaries between the till-covered plateau and the driftless areas on Jameson Land have been interpreted as caused by erosion (Möller *et al.* 1994). However, the eastern boundary may record a change from warm to cold basal conditions, or possibly, the terrestrial maximum limit of the main ice-sheet advance in Saalian time (Paper IV). If the latter is true, erratics on Jameson Land, Liverpool Land and other elevated areas along the East Greenland coast (Bretz 1935; Funder 1972; Funder & Hjort 1973) must have been brought there by one or several

earlier, even more extensive glaciations. Evidence of such glaciations is found for example in IRD records off southeast Greenland, where the largest IRD peaks are 1-4 Ma old (St. John & Kriesek 2002). It would perhaps also be easier to get the needed large increase in ice volume during longer cold periods and at a time before the efficient drainage system in the fjords was well developed.

The two advances from Liverpool Land have not previously been recognised and moraines in and along the Hurry Fjord have been thought to date from the Milne Land stade (cf. Table 3 in Paper IV), when glaciers briefly re-expanded along the East Greenland coast (Funder & Hjort 1973; Hjort 1979; Funder *et al.* 1998; Funder & Hansen 1999). Most of these moraines are undated and some of them, e.g. in the Gåseelv Valley (Lilliesköld & Salvigsen 1991) and at the mouth of the Hurry Fjord (Dowdeswell *et al.* 1994), may well belong to Liverpool Land mountain glaciation 2 (paper IV).

The nature of glacial cycles

Today both the Taymyr Peninsula and Jameson Land are situated in polar tundra environments (Köppen climate type ET; Henderson-Sellers & Robinson 1986), characterised by very low mean annual temperatures, little precipitation and continuous permafrost. During glacial periods the cold and the aridity were even more accentuated (Hahne & Melles 1999; Funder *et al.* 1998). For example, if precipitation is not reduced to almost zero during the Late Weichselian, ice-sheet models produce too much ice in the Kara Sea area to fit the geological record on the Taymyr Peninsula (M.J. Siegert *et al.* 1999; Siegert & Marsiat 2001). Due to the low temperatures, any ice that forms in these polar regions will initially be cold but may eventually, due to ice thickness, geothermal heat or friction, become warm-based.

The Weichselian glaciation in the Kara Sea region involved three glacial events of successively smaller amplitude (Fig. 2). The size reduction was a result of the successive westward shift of maximum glaciation with time, likely due to changes in atmospheric circulation (Velichko *et al.* 1997). Each glacial event called for the formation of an entire, new ice sheet, with the possible exception of the Middle Weichselian one (paper II; Hjort *et al.* 2002). During glacials the equilibrium line altitude (ELA) was probably lowered to sea level (M.J. Siegert *et al.* 1999; Grosswald & Hughes 2002) and ice sheets could form by ice accumulation on the exposed shelf (e.g. Barry *et al.* 1975) or from partly

grounded, thickening sea ice (Grosswald 1980; Hughes 1986).

A glaciation on East Greenland, on the other hand, requires ‘only’ the expansion of pre-existing glaciers, ice caps or the main ice sheet. Increased ice cover may be brought about by northward (increased precipitation) or southward (decreased temperatures) shifts of climatic zones, both leading to lowered ELA’s. The coastal glaciers and the Greenland ice sheet respond to different changes in climate and circulation and may expand and retreat out of phase (cf. Humlum 1999). A glacial cycle in the Scoresby Sund area may have started with both alpine glaciation in the coastal mountains and the expansion of outlet glaciers from the ice sheet (Fig. 4). During initial cooling there was still open water (a moisture source) close to the coast. Later, as sea ice expanded southwards, precipitation was reduced to glacial period levels and the coastal glaciers became less active, possibly changing to cold-based conditions. Similar switches in response to mass balance changes have been documented for glaciers on Svalbard (Glasser & Hambrey 2001). During the Saalian the ice sheet continued to expand, mainly due to low temperatures, while it remained relatively stationary during the Middle-Late Weichselian.

In both areas, Siberia and East Greenland, the early stages of a glacial cycle seem to have been the most dynamic, with extensive glacial advances of shelf-based ice, coastal glaciers and outlet glaciers from the ice sheet. Later, as the climate grew more severe (increased continentality and aridity), glaciations became more limited (Kara Sea) or static (Greenland). For example, the Late Weichselian ice advance onto the Taymyr Peninsula was probably caused by internal factors, rather than climatic (paper III), but nothing much happened with the relatively stable Greenland ice sheet in the Scoresby Sund area during the long Flakkerhuk stade (Hansen *et al.* 1999).

Deglaciations are usually more rapid than the growth of an ice sheet, mainly because accumulation is limited but ablation/calving is not (e.g. Letréguilly *et al.* 1991). The central parts of the Kara Sea ice sheets seem to have disappeared quickly, possibly aided by rising sea levels. In a strict sense, the marginal areas, e.g. the NTZ, have in fact not yet been deglaciated, as fossil glacier ice is still present under a thin cover of melted-out sediments (papers II, III). Glacier ice has also been reported from other areas covered by former Kara Sea ice sheets (Astakhov & Isayeva 1988; C. Siegert *et al.* 1999; Lokrantz & Ingólfsson submitted).

Conclusions

- The 750-km-long North Taymyr ice-marginal zone (NTZ) on the Taymyr Peninsula, Siberia, was formed along the front of ice sheets originating on the Kara Sea shelf. It records glacial events during the Early, Middle and Late Weichselian. The NTZ comprises large, ice-cored thrust-block moraines dotted with glaciokarst lakes, kames and small push moraines. At the distal margin are several deltas, subaqueous fans and sandur plains. Distal to the NTZ are laminated glaciolacustrine sediments deposited in lakes dammed by the ice sheets.

- The sedimentary record in the Ugleelv area on Jameson Land, East Greenland spans two glacial cycles. Glaciolacustrine sediments record a delta prograding from an outlet glacier in Scoresby Sund in the southwest. These sediments were later overridden by a glacier coming from the east (Liverpool Land) that deposited a local till and glaciofluvial sediments. After a period with periglacial conditions the Greenland ice sheet expanded over the area, depositing a till with regional distribution (Scoresby Sund glaciation). But the main impact was glaciofluvial erosion, including the erosion of the Ugleelv Valley, during deglaciation in the late Saalian. During the Weichselian a glacier advanced upvalley from the east, causing some erosion and deformation, and also depositing small end moraines. Aeolian sedimentation occurred in the Late Weichselian, showing there was no ice cover in the area at that time.

- Within my study areas, three types of ice-dynamic behaviour have been recognised. They are characterised by different sediment-landform associations, but share the trait of leaving few traces except for at their margins: (1) warm-based, ‘surge-like’ ice advance, with a margin that turned cold-based after reaching the maximum extent (the Kara Sea ice sheet); (2) mainly cold-based outer parts of an ice sheet, but with spatial and temporal changes in basal temperature (the Greenland ice sheet); and (3) mountain glaciation with valley glaciers (the Liverpool Land glaciers).

- The Kara Sea ice sheet was shelf/marine based and the ice advanced inward over land. It was unstable, experienced large shifts in areal extent and disappeared completely during interglacials. The glaciation of the Kara Sea had climatic causes, but internal ice dynamics may have governed ‘individual’ ice advances.

- The Greenland ice sheet is land-based and its ice advances towards the sea are strongly channelled by the fjords. Much due to the subglacial topography it is comparatively stable and the ice sheet remains during interglacials. The expansion or retreat of outlet glaciers in the fjords usually represents the major changes. Glaciations and ice advances are climatically driven.

- The most extensive Middle-Late Pleistocene glaciation took place during the Saalian in both the Kara Sea and Scoresby Sund areas. On Taymyr, three successively smaller glaciations occurred during the Weichselian, while the central Jameson Land plateau probably experienced only one minor Weichselian ice advance from the coastal mountains. See also Figs. 2 and 4.

Implications and ideas for the future

The work on Taymyr has delimited the eastern boundary of the Kara Sea ice sheet through time and, together with results from other QUEEN groups (Larsen *et al.* 1999; Thiede *et al.* 2001), disproved the idea of a great Late Weichselian Eurasian ice sheet (Peltier 1994; Grosswald & Hughes 2002). Another important part has been the documentation of the NTZ, which previously had only been outlined from satellite images (Kind & Leonov 1982; Isayeva 1984). So far, reliable absolute dates are relatively few and I would like to establish better chronological control of the glacial history of Taymyr. In determining the extent of the Late Weichselian Kara Sea ice sheet I find the recent results of marine-geological studies (Polyak *et al.* 2002a) very exciting and believe the combination of terrestrial and marine records will provide crucial answers. The sedimentary basins on Taymyr contain a long record of Quaternary environmental history (Kind & Leonov 1982) and it would also be possible to go further back in time and look at older glaciations, mainly south of the Byrranga Mountains, with a methodology similar to the one used in this study.

The Jameson Land results are roughly in agreement with previous studies (Möller *et al.* 1994; Funder *et al.* 1998) but provide more detail on the terrestrial glaciations in the otherwise fjord-dominated glacial record. To get a better idea of the ice-cover extent I would like to do detailed mapping of the plateau area, particularly of the glaciofluvial channels and till limits. It would also be interesting to look more closely at the late glacial–Holocene aeolian sediments and at what has usually been interpreted as ‘historical’ moraines. The possibilities

of establishing a good chronology are promising, since the aeolian deposits are excellent for luminescence dating and I also think that cosmogenic isotope dating could be fruitful, for example for the older glaciations in the plateau area.

Taken together, my work has given some insights into the course of glacial cycles and into the dynamic behaviour of ice sheets in two different polar areas. It would certainly be interesting to further compare these results with those from other formerly glaciated areas in various parts of the world, and also with different glaciological models.

Acknowledgements

First of all I would like to thank my supervisors, Per Möller, Christian Hjort and Lena Adrielsson for introducing me to the exciting area of polar research and for being a great supervising team. I am also grateful to my fellow PhD students, especially Gert Pettersson, Joachim Albrecht and Anna Broström, and everybody else in and outside the Department of Geology who in some way has helped me with my work. During fieldwork I enjoyed the company and help of Saskia Eriksson, Maksim Pavlov, Oleg Antonov, Dmitry Bolshiyakov, Per Bergenrud and Martin Adrielsson, in addition to my supervisors. Laila Johannesson is worth a ‘Thank you’ for all our discussions regarding both science and life as a PhD student over the past years. Special thanks go to my family for supporting me and, not least, for helping out with practicalities when necessary. Thanks also to Mark Benfer who checked the English language.

My position has been financed by the Faculty of Sciences at Lund University. The Taymyr project has been part of the European Union financed ‘Eurasian Ice Sheets’ programme (contract no. ENV4-CT97-0563) and the European Science Foundation’s QUEEN (Quaternary Environment of the Eurasian North) programme. Additional financial support has come from the Royal Swedish Academy of Sciences (KVA) and the Swedish Environmental Protection Agency (SNV). The logistics in Siberia were organised and to a large part paid for by the Swedish Polar Research Secretariat. The Jameson Land expedition was financed through grants from Helge Ax:son Johnson’s foundation, the Crafoord foundation, the Royal Physiographic Society in Lund, the Swedish Society for Anthropology and Geography (SSAG),

the Royal Swedish Academy of Sciences and the Faculty of Sciences at Lund University. The Danish Polar Centre and the Swedish Polar Research Secretariat provided logistic support. During my PhD studies I have received financial support for participating in conferences and courses from the Department of Geology, Lund University (Wahlbom's fund), the Nordic Academy for Advanced Study (NorFA) and Uppsala University (Knigge's foundation).

Svensk sammanfattning

Glacialgeologi och isdynamik vid randen av två istäckan, på Tajmyrhalvön i Sibirien och på Jameson Land, Östgrönland

Arktis utgör en viktig del i Jordens miljö och vad som händer där har stor påverkan på det globala klimatet, samtidigt som området är mycket känsligt för förändringar. Under kvartärtiden har stora isar flera gånger täckt Skandinavien, Ryssland, Sibirien och Nordamerika och täcker än idag stora delar av Grönland. Isarna har lämnat spår (landformer och sediment) efter sig och genom att undersöka dessa kan man ta reda på var och när istäckena fanns och dessutom hur de fungerade.

Den grundläggande orsaken till varför istäckan växer till och sedan försvinner är klimatet som förändras med hur Jordens bana förhåller sig till Solen, det vill säga hur solinstrålningen periodiskt varierar. En isframstöt har tidigare i allmänhet tolkats som att klimatet blivit överlag kallare, men andra faktorer, t.ex. närheten till öppet hav (som nederbördskillan), topografi och förhållandena vid isens botten, kan också påverka var isen finns och hur den rör sig. För att kunna tolka de geologiska arkiven rätt är det därför viktigt att man förstår samspillet mellan landskap, is och klimat.

Inom mitt avhandlingsarbete har jag undersökt den så kallade Norra Tajmyrisrandlinjen på Tajmyrhalvön i arktiska Sibirien och Ugleelvdalen på Jameson Land, Östgrönland, två geografiskt vitt skilda områden men som båda har varit nedisade och legat nära randen av stora istäckan. Syftet med mitt arbete har varit att rekonstruera nedisningshistorien i dessa områden, inklusive de miljöer som funnits och de processer som varit inblandade. Jag har undersökt landformer och sedimentlagerföljder,

studerat flyg- och satellitbilder samt använt mig av så kallade geografiska informationssystem för att få en helhetsbild av glacialgeologin i de två områdena. Åldersbestämningar har gjorts med hjälp av två olika metoder: optiskt stimulerad luminescens (OSL) och kol-14 (^{14}C).

Kontinentalsockeln i det grunda Karahavet, som torrläggs under istider då vatten binds i inlandsisarna, har varit nedisad tre gånger under den senaste istiden (Weichsel; 115 000 - 10 000 år före nutid). Varje is var mindre än den föregående. Istäckena ledde till att floderna på Tajmyrhalvön tvingades rinna söderut och, under de två äldre skedena, också till att sjöar dämades upp söder om iskanten. Den yngsta isframstötten, men troligen också de äldre, skedde snabbt och var bottenmältande. När isen nått sitt yttersta läge fick den en bottenfrusen kant där is och sediment trycktes samman och deformades. De äldsta delarna av Norra Tajmyrisrandlinjen bildades ursprungligen för ca 80 000 år sedan (Tidigweichsel), under isens reträtt från sin största utbredning. Senare isframstötter, för ca 65 000 år sedan (Mellanweichsel) och för mindre än 20 000 år sedan (Senweichsel), byggde på den ytterligare. Israndlinjen består av så kallade landformssystem bildade vid kanten av och ovanpå isen och domineras av 50-80 m höga och 2-3 km breda moränryggar. Stora områden underlagras fortfarande av gammal glaciäris och ovanpå denna håller än idag ett landskap med otaliga sjöar och sandkullar (kames) på att bildas i samband med att isen sakta smälter. Landformsystemet framför israndlinjen karakteriseras av sediment och landformer avsatta antingen under vatten eller på torra land, beroende på höjdläget och tidpunkten då de bildades.

Sedimentlagerföljden i Ugleelvdalen på Grönland omfattar tre moräner, samt issjö- och isälvsavlagringar och vindavsatta sediment. De äldsta delarna utgörs av ett delta som byggdes ut i en sjö uppdämd av en fjordglaciär i Scoresby Sund i väster, tidigt under den föregående istiden (Saale; mer än 130 000 år före nutid). Något senare rörde sig en glaciär från Liverpool Land i öster in över Jameson Land. Glaciären var bottenmältande och avsatte isälvsand och morän, bestående av lokalt material, samt små moränryggar. Därefter, under den senare delen av Saale, följde "den stora nedisningen" då den grönländska inlandsisen överkred området från väster. Isen var bottenfrusen men blev tidvis och på vissa ställen bottenmältande. Dess största påverkan på landskapet var genom smältvattenserrosion, bland annat nedskärning i Ugleelvdalen. Under större delen av Weichsel-isti-

den var Ugleelv-området isfritt, men vid ett tillfälle trängde en dalglaciär från Liverpool Land upp i dalgången. Förhållandena var mer föränderliga under de tidiga delarna av istidscyklerna än under de senare, då klimatet var extremt kallt och torrt.

Mina undersökningar tyder på att Karahavsisen och Grönlandsisen betar sig på olika sätt under en istidscykel. Karahavsisen, som är centrerad på kontinentalsockeln, förändras mycket i storlek och försvinner helt under värmeperioderna, då havet dessutom dränker sockeln. Enskilda isframstötter verkar styras av inre mekanismer snarare än av klimatförändringar. Den grönländska inlandsisen är däremot förhållandevis stabil, den finns av framför allt topografiska orsaker kvar under värmeperioderna (som idag) och reagerar främst på klimatförändringar genom utvidgning eller tillbakadragande av glaciärer i fjordarna.

References

- Adrielson, L. 2001: Subglacial hydrology and bed deformation dynamics during a Late Weichselian ice lobe surge in the Öresund strait area, south Scandinavia. *In: Volume of abstracts, Palaeo-ice stream international symposium, Aarhus, Denmark*. 7. Also at <http://www.inqua.au.dk/cog/abstracts.pdf> [18 July 2002].
- Arkhipov, S.A., Ehlers, J., Johnson, R.G. & Wright Jr, H.E. 1995: Glacial drainage towards the Mediterranean during the Middle and Late Pleistocene. *Boreas* 24, 196-206.
- Astakhov, V. 1992: The last glaciation in West Siberia. *Sveriges Geologiska Undersökning Ca 81*, 21-30.
- Astakhov, V. 1997: Late glacial events in the central Russian Arctic. *Quaternary International* 41/42, 17-25.
- Astakhov, V. 1998: The last ice sheet of the Kara Sea: terrestrial constraints on its age. *Quaternary International* 45/46, 19-28.
- Astakhov, V.I. & Isayeva, L.L. 1988: The 'Ice Hill': an example of 'retarded deglaciation' in Siberia. *Quaternary Science Reviews* 7, 29-40.
- Barry, R.G., Andrews, J.T. & Mahaffy, M.A. 1975: Continental ice sheets – conditions for growth. *Science* 190, 979-981.
- Berger, A., Imbrie, J., Hays, J., Kukla, G. & Salzman, B. 1984: *Milankovitch and climate: understanding the response to astronomical forcing*. Reidel, Dordrecht, Netherlands.
- Berger, G.W., Melles, M. & Raab, A. 2002: Limiting extent of ice sheets in the Russian high-arctic during isotope stages 2-3 from IRSL dating of lake sediments, Taymyr Peninsula. *Poster abstract, ARCSS workshop, Seattle, USA*. Also at http://siempre.arcus.org/4DACTION/wi_pos_displayAbstract/4/305 [10 July 2002].
- Bolshiyarov, D.Yu. & Makeev, V.M. 1995: *Severnaya Zemlya archipelago: glaciation and environmental history*. Gidrometeoizdat, St Petersburg. 214 pp. (in Russian)
- Bretz, J.H. 1935: Physiographic studies in East Greenland. *In Boyd, L.A. (ed.): The fiord region of East Greenland*. American Geographical Society, Special publication 18; 159-267.
- Clark, P.U. 1994: Unstable behavior of the Laurentide Ice Sheet over deforming sediment and its implications for climate change. *Quaternary Research* 41, 19-25.
- Clark, P.U., Alley, R.B. & Pollard, D. 1999: Northern Hemisphere ice-sheet influences on global climate change. *Science* 286, 1104-1111.
- Dowdeswell, J.A., Uenzelmann-Neben, G., Whittington, R.J. & Marienfeld, P. 1994: The Late Quaternary sedimentary record in Scoresby Sund, East Greenland. *Boreas* 23(4), 294-310.
- Dyke, A.S. 1993: Landscapes of cold-centred Late Wisconsinan ice caps, Arctic Canada. *Progress in Physical Geography* 17(2), 223-247.
- Dyke, A.S. & Savelle, J.M. 2000: Major end moraines of Younger Dryas age on Wollaston Peninsula, Victoria Island, Canadian Arctic: implications for paleoclimate and for formation of hummocky moraine. *Canadian Journal of Earth Sciences* 37, 601-619.
- Eastman, J.R. 1997: *Idrisi for Windows*, Version 2. Clark Laboratories, Clark University, Worcester, MA.
- England, J., Smith, I.R. & Evans D.J.A. 2000: The last glaciation of east-central Ellesmere Island, Nunavut: ice-dynamics, deglacial chronology, and sea level change. *Canadian Journal of Earth Sciences* 37, 1355-1371.
- Funder, S. 1972: Remarks on the Quaternary geology of Jameson Land and adjacent areas, Scoresby Sund, East Greenland. *Rapport Grönlands geologiske undersøgelse* 48, 93-98.
- Funder, S. & Hansen, L. 1999: The Greenland ice sheet - a model for its culmination and decay during and after the last glacial maximum. *Bulletin of the Geological Society of Denmark* 42, 137-152.
- Funder, S. & Hjort, C. 1973: Aspects of the

- Weichselian chronology in central East Greenland. *Boreas* 2(2), 69-84.
- Funder, S., Hjort, C., Landvik, J.Y., Nam, S-I., Reeh, N. & Stein, R. 1998: History of a stable ice margin – East Greenland during the middle and upper Pleistocene. *Quaternary Science Reviews* 17, 77-123.
- Funder, S., Riazanova, M., Rydlevski, A. & Seidenkrantz, M.-S. 1999: Late Quaternary events in northern Siberia – preliminary results of field work on coastal Taymyr. In: *Abstract volume of the third QUEEN workshop, Øystese, Norway*; 8. Also at <http://www.geomar.de/~hbauch/king/html/99abst.html> [2 July 2002].
- Glasser, N.F. & Hambrey, M.J. 2001: Styles of sedimentation beneath Svalbard valley glaciers under changing dynamic and thermal regimes. *Journal of the Geological Society of London* 158(4), 697-707.
- GLOBE Task Team and others (Hastings, D.A., Dunbar, P.K., Elphinstone, G.M., Bootz, M., Murakami, H., Maruyama, H., Masaharu, H., Holland, P., Payne, J., Bryant, N.A., Logan, T.L., Muller, J.-P., Schreier, G. & MacDonald, J.S. (Eds.)), 1999: *The Global Land One-kilometer Base Elevation (GLOBE) Digital Elevation Model, Version 1.0*. National Oceanic and Atmospheric Administration, National Geophysical Data Center, Colorado, U.S.A. Digital data base on the World Wide Web, <http://www.ngdc.noaa.gov/seg/topo/globe.shtml> [14 June 1999].
- Grosswald, M.G. 1980: Late Weichselian ice sheet of northern Eurasia. *Quaternary Research* 13, 1-32.
- Grosswald, M.G. & Hughes, T.J. 2002: The Russian component of an Arctic Ice Sheet during the Last Glacial Maximum. *Quaternary Science Reviews* 21, 121-146.
- Hahne, J. & Melles, M. 1999: Climate and vegetation history of the Taymyr Peninsula since Middle Weichselian time – palynological evidence from lake sediments. In Kassens, H., Bauch, H.A., Dmitrenko, I.A., Eicken, H., Hubberten, H.W., Melles, M., Thiede, J. & Timokhov, L.A. (eds.): *Land-ocean systems in the Siberian Arctic: Dynamics and history*. Springer-Verlag, Berlin; 407-423.
- Ham, N.R. & Attig, J.W. 1996: Ice wastage and landscape evolution along the southern margin of the Laurentide Ice Sheet, north-central Wisconsin. *Boreas* 25, 171-186.
- Hansen, L., Funder, S., Murray, A.S. & Mejdahl, V. 1999: Luminescence dating of the last Weichselian glacier advance in East Greenland. *Quaternary Geochronology* 18, 179-190.
- Henderson-Sellers, A. & Robinson, P.J. 1986: *Contemporary climatology*. Longman Scientific & Technical, Harlow. 439 pp.
- Henriksen, M., Mangerud, J., Maslenikova, O., Matiouchkov, A. & Tveranger, J. 2001: Weichselian stratigraphy and glaciotectonic deformation along the lower Pechora River, Arctic Russia. *Global and Planetary Change* 31, 297-319.
- Hjort, C. 1979: Glaciation in northern East Greenland during the Late Weichselian and Early Flandrian. *Boreas* 8, 281-296.
- Hjort, C., Möller, P. & Alexanderson, H. 2002: Weichselian glaciation of the Taymyr Peninsula, Siberia. In Ehlers, J. & Gibbard, P.L. (eds): *Quaternary glaciations – extent and chronology. Vol. 1, Europe*. Quaternary Book Series, Elsevier, Amsterdam; 000-000/in press
- Hughes, T.J. 1986: The marine ice transgression hypothesis. *Geografiska Annaler* 69A(2), 237-250.
- Humlum, O. 1999: Late-Holocene climate in central West Greenland: meteorological data and rock-glacier isotope evidence. *The Holocene* 9(4), 581-594.
- Isayeva, LL. 1984: Late Pleistocene glaciation of north-central Siberia. In Velichko, A.A. (ed.): *Late Quaternary environments of the Soviet Union*. Univ. Minnesota Press, Minneapolis; 21-30.
- Johnsen, S.J., Clausen, H.B., Dansgaard, W., Gundestrup, N.S., Hansson, M., Jonsson, P., Steffensen, J.P. & Sveinbjörnsdóttir, A.E. 1992: A “deep” ice core from East Greenland. *Meddelelser om Grønland, Geoscience* 29, 3-22.
- Kind, N.V. & Leonov, B.N. (eds). 1982: *Antropogen Taymyra*. Nauka, Moscow. 181 pp. (in Russian)
- Kleiber, H.P., Niessen, F. & Weiel, D. 2001: The Late Quaternary evolution of the western Laptev Sea continental margin, Arctic Siberia – implications from sub-bottom profiling. *Global and Planetary Change* 31, 105-124.
- Kleman, J., Borgström, I., Robertsson, A.-M. & Lilliesköld, M. 1992: Morphology and stratigraphy from several deglaciations in the Transtrand Mountains, western Sweden. *Journal of Quaternary Science* 7, 1-17.
- Knies, J., Nowaczyk, N., Muller, C., Vogt, C. & Stein, R. 2000: A multiproxy approach to reconstruct the environmental changes along the Eurasian continental margin over the last 150 000 years. *Marine Geology* 163, 317-344

- Knies, J., Kleiber, H.-P., Matthiessen, J., Müller, C. & Nowaczyk, N. 2001: Marine ice-rafted debris records constrain maximum extent of Saalian and Weichselian ice-sheets along the northern Eurasian margin. *Global and Planetary Change* 31, 45-64.
- Lambeck, K. 1995: Constraints on the Late Weichselian ice sheet over the Barents Sea from observations of raised shorelines. *Quaternary Science Reviews* 14, 1-16.
- Lambeck, K. 1996: Limits on the areal extent of the Barents Sea ice sheet in Late Weichselian time. *Global and Planetary Change* 12, 41-51.
- Lambeck, K., Smither, C. & Johnston, P. 1998: Sea-level change, glacial rebound and mantle viscosity for northern Europe. *Geophysical Journal International* 134, 102-144.
- Larsen, E., Funder, S. & Thiede, J. (eds.) 1999: Late Quaternary history of northern Russia and adjacent shelves. *Boreas* 28(1), 242 pp.
- Letréguilly, A., Huybrechts, P. & Reeh, N. 1991: Steady-state characteristics of the Greenland ice sheet under different climates. *Journal of Glaciology* 37(125), 149-157.
- Lilliesköld, M. & Salvigsen, O. 1991: Glacial features and stratigraphy on eastern Jameson Land. In Möller, P., Hjort, C. & Ingólfsson, Ó. (eds.): *The last interglacial-glacial cycle: preliminary report on the PONAM fieldwork on Jameson Land and Scoresby Sund, East Greenland*. LUNDQUA Report 33; 111-116.
- Lokrantz, H. & Ingólfsson, O. submitted: Origin of a massive ground ice body on Yugorski Peninsula, Arctic Russia: buried glacier ice or intrasedimental segregation ice? In Lokrantz, H. 2002: *Pleistocene stratigraphy, ice sheet history and environmental development in the southern Kara Sea, Arctic Russia*. Göteborg University, Earth Sciences Centre, Thesis A77; appendix V.
- Mangerud, J., Svendsen, J.I. & Astakhov, V.I. 1999: Age and extent of the Barents and Kara ice sheets in Northern Russia. *Boreas* 28(1), 46-80.
- Mangerud, J., Astakhov, V., Jakobsson, M. & Svendsen, J.I. 2001: Huge ice-age lakes in Russia. *Journal of Quaternary Science* 16(8), 773-777.
- Manley, W.F., Lokrantz, H., Gataullin, V., Ingólfsson, Ó., Forman, S.L. & Andersson, T. 2001: Late Quaternary stratigraphy, radiocarbon chronology, and glacial history at Cape Shpindler, southern Kara Sea, Arctic Russia. *Global and Planetary Change* 31, 239-254.
- Mooers, H.D. 1990: A glacial-process model: The role of spatial and temporal variations in glacier thermal regime. *Geological Society of America Bulletin* 102, 243-251.
- Möller, P., Hjort, C., Adrielsson, L. & Salvigsen, O. 1994: Glacial history of interior Jameson Land, East Greenland. *Boreas* 23, 320-348.
- Möller, P., Bolshiyarov, D.Yu. & Bergsten, H. 1999a: Weichselian geology and palaeoenvironmental history of the central Taymyr Peninsula, Siberia, indicating no glaciation during the last global glacial maximum. *Boreas* 28(1), 92-114.
- Möller, P., Bolshiyarov, D.Y., Jansson, U. & Schneider, G.V. 1999b: The "Ledianaya gravel event" – a marker of the last glacioisostatic-induced depression along the Byrranga Mountains and south thereof. In: *Abstract volume of the third QUEEN workshop, Øystese, Norway*; 41. Also at <http://www.geomar.de/~hbauch/king/html/99abst.html> [2 July 2002].
- Möller, P., Fedorov, G., Jönsson, C. & Pavlov, M. 2000: The Cape Chelyuskin area on northern Taymyr – outside the Weichselian glaciations? In: *Abstract volume of the fourth QUEEN workshop, Lund, Sweden*; 35. Also at <http://www.geomar.de/~hbauch/king/html/00abst.html> [2 July 2002].
- Niessen, F., Stein, R. & Dittmers, K. 2002: LGM glaciation and Siberian river runoff in the Kara Sea: results from the RV "Boris Petrov" expedition 2001 (SIRRO-project). *Terra Nostra* 2002(3), 78.
- Ó Cofaigh, C., Lemmen, D.S., Evans, D.J.A. & Bednarski, J. 1999: Glacial landform-sediment assemblages in the Canadian High Arctic and their implications for late Quaternary glaciation. *Annals of Glaciology* 28, 195-201.
- Oerlemans, J. 1982: Glacial cycles and ice-sheet modelling. *Climatic change* 4, 353-374.
- Pavlidis, Yu.A., Dunayev, N.N. & Shcherbakov, F.A. 1997: The late Pleistocene of the arctic Eurasian shelves. *Quaternary International* 41/42, 3-9.
- Peltier, W.R. 1994: Ice-age paleotopography. *Science* 265(5169), 195-201.
- Polyak, L., Forman, S.L., Herlihy, F.A., Ivanov, G. & Krinitsky, P. 1997: Late Weichselian deglacial history of the Svyatova (Saint) Anna Trough, northern Kara Sea, Arctic Russia. *Marine Geology* 143, 169-188.
- Polyak, L., Gataullin, V. & Hidrolog-2001 shipboard party 2002a: New data on the glacial history of the northern Kara Sea. In: *Abstract volume of the sixth QUEEN workshop, Spiez, Switzerland*; 44. Also at <http://www.geomar.de/~hbauch/king/html/02abst.html> [2 July 2002].

- Polyak, L., Levitan, M., Khusid, T., Merklin, L. & Mukhina, V. 2002b: Variations in the influence of riverine discharge on the Kara Sea during the last deglaciation and the Holocene. *Global and Planetary Change* 32, 291-309.
- Raab, A., Melles, M. & Hubberten, H.-W. 2001: Weichselian and Holocene environmental history of Severnaya Zemlya, northern central Siberia, as inferred from Changeable Lake sediments. In: *EUG XI, Strasbourg, France, Journal of Conference Abstracts* 6(1); 216. Also at <http://www.campublic.co.uk/EUGXI/EVO6.pdf> [10 July 2002].
- Rutter, N. 1995: Problematic ice sheets. *Quaternary International* 28, 19-37.
- Sharpe, D.R. 1988: Late glacial landforms of Wollaston Peninsula, Victoria Island, Northwest Territories: product of ice-marginal retreat, surge, and mass stagnation. *Canadian Journal of Earth Sciences* 25, 262-279.
- Siegert, C., Derevyagin, A.Y., Shilova, G.N., Hermichen, W.D. & Hiller, A. 1999: Paleoclimatic indicators from permafrost sequences in the eastern Taymyr lowland. In Kassens, H., Bauch, H.A., Dmitrenko, I.A., Eicken, H., Hubberten, H.W., Melles, M., Thiede, J. & Timokhov, L.A. (eds.): *Land-ocean systems in the Siberian Arctic: Dynamics and history*. Springer-Verlag, Berlin; 477-499.
- Siegert, M.J., Dowdeswell, J.A. & Melles, M. 1999: Late Weichselian glaciation of the Russian High Arctic. *Quaternary Research* 52, 273-285.
- Siegert, M.J. & Marsiat, I. 2001: Numerical reconstructions of LGM climate across the Eurasian Arctic. *Quaternary Science Reviews* 20, 1595-1605.
- St. John, K.E.K. & Krissek, L.A. 2002: The late Miocene to Pleistocene ice-rafting history of southeast Greenland. *Boreas* 31(1), 28-35.
- Stein, R., Nam, S.-I., Grobe, H. & Hubberten, H. 1996: Late Quaternary glacial history and short-term ice-rafted debris fluctuations along the East Greenland continental margin. In Andrews, J.T., Austin, W.E.N., Bergsten, H. & Jennings, A.E. (eds.): *Late Quaternary palaeoceanography of the North Atlantic margins*. Geological Society Special Publication 111; 135-151.
- Svendsen, J.I., Astakhov, V.I., Bolshiyakov, D.Yu., Demidov, I., Dowdeswell, J.A., Gataullin, V., Hjort, C., Hubberten, H.W., Larsen, E., Mangerud, J., Melles, M., Möller, P., Saarnisto, M. & Siegert, M. 1999: Maximum extent of the Eurasian ice sheets in the Barents and Kara Sea region during the Weichselian. *Boreas* 28(1), 234-242.
- Svendsen, J.I. (ed.) 2001: *Eurasian Ice Sheets*. Final report 1998-2000 to the Commission of the European Communities' Environment and Climate Research Programme.
- Thiede, J., Bauch, H.A., Hjort, C. & Mangerud, J. (eds.) 2001: The late Quaternary stratigraphy and environments of northern Eurasia and the adjacent seas – new contributions from QUEEN. *Global and Planetary Change* 31. 474 pp.
- Tveranger, J., Astakhov, V., Mangerud, J. & Svendsen, J.I. 1999: Surface form of the south-western sector of the last Kara Sea ice sheet. *Boreas* 28(1), 81-91.
- Velichko, A.A., Kononov, Yu.M. & Faustova, M.A. 1997: The last glaciation of Earth: size and volume of the ice sheets. *Quaternary International* 41/42, 43-51.
- Wilch, E. & Hughes, T.J. 2000: Calculating basal thermal zones beneath the Antarctic ice sheet. *Journal of Glaciology* 46(153), 297-310.
- Wysota, W. 1999: Ice sheet maximum limit of the Vistulian Glaciation in the mid-eastern Chełmno-Dobrzyń Lakeland, northern Poland. *Geological Quarterly* 43(2), 189-202.

