



# LUND UNIVERSITY

## Rapidly Emerging Role of Whole Body Vibration Therapy in the Management of Neurologic Diseases Besides Polio Response

Brogårdh, Christina; Flansbjerg, Ulla-Britt; Lexell, Jan

*Published in:*  
Archives of Physical Medicine and Rehabilitation

*DOI:*  
[10.1016/j.apmr.2010.11.019](https://doi.org/10.1016/j.apmr.2010.11.019)

2011

[Link to publication](#)

*Citation for published version (APA):*  
Brogårdh, C., Flansbjerg, U.-B., & Lexell, J. (2011). Rapidly Emerging Role of Whole Body Vibration Therapy in the Management of Neurologic Diseases Besides Polio Response. *Archives of Physical Medicine and Rehabilitation*, 92(4), 677-678. <https://doi.org/10.1016/j.apmr.2010.11.019>

*Total number of authors:*  
3

### General rights

Unless other specific re-use rights are stated the following general rights apply:  
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117  
221 00 Lund  
+46 46-222 00 00





LUND UNIVERSITY  
Faculty of Medicine

---

# LUP

*Lund University Publications*

Institutional Repository of Lund University

---

This is an author produced version of a paper published in Archives of Physical Medicine and Rehabilitation. This paper has been peer-reviewed but does not include the final publisher proof-corrections or journal pagination.

Citation for the published paper:  
Christina Brogårdh, Ulla-Britt Flansbjer, Jan Lexell

"Rapidly Emerging Role of Whole Body Vibration Therapy in the Management of Neurologic Diseases Besides Polio Response"

Archives of Physical Medicine and Rehabilitation  
2011 92(4), 677 - 678

<http://dx.doi.org/10.1016/j.apmr.2010.11.019>

Access to the published version may require journal subscription.

Published with permission from: Elsevier Science Inc

Respon to Ms. Ref. No.: ARCHIVES-PMR-D-10-00888R1  
Title: Rapidly emerging role of whole body vibration therapy in the management of neurological diseases besides polio

**Christina Brogårdh, RPT, PhD<sup>1,2</sup>, Ulla-Britt Flansbjer, RPT, PhD<sup>1,2</sup>  
and Jan Lexell, MD, PhD<sup>1,2</sup>**

From <sup>1</sup>Department of Rehabilitation Medicine, Skåne University Hospital, Lund, and  
<sup>2</sup>Division of Rehabilitation Medicine, Department of Clinical Sciences, Lund, Lund  
University, Lund, Sweden

Corresponding author: Christina Brogårdh, RPT, PhD, Department of Rehabilitation  
Medicine, Skåne University Hospital, SE-221 85 Lund, Sweden.  
E-mail: christina.brogardh@skane.se

To Editor,

We thank xxx et al. for their comments on our brief report regarding whole-body vibration (WBV) training in persons with late effects of polio (1). Their overall conclusion that WBV therapy has a rapidly emerging role in the management of neurological disorders is equally thought provoking. Despite being on the market for over a decade and tested in different neurological disorders, there are not yet any clear results regarding positive effects of WBV training. Even though some studies have reported improvements (2,3), others have not found it to be better than traditional interventions (3-6). The reported effects are usually small in comparison with other established interventions and often within the limits of normal variations or measurement errors. We have performed assessments of reliability (7-10) of muscle strength and gait performance in patients with late effects of polio and stroke and have shown that increases on group level must be at least 7%, and often greater, to be considered clinically real improvements. Improvements reported after WBV training are often below these values and therefore would not indicate a clinically real improvement.

Furthermore, there are few randomized controlled trials (RCT) of WBV training in patients with neurological disorders. Those that have used such structured design often report no or small effects. The authors refer to a randomized cross-over study by Schyns et al. (5) and state that balance improved and muscle spasm was attenuated after WBV training in patients with multiple sclerosis (MS). However, balance was not primarily assessed in that study and other outcomes measures, such as gait performance, were not significant. In addition, the authors of that study themselves concluded that "...there is limited evidence that the addition of whole body vibration provides any additional improvements". A recently published RCT of patients with MS (6) also reported that WBV training over 20 weeks had no effects on leg muscle performance and functional capacity. In patients with Parkinson's disease (PD) (3) and stroke (4), WBV training was not found to be more effective than conventional training regarding gait performance (3), balance or activities of daily living (4). In our opinion, studies so far have not been able to show that WBV training is a viable treatment for neurological disorders.

An important aspect of the design of WBV training studies is the choice of outcome measures. Some studies have primarily assessed body functions for example muscle strength and balance but there is little evidence of transfer of gains to activities of daily living (for example gait performance) after WBV training. From a rehabilitation perspective, an

intervention is considered to be effective when it reduces not only impairments but also activity limitations and subsequently participation restrictions.

We do agree with xxx et al. that there is insufficient evidence of the effects of WBV training and therefore a need for larger RCTs in patients with neurological disorders, using sound reliable outcome measurements. We have recently completed a double blind RCT of WBV training with two different amplitudes in chronic stroke patients and the findings will be reported at upcoming international conferences during 2011.

## References

1. Brogårdh C, Flansbjerg UB, Lexell J. No effects on muscle strength and gait performance after whole-body vibration training in persons with late effects of polio: a pilot study. *Arch Phys Med Rehabil* 2010; 91:1474- 7.
2. Wunderer K, Schabrun SM, Chipchase LS. Effects of whole body vibration on strength and functional mobility in multiple sclerosis. *Physiother Theory Pract* 2010; 26: 374-84.
3. Ebersbach G, Edler D, Kaufhold O, Wissel J. Whole body vibration versus conventional physiotherapy to improve balance and gait in Parkinson's disease. *Arch Phys Med Rehabil* 2008; 89: 399-403.
4. van Nes IJW, Latour H, Schils F, Meijer R, van Kuijk A, Geurts ACH. Long-term effects of 6-week whole-body vibration on balance recovery and activities of daily living in the postacute phase of stroke. *Stroke* 2006; 37: 2331-35.
5. Schyns F, Paul L, Finlay K, Ferguson C, Noble E. Vibration therapy in multiple sclerosis: a pilot study exploring its effects on tone, muscle force, sensation and functional performance. *Clin Rehabil* 2009; 23: 771-81.
6. Broekmans T, Roelants M, Alders G, Feys P, Thijs H, Eijnde B. Exploring the effects of a 20-week whole-body vibration training programme on leg muscle performance and function in persons with multiple sclerosis. *J Rehabil Med* 2010; 42: 866-72.
7. Flansbjerg U-B, Lexell J. Reliability of knee extensor and flexor muscle strength measurements in persons with late effects of polio. *J Rehabil Med* 2010; 42: 588-92.
8. Flansbjerg U-B, Lexell J. Reliability of gait performance tests in individuals with late effects of polio. *PM&R* 2010; 2: 125-31.
9. Flansbjerg UB, Holmbäck AM, Downham D, Patten C, Lexell J. Reliability of gait performance tests in men and women with hemiparesis after stroke. *J Rehabil Med* 2005; 37: 75-82.

10. Flansbjer UB, Holmbäck AM, Downham D, Lexell J. What change in isokinetic knee muscle strength can be detected in men and women with hemiparesis after stroke? *Clin Rehabil* 2005; 19: 514-22.