

LUND UNIVERSITY

Relativistic Coulomb excitation of 88Kr

Moschner, K.; Blazhev, A.; Jolie, J.; Warr, N.; Boutachkov, P.; Bednarczyk, P.; Sieja, K.; Algora, A.; Ameil, F.; Bentley, M. A.; Brambilla, S.; Braun, N.; Camera, F.; Cederkäll, J.; Corsi, A.; Danchev, M.; Dijulio, D.; Fahlander, C.; Gerl, J.; Giaz, A.; Golubev, P.; Górska, M.; Grebosz, J.; Habermann, T.; Hackstein, M.; Hoischen, R.; Kojouharov, I.; Kurz, N.; Mărginean, N.; Merchán, E.; Möller, T.; Naqvi, F.; Nara Singh, B. S.; Nociforo, C.; Pietralla, N.; Pietri, S.; Podolyák, Zs; Prochazka, A.; Reese, M.; Reiter, P.; Rudigier, M.; Rudolph, D.; Sava, T.; Schaffner, H.; Scruton, L.; Taprogge, J.; Thomas, T.; Weick, H.; Wendt, A.; Wieland, O.

Published in: Physical Review C - Nuclear Physics

DOI: 10.1103/PhysRevC.94.054323

2016

Document Version: Publisher's PDF, also known as Version of record

Link to publication

Citation for published version (APA):

Moschner, K., Blazhev, A., Jolie, J., Warr, N., Boutachkov, P., Bednarczyk, P., Sieja, K., Algora, A., Ameil, F., Bentley, M. A., Brambilla, S., Braun, N., Camera, F., Cederkäll, J., Corsi, A., Danchev, M., Dijulio, D., Fahlander, C., Gerl, J., ... Wollersheim, H. J. (2016). Relativistic Coulomb excitation of 88Kr. *Physical Review C - Nuclear Physics*, *94*(5), Article 054323. https://doi.org/10.1103/PhysRevC.94.054323

Total number of authors: 51

Creative Commons License: Unspecified

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

• Users may download and print one copy of any publication from the public portal for the purpose of private study

or research.

You may not further distribute the material or use it for any profit-making activity or commercial gain

You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: https://creativecommons.org/licenses/

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Download date: 07. Jul. 2025

LUND UNIVERSITY

PO Box 117 221 00 Lund +46 46-222 00 00

Relativistic Coulomb excitation of ⁸⁸Kr

K. Moschner,¹ A. Blazhev,¹ J. Jolie,¹ N. Warr,¹ P. Boutachkov,^{2,3} P. Bednarczyk,⁴ K. Sieja,⁵ A. Algora,⁶ F. Ameil,² M. A. Bentley,⁷ S. Brambilla,⁸ N. Braun,¹ F. Camera,⁸ J. Cederkäll,⁹ A. Corsi,⁸ M. Danchev,¹⁰ D. DiJulio,⁹ C. Fahlander,⁹ J. Gerl,² A. Giaz,⁸ P. Golubev,⁹ M. Górska,² J. Grebosz,⁴ T. Habermann,² M. Hackstein,¹ R. Hoischen,^{2,9} I. Kojouharov,² N. Kurz,² N. Mărginean,¹¹ E. Merchán,² T. Möller,³ F. Naqvi,^{1,2} B. S. Nara Singh,⁷ C. Nociforo,² N. Pietralla,³ S. Pietri,² Zs. Podolyák,¹² A. Prochazka,² M. Reese,³ P. Reiter,¹ M. Rudigier,^{1,12} D. Rudolph,⁹ T. Sava,¹¹ H. Schaffner,² L. Scruton,⁷ J. Taprogge,¹ T. Thomas,¹ H. Weick,² A. Wendt,¹ O. Wieland,⁸ and H.-J. Wollersheim²

²GSI Helmholtzzentrum für Schwerionenforschung GmbH, Darmstadt, Germany

³Institut für Kernphysik, Technische Universität Darmstadt, Darmstadt, Germany

⁴Institute of Nuclear Physics PAN, Kraków, Poland

⁵Université de Strasbourg, IPHC, CNRS, UMR7178, Strasbourg, France

⁶Instituto de Fisica Corpuscular, CSIC-Universidad de Valencia, Valencia, Spain

⁷Department of Physics, University of York, York, United Kingdom

⁸Dipartimento di Fisica, Università de Milano, and INFN Sezione Milano, Milano, Italy

⁹Department of Physics, Lund University, Lund, Sweden

¹⁰Faculty of Physics, University of Sofia, Sofia, Bulgaria

¹¹Horia Hulubei NIPNE, Bucharest, Romania

¹²Department of Physics, University of Surrey, Guildford, United Kingdom

(Received 10 February 2015; revised manuscript received 15 February 2016; published 28 November 2016)

To investigate the systematics of mixed-symmetry states in N = 52 isotones, a relativistic Coulomb excitation experiment was performed during the PreSPEC campaign at the GSI Helmholtzzentrum für Schwerionenforschung to determine E2 transition strengths to 2^+ states of the radioactive nucleus ⁸⁸Kr. Absolute transition rates could be measured towards the first and third 2^+ states. For the latter a mixed-symmetry character is suggested on the basis of the indication for a strong M1 transition to the fully symmetric 2_1^+ state, extending the knowledge of the N = 52 isotones below Z = 40. A comparison with the proton-neutron interacting boson model and shell-model predictions is made and supports the assignment.

DOI: 10.1103/PhysRevC.94.054323

I. INTRODUCTION

The proton-neutron interacting boson model (IBM-2) [1] represents an algebraic formulation of a truncated shell model appropriate for the description of quadrupole collectivity. Because the IBM-2 is based on nucleon pairs formed by either two valence neutrons or two valence protons, it allows also the study of proton-neutron correlations, albeit of nucleon pairs approximated by bosons.

Dynamical symmetries, connected to the equations describing the dynamics of a quantal system, lead to conserved quantities denoted by quantum numbers and stringent selection rules which permit one to test the goodness of these quantum numbers. Furthermore, they give rise to a complete set of basis states, which can be constructed analytically. In the IBM-2 several dynamical symmetries can be constructed. Here we focus on those Hamiltonians that do not change upon exchange of the proton boson and neutron boson labels. Those can be classified using a new symmetry: *F*-spin symmetry [1]. A detailed account of the dynamical symmetries of these Hamiltonians is given in Ref. [2].

The concept of *F* spin in the IBM-2 extends the isospin formalism to the IBM-2 boson systems. These bosons are considered to be "elementary" particles that form a doublet with projection $F_z = +1/2$ (proton boson) and $F_z = -1/2$ (neutron boson). For the description of a given nucleus with fixed numbers of proton and neutron bosons, N_{π} and N_{ν} ,

respectively, the z component of the F spin is a good quantum number; it is $F_z = (N_{\pi} - N_{\nu})/2$. For a given total boson number, $N = N_{\pi} + N_{\nu}$, the total F-spin quantum number can take values between F_z and $F_{\text{max}} = N/2$.

While the low-lying collective states have $F = F_{\text{max}}$ and are symmetric with respect to the pairwise exchange of boson Fspin labels, the IBM-2 predicts entire new classes of collective states with F-spin quantum numbers $F < F_{\text{max}}$. Their wave functions contain at least one pair of proton and neutron bosons that is antisymmetric under the exchange of the proton and neutron labels. These states are called mixed-symmetry (MS) states.

In the mid-1980s the first MS state, a collective $J^{\pi} = 1^+$ excitation called scissors mode, was discovered in deformed gadolinium nuclei [3] by the research group of Richter. The IBM predicted the scissors mode [4] as one representative of the class of collective MS states in which valence neutron and proton pairs move out of phase. Scissors states were subsequently observed in many deformed nuclei and intensive theoretical studies of such states were performed [5,6].

According to the IBM-2 approach the whole class of MS states is formed by quadrupole collectivity. The building block of all MS excitations is therefore the MS 2_{ms}^+ state. It should occur in spherical nuclei as the lowest, one-phonon MS state on which collective multiphonon structures can be built. The existence of such a multiphonon structure with MS character was discovered in the nuclide ⁹⁴Mo [7,8] and can be explained

by the *F*-spin symmetric O(6) limit of the IBM-2 [2]. Recent measurements on ⁹⁴Mo and other N = 52 isotones confirm [9–15] the initial findings as a general phenomenon [16]. Here, the evolution of MS states in N = 52 isotones can be tracked over different proton shells. This enables the investigation of the variation of the proton-neutron interaction as a function of the nuclear valence space [17].

Unfortunately, the light N = 52 isotones below Z = 40 are unstable nuclei. No data on absolute transition rates of MS states existed prior to the present study. Absolute electromagnetic transition rates to low-lying states are, however, necessary information to uniquely identify MS states. Therefore, they are prerequisite for the systematic understanding of the proton-neutron nonsymmetric building blocks of nuclear collectivity.

It was the aim of this study to determine the quadrupole collectivity and to identify the lowest MS state of neutron-rich ⁸⁸Kr by using relativistic Coulomb excitation with a radioactive ion beam (RIB). Coulomb excitation is very well suited also for the population of one-phonon MS states [10,18,19]. The single-step Coulomb excitation reaction populates in a very clean way the excited 2⁺ states. The desired $B(E2; 2_{ms}^+ \rightarrow 0_1^+)$ and $B(M1; 2_{ms}^+ \rightarrow 2_1^+)$ values can be deduced from the measured cross sections and branching ratios.

Recently, Mücher *et al.* were able to measure the $B(E2; 0_1^+ \rightarrow 2_1^+)$ in ⁸⁸Kr at REX-ISOLDE [14]. In that experiment it was also attempted to measure nonyrast states of ⁸⁸Kr, in particular the 2_3^+ state. However, using low-energy Coulomb excitation the cross section for the excitation of the MS state was more than three orders of magnitude smaller than the one for excitation of the 2_1^+ state. The result was that it was not possible to measure the B(E2) of the 2_3^+ state. Using the predicted $B(E2; 0_1^+ \rightarrow 2_{ms}^+)$ value, which is a factor of 10 weaker than the $B(E2; 0_1^+ \rightarrow 2_1^+)$ value [16], the cross section for Coulomb excitation using a ⁸⁸Kr beam at intermediate energy was calculated to be about two orders of magnitude higher than for the above-mentioned REX-ISOLDE measurement [20]. Thus, the PreSPEC setup at the GSI Helmholtzzentrum für Schwerionenforschung was chosen to obtain transition strengths for the possible MS candidate state in ⁸⁸Kr at 2216 keV. Figure 1 shows the relevant part of the level scheme of ⁸⁸Kr.

II. EXPERIMENTAL DETAILS

A ²³⁸U primary beam at 650 MeV/nucleon was provided by the GSI accelerator complex. The ⁸⁸Kr isotopes were produced by in-flight fission on a 0.66 g/cm² thick ⁹Be primary target. The nuclei of interest were selected by the fragment separator (FRS) [22] utilizing the $B\rho$ - ΔE - $B\rho$ method. For identification purposes the fission fragments were tracked by the FRS beam detectors in the middle and final focal plane on an eventby-event basis. By measurement of the time of flight (TOF) through the separator as well as the energy loss in a multiple sampling ionization chamber the composition of the secondary beam could be obtained. The FRS particle identification plot in terms of atomic number Z and mass-over-charge ratio A/Qis shown in Fig. 2. A clean separation of the isotope of interest was provided by full width at half maximum resolutions of



FIG. 1. Partial level scheme of 88 Kr illustrating the lowest lying 2^+ states and the known transitions between them. The thickness of the arrows expresses the branching ratios of the transitions adopted from Ref. [21]. Transitions that remained unobserved in this work are drawn with dashed grey lines. Energy labels are in keV.

0.96% in A/Q and 0.91% in Z. A total of 2.0×10^{6} ⁸⁸Kr nuclei could be identified during the experiment.

Coulomb excitation of the 88Kr secondary beam at an energy of 128 MeV/nucleon took place on a 0.386 g/cm² thick, secondary ¹⁹⁷Au target. Deexcitation γ rays were detected by the PreSPEC Ge-detector array in the fast-beam configuration [23]. The array consisted of 15 EUROBALL Cluster detectors [24] placed under forward angles. The detection efficiency is enhanced due to the Lorentz boost of the γ -ray angular distribution at ion velocities of $\beta =$ $v/c \sim 0.5$. The measurement was carried out using particle- γ coincidences to trigger the data acquisition. The ions after the secondary target were identified by the Lund-York-Cologne Calorimeter (LYCCA) [25], which acts as a $\Delta E - E$ telescope. Additionally, it provides the TOF and trajectory of the excited nuclei of interest after the target. This is particularly important to determine the velocity and scattering angles needed for a proper Doppler correction of the measured γ rays, as well as to ensure selection of safe Coulomb excitation events.



FIG. 2. FRS particle identification plot for the ⁸⁸Kr setting.

III. ANALYSIS

For the analysis of the Coulomb excitation, the same mass and charge numbers for ingoing and outgoing particles at the secondary target were selected. This was done by selecting events with proper A/Q and Z measured by the FRS particle detectors, as well as proper TOF, ΔE , and E determined by LYCCA. Further conditions on the scattering angle of ⁸⁸Kr and prompt time in the particle- γ coincidences were applied to select safe Coulomb excitation of the relativistic projectile and enhance the peak-to-background ratio of the obtained γ -ray spectra, respectively. By applying an event-by-event Doppler correction the $2^+_1 \rightarrow 0^+_1 \gamma$ transition of ⁸⁸Kr at 775 keV could be clearly identified in the total γ -ray spectrum.

The EUROBALL Cluster detectors forming the PreSPEC array are arranged in two rings at mean polar angles of 16.3° and 34.7° comprising 5 and 10 cluster detectors, respectively. In the above-mentioned total γ -ray spectrum (both rings) for ⁸⁸Kr no clear evidence for Coulomb excitation of the 2^+_3 state could be observed. By separately considering detectors of the inner or outer rings, the spectra shown in Fig. 3 could be extracted.

While no clear peak structure in addition to the $2_1^+ \rightarrow 0_1^+$ transition is observed in the spectrum for the inner ring shown in Fig. 3(a), in the spectrum for the outer ring shown in Fig. 3(b) the region around 1441 keV exhibits significant excess of detected γ -ray events. This energy region corresponds exactly to the energy of the $2_3^+ \rightarrow 2_1^+$ transition in ⁸⁸Kr suggesting the Coulomb excitation of the 2_3^+ state. Other transitions in the spectrum are not observed. The rather broad and slightly asymmetric shape of the peak at 1441 keV is interpreted to originate from a very short lifetime of a few tens of femtoseconds, making it decay mainly while slowing down in the secondary target. Such fast transitions (short lifetimes) are expected for MS states due to high $B(M1; 2_{ms}^+ \rightarrow 2_1^+)$ values [16].



FIG. 3. Doppler-corrected γ -ray energy spectra for ⁸⁸Kr for the (a) inner and (b) outer ring of germanium detectors in the PreSPEC setup. The spectra are drawn with their statistical errors. Fits to the observed transitions and the 1σ observation limit in case of the $2_3^+ \rightarrow 2_1^+$ in the inner ring and a background function are indicated with the red (gray) line (see text for details).

The nonobservation in the inner ring can be understood considering the lower total efficiency due to fewer detectors in this group and the higher noise level due to more atomic background in close vicinity to the beam pipe.

Determining the peak areas in the spectrum for the outer ring for both transitions of ⁸⁸Kr led to values of 81(13) counts in the $2_1^+ \rightarrow 0_1^+$ and 29(10) counts in the $2_3^+ \rightarrow 2_1^+$ 2_1^+ case. Considering the branching ratio of the $2_3^+ \rightarrow 2_1^+$ transition of 85.4(2)% [21] and an energy-dependent increase of detection efficiency of 35% for the lower lying line, we were able to extract the value for $B(E2; 0_{g.s.}^+ \rightarrow 2_3^+)$ of ⁸⁸Kr by comparing the effective count rates. Normalizing to the value $B(E2; 0_{g.s.}^+ \rightarrow 2_1^+) = 0.093(9) e^2 b^2$ given in Ref. [14] we calculated a value of $B(E2; 0_{g.s.}^+ \rightarrow 2_3^+) = 0.04(2) e^2 b^2$ or $B(E2 \downarrow) = 3.4(17)$ W.u. Although small, this value exceeds the typical transition strength of $2_{ms}^+ \rightarrow 0_1^+$ transition of about 1 W.u. A similar value of 3.7(8) W.u. was also observed for ⁹²Zr [12].

IV. RESULTS

The decay from the 2_3^+ state to the 2_1^+ state is most likely a nearly pure *M*1 transition. Figure 4 shows the experimentally obtained, normalized γ -ray intensities extracted for both rings with their 1 σ error margins, including a 1 σ upper limit for the 1441 keV transition due to the nonobservation in the inner ring [see Fig. 3(a)]. The experimental data was least-squares fitted with theoretical angular distribution functions calculated using the computer program DWEIKO [20]. The best description of the data (indicated by red lines in Fig. 4) results in a multipole mixing ratio $\delta = 0.08^{+0.09}_{-0.05}$ and indicates the preference for the *M*1 character for the $2_3^+ \rightarrow 2_1^+$ transition.

Assuming the determined M1 character a collective $B(M1; 2_3^+ \rightarrow 2_1^+)$ of 0.6(3) μ_N^2 is obtained in agreement with the expectations for a MS character of the 2_3^+ state. For the E2 transition strength we obtain a value of $B(E2; 2_3^+ \rightarrow 2_1^+) = 1.0^{+8.7}_{-0.9}$ W.u., which is in agreement within the error



FIG. 4. Observed intensities with their statistical uncertainties (data points) and calculated angular distributions for the $2_1^+ \rightarrow 0_1^+$ with pure *E*2 character (dashed line) and the $2_3^+ \rightarrow 2_1^+$ (solid line) transitions. The angular range of the inner and outer rings of germanium detectors are marked with gray color. A 1 σ observation limit is indicated for the $2_3^+ \rightarrow 2_1^+$ in the inner ring.

TABLE I. Deduced experimental transition strengths from the present data in comparison to model calculations in the IBM-2.

Transition	σL	$B(\sigma L)_{\text{expt}}$	$B(\sigma L)_{O(6)}$	$B(\sigma L)$ [26,27]
$2^+_2 \rightarrow 0^+_{g,g}$	E2	≼1.0 W.u.	forbidden	0.2 W.u.
$2^{\tilde{+}}_{3} \rightarrow 0^{\tilde{+}}_{g,s}$	E2	3.4(17) W.u.	0.8 W.u.	0.5 W.u.
$2^+_3 \rightarrow 2^+_1$	E2	$1.0^{+8.7}_{-0.9}$ W.u.	forbidden	0.6 W.u.
$2^+_3 \rightarrow 2^+_1$	M1	$0.6(3) \mu_N^2$	$0.30 \ \mu_N^2$	$0.19 \ \mu_N^2$

bars both with the theoretical values (see Table I) and the value of B(E2) = 5(3) W.u. obtained for the ⁹⁴Mo case [9].

In Fig. 3 there is no clear indication of an 802 keV line related to the $2_2^+ \rightarrow 2_1^+$ transition of ⁸⁸Kr. However, excitation of this state cannot be fully excluded due to the high statistical uncertainty of the γ -ray background. Estimating the sensitivity limit for clear identification of the γ -ray transitions to be ten counts, an upper limit of $B(E2; 0_{g.s.}^+ \rightarrow 2_2^+) \leq 0.012 e^2 b^2$ for the excitation of this state follows. Table I lists the adopted values for the transition rates obtained from the present measurement and analysis.

V. DISCUSSION

Using the bare values for the boson g factors $g_n = 0$ and $g_p = 1.0 \mu_N^2$ and effective quadrupole charges of $e_n = 0$ and $e_p = 0.07$ eb obtained in IBM-2 calculations of the even-even ^{88–96}Kr isotopes [26,27], Table I gives the theoretical values in the *F*-spin symmetric O(6) limit [2] assuming ⁷⁸Ni as an inert core. Also given are the values obtained in Refs. [26,27] using IBM-2 parameters based on self-consistent mean-field calculations using a microscopic Gogny energy-density functional. In this calculation the 2_3^+ state is a mixed state with only 57% MS character. Better agreement of the experimental values is obtained for the O(6) limit which indicates a pure MS character for the 2_3^+ state.

Figure 5 shows the systematics for the even-even N = 52 isotones including our new data which fit the general behavior of the lowest MS states in this region. Although the dominant proton orbit changes from $g_{9/2}$ above 92 Zr to the upper part of the *pf* shell in ⁸⁸Kr, the collectivity of the lowest MS state remains at the same level. However, it should be noted that all experimentally known $B(M1; 2_{ms}^+ \rightarrow 2_1^+)$ in the N = 52 chain exceed the O(6) expectations.

For ⁹⁴Mo a good description of the *M*1 transition strength was obtained by a shell-model approach using the surface delta interaction (SDI) as the residual interaction and ⁸⁸Sr as an inert core [17]. Using the same model space, SDI parameters, and effective *g* factors a similar good agreement for the $B(M1; 2_{ms}^+ \rightarrow 2_1^+)$ value of one-phonon MS states in ⁹²Zr and ⁹⁶Ru is achieved. As pointed out in Ref. [12] these calculations show less good agreement with the experimentally obtained *g* factors for low-lying states in the N = 52isotones.

To obtain a microscopic description below Z = 40, the shell-model valence space has to be enlarged with respect to the proton pf orbitals. Such an approach based on a ⁷⁸Ni core



FIG. 5. (a) Excitation energies, (b) reduced B(E2), and (c) B(M1) transition strengths from the 2_1^+ and 2_{ms}^+ states in N = 52 isotones [13–16,28,29] together with the newly determined values for ⁸⁸Kr in red. Additionally shown are the results from shell-model calculations using the SDI [17] (green lines) and from new calculations based on the work of Ref. [30] (red lines).

and an extended model space $(1f_{5/2}, 2p_{1/2}, 2p_{3/2}, 1g_{9/2})$ for protons and $(2d_{5/2}, 3s_{1/2}, 2d_{3/2}, 1g_{7/2}, 1h_{11/2})$ for neutrons was presented in Ref. [30] and reproduced spectroscopic data of mixed-symmetry states in zirconium isotopes as well as yrast excitations in ⁸⁸Kr [31]. With the same effective interaction but slightly changed effective charges of $e_n = 0.7$ eb, $e_p = 1.7$ eb, orbital g factors $g_l^n = 0$, $g_l^p = 1.0 \mu_N^2$, and a quenching factor of 0.75 for the bare nucleon spin g factors, we obtained the theoretical energies and transition strengths for the 2_1^+ and 2_{ms}^+ states in ⁸⁸Kr, ⁹⁰Sr, and ⁹²Zr shown in Fig. 5. These shell-model calculations also predict mixed-symmetry states at the correct energies with clearly enhanced $B(M1; 2_{ms}^+ \rightarrow 2_1^+)$ values and E2 transitions to the ground state in good agreement with the experimentally observed strengths. However, the B(M1)strengths are underestimated for ⁸⁸Kr and ⁹²Zr. Also, the wavefunction analysis suggests that the identified states might not fully be the mixed-symmetric one-phonon 2_{ms}^+ states as pointed out for ⁹²Zr in Ref. [30]. The discrepancy between experiment and theory for the MS candidate state in ⁹⁰Sr at 1892 keV is larger and inverted (see Fig. 5). Here, the magnitude of the experimental transition strengths stems from a single lifetime measurement of $\tau = 3(2)$ ps using the $\beta \gamma \gamma$ fast timing method reported in Ref. [28]. A confirmation of this experimental lifetime is needed.

VI. SUMMARY

In conclusion, we obtained $B(E2; 0_{g.s.}^+ \rightarrow 2_3^+)$ and $B(M1; 2_3^+ \rightarrow 2_1^+)$ values as well as an upper limit for the $B(E2; 0_{g.s.}^+ \rightarrow 2_2^+)$ value from relativistic Coulomb excitation of the radioactive nucleus ⁸⁸Kr. The strong *M*1 character of the $2_3^+ \rightarrow 2_1^+$ transition indicates the mixed-symmetric nature of the $2_3^+ \rightarrow 2_1^+$ state, which is supported by model calculations in the proton-neutron interacting boson model. New shell-model calculations for the N = 52 isotones ⁹²Zr, ⁹⁰Sr, and ⁸⁸Kr with

- A. Arima, T. Otsuka, F. Iachello, and I. Talmi, Phys. Lett. B 66, 205 (1977).
- [2] P. Van Isacker, K. Heyde, J. Jolie, and A. Sevrin, Ann. Phys. (NY) 171, 253 (1986).
- [3] D. Bohle, A. Richter, W. Steffen, A. E. L. Dieperink, N. Lo Iudice, F. Palumbo, and O. Scholten, Phys. Lett. B 137, 27 (1984).
- [4] F. Iachello, Nucl. Phys. A **358**, 89 (1981).
- [5] A. Richter, Prog. Part. Nucl. Phys. 34, 261 (1995).
- [6] K. Heyde, P. von Neumann-Cosel, and A. Richter, Rev. Mod. Phys. 82, 2365 (2010).
- [7] N. Pietralla, C. Fransen, D. Belic, P. von Brentano, C. Frießner, U. Kneissl, A. Linnemann, A. Nord, H. H. Pitz, T. Otsuka, I. Schneider, V. Werner, and I. Wiedenhöver, Phys. Rev. Lett. 83, 1303 (1999).
- [8] N. Pietralla, C. Fransen, P. von Brentano, A. Dewald, A. Fitzler, C. Frießner, and J. Gableske, Phys. Rev. Lett. 84, 3775 (2000).
- [9] C. Fransen, N. Pietralla, Z. Ammar, D. Bandyopadhyay, N. Boukharouba, P. von Brentano, A. Dewald, J. Gableske, A. Gade, J. Jolie, U. Kneissl, S. R. Lesher, A. F. Lisetskiy, M. T. McEllistrem, M. Merrick, H. H. Pitz, N. Warr, V. Werner, and S. W. Yates, Phys. Rev. C 67, 024307 (2003).
- [10] N. Pietralla, C. J. Barton, R. Krucken, C. W. Beausang, M. A. Caprio, R. F. Casten, J. R. Cooper, A. A. Hecht, H. Newman, J. R. Novak, and N. V. Zamfir, Phys. Rev. C 64, 031301 (2001).
- [11] H. Klein, A. F. Lisetskiy, N. Pietralla, C. Fransen, A. Gade, and P. von Brentano, Phys. Rev. C 65, 044315 (2002).
- [12] V. Werner et al., Phys. Lett. B 550, 140 (2002).
- [13] C. Fransen, A. Blazhev, A. Dewald, J. Jolie, D. Mücher, O. Möller, and T. Pissulla, AIP Conf. Proc. **1090**, 529 (2009).
- [14] D. Mücher et al., AIP Conf. Proc. 1090, 587 (2009).
- [15] A. Hennig, T. Ahn, V. Anagnostatou, A. Blazhev, N. Cooper, V. Derya, M. Elvers, J. Endres, P. Goddard, A. Heinz, R. O. Hughes, G. Ilie, M. N. Mineva, P. Petkov, S. G. Pickstone, N. Pietralla, D. Radeck, T. J. Ross, D. Savran, M. Spieker, V. Werner, and A. Zilges, Phys. Rev. C 92, 064317 (2015).

a ⁷⁸Ni core also reproduce the measured transition strengths of the states in question.

ACKNOWLEDGMENTS

Parts of this research were carried out at the UNILAC/SIS accelerators at GSI Darmstadt, a member of the Helmholtz Association (HGF). This work was supported by the German BMBF under Contracts No. 06KY9136I, No. 05P12PKFNE, and No. 05P12RDFN8, HIC for FAIR, and the Swedish Research Council.

- [16] N. Pietralla, P. von Brentano, and A. F. Lisetskiy, Prog. Part. Nucl. Phys. 60, 225 (2008).
- [17] A. F. Lisetskiy, N. Pietralla, C. Fransen, R. Jolos, and P. von Brentano, Nucl. Phys. A 677, 100 (2000).
- [18] G. Rainovski, N. Pietralla, T. Ahn, C. J. Lister, R. V. F. Janssens, M. P. Carpenter, S. Zhu, and C. J. Barton, Phys. Rev. Lett. 96, 122501 (2006).
- [19] M. Danchev, G. Rainovski, N. Pietralla, A. Gargano, A. Covello, C. Baktash, J. R. Beene, C. R. Bingham, A. Galindo-Uribarri, K. A. Gladnishki, C. J. Gross, V. Y. Ponomarev, D. C. Radford, L. L. Riedinger, M. Scheck, A. E. Stuchbery, J. Wambach, C. H. Yu, and N. V. Zamfir, Phys. Rev. C 84, 061306 (2011).
- [20] C. Bertulani, C. Campbell, and T. Glasmacher, Comput. Phys. Commun. 152, 317 (2003).
- [21] E. McCutchan and A. Sonzogni, Nucl. Data Sheets 115, 135 (2014).
- [22] H. Geissel *et al.*, Nucl. Instrum. Methods Phys. Res. B 70, 286 (1992).
- [23] H. Wollersheim *et al.*, Nucl. Instrum. Methods Phys. Res. A 537, 637 (2005).
- [24] J. Eberth et al., Prog. Part. Nucl. Phys. 28, 495 (1992).
- [25] P. Golubev *et al.*, Nucl. Instrum. Methods Phys. Res. A **723**, 55 (2013).
- [26] M. Albers *et al.*, Phys. Rev. Lett. **108**, 062701 (2012); **109**, 209904 (2012); **114**, 189902 (2015).
- [27] M. Albers *et al.*, Nucl. Phys. A **899**, 1 (2013); **947**, 260 (2016).
- [28] H. Mach, F. Wohn, G. Molnár, K. Sistemich, J. C. Hill, M. Moszyński, R. Gill, W. Krips, and D. Brenner, Nucl. Phys. A 523, 197 (1991).
- [29] C. Fransen (unpublished).
- [30] K. Sieja, F. Nowacki, K. Langanke, and G. Martínez-Pinedo, Phys. Rev. C 79, 064310 (2009).
- [31] M. Czerwiński, T. Rzaca-Urban, K. Sieja, H. Sliwinska, W. Urban, A. G. Smith, J. F. Smith, G. S. Simpson, I. Ahmad, J. P. Greene, and T. Materna, Phys. Rev. C 88, 044314 (2013).