



LUND UNIVERSITY

The risk for depression in patients with ankylosing spondylitis: a population-based cohort study

Meesters, Jorit J. L.; Bremander, Ann; Bergman, Stefan; Petersson, Ingemar; Turkiewicz, Aleksandra; Englund, Martin

Published in:
Arthritis Research and Therapy

DOI:
[10.1186/s13075-014-0418-z](https://doi.org/10.1186/s13075-014-0418-z)

2014

[Link to publication](#)

Citation for published version (APA):

Meesters, J. J. L., Bremander, A., Bergman, S., Petersson, I., Turkiewicz, A., & Englund, M. (2014). The risk for depression in patients with ankylosing spondylitis: a population-based cohort study. *Arthritis Research and Therapy*, 16(418), 1-5. <https://doi.org/10.1186/s13075-014-0418-z>

Total number of authors:
6

General rights

Unless other specific re-use rights are stated the following general rights apply:
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

RESEARCH ARTICLE

Open Access

The risk for depression in patients with ankylosing spondylitis: a population-based cohort study

Jorit JL Meesters^{1,2}, Ann Bremander^{3,4,5}, Stefan Bergman^{3,4}, Ingemar F Petersson^{1,3,6}, Aleksandra Turkiewicz⁶ and Martin Englund^{1,6,7*}

Abstract

Introduction: Depression is frequent in ankylosing spondylitis (AS) patients. However, epidemiological data about the potential increase in risk are lacking. This study compares the rate of doctor-diagnosed depression in a well defined cohort of AS patients to the general population seeking care.

Methods: The Skåne Healthcare Register comprises healthcare data of each resident in Region Skåne, Sweden (population 1.2 million), including ICD-10 diagnoses. Using physician coded consultation data from years 1999 to 2011, we calculated depression consultation rates for all AS patients. We obtained standardized depression-rate ratios by dividing the observed depression rate in AS patients by the expected rate based on the corresponding age- and sex-specific rates of depression in the general population seeking care. A ratio >1 equals a higher rate of depression among AS patients.

Results: The AS cohort consisted of 1738 subjects (65% men) with a mean age of 54 years. The reference population consisted of 967,012 subjects. During the 13-year observation period 10% (n = 172) of the AS cohort had a doctor-diagnosed depression compared to 6% (n = 105) to be expected. The standardized estimate of depression-rate ratio was 1.81 (95% confidence interval 1.44 to 2.24) in women men and 1.49 (1.20 to 1.89) in men.

Conclusions: The rate of doctor-diagnosed depression is increased about 80% in female and 50% in male AS patients. Future challenges are to timely identify and treat the AS patients who suffer from depression.

Introduction

The prevalence of ankylosing spondylitis (AS) in the Western world is estimated to be 0.1 to 0.2%, with an estimated incidence of 7/100,000 per year [1-3]. Due to the nature of the disease, AS typically affects physical function, quality of life [4,5], and work productivity [6,7]. Comorbid conditions, including uveitis, inflammatory bowel and cardiovascular disease, contribute to the prognosis and total burden of AS [8].

Clinical depression has also been reported to be more frequent in chronic diseases [9], rheumatic diseases in general [10] and AS [11-13]. However, an estimate of the actual increase in risk, and its age- and/or gender-

dependency, of depression in AS patients is lacking. Therefore, our objective was to study consultation rates of doctor-diagnosed depression in female and male AS patients compared to the general population seeking healthcare in a population-based open cohort study. We hypothesized that the consultation rates of depression are significantly higher in AS patients.

Methods

The Skåne healthcare register (SHR)

Detailed information about the SHR is provided in previous articles [8,14]. In brief, each single healthcare consultation in this region generates data entries by the healthcare provider that are transferred to central databases by the patient's unique 10-digit personal identification number. Data includes information about date of consultation and international classification of disease (ICD)-10 diagnostic codes

* Correspondence: martin.englund@med.lu.se

¹Epidemiology and Register Centre South Skåne, Skåne University Hospital, Lund, Sweden

⁶Orthopaedics, Clinical Sciences Lund, Lund University, Lund, Sweden

Full list of author information is available at the end of the article

by the responsible physician, drawn from the patients' computer-based medical records.

The Swedish Population Register

The Swedish Population Register is the civil registration of vital events (for example, births, deaths, removal) of Swedish inhabitants. For this study, we linked SHR data via the personal identification number to add information on date of death and residency.

Inclusion criteria for the AS cohort

We required all AS cases to be at least 20 years of age and to be resident in Skåne Region by the first healthcare visit in the SHR between 1999 and 2011. Further, our case definition required a clinic visit with a diagnosis of AS (ICD-10 code M45) made by a physician during the years 1999 to 2011 (13 calendar years), at least once at a rheumatology clinic or internal medicine clinic, or at least twice at any other clinic including general practice. The date of fulfilling the above definition marked the beginning of the exposure time. A previous validation of spondyloarthritis at large, showed 98% of patients identified via the register to be valid spondyloarthritis patients [2].

Inclusion criteria for the reference population

In this article the general population seeking healthcare is regarded as the reference population. For the AS cohort, we required all subjects in the reference population to be residents in Skåne Region and aged 20 years or older by the first healthcare visit registered in the SHR between 1999 and 2011. Then we required all of the subjects to have had at least one clinic visit between 1 January 1999 and 31 December 2011, with any diagnosis by any physician in the SHR, to minimize confounding.

Outcome

For the reference population we calculated each single subject's person-time from the first healthcare visit until one of the following four events had occurred for the first time within the observation period from the first visit until 31 December 2011, in the following order of priority: 1) occurrence of doctor-diagnosed depressive episode (ICD-10 code F32) or recurrent depressive disorder (F33); 2) death; 3) end of the observation period if the subject was still a resident in the county by that time; or 4) date of relocation outside of the region. The same procedure was applied to the AS patients, where person-time was calculated from the date of fulfilling the AS definition.

Statistical analysis

Observed depression rate for AS patients

We calculated observed consultation rates for doctor-diagnosed depression (depression rates) among patients

fulfilling our AS case criteria. The total number of AS patients having had the diagnosis within the study period formed the numerator of the rate, and the AS patients' added person-time formed the denominator of the rate. The age of all subjects was defined as the age at the mean follow-up time. We calculated rates for each age and sex stratum (using 5-year age categories).

Expected depression rate for AS patients

We calculated the expected doctor-diagnosed depression rate as if the rate in the general population seeking healthcare prevailed, using age and sex of the AS patients as the standard. Hence, using data from the SHR, we calculated the sum of each resident's person-time and counted all subjects diagnosed with the first occurrence of depression during the study period. We then calculated the expected depression rate by multiplying the stratum-specific rates in the reference population with the appropriate age and sex weightings from the AS cohort. Then we added the stratum-specific products to obtain the age- and sex-standardized rate.

Standardized depression-rate ratio and absolute rate difference

We calculated age- and sex-standardized depression-rate ratio by dividing the observed morbidity rate in the AS cohort by their expected morbidity rate. A rate ratio >1 equals a higher depression rate in the AS cohort than in the general population seeking healthcare. We calculated 95% CI for the rate ratios using the method described by Breslow [15]. We also calculated the absolute depression-rate difference (observed rate minus expected rate) to provide results of excess doctor-diagnosed depression on the absolute scale as well.

Gender differences

To evaluate the difference in depression-rate ratio in women and men for AS patients and the reference population, respectively, we also calculated the ratio of these rate ratios with their 95% CI.

Diabetes mellitus

To facilitate interpretation of the results obtained for the AS patients, we also evaluated patients with another chronic disease reported to be associated with increased risk of depression, insulin-dependent and non-insulin-dependent diabetes mellitus (ICD-10 code E10 or E11) [16]. Our case definition for diabetes required a clinic visit, with a physician's diagnosis of diabetes at least once at an endocrinology or internal medicine clinic, or at least twice at any other clinic, including general practice.

Ethics statement

The study was conducted according to the Declaration of Helsinki and was approved by the Ethical Review Board of Lund University, Sweden. The population in the Skåne region was informed of the study via regional news press and offered the opportunity to opt out, a process sanctioned by the Ethical Review Board. Thus, no individually signed informed consent forms were required according to the decision from the Ethical Review Board (a common procedure in Sweden for population-based register studies).

Results

The number of residents (aged ≥ 20 years) in Skåne Region who fulfilled our AS case criteria during our 13-year study period was 1,738 (64.4% men), mean age (SD) 54.5 (14.3) years (1,275 AS patients (73%) had been diagnosed at a rheumatology or internal medicine clinic). The total adult general population seeking healthcare during the same period consisted of 967,012 subjects (47.6% men).

During the study period 10% ($n = 172$) of the AS cohort had at least one episode of doctor-diagnosed depression compared to the expected 6% ($n = 105$) based on age- and sex-standardized data from the reference population. The standardized depression-rate ratio in subjects with AS (Table 1) was significantly elevated (1.63; 95% CI 1.40, 1.89). The standardized depression-rate ratio in women (1.81; 95% CI 1.44, 2.24) and in men (1.49; 95% CI 1.20, 1.89) were both elevated. However, the age-standardized ratio between these rate ratios (women versus men) was not significantly different (1.20; 95% CI 0.89, 1.62), implying no substantial interaction between AS and sex for the risk of doctor-diagnosed depression.

The absolute rate difference (observed minus expected) for doctor-diagnosed depression in subjects with AS was 649 per 100,000 person-years. The absolute rate difference

was greater in women (1,138 per 100,000 person-years) compared to men (421 per 100,000 person-years). However, this difference is explained by the much higher observed depression rate in women in general compared to men (Table 1). The rate of doctor-diagnosed depression in women in the reference population was highly elevated compared to men (standardized depression rate ratio = 1.66; 95% CI 1.63, 1.68), that is, women had a 66% higher increased rate than men. The standardized depression-rate ratio in subjects with diabetes mellitus was significantly elevated (1.67; 95% CI 1.63, 1.71).

Discussion

In a cohort study design, using comprehensive population-based epidemiologic data based on physicians' diagnostic coding, our hypothesis that consultation rates of depression are significantly higher in AS patients was confirmed. We found an 80% increased rate of doctor-diagnosed depression in women with AS and 50% in men with AS compared to the general population seeking care. The higher rate in women is due to general sex differences and thus is not so much AS-specific. Although these findings come as no surprise, population-based epidemiologic estimates of the actual increase in risk and potential age and gender aspects are lacking. These findings enhance our understanding of the total burden for AS patients and may contribute to improved health assessment and treatment planning [17].

Prior studies have suggested a relationship between depression and rheumatic diseases [10] and AS [11,13]. In the present study we found evidence that the risk for depression is higher in AS patients, and the increase is in line with patients with diabetes mellitus. We further found the absolute rate of doctor-diagnosed depression to be much higher in women than men, both in the general population and in AS patients. These findings are in

Table 1 Doctor-diagnosed depression in ankylosing spondylitis (AS) patients ($n = 1,738$) compared with the general population seeking healthcare ($n = 967,012$)

	Observed			Expected		Absolute rate differences ²	Rate ratio ³	95% CI
	Person time (years)	Number of cases ¹	Depression rate ²	Number of cases	Depression rate ²			
Men, all	6,983	89	1,275	60	853	421	1.49	1.20, 1.89
≤50 years	2,879	42	1,376	27	933	443	1.47	1.10, 1.93
>50 years	4,104	47	1,155	31	759	396	1.52	1.07, 2.10
Women, all	3,259	83	2,547	46	1,409	1,138	1.81	1.44, 2.24
≤50 years	1,274	39	2,825	20	1,546	1,279	1.83	1.35, 2.42
>50 years	1,984	44	2,244	25	1,260	984	1.78	1.24, 2.48
Total, all	10,242	172	1,679	105	1,030	649	1.63	1.40, 1.89
≤50 years	4,153	83	1,825	47	1,123	702	1.63	1.32, 1.98
>50 years	6,089	89	1,512	56	923	588	1.64	1.28, 2.06

¹AS patients diagnosed with depression (ICD-10 code F32 or F33) in 1999 to 2011. ²Per 100,000 person-years. ³Bold figures indicate statistically significant results.

line with literature regarding prevalence of depression in the general population [18] and in patients with rheumatic diseases [10].

In our study, we did not include person-time for residents not seeking healthcare in the study time frame. Including such person-time would elevate the rate ratios even further. This decision was made to compensate for bias caused by tendency to consult (irrespective of reason) being higher in patients with previous consultations. Still, we cannot fully exclude the possibility of residual doctors' work-up bias towards patients with a chronic disease such as AS. However, we find this unlikely because depression is not a typical diagnosis for which patients are screened, unlike hypertension, hypercholesterolemia, or diabetes. Furthermore, we included both prevalent and incident disease, both for the AS diagnosis and for depression, and data have not been controlled for possible confounding factors other than age and sex, and propensity to seek care. Further, in this study we could not relate the risk to disease duration, disease severity or other covariates, therefore residual confounding is possible. Moreover, the use of diagnostic codes based on physicians coding in everyday practice might pose a threat to the validity. However, the accuracy of the diagnostic coding from the SHR has previously been validated for spondyloarthritis, as well as specifically for psoriatic arthritis and psoriasis, with positive predictive values consistently exceeding 81% [19]. The diagnostic coding of depression could not be externally verified, for example, by review of medical records, due to restrictions in the approval of our institutional review board.

Conclusions

Based on our findings, we conclude that the consultation rate for depression is increased by more than 60% in AS patients compared to the background population seeking care, with women having somewhat larger increase in risk. One future challenge is to timely identify AS patients with depression. Research should focus on finding out which patient-reported outcome measures are most suitable to identify a depressive disorder in AS patients. The Hospital Anxiety and Depression Scale, the Hamilton Rating Scale for Depression, the Beck Depression Inventory, or the Center for Epidemiologic Studies Depression Scale [10] could be considered, as these have previously been used to assess depression in studies of rheumatic disorders.

Abbreviations

AS: ankylosing spondylitis; ICD: International classification of diseases; SHR: Skåne healthcare register.

Competing interests

IP, SB and ME received payment from Pfizer for lectures in clinical epidemiology. The other authors declare that they have no competing interests.

Authors' contributions

JM participated in the analysis and interpretation of data, and the draft of the manuscript. AB participated in the design of the study, the interpretation of data and helped to draft the manuscript revising it critically for important intellectual content. SB participated in the design of the study, the interpretation of data and helped to draft the manuscript revising it critically for important intellectual content. IP participated in the design of the study, the interpretation of data and helped to draft the manuscript revising it critically for important intellectual content. AT participated in the acquisition, analysis and interpretation of data, and (helped to) draft the manuscript revising it critically for important intellectual content. ME participated in the design of the study, the acquisition, analysis and interpretation of the data, and the draft of the manuscript. All authors read and approved the final manuscript.

Acknowledgements

We would like to acknowledge funding support from Swedish Research Council, Medical Faculty Lund University, and Region Skåne.

Author details

¹Epidemiology and Register Centre South Skåne, Skåne University Hospital, Lund, Sweden. ²Department of Orthopaedics, Leiden University Medical Centre, Leiden, the Netherlands. ³Department of Clinical Sciences, Lund, Section of Rheumatology, Lund University, Lund, Sweden. ⁴Spenshult, Research and Development Center, Oskarström, Sweden. ⁵School of Business and Engineering, Department of Exercise Physiology, Biomechanics and Health, Halmstad University, Halmstad, Sweden. ⁶Orthopaedics, Clinical Sciences Lund, Lund University, Lund, Sweden. ⁷Clinical Epidemiology Research & Training Unit, Boston University School of Medicine, Boston, MA, USA.

Received: 26 November 2013 Accepted: 1 August 2014

Published online: 18 August 2014

References

1. Bakland G, Nossent HC, Gran JT: **Incidence and prevalence of ankylosing spondylitis in Northern Norway.** *Arthritis Rheum* 2005, **53**:850–855.
2. Haglund E, Bremander AB, Petersson IF, Strombeck B, Bergman S, Jacobsson LT, Turkiewicz A, Geborek P, Englund M: **Prevalence of spondyloarthritis and its subtypes in southern Sweden.** *Ann Rheum Dis* 2011, **70**:943–948.
3. Carter ET, McKenna CH, Brian DD, Kurland LT: **Epidemiology of Ankylosing spondylitis in Rochester, Minnesota, 1935–1973.** *Arthritis Rheum* 1979, **22**:365–370.
4. Dagfinrud H, Mengshoel AM, Hagen KB, Loge JH, Kvien TK: **Health status of patients with ankylosing spondylitis: a comparison with the general population.** *Ann Rheum Dis* 2004, **63**:1605–1610.
5. Singh JA, Strand V: **Spondyloarthritis is associated with poor function and physical health-related quality of life.** *J Rheumatol* 2009, **36**:1012–1020.
6. Chorus AM, Miedema HS, Boonen A, van der Linden S: **Quality of life and work in patients with rheumatoid arthritis and ankylosing spondylitis of working age.** *Ann Rheum Dis* 2003, **62**:1178–1184.
7. Strombeck B, Englund M, Bremander A, Jacobsson LT, Kedza L, Kobelt G, Petersson IF: **Cost of illness from the public payers' perspective in patients with ankylosing spondylitis in rheumatological care.** *J Rheumatol* 2010, **37**:2348–2355.
8. Bremander A, Petersson IF, Bergman S, Englund M: **Population-based estimates of common comorbidities and cardiovascular disease in ankylosing spondylitis.** *Arthritis Care Res (Hoboken)* 2011, **63**:550–556.
9. Wells KB, Golding JM, Burnam MA: **Psychiatric disorder in a sample of the general population with and without chronic medical conditions.** *Am J Psychiatry* 1988, **145**:976–981.
10. Geenen R, Newman S, Bossemma ER, Vriezckolk JE, Boelen PA: **Psychological interventions for patients with rheumatic diseases and anxiety or depression.** *Best Pract Res Clin Rheumatol* 2012, **26**:305–319.
11. Martindale J, Smith J, Sutton CJ, Grennan D, Goodacre L, Goodacre JA: **Disease and psychological status in ankylosing spondylitis.** *Rheumatology (Oxford)* 2006, **45**:1288–1293.
12. Hyphantis T, Kotsis K, Tsifetaki N, Creed F, Drosos AA, Carvalho AF, Voulgari PV: **The relationship between depressive symptoms, illness perceptions and quality of life in ankylosing spondylitis in comparison to rheumatoid arthritis.** *Clin Rheumatol* 2013, **32**:635–644.

13. Barlow JH, Macey SJ, Struthers GR: **Gender, depression, and ankylosing spondylitis.** *Arthritis Care Res* 1993, **6**:45–51.
14. Englund M, Joud A, Geborek P, Felson DT, Jacobsson LT, Petersson IF: **Prevalence and incidence of rheumatoid arthritis in southern Sweden 2008 and their relation to prescribed biologics.** *Rheumatology (Oxford)* 2010, **49**:1563–1569.
15. Breslow NE, Day NE: **Statistical methods in cancer research. Volume II—The design and analysis of cohort studies.** *IARC Sci Publ* 1987, **82**:1–406.
16. Kan C, Silva N, Golden SH, Rajala U, Timonen M, Stahl D, Ismail K: **A systematic review and meta-analysis of the association between depression and insulin resistance.** *Diabetes Care* 2013, **36**:480–489.
17. Kremer JM, Greenberg J: **Interpreting registry-derived drug studies: does societal context matter?** *Arthritis Rheum* 2009, **60**:3155–3157.
18. Task Force on Major and Chronic Diseases: *Major and Chronic Diseases, Report 2007.* Luxembourg, Luxembourg: European Committee; 2008.
19. Lofvendahl S, Theander E, Svensson A, Carlsson KS, Englund M, Petersson IF: **Validity of diagnostic codes and prevalence of physician-diagnosed psoriasis and psoriatic arthritis in southern sweden - a population-based register study.** *PLoS One* 2014, **9**:e98024.

doi:10.1186/s13075-014-0418-z

Cite this article as: Meesters *et al.*: The risk for depression in patients with ankylosing spondylitis: a population-based cohort study. *Arthritis Research & Therapy* 2014 **16**:418.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

