

LUND UNIVERSITY

48Ca+249Bk Fusion Reaction Leading to Element Z=117: Long-Lived α-Decaying 270Db and Discovery of 266Lr

Khuyagbaatar, J.; Yakushev, A.; Düllmann, Ch.E.; Ackermann, D.; Andersson, L.-L.; Asai, M.; Block, M.; Boll, R.A.; Brand, H.; Cox, D.M.; Dasgupta, M.; Derkx, X.; Di Nitto, A.; Eberhardt, K.; Even, J.; Evers, M.; Fahlander, Claes; Forsberg, Ulrika; Gates, J.M.; Gharibyan, N.; Golubev, Pavel; Gregorich, K.E.; Hamilton, J.H.; Hartmann, W.; Herzberg, R.-D.; Heßberger, F.P.; Hinde, D.J.; Hoffmann, J.; Hollinger, R.; Hübner, A.; Jäger, E.; Kindler, B.; Kratz, J.V.; Krier, J.; Kurz, N.; Laatiaoui, M.; Lahiri, S.; Lang, R.; Lommel, B.; Maiti, M.; Miernik, K.; Minami, S.; Mistry, A.; Mokry, C.; Nitsche, H.; Omtvedt, J.P.; Pang, G.K.; Papadakis, P.; Renisch, D.; Roberto, J.

Published in: Physical Review Letters

10.1103/PhysRevLett.112.172501

2014

Link to publication

Citation for published version (APA):

Khuyagbaatar, J., Yakushev, A., Düllmann, C. E., Ackermann, D., Andersson, L.-L., Asai, M., Block, M., Boll, R. A., Brand, H., Cox, D. M., Dasgupta, M., Derkx, X., Di Nitto, A., Eberhardt, K., Even, J., Evers, M., Fahlander, C., Forsberg, U., Gates, J. M., ... Yakusheva, V. (2014). 48Ca+249Bk Fusion Reaction Leading to Element Z=117: Long-Lived α-Decaying 270Db and Discovery of 266Lr. *Physical Review Letters*, *112*(17), Article 172501. https://doi.org/10.1103/PhysRevLett.112.172501

Total number of authors:

General rights

Unless other specific re-use rights are stated the following general rights apply:
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

• Users may download and print one copy of any publication from the public portal for the purpose of private study

You may not further distribute the material or use it for any profit-making activity or commercial gain
You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: https://creativecommons.org/licenses/

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Download date: 17, Oct. 2025

LUND UNIVERSITY

PO Box 117 221 00 Lund +46 46-222 00 00 3

48 Ca + 249 Bk Fusion Reaction Leading to Element Z=117: Long-Lived lpha-Decaying 270 Db and Discovery of 266 Lr

J. Khuyagbaatar, ^{1,2,*} A. Yakushev, ² Ch. E. Düllmann, ^{1,2,3} D. Ackermann, ² L.-L. Andersson, ¹ M. Asai, ⁴ M. Block, ² R. A. Boll, H. Brand, D. M. Cox, M. Dasgupta, X. Derkx, A. Di Nitto, K. Eberhardt, J. Even, M. Evers, C. Fahlander, U. Forsberg, J. M. Gates, N. Gharibyan, P. Golubev, K. E. Gregorich, J. H. Hamilton, W. Hartmann, R.-D. Herzberg, F. P. Heßberger, D. J. Hinde, J. Hoffmann, R. Hollinger, A. Hübner, E. Jäger, B. Kindler, J. V. Kratz, J. Krier, N. Kurz, M. Laatiaoui, S. Lahiri, R. Lang, B. Lommel, M. Maiti, K. Miernik, S. Minami, M. Maiti, M. Maiti, M. Maiti, M. Miernik, R. Miernik, M. Maiti, M. Mai A. Mistry, ⁶ C. Mokry, ^{1,3} H. Nitsche, ⁹ J. P. Omtvedt, ¹³ G. K. Pang, ⁹ P. Papadakis, ^{6,14} D. Renisch, ³ J. Roberto, ⁵ D. Rudolph, ⁸ J. Runke, ² K. P. Rykaczewski, ⁵ L. G. Sarmiento, ⁸ M. Schädel, ^{2,4} B. Schausten, ² A. Semchenkov, ¹³ D. A. Shaughnessy, ¹⁰ P. Steinegger, ^{15,16} J. Steiner, ² E. E. Tereshatov, ¹⁰ P. Thörle-Pospiech, ^{1,3} K. Tinschert, ² T. Torres De Heidenreich, ² D. M. Schädel, ^{2,4} B. Schausten, ³ A. Semchenkov, ¹⁵ D. A. Shaughnessy, ¹⁰ P. Thörle-Pospiech, ¹⁷ T. Torres De Heidenreich, ² D. M. Schädel, ^{2,4} B. Schausten, ² A. Semchenkov, ¹³ D. A. Shaughnessy, ¹⁰ P. Thörle-Pospiech, ^{1,3} K. Tinschert, ² T. Torres De Heidenreich, ² D. M. Schädel, ^{2,4} B. Schausten, ³ A. Semchenkov, ¹⁵ D. A. Shaughnessy, ¹⁰ P. Thörle-Pospiech, ¹⁷ P. Thörle-Pospiech, ¹⁷ P. Thörle-Pospiech, ¹⁷ P. Thörle-Pospiech, ¹⁸ P. Thörle-Pospiec N. Trautmann, A. Türler, 15,16 J. Uusitalo, 14 D. E. Ward, 8 M. Wegrzecki, 17 N. Wiehl, 1,3 S. M. Van Cleve, 5 and V. Yakusheva 1 ¹Helmholtz Institute Mainz, 55099 Mainz, Germany ²GSI Helmholtzzentrum für Schwerionenforschung, 64291 Darmstadt, Germany ³Johannes Gutenberg-Universität Mainz, 55099 Mainz, Germany ⁴Advanced Science Research Center, Japan Atomic Energy Agency, Tokai, Ibaraki 319-1195, Japan ⁵Oak Ridge National Laboratory, Oak Ridge, Tennessee 37831, USA ⁶University of Liverpool, Liverpool L69 7ZE, United Kingdom ⁷The Australian National University, Canberra, Australian Capital Territory 0200, Australia ⁸Lund University, 22100 Lund, Sweden ⁹Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA ¹⁰Lawrence Livermore National Laboratory, Livermore, California 94551, USA ¹Vanderbilt University, Nashville, Tennessee 37235, USA ¹²Saha Institute of Nuclear Physics, Kolkata 700064, India ¹³University of Oslo, 0315 Oslo, Norway ¹⁴University of Jyväskylä, 40351 Jyväskylä, Finland ¹⁵Paul Scherrer Institute, 5232 Villigen, Switzerland ¹⁶University of Bern, 3012 Bern, Switzerland

The superheavy element with atomic number Z=117 was produced as an evaporation residue in the 48 Ca + 249 Bk fusion reaction at the gas-filled recoil separator TASCA at GSI Darmstadt, Germany. The radioactive decay of evaporation residues and their α -decay products was studied using a detection setup that allowed measuring decays of single atomic nuclei with half-lives between sub- μ s and a few days. Two decay chains comprising seven α decays and a spontaneous fission each were identified and are assigned to the isotope $^{294}117$ and its decay products. A hitherto unknown α -decay branch in 270 Db (Z=105) was observed, which populated the new isotope 266 Lr (Z=103). The identification of the long-lived ($T_{1/2}=1.0^{+1.9}_{-0.4}$ h) α -emitter 270 Db marks an important step towards the observation of even more long-lived nuclei of superheavy elements located on an "island of stability."

¹⁷Institute of Electron Technology, 02-668 Warsaw, Poland (Received 22 February 2014; published 1 May 2014)

DOI: 10.1103/PhysRevLett.112.172501 PACS numbers: 27.90.+b, 23.60.+e, 25.70.Gh

The existence of superheavy elements (SHE) was predicted in the late 1960s as one of the first harvests of the macroscopic-microscopic theory of the atomic nucleus [1,2]. Shell effects were predicted to be pronounced at Z=114 and N=184. Modern theoretical approaches [3,4] confirm this concept. Very long half-lives for these nuclei forming an "island of stability" are expected. In the past 50 years, a wealth of both theoretical and experimental work has been devoted to SHE [5,6] (and references therein). To date, nuclei associated with the "island of stability" can be accessed preferentially in 48 Cainduced fusion reactions with actinide targets [6].

Successful use of these reactions was pioneered at the Dubna Gas-Filled Recoil Separator (DGFRS) at the Flerov Laboratory of Nuclear Reactions in Dubna, Russia [6]. Synthesis of nuclei of SHE with Z up to 118 has been reported [7]. Identification was achieved by detecting their subsequent α -decay chain, which always ended with spontaneous fission (SF) [6,7]. Independent experiments at other laboratories have agreed with findings at the DGFRS [8–16]. Discovery of elements with Z=114 and Z=116 was approved by the IUPAC-IUPAP Joint Working Party [17], and recently they were named flerovium (Fl) and livermorium (Lv), respectively [18].

The identification and assignment of nuclear α -decay chains of the odd-Z elements 115 [7,16,19] and 117 [20–23] is more difficult compared to that of the even-Zones. They include more complex α -decay patterns and often longer half-lives, $T_{1/2}$. Moreover, odd nucleons significantly hinder the SF decay of these nuclei, essentially leading to long partial SF half-lives. Thus, the reliability of the assignment of members of such chains is often reduced, because the probability to observe random correlations of unwanted background present in the data set increases with increasing the correlation search time, Δt , between two genuine chain members [5,6]. The longest half-lives detected for α -decaying nuclei of SHE are about 2 min, for ²⁶⁹Sg [12] and ²⁷¹Sg [7]. A successful closer approach towards the center of the "island of stability" depends on a reliable identification of single long-lived nuclei of SHE. An attractive option is to advance setups suitable for registering various nuclear decay modes. Also, detection of their ions stored in, e.g., Penning traps, has been suggested [24,25].

Here, we report on results of the 48 Ca + 249 Bk fusion reaction leading to the SHE with Z=117. First reports on the observation of two isotopes 293,294 117 produced in the 249 Bk (48 Ca, 3–4n) reactions came from a collaboration working at the DGFRS [20,21]. Recently, further data were reported, again from the DGFRS [22,23]. This reaction is unique due to the short half-life of 249 Bk [$T_{1/2}=330(4)$ d, [26]], necessitating dedicated production at Oak Ridge National Laboratory [21] and safe handling of the highly radioactive material. This severely limits the number of independent experiments dedicated to the production of element Z=117.

The experiment was performed at the gas-filled TransActinide Separator and Chemistry Apparatus (TASCA) at GSI [27]. A pulsed (5-ms-long pulses, 50 s⁻¹ repetition rate) ⁴⁸Ca¹⁰⁺ beam was accelerated to energies of 268.3, 270.2, and 274.1 MeV. Targets in ²⁴⁹Bk₂O₃ form were electrodeposited [28] on 2.20(11) μ m thick Ti backings, which faced the beam. Four targets with an area of 6 cm² each were mounted on a wheel, which rotated synchronously with the beam macrostructure [29]. The $\approx 0.48 \text{ mg/cm}^2$ thick target consisted of ²⁴⁹Bk (62%) and its decay product ²⁴⁹Cf (38%) at the beginning of the experiment. The average beam intensity was $4.7 \times 10^{12} \text{ s}^{-1}$. The beam energies in the center of the target were estimated to be $E_{lab} = 252.1(21)$, 254.0(21), and 258.0(21) MeV [30]. Beam doses of 2.3, 4.9, and 3.9×10^{18} were accumulated during 6.3, 10.5, and 10.4 d, respectively. At these beam energies, the evaporation of three or four neutrons is expected from the ²⁹⁷117 compound nuclei at excitation energies of $E^* = 37.8-41.3$, 39.4–42.9, and 42.7–46.2 MeV [31] populating the evaporation residues (ER) ²⁹⁴117 and ²⁹³117 [20–23], respectively.

Compared to the experiments on ^{288,289}Fl performed in 2009 [11,13], TASCA was significantly upgraded for the current experiment [32]. According to the predicted

average charge state (\approx 6.8) of ^{293,294}117 ions moving in helium gas at 0.8 mbar pressure [33,34], the TASCA magnetic system was set to center ions with a magnetic rigidity ($B\rho$) of 2.20 Tm into the focal plane.

The detection system of TASCA consisted of a multiwire proportional counter (MWPC) and a focal plane detector box (FPDB). The latter consisted of a doublesided silicon strip detector (DSSSD)-based implantation detector (hereafter: stop detector), with eight DSSSDs mounted perpendicular in the backward hemisphere of the stop detector to form a five-sided box configuration (box detectors). The FPDB detection efficiency for α particles emitted from implanted nuclei was estimated to be about 80% based on data from test experiments. The stop detector comprised 144 vertical (X) and 48 horizontal (Y) strips on the front and back sides, respectively. Two adjacent single-sided Si-strip detectors were mounted directly behind the stop detector to register particles passing through the stop detector (veto detector). The efficiency of TASCA to guide ER to the stop detector was estimated to be 47(5)% [35]. The average total triggering counting rate was $\approx 700 \text{ s}^{-1}$, distributed over the almost 7000 pixels of the stop detector.

All preamplifier signals from the Y strips of the stop detector were digitized by 60 MHz-sampling ADCs and stored in 50 μ s-long traces. All other signals, i.e., those from the X strips of the stop detector, from box and veto detectors, and from the MWPC were processed with analog electronics and peak-sensing ADCs with a dead time of about 35 μ s. The stop detector's analog and digital branches, which independently initialized the data storage, allowed determining the energy of events in two independent ways. Energy calibrations of stop and box detectors were made using an external α source and α decays of implanted nuclei.

Different position and time correlation analyses between ER-, α -, and SF-like events were carried out. Promising candidates for decay chains were selected from results of analyses searching for ER- α - α ($\Delta t_{\rm ER-}\alpha$ < 1 s, Δt_{α - α < 20 s) and ER- α - α -SF ($\Delta t_{\rm ER-}\alpha$ < 20 s, Δt_{α - α < 200 s, Δt_{α -SF < 500 s) correlations. The search conditions were (i) ER-like events: energy of 3–20 MeV, coincident with a signal from the MWPC; (ii) first and second α -like events: 8.5–12 and 8–11 MeV, respectively; (iii) SF-like events: energies > 50 MeV. Alpha and SF-like events were required to be without coincident MWPC signals.

In total, four chains of two different kinds, two long and two shorter ones with significantly different decay properties, all terminated by SF, were identified. All observed chains will be presented in detail in [36]. Here, we focus on the two long decay chains, which are shown in Fig. 1. The chains were detected at $E_{\rm lab} = 254.0$ (chain 1) and $E_{\rm lab} = 258.0$ MeV (chain 2) in strips X = 103 and 111, respectively, which are on the right half (higher $B\rho$) of the stop detector. In this area, the implantation rate, which is




FIG. 1 (color online). Decay chains assigned to $^{294}117$. Experimental energies in MeV and Δt of all events together with their digitized traces are shown. Energies of reconstructed α particles are marked by an asterisk (*). Boxes with black triangles are corresponding to events observed during beam-off periods. The measured fission fragment energies are from stop + box detectors. Theoretically predicted [38] α -decay energies (in MeV), partial α , and unhindered SF half-lives are given on the left.

overall dominated by transfer products that have lower $B\rho$ than the ER, is significantly lower than the average over the whole stop detector [37].

Chain 1 was found in pixel X = 103 and Y = 41 (Fig. 1). Five α decays occurred during beam-off periods, where the counting rate of α -like events within the energy range of 6–12 MeV was 8×10^{-5} s⁻¹ in this pixel. This number is based on seventy-four events detected in all beam-off periods during the complete 254.0 MeV run. A small stop detector signal of a fourth member (α_3) of the chain was stored only by the digital branch: an energy of 0.6 MeV was registered in strip Y = 41 with $\Delta t = 5.96$ s after α_2 , in a beam-off period. A coincident signal was registered in a box detector and a full energy of 9.3(3) MeV was reconstructed. The counting rate for such events with reconstructed full energies of 6-12 MeV in strip Y = 41with missing X signal was 4×10^{-4} s⁻¹ during beam-off periods. One member, α_6 , of chain 1 was detected during a beam-on period, where the counting rate was 1×10^{-3} s⁻¹. Only one SF-like event was detected in pixel (103,41) during the entire 254.0 MeV run. The counting rate of ERlike events was 4×10^{-3} s⁻¹. A probability of 5×10^{-15} for the observation of random sequences ER- α_{1-7} -SF similar to chain 1 was calculated, according to [39].

Chain 2 occurred in pixel X = 111 and Y = 19(Fig. 1). In all beam-off periods during the 258.0 MeV run, fifty-nine α -like events within the energy range of 6-12 MeV were detected, resulting in a counting rate of 7×10^{-5} s⁻¹; six of these are members of chain 2. Only one SF-like event was found in this pixel during the entire 258.0 MeV run; it occurred 29 h after the last α decay. A signal with an energy $E_{\rm stop}=4.64~{\rm MeV}$ was detected during a beam-off period in between the α_2 and α_4 members of the chain and attributed to the α_3 member. It was likely emitted into the backward direction under a rather shallow angle, thus leaving a relatively large fraction of its energy in the stop detector, but missed detection in the box detector. In total, only twenty-nine events with energies from 3-6 MeV were registered during the beamoff periods in the entire 258.0 MeV run, supporting the nonrandom origin of this event. The probability to observe random sequences similar to chain 2 was estimated to be 5×10^{-16} .

The decay properties of the corresponding members of both chains are compatible with one common origin [39]. The high energies exceeding 10 MeV for the first and second α particles and their correlation times point to an origin from heavy nuclei; however, the observed sequence ER- α_1 - α_2 is not compatible with that of any known nucleus with $Z \le 116$ [7,26]. On the other hand, high energetic α like events, if registered solely with analog electronics, can result from the summing of overlapping low-energy signals [40]. The registered traces of all members of both chains are also presented in Fig. 1. All signals from α particles are recorded as single-peak traces, and the deduced energies from the peak amplitudes agree with those from the analog branch within the energy resolution. Our data allow excluding pile-up events as the source of any member of the two decay chains. Thus, these chains are assigned to originate from the radioactive decay of ER with Z > 116, produced in the fusion of ⁴⁸Ca either with ²⁴⁹Bk or ²⁴⁹Cf, namely, ^{293,294}117 or ^{293,294}118. We attribute these chains to ²⁹⁴117, where the odd proton and neutron are responsible [3,38] for the longest ever measured nuclear α -decay chain originating from SHE [7–16,19].

As is visible in the trace of the ER of chain 1, a small signal with an energy of about 1.4 MeV was observed 1.5 μ s after the ER implantation. The probability to observe a signal with an energy of 1–2 MeV presumably due to heavy charged particles like He-gas atoms or protons in the trace of an ER-like event is less than 1%. The absence of such a signal in the trace of the ER of chain 2, where the ER was implanted deeper into the stop detector, leads us to interpret this signal as not being due to a charged heavy particle from a true member of chain 1 (e.g., an escaped α particle). However, various scenarios leading to the observation of such a signal (e.g., an isomer decaying by cascades of conversion electrons) are being considered [36].

The theoretical calculation of decay properties of odd-odd nuclei is complicated by the scarce information on their structure [3]. However, reliable predictions of the α -decay properties of nuclei originating from ²⁹⁴117 have been given in [38], which satisfyingly agreed with results from the DGFRS [20–23]. The predictions from [38] are included in Fig. 1. Best agreement between the experimental and theoretical α -decay properties is observed for 294 117, 290 115, 286 113, and 270 Db. The experimental T_{SF} of 266 Lr is about 3×10^3 times longer than the theoretical prediction, which can be explained by the presence of odd nucleons. Using the same hindrance factor as a rough approximation, T_{SF} for ²⁸²Rg, ²⁷⁸Mt, ²⁷⁴Bh, and ²⁷⁰Db can be estimated to 1.2 h, 1.5 min, 1.3 h, and 12 h, respectively. These T_{SF} values are larger than the predicted T_a , which points to the dominance of α decay, in accordance with experimental observations. A possible SF branch of ≈8% can be estimated for ²⁷⁰Db based on the estimated value for $T_{\rm SF}$ and the experimental T_{α} .

A summary of the present experimental data together with results reported from groups working at the DGFRS [20–23] is given in Table I. Comparing these data, there is good agreement in most cases. A new finding in our work is the identification of α decay in ²⁷⁰Db and the new nucleus

TABLE I. Decay properties of the nuclei originating from ²⁹⁴117 observed in this work, reported from DGFRS [23] and from the combined data sets. Half-life uncertainties corresponding to the 68% confidence level are given, according to [39].

	E_{α} (MeV) $T_{1/2}$		Decay mode ^a	$N_{lpha}/N_{ m SF}^{\scriptscriptstyle m b} \ T_{1/2}$
Nuclei	This work	DGFRS [23]	Combined	
²⁹⁴ 117	11.05(4)	10.81–10.97	α	5/-
	51^{+94}_{-20} ms	50^{+60}_{-18} ms	100%	51^{+41}_{-16} ms
²⁹⁰ 115	10.31(4)	9.78-10.28	α	5/-
	$1.3^{+2.3}_{-0.5}$ s	$0.24^{+0.28}_{-0.09}$ s	100%	$0.75^{+0.61}_{-0.23}$ s
²⁸⁶ 113	9.3(3)	9.61-9.75	α	6/-
	$2.9^{+5.3}_{-1.1}$ s	13^{+12}_{-4} s	100%	$7.9^{+5.5}_{-2.3}$ s
282 Rg	9.05(3), 8.86(3)	9.01(5)	α	6/-
	$3.1^{+5.7}_{-1.2}$ min	$1.0^{+1.0}_{-0.3}$ min	100%	2.1 ^{+1.4} _{-0.6} min
^{278}Mt	9.45(3)	9.38-9.55	α	5/-
	$3.6^{+6.5}_{-1.4}$ s	$5.2^{+6.2}_{-1.8}$ s	100%	$4.4^{+3.6}_{-1.4}$ s
²⁷⁴ Bh	8.84(3)	8.76(5)	α	5/-
	30^{+54}_{-12} s	54^{+65}_{-19} s	100%	42^{+34}_{-13} s
²⁷⁰ Db	7.90(3)	SF	$lpha/\mathrm{SF}^{\scriptscriptstyle\mathrm{c}}$	2/1°
	$1.0^{+1.9}_{-0.4}$ h	17^{+15}_{-6} h	83/17%	$1.1^{+1.5}_{-0.4}$ h
266 Lr	SF	(not observed)	SF	-/2
	11^{+21}_{-5} h		100%	11^{+21}_{-5} h

^aCorresponding α and/or SF branching ratios are given.

²⁶⁶Lr. In total four decay chains assigned to ²⁹⁴117 were reported from the DGFRS. All chains were assigned to be terminated by SF of 270 Db. The measured Δt were 33, 38, 24, and 1.2 h [20,22,23]. Respecting our data, which indicate the half-lives of $^{270}\mathrm{Db}$ and $^{266}\mathrm{Lr}$ to be $1.0^{+1.9}_{-0.4}$ and 11^{+21}_{-5} h, respectively, it appears that the last one of the DGFRS chains, where a significantly shorter Δt was recorded than in the three first ones, was indeed terminated by SF of 270 Db. In the other cases, the α decay of 270 Db may have remained unidentified (as considered in [20–22]), while the chains terminated by SF of ²⁶⁶Lr. If so, a branching ratio for SF, of about 1/6 = 17% is deduced for ²⁷⁰Db. This is in agreement with the estimate given above within the uncertainty of low statistics [39].

The partial SF half-life (6.5 h) of ²⁷⁰Db is comparable to those of 267 Db ($T_{1/2} \approx 1.4$ h, SF) [19] and 268 Db ($T_{1/2} \approx 25$ h, SF) [19], which lie in the vicinity of the neutron shell closure at N = 162. The half-life of the new nucleus 266 Lr with N = 163 is comparable to those of the heavier isotones 267 Rf ($T_{1/2} \approx 1.3$ h, SF) [7] and 268 Db. This suggests the stabilizing influence of the N = 162shell closure to extend towards lower Z at least down to Z = 103, which is also seen in the low E_{α} value of ²⁷⁰Db.

The cross sections for the production of $^{294}117$ were evaluated to $0.7^{+1.6}_{-0.6}$ and $0.9^{+2.0}_{-0.7}$ pb at $E_{\rm lab} = 252.0$ –256.1 and 255.9-260.0 MeV, respectively. These are consistent with values reported from the DGFRS. At $E_{\rm lab} = 250.0$ -254.1 MeV no events were observed, corresponding to an upper cross section limit of 2.9 pb [39].

In conclusion, we observed the nuclear decay of two atoms of element Z=117 and its daughter products, synthesized in the ${}^{48}{\rm Ca}+{}^{249}{\rm Bk}$ reaction. The nuclei ²⁹⁴117, ²⁹⁰115, ²⁸⁶113, ²⁸²Rg, ²⁷⁸Mt, ²⁷⁴Bh, ²⁷⁰Db were identified by their α decays and ²⁶⁶Lr by its SF decay, which terminated the decay chains. Results of the present work confirm previously reported data [20-23] on the decay chains assigned to ²⁹⁴117. In addition, we report a previously unknown α branch in ²⁷⁰Db, which populated the new SF unstable nucleus ²⁶⁶Lr. Our data confirm the perseverance of the N = 162 shell closure towards lighter Z, at least down to Z = 103. ²⁷⁰Db is the most long-lived α -decaying nucleus above No (Z=102). The decay chain members from $^{290}115$ to 266 Lr all decay with $T_{1/2} \gtrsim 1$ s, which opens prospects for their chemical investigation and off-line studies. Our experimental data show the sensitivity of TASCA for the identification of the radioactive decay of single nuclei with half-lives from sub-\mu s up to about 1 d. Future studies of SHE designed to allow a closer approach towards the center of the "island of stability" may require the safe measurement of even longer half-lives.

We are grateful for GSI's ECR ion source and UNILAC staff, and the Experimental Electronics department for their continuous support of the experiment. This work was financially supported in part by the German (05P12UMFNE), the Helmholtz association

Number of α and SF events used for the calculations of $T_{1/2}$.

^cOne SF event out of four reported in [23], see text for discussion.

(VH-NG-723), the Australian and Swedish Research Councils, the U.S. Department of Energy by LLNL (DE-AC52-07NA27344), the Laboratory Directed Research and Development Program at LLNL (11-ERD-011) and the Helmholtz Institute Mainz. This work was co-sponsored by the Office of Science, U.S. Department of Energy, and supported under U.S. DOE Grant No. DE-AC05-00OR22725, and at Vanderbilt DOE Grant No. DE-FG05-88ER40407, and the UK Science and Technology Funding Council (STFC). J. M. G., K. E. G., and H. N. were supported by the U.S. Department of Energy, Office of Science, Nuclear Physics, Low Energy Physics Program, through the Lawrence Berkeley National Laboratory under Contract No. DE-AC02-05CH11231.

- *J.Khuyagbaatar@gsi.de
- [†]Indian Institute of Technology Roorkee, Roorkee 247667, India.
- [1] A. Sobiczewski, F. A. Gareev, and B. N. Kalinkin, Phys. Lett. **22**, 500 (1966).
- [2] W. D. Myers and W. J. Swiatecki, Nucl. Phys. 81, 1 (1966).
- [3] A. Sobiczewski and K. Pomorski, Prog. Part. Nucl. Phys. 58, 292 (2007).
- [4] S. Ćwiok, P.-H. Heenen, and W. Nazarewicz, Nature (London) 433, 705 (2005).
- [5] S. Hofmann and G. Münzenberg, Rev. Mod. Phys. 72, 733 (2000).
- [6] Yu. Ts. Oganessian, J. Phys. G 34, R165 (2007).
- [7] Yu. Ts. Oganessian, Radiochim. Acta **99**, 429 (2011).
- [8] S. Hofmann et al., Eur. Phys. J. A 32, 251 (2007).
- [9] L. Stavsetra, K. Gregorich, J. Dvorak, P. Ellison, I. Dragojević, M. Garcia, and H. Nitsche, Phys. Rev. Lett. 103, 132502 (2009).
- [10] R. Eichler et al., Radiochim. Acta 98, 133 (2010).
- [11] Ch. E. Düllmann et al., Phys. Rev. Lett. 104, 252701 (2010).
- [12] P. A. Ellison et al., Phys. Rev. Lett. 105, 182701 (2010).
- [13] J. M. Gates et al., Phys. Rev. C 83, 054618 (2011).
- [14] S. Hofmann et al., Eur. Phys. J. A 48, 62 (2012).
- [15] A. Yakushev et al., Inorg. Chem. 53, 1624 (2014).
- [16] D. Rudolph et al., Phys. Rev. Lett. 111, 112502 (2013).
- [17] R. C. Barber, P. J. Karol, H. Nakahara, E. Vardaci, and E. W. Vogt, Pure Appl. Chem. 83, 1485 (2011).
- [18] R. D. Loss and J. Corish, Pure Appl. Chem. **84**, 1669 (2012).
- [19] D. Rudolph et al., Acta Phys. Pol. B 45, 263 (2014).

- [20] Yu. Ts. Oganessian et al., Phys. Rev. Lett. 104, 142502 (2010).
- [21] Yu. Ts. Oganessian *et al.*, Phys. Rev. C **83**, 054315 (2011).
- [22] Yu. Ts. Oganessian *et al.*, Phys. Rev. Lett. **109**, 162501 (2012).
- [23] Yu. Ts. Oganessian et al., Phys. Rev. C 87, 054621 (2013).
- [24] M. Block et al., Nature (London) 463, 785 (2010).
- [25] E. Minaya Ramirez et al., Science 337, 1207 (2012).
- [26] http://www.nndc.bnl.gov/ensdf/.
- [27] A. Semchenkov *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. B **266**, 4153 (2008).
- [28] J. Runke et al., J. Radioanal. Nucl. Chem. 299, 1081 (2014).
- [29] E. Jäger, H. Brand, Ch. E. Düllmann, J. Khuyagbaatar, J. Krier, M. Schädel, T. Torres, and A. Yakushev, J. Radioanal. Nucl. Chem. 299, 1073 (2014).
- [30] J. F. Ziegler, Nucl. Instrum. Methods Phys. Res., Sect. A 219, 1027 (2004).
- [31] W. D. Myers and W. J. Swiatecki, Nucl. Phys. A601, 141 (1996).
- [32] A. Yakushev et al. (to be published).
- [33] J. Khuyagbaatar *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **689**, 40 (2012).
- [34] J. Khuyagbaatar, V. P. Shevelko, A. Borschevsky, Ch. E. Düllmann, I. Yu. Tolstikhina, and A. Yakushev, Phys. Rev. A 88, 042703 (2013).
- [35] K. E. Gregorich, Nucl. Instrum. Methods Phys. Res., Sect. A **711**, 47 (2013).
- [36] J. Khuyagbaatar et al. (to be published).
- [37] See Supplemental Material at http://link.aps.org/supplemental/10.1103/PhysRevLett.112.172501, Fig. 1, showing the distribution of implantation counting rates along the *X* strips of the stop detector.
- [38] A. Sobiczewski, Acta Phys. Pol. B 41, 157 (2010).
- [39] K.-H. Schmidt, C.-C. Sahm, K. Pielenz, and H.-G. Clerc, Z. Phys. A 316, 19 (1984).
- [40] See supplemental Material at http://link.aps.org/supplemental/10.1103/PhysRevLett.112.172501, Fig. 2, showing an example for a spectrum containing events from the decay-chain 222 Pa $^{-218}$ Ac, where energies of both α 's are summed up due to the short $T_{1/2}=1~\mu s$ of the daughter. Only an analysis of digitized data, as they are simultaneously recorded at TASCA, allows resolving pile-up events and producing clear spectra for mother and daughter. While the analog spectrum contains a large number of entries within wide energy range 7–18 MeV, the spectra produced from the digitized data clearly show peaks.