



# LUND UNIVERSITY

## **Individual whole-body concentration of (137)Cesium is associated with decreased blood counts in children in the Chernobyl-contaminated areas, Ukraine, 2008-2010.**

Lindgren, Anna; Stepanova, Eugenia; Vdovenko, Vitaliy; McMahon, Daria; Litvinetz, Oksana; Leonovich, Elena; Karmaus, Wilfried

*Published in:*

Journal of Exposure Science & Environmental Epidemiology

*DOI:*

[10.1038/jes.2013.60](https://doi.org/10.1038/jes.2013.60)

2015

[Link to publication](#)

*Citation for published version (APA):*

Lindgren, A., Stepanova, E., Vdovenko, V., McMahon, D., Litvinetz, O., Leonovich, E., & Karmaus, W. (2015). Individual whole-body concentration of (137)Cesium is associated with decreased blood counts in children in the Chernobyl-contaminated areas, Ukraine, 2008-2010. *Journal of Exposure Science & Environmental Epidemiology*, 25(3), 334-342. <https://doi.org/10.1038/jes.2013.60>

*Total number of authors:*

7

### **General rights**

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117  
221 00 Lund  
+46 46-222 00 00

**Individual whole body concentration of <sup>137</sup>Cesium is associated with decreased blood counts in children in the Chernobyl contaminated areas, Ukraine, 2008-2010.**

**Authors:** Anna Lindgren<sup>1</sup>, Eugenia Stepanova<sup>2</sup>, Vitaliy Vdovenko<sup>2</sup>, Daria McMahon<sup>3</sup>, Oksana Litvinetz<sup>2</sup>, Elena Leonovich<sup>2</sup>, Wilfried Karmaus<sup>4</sup>

<sup>1</sup>Department of Occupational and Environmental Medicine, Lund University, Lund, Sweden.

<sup>2</sup>Research Center for Radiation Medicine, Academy of Medical Sciences of Ukraine, Kyiv, Ukraine.

<sup>3</sup>Department of Epidemiology and Biostatistics, Norman J. Arnold School of Public Health, University of South Carolina, Columbia, South Carolina, USA.

<sup>4</sup> Division of Epidemiology, Biostatistics and Environmental Health, School of Public Health, University of Memphis, Tennessee, USA.

**Corresponding author**

Wilfried Karmaus, MD, Dr.med., MPH

Division of Epidemiology, Biostatistics and Environmental Health, School of Public Health, University of Memphis, 301 Robison Hall, Memphis, TN 38152, USA.

Tel: (+1) 901-678-2491

Fax: (+1) 901-678-0372

Email: karmaus1@memphis.edu

## **Abstract**

Narodichesky region, Zhitomir Oblast, Ukraine, is situated approximately 80 km from the Chernobyl Nuclear Power Plant, which exploded in 1986 and polluted the environment. A previous study found that children living in villages with high activity of  $^{137}\text{Cs}$  in the soil had decreased levels of hemoglobin, erythrocytes and thrombocytes. These findings motivated the present study which used a more comprehensive exposure assessment, including individual whole-body concentrations (WBC) of  $^{137}\text{Cs}$  (Bq/kg). This cross-sectional sample examined between 2008-2010, included 590 children in the age 0-18 years.

Children with higher individual  $\log(\text{WBC})$  activity in the body, had significantly decreased hemoglobin, erythrocyte and thrombocyte counts. The effect of  $\log(\text{WBC})$  on decreased thrombocyte count was only seen in children older than 12 years. The average village activity of  $^{137}\text{Cs}$  ( $\text{kBq}/\text{m}^2$ ) in soil was associated with decreased blood counts only indirectly, through  $^{137}\text{Cs}$  in the body as an intermediate variable.

Children in this study were born at least four years after the accident and thus exposed to low doses of ionizing radiation from  $^{137}\text{Cs}$ . This cross-sectional study indicates that low levels may be associated with decreased blood counts, but we cannot exclude that these results are due to residual confounding factors.

## **Key words**

child exposure/health, epidemiology, personal exposure, radiation

## **Running short title**

$^{137}\text{Cs}$  and blood counts in children in Ukraine

## **Ethical permission**

The study was approved by the Committee on Bioethics of the Research Center for Radiation Medicine, Academy of Medical Sciences of Ukraine (Report #1, August 31, 2007) and by the Office of Research Compliance at the University of South Carolina.

## **BACKGROUND**

The Narodichesky region in Zhitomir Oblast, Ukraine, is one of the most contaminated populated areas around the Chernobyl Nuclear Power Plant, which exploded in 1986. The population in the contaminated areas is still exposed to ionizing radiation from the accident, mainly from the long-lived radioactive isotope  $^{137}\text{Cs}$  (1), which has a physical half-life of 30 years. A previous longitudinal follow-up (1993-1998) found that children living in villages with high activity of  $^{137}\text{Cs}$  in the soil had decreased levels of hemoglobin, erythrocytes and thrombocytes (2). These findings motivated the present study, which used a more comprehensive exposure assessment, including individual whole-body concentrations (WBC) of  $^{137}\text{Cs}$  (Bq/kg).

Radiation exposure in the years after the accident has been declining not only because of physical decay of the radionuclides, but also because of vertical migration deeper into the soil, which shields off the exposure, and because of countermeasures to reduce the doses (3). The average total (external and internal) radiation dose from Chernobyl for the population in the Narodichesky region has been estimated to be 17 mSv for the single year 1986, 13 mSv in cumulative dose for 1987-90, and only 3.2 mSv in cumulative dose 2001-05 (1). The average total dose from the accident for the single year 2008 in the Narodichesky region was

estimated to be 0.13-2.2 mSv, depending on village (4). The exposure is thus small in children born today compared to early after the accident.

External exposure from soil is only a part of the total dose, in the Narodichesky region it was approximately 70% of total exposure during 1986-2005 (1). In addition to external exposure radiating directly from radioactive deposition in the soil, people are internally exposed from ingestion of contaminated food; high activity of  $^{137}\text{Cs}$  has been found in milk, meat and especially forest products such as mushrooms, wild berries, game meat, and fish from local ponds (3, 5, 6). Radioactivity from inhaled dust was an additional major source of internal exposure in the first years after the accident, but is an insignificant pathway today (7). The levels of food contamination do not necessarily decrease as steadily as the soil radiation, since it's not affected by the vertical migration, but rather fluctuate from year to year depending on the harvest. The internal exposure from food makes up an increasing proportion of the total exposure in later years in many areas (7, 8).

Generally, a high level of  $^{137}\text{Cs}$  deposition in soil can be expected to also contribute to a high level of  $^{137}\text{Cs}$  in the body, and such a correlation has been reported (9), but there are also studies which have found no correlation (10, 11). The lack of agreement depends on many factors such as the soil-transfer to vegetation, dietary habits, and countermeasures applied to protect the population.

$^{137}\text{Cs}$  is an alkali metal which has very similar characteristics to Potassium and is homogeneously spread in the body. The average biological half-life in the body is 100 days before excretion. The uptake is rapid and complete both by inhalation and ingestion (12). Considering the homogenous spread of moderate amounts of  $^{137}\text{Cs}$  in the body, organ-specific doses resulting from  $^{137}\text{Cs}$  in the body can in the Zhitomir region be expected to be low.

Although the acute detrimental effects of high-dose radiation on the hematopoietic system are well known, both from Chernobyl (13), and other radiation accidents (14), there is little information on long-term low dose radiation effects on the hematopoietic system. The Techa river incident, a plutonium processing facility established in 1948 with radioactive waste leakage into the river, caused decreased levels of thrombocytes, erythrocytes and hemoglobin, in addition to a range of other symptoms which together were labeled "chronic radiation syndrome" (15, 16). However, in residents of the areas along the Techa river, the accumulated bone-marrow doses from internal  $^{137}\text{Cs}$  and  $^{90}\text{Sr}$  were up to 3-4 Sv, considerably higher than the total accumulated doses in the Narodichesky region.

Regarding health effects in the Chernobyl area, in addition to our previous study, we have identified six published articles on long-term effects on red blood cells, but these are in people born before the accident or irradiated in utero (17-22). Stepanova et al (2006) found more transitory, prehemolytic and degenerative forms of erythrocytes in children irradiated in utero, in comparison with control children.

A study with cross-sectional comparisons in 1986, 1992 and 1998 found no difference in mean hemoglobin, erythrocyte or thrombocyte counts in children residing in settlements with  $^{137}\text{Cs}$  soil activity of 37-555 kBq/m<sup>2</sup> compared to children living in settlements with less than 37 kBq/m<sup>2</sup> (19). However, this study only describe means in the two exposure groups without statistical adjustment for any factors, which is typical for many of the studies of health effects from Chernobyl (23). In contrast, in a repeated measurement study (1993-1998), which included children born before and after the accident, we found decreased hemoglobin, erythrocyte and thrombocyte counts in children residing in villages with  $^{137}\text{Cs}$  soil activity of 266-310 kBq/m<sup>2</sup> and 350-879 kBq/m<sup>2</sup> compared to 29-112 kBq/m<sup>2</sup> (2).

The purpose of the analysis at hand is to investigate whether our previous findings can be replicated and expanded upon, by using more comprehensive exposure assessment including not only average village soil activity of  $^{137}\text{Cs}$  but also individual measurements of whole body concentration of  $^{137}\text{Cs}$ .

## **METHODS**

### **Study population**

Settlements in the Narodichesky region (Zhitomir Oblast, Ukraine), are located approximately 80 km from the Chernobyl Nuclear Power Plant. In 2001, a population census determined that this region had approximately 11 400 inhabitants, including 2000 children. The study area has been extensively described in previous publications (2, 24). The lifestyle in the Narodichesky region is similar in all villages, with predominantly locally homegrown food supply and little migration. All children in Narodichesky region are provided with free breakfast and lunch at school. The menu at school has to meet standards of sanitary hygiene and follow the provision of The Ukrainian Government (25). The meals served in school are prepared from products imported from areas free from contamination with radionuclides. Meals consumed at home, at the 5 o'clock snack and supper, are prepared from local produce (milk, meat, fish, vegetables, berries, mushrooms and fruit).

### **Study selection and missing data**

This cross-sectional study included 590 children in the age 0-18 years; 566 (age 2-18) had complete exposure measurements on  $^{137}\text{Cs}$  whole body concentrations and residential  $^{137}\text{Cs}$  soil estimates, and were used in the analyses.

We examined predominantly children of school age. In addition, some preschool children were assessed upon request of their parents. On the day of examination, all school children present in a class were transported to the Central Hospital of Narodichi, by their teachers. Parents were notified of the forthcoming examinations in advance, and their verbal permission was requested for young children (0-6 years). Parents of all school children gave their written permission for the examination.

We excluded from the analysis 23 children with missing values of  $^{137}\text{Cs}$  WBC and one child with missing information on  $^{137}\text{Cs}$  in soil. Missing WBC was found exclusively in children 0-4 years of age, since these children had difficulties to comply with the exposure assessment.

### **Exposure assessment**

**Village average soil activity of  $^{137}\text{Cs}$ .** The average  $^{137}\text{Cs}$  soil activity per village for 2008 was obtained from a report published by the Ukrainian Ministry of Health (4). The Ministry publishes yearly estimates for the purpose of monitoring the population exposure. The report stated that the estimates of  $^{137}\text{Cs}$  in soil for 2008 were based on soil measurements conducted in 1992 and that a decay function was applied to obtain the estimates for 2008. The estimated average external effective dose for each village in 2008, was also obtained from this publication.

**Individual measurements of whole body concentration of  $^{137}\text{Cs}$ .** Whole-body concentration of  $^{137}\text{Cs}$  was measured for all children in 2008-2010 using a gamma-spectrometer (Whole Body Counter SCRINNER – 3M, designed and produced by INECO (Ukrainian Institution of Human Ecology, Academy of Technological Sciences).

”SCRINNER-3M” is used in 12 regional medical centers of the Ukraine for mass screening of



residents of contaminated areas (26), and is also used in Belarus, at the Whole Body Counter Laboratory at the Institute of Radiation Safety (BELRAD) (27).

The whole-body counter is designed as a standard chair (with lead shielding in the back and seat of the chair for local protection from background radiation). The single detector of the "SCRINNER-3M" is incorporated into the back of the chair and detects  $\gamma$ -rays from the top of the head to the knee level. The angle between the back of a chair and the seat is 100 degrees, and the distance from the back and the base of the chair to detector is 40 cm. Children were sitting in upright position during the measurements.

The "SCRINNER-3M" is equipped with Pb collimator: thickness 50mm, coaxial scintillation detector NaI(Tl):  $\varnothing 150 \times 100$ mm. Prior to the measurements, "SCRINNER -3M" is calibrated using phantoms in 6 age groups, composed from taurpaline filled with dried peas with known radionuclide content. Additional information about the phantoms is presented in Supplemental file 1 (S1). IAEA has made intercomparisons of chair type whole body counters with other whole body counters, and also of "dried pea phantoms" (28). The general characteristics of the "SCRINNER-3M" in calibrated state is given in Table I. The minimum detection activity (MDA) for 3 minutes exposition on a 70kg adult phantom (with a maximum error of 30%) was 340Bq. The measurements were conducted according with the recommendations provided by the Research Center of Radiation Medicine Academy of Medical Sciences of Ukraine (29, 30). The protocol has been published (31). Each measurement continued until its error dropped to the level of 30% or lower as required by the standards of International Commission on Radiological Protection (ICRP). Under these measurement conditions, the required measurement error for most children was obtained within 3 minutes, and only a few children required up to 5 minutes. Calculations of measured activity (Bq) accounted for child's weight by including calibration coefficients. To further

reduce residual confounding by weight, we took further account of child's weight by using WBC (Bq/kg) as the unit for our regression analyses, instead of the measured activity (Bq).

Every child had only one measurement of the WBC of  $^{137}\text{Cs}$ . Almost all of the children, 561 out of 566, had this measurement in 2009. The measurements were predominately conducted in three months: April (n=255), September (n=209), and December (n=55). A few measurements were done in others months. Since levels of  $^{137}\text{Cs}$  are known to potentially vary by season due to different food consumption during different seasons (i.e., mushrooms in the autumn) (32, 33), we adjusted for season in the final multivariable statistical analyses.

### **Calculations of individual internal effective dose from $^{137}\text{Cs}$**

The internal dose calculations were based on the age-dependent committed dose coefficients for ingestion  $e$  (nSv/Bq), from ICRP publication 72 (34). Based on the point-measurements of  $^{137}\text{Cs}$  in the body, a constant steady-state of the measured activity was assumed for the whole year. A biological half-life of 100 days was assumed, with an effective half-life of 99 days.

The intake in Bq to remain in steady state was first calculated for one day:  $(\ln(2)/99) \times 137\text{Cs}$  (Bq). The doses were then calculated as:

Daily intake to remain in steady state (Bq)  $\times e$  (nSv/Bq)  $\times 365$ , and converted to mSv/year.

### **Medical assessment and questionnaires**

Medical assessments followed the same protocol as the previous studies (2, 23). Blood was collected in tubes containing EDTA. A blood count, including erythrocytes (red blood cell count; RBC), leukocytes, thrombocytes (Plt) and hemoglobin (Hb), was conducted using Sysmex model F-800 (TOA Medical Electronics Company, Kobe, Japan). Normal blood smears were stained using the standardized azure B-eosin GIEMSA Y Romanowsky

procedure and the cells were counted. We calculated the Color Parameter (CP) to classify anemia. CP is an old parameter equal to Mean Corpuscular Hemoglobin (MCH)\*0.03 with  $MCH = Hb \text{ (g/L)} / \text{number of erythrocytes (x10}^{12}\text{cells/L)}$  (35).

CP <0.8 (hypochrom) may be indicative of iron-deficiency anemia, CP = 0.8-1.0 (normochrom) is considered normal range, and CP >1.0 (hyperchrom) may be indicative of B12/folate-deficiency anemia. The reference values for hemoglobin, erythrocytes, and thrombocytes are provided in (Table II) (36, 37).

More extensive confounder information was gathered in this study compared to our previous investigation. Information on potential confounders such as environmental tobacco smoke, active tobacco smoking, and type of fuel used for residential heating or cooking, was collected during interviews. Medical assessments, interviews, and the measurements of WBC of  $^{137}\text{Cs}$ , were performed on the same day.

### **Statistical methods**

Multiple Linear Regression analyses were performed using R (R Development Core Team, 2012), version 2.15.0 to estimate the associations of  $^{137}\text{Cs}$  in soil and WBC with blood counts. The outcome variables were hemoglobin, erythrocyte count, and thrombocyte count. The exposure variables were log(WBC) and  $^{137}\text{Cs}$  in soil ( $\text{kBq/m}^2$ ). The log transformation of WBC ( $\text{Bq/kg}$ ) was done to fulfill the model assumptions of linearity and constant variance. We adjusted all estimates for sex of the child, age (continuous), season (spring, summer, autumn, winter), environmental tobacco smoke (ETS) (Yes/No), active tobacco smoking (0 cigarettes/day, 1-10 cigarettes/day, >10 cigarettes/day) and use of coal/wood for cooking (Yes/No). These variables were deemed as potential confounders, and kept in the model regardless of their statistical significance, for model consistency. All 566 children had

complete information on these variables. None of the 566 observed values were excluded from the analyses, despite a few extreme values on WBC. A few extreme values were expected since the  $^{137}\text{Cs}$  intake may be highly variable.

Effects of interaction between  $^{137}\text{Cs}$  in soil and age, and  $\log(\text{WBC})$  and age on blood counts, were tested by partial t-test with age as a continuous variable. If the interaction terms were significant we presented estimates stratified by age. As a sensitivity analysis we included CP as an additional categorical variable (hypochrom, normochrom, hyperchrom), in the model which estimated effects of  $^{137}\text{Cs}$  in soil and  $\log(\text{WBC})$  on Hb. The purpose was to see if the effect estimates changed when adjusting for hypochrom and hyperchrom erythrocytes, which are often caused by nutritional deficiencies. The significance level for all analyses was  $\alpha=0.05$ .

Path-analysis, i.e. Covariance Analysis of Linear Structural Equations (28), was performed using SAS version 9.3. Path-analysis is a method used to decompose sources of correlation to see how much of the total effects of a variable is due to its "direct effect" (partial correlation with the outcome after adjustment for other variables), and "indirect effect" (correlation with variables which have a partial correlation with the outcome). Path-analysis can be used to test the assumption that a variable is an intermediate step in a chain of responses. If it is, the exposure variable should have a direct effect on the assumed intermediate variable, which should have a direct effect on the outcome.

Based on results from our previous study, we hypothesized that  $^{137}\text{Cs}$  in soil 1) has a direct effect on blood counts, and 2) also has an indirect effect on blood counts through WBC Bq/kg as an intermediate variable. The full model that we first specified is included in the Supplemental file 4 (S4). Blom transformation was used to standardize the variables. The model was modified to fulfill the criteria for model fit, with non-significant Chi-square test,

Root Mean Square Residual (RMSR) close to zero, goodness of fit index (GFI) and adjusted goodness of fit index (AGFI) > 0.98, indicating a good fit (38). The final fitted path models supported by the data, are presented in Supplemental file 2, S2.

## **RESULTS**

### **Population**

Fifty percent of the children were recruited from the small town Narodichi, and the rest from 27 different rural villages (Table III). Five children were living outside the Narodichesky region.

All children were born after the accident, the oldest child in 1990 (Table IV). The majority (73.5%) of the children were between the ages of 12-18 at the time of examination. Just under half of the sample (45%) was exposed to environmental tobacco smoke (ETS), but only 5% of the children were active daily tobacco smokers (Table IV).

### **Exposure description of WBC and $^{137}\text{Cs}$ kBq/m<sup>2</sup>**

The individual WBC Bq/kg had a right-skewed distribution, with few observations above 300 Bq/kg (Fig 1a). The average activity of  $^{137}\text{Cs}$  in soil of the village had a left-skewed distribution with approximately half of the population living in a village with >200 kBq/m<sup>2</sup> (Fig 1b, Table III). Overall, the individual WBC had moderate positive correlation with the activity of  $^{137}\text{Cs}$  in soil (Spearman's rank correlation,  $r_s = 0.508$   $p < 0.001$ ; Fig 1c).

The WBC was highest for measurements performed in the spring (Figure 2). The mean internal effective dose was 0.14 mSv/year (Table III). There were 5 children with an estimated effective internal dose >1mSv/year.

## Blood markers

Overall, many children appeared to be low in hemoglobin concentrations and erythrocyte counts. 21.7% of the children were below the reference values for Hb, and 65.4% were below the reference values for erythrocytes (RBC) for their age. Hemoglobin values and erythrocyte counts below reference levels were more common in the older children (Table II). For thrombocyte counts (PLT), only 0.7% of the children were below the reference values for their age.

Plots of the bivariate relationships between individual WBC and log(WBC) with blood counts, suggest negative associations between WBC and Hb, RBC:s and PLT:s (Fig 3). Although less apparent, similar relationships were observed between  $^{137}\text{Cs}$  activity in soil (kBq/m<sup>2</sup>) and Hb, RBC:s and PLT:s (Fig 4). The log transformed WBC (Bq/kg), log(WBC), had a more linear relationship with the outcomes, and was used for the multivariate analyses

In multivariate analyses, log(WBC), was associated with the hemoglobin level (Table V). One unit increase in log(WBC) decreased hemoglobin by -5.94 g/L, 95%CI [-6.75, -5.13]. One unit increase in  $^{137}\text{Cs}$  activity in residential soil was associated with decreased Hb, when adjusting for confounders other than individual log(WBC), but when adjusting for log(WBC) the effect was small and statistically insignificant (Table III). There was no interaction between log(WBC) or  $^{137}\text{Cs}$  in soil and age, regarding the effects on Hb.

The log(WBC), was also associated with erythrocyte count (RBC; Table V). One unit increase in log(WBC) decreased the RBC by  $-0.238 \times 10^{12}/\text{L}$ , 95%CI [-0.27, -0.21]. One unit increase in  $^{137}\text{Cs}$  activity in soil was also associated with decreased RBC when adjusting for confounders other than individual log (WBC), but when adjusting for log (WBC) the effect

was very small and instead associated with increased RBC (Table V). There was no interaction between log(WBC) or  $^{137}\text{Cs}$  in soil and age, regarding the effects on RBC.

Regarding the effects on thrombocytes, there was a significant interaction between log(WBC) and age (p-value=0.004), when using age as a continuous variable, but nonsignificant for  $^{137}\text{Cs}$  in soil and age. We therefore show separate results for age < 12 years and 12-18 years. In children 12 years and older, log(WBC) was associated with thrombocyte count (Table V). One unit increase in log(WBC) decreased the count by  $-15.99 \times 10^9/\text{L}$ , 95% CI [-21.37, -10.62]. In children 12 years and older, one unit increase in  $^{137}\text{Cs}$  activity in soil was also associated with decreased thrombocytes when adjusting for confounders other than individual log(WBC), but when adjusting for log(WBC) the effect was very small and statistically insignificant (Table V). In children 2-11 years, there was no statistical association between log(WBC) or  $^{137}\text{Cs}$  in soil with the thrombocyte count.

### **Sensitivity analysis using Color Parameter**

There were 18 children with color parameter <0.8, indicating low levels of hemoglobin per erythrocyte; 527 children had a normal CP=0.8- 1.0, and 21 children had a color parameter >1.0 indicating high levels of hemoglobin per erythrocyte. Adjusting for CP as a categorical variable with three groups (CP< 0.8, CP=0.8-1.0, CP >1.0) did not change the effect estimate for log(WBC) on Hb, which changed by less than 2%. The effect of  $^{137}\text{Cs}$  activity in soil remained small and statistically insignificant after adjustment for CP. The fact that adjustment for CP did not change the results, decreases the likelihood (but does not fully exclude) that iron-deficiency anemia or B<sub>12</sub>/folate-deficiency anemia (which typically exhibits abnormal CP), are confounders in this study.

## **Path analysis investigating $^{137}\text{Cs}/\text{kg}$ as an intermediate variable between $^{137}\text{Cs}$ in soil and Hb, RBC and PLT**

The partial correlations supports our hypothesis that residential soil activity of  $^{137}\text{Cs}$  had an indirect effect on blood counts through a direct effect of WBC (Bq/kg). There was no support for any direct effects from  $^{137}\text{Cs}$  in soil. The graphs of the final fitted models supported by the path analysis can be seen in the Supplemental file 2 (S2).

## **DISCUSSION**

The main strength of the study was, compared to our previous study, that the village levels of  $^{137}\text{Cs}$  exposure were complemented by assessment of individual levels of  $^{137}\text{Cs}$  WBC, which can be expected to decrease the amount of measurement error. We also gathered individual information on risk factors such as active tobacco smoking, environmental tobacco smoke, and household fuels such as coal and wood, which had only marginal influence on the risk estimates. Since the residents in the investigated areas are poor, anemia due to poor nutrition is not unlikely, and we hypothesized this could confound or dilute the effects of radiation. Our results, however, show that most children had normal CP, which indicates that iron or  $\text{B}_{12}$ /folate deficiency due to poor nutrition is not a main cause of anemia in this area, and our sensitivity analysis shows that having low/high CP was not associated with  $^{137}\text{Cs}$  exposure. These findings speak against that the low blood counts are due to nutritional deficiencies. However, we cannot exclude completely that a combination of iron-deficiency and  $\text{B}_{12}$ -may exist. The combination of both would result in a normal CP. Hence, the absence of other biochemical parameters such as ferritin and  $\text{B}_{12}$  is a limitation of our study. The expected effect of long-term exposure to  $^{137}\text{Cs}$  on red blood cells is most likely related to activation of



free radical oxidation in these cells. It is also possible, that long-term exposure  $^{137}\text{Cs}$  may directly affect bone marrow and lead to decreased production of blood cells and development of normochrom (normocytic) anemia, which is also what was mainly observed in this population.

Our results show that children of the Narodichesky region with high levels of log transformed WBC of  $^{137}\text{Cs}$ , had decreased levels of hemoglobin, and erythrocyte and thrombocyte counts. The average village level of  $^{137}\text{Cs}$  in soil was not associated with decreased blood counts after adjustment for individual body burden of  $^{137}\text{Cs}$ . The effects of log(WBC) on decreased thrombocyte counts were only seen in children 12 years and older.

Our results, supported by a path analysis, suggest that internal exposure of  $^{137}\text{Cs}$  affects blood counts, and that  $^{137}\text{Cs}$  in soil has only indirect effects on the blood counts via internal exposure to  $^{137}\text{Cs}$  as an intermediate variable, but no direct effect due to external exposure was detected. However this finding should be interpreted with caution and cannot be fully attributed to internal dose being more important than external exposure, since it may also be influenced by measurement errors of the external exposure. The external exposure was assessed on average village level, while internal  $^{137}\text{Cs}$  was measured on an individual level, i.e. a larger measurement error can be expected to dilute the effects of external exposure. A strength was; however, that there was a high variability in the level of  $^{137}\text{Cs}$  soil contamination between the villages.

The level of external exposure has been decreasing for each year since the accident, but the ranking of which the population in the villages receives high or low external exposure is identical since 1992 (4). The average soil contamination for the village of a child can thus be used as a proxy of both cumulative and current exposure of that child. The population is stable

and rarely relocates to places with different external exposure levels, which further strengthens the validity as a proxy for long-term exposure.

Internal exposure to  $^{137}\text{Cs}$  measured by WBC is a point measurement primarily reflecting current intake, but it may also have long-term predictive value if dietary habits and levels of dust inhalation are stable. The very highest  $^{137}\text{Cs}$ -whole body concentrations are likely to result from very recent intakes of contaminated food. The fact that we adjusted for season and also took the logarithm of WBC, shrinks the values of the peak whole body counts and probably makes the counts more representative of average long-term intake.

The cross-sectional nature of the study, however, was a limitation. To study dose-effect relationships, one needs to have information about the total cumulative dose for the entire period of residence in the Narodichesky region.

Since WBC reflects dietary intake, the most likely potential confounding factors would be nutritional deficiencies, or environmental contaminants correlated with  $^{137}\text{Cs}$ . There may be a possibility of environmental confounding or partial effects due to other long-lived radionuclides released from the Chernobyl accident, such as  $^{90}\text{Sr}$ ,  $^{239}\text{Pu}$  and  $^{241}\text{Am}$ . These radionuclides were not as widely geographically distributed as  $^{137}\text{Cs}$ , but mainly deposited within 100km of the Chernobyl reactor, due to dispersal with larger particle size (3). However, these radionuclides have a relatively high deposition in Zhitomir Oblast (1, 39), but have not been measured to the same extent as  $^{137}\text{Cs}$ . They are expected to make a small contribution to the total radiation dose in the population, but they are prone to bone deposition and are also accumulated in the bone due to very long physical and biological half-life (12). Especially Plutonium and Americium are concentrated in the endosteum (12), which is also where stem cell production in the bone marrow is concentrated (40). In addition, a large amount of lead (Pb) was dumped on the burning Chernobyl reactor to extinguish the

fire, and since lead is a known cause of anemia the possibility of confounding due to lead has to be considered. However, a study conducted in areas 30-100km from Chernobyl, found levels of lead in soil to be low, between 2-27 mg/kg, which equals background values in these soil types (41).

The present study supports the previous finding that children living in villages with high level of  $^{137}\text{Cs}$  in soil had decreased levels of hemoglobin, red blood cells and thrombocytes (2). The previous study showed that thrombocytes were less affected in later years which is also in agreement with our current finding that thrombocytes are only affected by WBC in older children. A previous study by Babeshko et al, did not find any difference in blood cells count depending on village level of  $^{137}\text{Cs}$  in soil, but this may depend on the simple comparison of means between the two groups of exposure, without adjustment for other factors (19).

Previous studies in Zhitomir Oblast have found morphologic changes in red blood cells in children irradiated in utero and living in the exposed areas (22). This is not directly comparable since our children are born at least a few years after the accident, and thus been exposed to lower doses of radiation.

Studies in Kyrgyzstan have found decreased Hb and erythrocytes in children and adults living in close proximity to radioactive waste dumps, compared to control groups (42,43). However, the exact levels of radioactive exposure for these individuals were not presented.

The current total doses from the Chernobyl accident in Zhitomir Oblast have been estimated to be low, on average 0.13-2.2 mSv/year in 2008 (4), and this agreed well with the doses observed in our study. For some children, the levels were above the  $1\text{mSv y}^{-1}$ , which is the reference level as recommended by the ICRP for population exposure from a non-natural source (44), but well below the  $20\text{mSv y}^{-1}$  which is the ICRP reference level for occupational exposure.

After the Techa river incident, in which people were exposed to accumulated bone marrow doses of up to 3-4Sv, hematopoiesis recovery was seen after reduction in red bone marrow dose rates to 100 mSv/year or lower (16). However, the dose reconstruction is retrospective in these studies, and the high uncertainty in the dose estimates can be expected to have diminished the ability to find effects from lower doses.

The ability to find effects of low-level exposure in epidemiological studies is highly dependent on the precision of the exposure assessment. WHO concluded that there is a complete lack of analytical studies of long-term effects from Chernobyl (23). This is the only study on long-term effects from  $^{137}\text{Cs}$  on blood cells, which has used individual exposure assessment. Our findings suggest that the radiation exposure from fallout after the Chernobyl accident may still affect children in the contaminated areas. However, possible confounding from other radionuclides or heavy metals released from Chernobyl, cannot be excluded and should be considered in future studies.

Although an association between low erythrocytes/Hb/thrombocytes and high  $^{137}\text{Cs}$  was detected, the blood counts were only mildly decreased. No major symptoms should be expected of these subnormal blood counts observed in the children, except possible tiredness in the children with most apparent anemia. However, even if mildly decreased blood counts do not markedly affect a healthy child under normal circumstances, the resources to cope with diseases or hard physical exercise may be diminished, especially if the subnormal blood counts reflect underlying disease with diminished cell production capacity.

Children in this study were born at least four years after the accident and thus exposed to low doses of ionizing radiation from  $^{137}\text{Cs}$ . This cross-sectional study indicates that these levels may be associated with decreased blood counts, but we cannot exclude that our results are due

to residual confounding factors. Since WBC reflects dietary intake, the most likely potential confounding factors could be nutritional deficiencies, or environmental factors causing anemia/decreased blood count. Future studies are needed to investigate the possible role of other radionuclides which may be correlated with  $^{137}\text{Cs}$ , and possibly other heavy metals, in addition to  $^{137}\text{Cs}$ .

### **Acknowledgements**

Anna Lindgren received a scholarship from the Swedish Council for Working Life and Social Research (FAS) (2011-1769), for conducting research at Arnold School of Public Health. Financial support for the health screenings was provided by the National Academy of Medical Sciences of Ukraine, Kyiv, Ukraine. This research was partially supported by a grant from the United States Civilian Research and Development Foundation (UKB1-2929-KV-08).

### **Competing interests**

The authors have no competing interests.

### **Supplementary information**

Supplementary information is available at Journal of Exposure Science & Environmental Epidemiology's website

**Supplemental File S1.** Phantoms used for calibration of the whole body counter. Word-document.

**Supplemental File S2.** Path analysis. Code and graphs for the path analysis testing the assumption of  $^{137}\text{Cs}$  in the body being an intermediate variable between  $^{137}\text{Cs}$  in soil and low blood counts. Word-document.

Table I. General characteristics, WBC “SCRINNER-3M.

Multichannel Pulse Analyzer (MCA)	ADC (1024 ch), interface serial RS-232
Detector	NaI(Tl) Ø150x100 mm
Collimator	Pb, thickness 50 mm
Measured radioisotope	<sup>137</sup> Cs
Registration energy, keV	662
MCA step, keV/Ch	2
Energy resolution, %	10.2
Registration efficiency for 70kg adult, Bq/cpm	6,5
Background attenuation factor for 70kg adult	0.77
Background counts, cpm	2200
Minimum detection activity (MDA, sensitivity level for 3 min exposition of 70 kg adult), Bq	340

Table II. Ukrainian reference levels for Hb, RBC and PLT, and prevalence of children with current values below reference levels.

	Reference level	(n below)/(total n )	% below
Hb <sup>1</sup> (g/L)	0-4 years $\geq 110$	1/8	12.5%
	5-7 years $\geq 115$	5/25	20%
	8 and older $\geq 120$	117/533	22%
Total		123/566	21.7%
RBC <sup>2</sup> ( $\times 10^{12}$ cells/L)	2-5 years $\geq 3.9$	4/16	25%
	6-11 years $\geq 4.0$	26/134	19.4 %
	12 to 17 years $\geq 4.5$	340/416	81.7%
Total		370/566	65.4%
PLT <sup>3</sup> ( $\times 10^9$ cells/L)	All ages $\geq 150$	4/566	0.7%

<sup>1</sup> Shabalov, N.P., 2001 (35).

<sup>2</sup> Nicholson JF, Pesce M, 2000 (36).

<sup>3</sup> Stepanova, E et al, 2008 (2).



Table III. Description of current  $^{137}\text{Cs}$  exposure level for villages included in the study. Official residential soil activity and external doses of  $^{137}\text{Cs}$  for each village, and measured individual whole body concentration and internal dose of  $^{137}\text{Cs}$  (n=566).

		External exposure (official estimates for year 2008)			Internal exposure (point-measures, 2008-2010)		
		Official average surface density of $^{137}\text{Cs}$ in soil <sup>1</sup>	Official average external effective dose <sup>1</sup>	$^{137}\text{Cs}$ WBC, measured in this study Mean	Internal effective dose calculated in this study Mean Range <sup>2</sup>		
Village	N	%	(kBq/m <sup>2</sup> )	(mSv/year)	$^{137}\text{Cs}$ (Bq/kg)	(mSv/year)	
Jitomir <sup>3</sup>	1	0.2	-	-	18	0.03	-
Kiev <sup>3</sup>	1	0.2	-	-	17	0.02	-
Brodnik	16	2.8	41	0.11	27	0.03	(0.02-0.06)
Liplanchina	1	0.2	53	0.15	32	0.06	-
Vyasovka	8	1.4	54	0.15	31	0.04	(0.01-0.06)
Rubejevka	2	0.4	55	0.15	47	0.02	(0.02-0.03)
Radcha	3	0.5	57	0.16	33	0.04	(0.03-0.04)
Budo-Golubevi	2	0.4	58	0.16	131	0.15	(0.09-0.20)
N. Radcha	2	0.4	72	0.20	24	0.04	(0.04-0.05)
Klochki	2	0.4	73	0.20	16	0.04	(0.03-0.04)
Norinci	15	2.7	77	0.21	26	0.04	(0.02-0.09)
Guto-Mariatyn	6	1.1	80	0.22	61	0.07	(0.02-0.14)
Babinichi	10	1.8	87	0.24	68	0.06	(0.02-0.10)
Laski	19	3.4	88	0.24	15	0.03	(0.005-0.05)
Jerev	14	2.5	90	0.25	44	0.06	(0.03-0.08)
Bolotnica	17	3	91	0.25	30	0.05	(0.01-0.21)
Latashi	5	0.9	92	0.26	16	0.03	(0.02-0.05)
S.Dorogin	4	0.7	114	0.32	30	0.06	(0.02-0.15)
Zakusily	33	5.8	114	0.31	72	0.09	(0.03-0.22)
N. Dorogin	2	0.4	117	0.32	20	0.04	(0.03-0.04)
Zalesie	2	0.4	137	0.38	91	0.13	(0.07-0.18)
Suharevka	15	2.7	145	0.40	95	0.11	(0.04-0.21)
Moteyki	1	0.2	155	0.43	201	0.38	-
Megeliska	28	4.9	163	0.45	125	0.12	(0.03-0.25)
Selec	23	4.1	214	0.59	135	0.23	(0.03-1.36)
Narodichi	284	50.2	242	0.67	123	0.17	(0.009-1.41)
Basar	49	8.7	252	0.70	138	0.16	(0.01-0.34)
Lubarka	1	0.2	274	0.76	139	0.27	-

<sup>1</sup>Ministry of Health of the Ukraine (2009). Certification doses for the settlements of Ukraine located in contaminated territories after the Chernobyl accident. Data for 2008. Report #13. Kiev.

<sup>2</sup>Range (Min, Max), is not presented for the villages where only one child was measured.

<sup>3</sup>There are no official estimates of the external effective dose from  $^{137}\text{Cs}$  in Jitomir and Kiev, since these areas are considered non-contaminated.

Table IV. Characteristics of the study population (n=566), and distribution of age, exposure and outcome variables.

		N	%
Sex	Female	266	47
	Male	300	53
Age group (at time of examination)	<6	16	2.8
	6-11	134	23.7
	12-18	416	73.5
Birth year	2002-2007	29	5
	1996-2001	230	40.6
	1990-1995	307	54.2
Environmental tobacco smoke (ETS)	No	313	55
	Yes	253	45
Active daily tobacco smoking, (cigarettes/day)	No	538	95
	1-10	23	4
	>10	5	0.9
Use of coal/wood for cooking	No	403	71
	Yes	163	29
Activity of <sup>137</sup> Cs in residential soil (kBq/m <sup>2</sup> )	<116	161	28.4
	116-164	48	8.4
	>165	357	63.1
	Mean	Median	Range
Age (years)	12.9	13	2-18
Weight (kg)	47.4	48	12.5-99.7
Activity of <sup>137</sup> Cs in residential soil (kBq/m <sup>2</sup> )	189.4	242	10-274
Activity of <sup>137</sup> Cs in the body (Bq)	4447	3404	185-42480
WBC <sup>137</sup> Cs (Bq/kg)	100.5	77.1	4-916
Log (WBC)	4.2	4.3	1.4-6.8
Effective internal dose (mSv/year)	0.14	0.10	0.005-1.41
Hemoglobin (g/L)	125.5	124	96-155
Erythrocyte cell count (RBC) x 10 <sup>12</sup> cells/L	4.2	4.1	3.1-5.2
Thrombocyte count x 10 <sup>9</sup> platelets/L	250.5	249	138-422

Table V. Linear relationship between the whole body concentration (WBC) of  $^{137}\text{Cs}$  (Bq/kg) and the activity of  $^{137}\text{Cs}$  in residential soil ( $\text{kBq/m}^2$ ) and blood counts (n=566).

	Estimate	SE	p-value
<u>Hemoglobin (g/L)</u>			
Log(WBC) <sup>1</sup>	-5.940	0.411	<0.001***
Cs in soil ( $\text{kBq/m}^2$ ) <sup>1</sup>	-0.00241	0.00621	0.6978
<u>Red Blood Cells (<math>\times 10^{12}</math> cells/L)</u>			
Log(WBC) <sup>1</sup>	-0.238	0.0148	<0.001***
Cs in soil ( $\text{kBq/m}^2$ ) <sup>1</sup>	0.000548	0.000224	0.0149*
<u>Platelets (<math>\times 10^9</math> plt/L)</u>			
<i>Age &lt; 12</i>			
Log(WBC) <sup>2</sup>	-4.7479	5.348	0.3762
Cs in soil ( $\text{kBq/m}^2$ ) <sup>2</sup>	-0.08901	0.06682	0.1850
<i>Age <math>\geq 12</math></i>			
Log(WBC) <sup>1</sup>	-15.99	2.735	<0.001***
Cs in soil ( $\text{kBq/m}^2$ ) <sup>1</sup>	-0.052	0.0446	0.24387

<sup>1</sup> Adjusted for Cs in soil ( $\text{kBq/m}^2$ ), log(Cs Bq/kg), age, sex, exam season, ETS, active tobacco smoking, use of coal/wood for cooking.

<sup>2</sup> Not adjusted for active smoking, since no children below 12 were smokers

## REFERENCES

1. Ministry of Ukraine of Emergencies and Affairs of population protection from the Consequences of Chernobyl Catastrophe. 20 years after Chernobyl Catastrophe. Future Outlook. National Report of Ukraine, 2006. [in English] Available at: [http://chernobyl.undp.org/english/docs/ukr\\_report\\_2006.pdf](http://chernobyl.undp.org/english/docs/ukr_report_2006.pdf) Accessed 05/31/2013.
2. Stepanova E, Karmaus W, Naboka M, Vdovenko V, Mousseau T, Shestopalov VM, et al. Exposure from the Chernobyl accident had adverse effects on erythrocytes, leukocytes, and platelets in children in the Narodichesky region, Ukraine: a 6-year follow-up study. *Environmental health*. 2008;7:21.
3. IAEA. Chernobyl's Legacy: Health, Environmental and Socio-economic Impacts and Recommendations to the Governments of Belarus, the Russian Federation and Ukraine. Austria, 2006.
4. Ministry of Health of the Ukraine. Certification doses for the settlements of Ukraine located in contaminated territories after the Chernobyl accident. Data for 2008. Report no 13. Kyiv, 2009. [in Ukrainian]
5. Korzun VN KL, Stepanova E.I, Torbin VF. Ionizing radiation and nutrition in children. *Chernobilinterinform*: Kyiv, Ukraine, 1997.
6. Travnikova IG, Bazjukin AN, Bruk GJ, Shutov VN, Balonov MI, Skuterud L, et al. Lake fish as the main contributor of internal dose to lakeshore residents in the Chernobyl contaminated area. *Journal of environmental radioactivity*. 2004;77(1):63-75.

7. Ministry of Ukraine of Emergencies and Affairs of Population Protection from the Consequences of Chernobyl Catastrophe. Radiological status of the territories assigned to zones of radioactive contamination (by districts). Kyiv, 2008. [in Ukrainian]

8. Ministry of Ukraine of Emergencies. 25 years after the Chernobyl accident. Safety for the Future. National Report of Ukraine. Kyiv, 2011. [in English] Available at: [http://www.kavlinge.se/download/18.2b99484f12f775c8dae80001245/25\\_Chornobyl\\_angl.pdf](http://www.kavlinge.se/download/18.2b99484f12f775c8dae80001245/25_Chornobyl_angl.pdf) Accessed 05/31/2013.

9. Sharifov VF, Koulikova NV, Voropai LV, Kroupnik TA, Masyakin VB, Cot VA, et al. Finding of the Chernobyl Sasakawa Health and Medical Cooperation Project: 137 Cs concentrations among children around Chernobyl. In: Yamashita S, Shibata Y, editor. Chernobyl: A Decade Proceedings of the Fifth Chernobyl Sasakawa Medical Cooperation Symposium, Kiev, Ukraine, 14–15 October 1996. Amsterdam, Lausanne, New York: Elsevier; 1997. pp. 39–44.

10. Zvonova IA, Jesko TV, Balonov MI, Danilova JO, Wallström E, Alpsten M, et al. 134Cs and 137Cs whole-body measurements and internal dosimetry of the population living in areas contaminated by radioactivity after the chernobyl accident. Radiation protection dosimetry. 1995;62(4):213-21.

11. Bernhardsson C, Zvonova I, Rääf C, Mattson S. Measurements of long-term external and internal radiation exposure of inhabitants of some villages of the Bryansk region of Russia after the Chernobyl accident. Science of the Total Environment. 2011(409):4811-7.

12. ICRP. Limits for intakes of radionuclides by workers. Publication No 30. Annals of the ICRP. 1979; 2(3/4).
13. Mettler FA, Jr., Gus'kova AK, Gusev I. Health effects in those with acute radiation sickness from the Chernobyl accident. Health physics. 2007;93(5):462-9.
14. Baverstock KF, Ash PJ. A review of radiation accidents involving whole body exposure and the relevance to the LD50/60 for man. The British journal of radiology. 1983;56(671):837-44.
15. Akleyev AV, Kossenko MM, Silkina LA, Degteva MO, Yachmenyov VA, Awa A, et al. Health effects of radiation incidents in the southern Urals. Stem cells. 1995;13 Suppl 1:58-68.
16. Akleyev AV. Chronic radiation syndrome among residents of the Techa riverside villages. Radiation protection dosimetry. 2012;151(4):689-95.
17. Fenech M, Perepetskaya G, Mikhalevich L. A more comprehensive application of the micronucleus technique for biomonitoring of genetic damage rates in human populations--experiences from the Chernobyl catastrophe. Environmental and molecular mutagenesis. 1997;30(2):112-8.
18. Livingston GK, Jensen RH, Silberstein EB, Hinnefeld JD, Pratt G, Bigbee WL, et al. Radiobiological evaluation of immigrants from the vicinity of Chernobyl. International journal of radiation biology. 1997;72(6):703-13.

19. Babeshko VG, Bruslova EM, Tsvyetskova NM, Dzurynska EN, Mochanova NJ, Talko VV et al. Hemopoietic system. In: Babeshko V, Voizianov A, Bazyka D (eds). Health effects of Chernobyl accident. Dia Ltd: Kyiv, Ukraine, 2003, pp. 412-26.
20. Vartanian LS, Gurevich S, Kozachenko AI, Nagler LG, Burlakova EB. [Age-related peculiarities of effect of low dose ionizing radiation on blood antioxidant enzyme system status in Chernobyl's accident liquidation participant]. [Advances in gerontology] Gerontologicheskoe obshchestvo. 2004;14:48-54.
21. Spirichev VB, Donchenko GV, Blazheevich NV, Parkhomenko Iu M, Aleinik SI, Golubkina NA, et al. [To the 20th anniversary of the Chornobyl accident study of vitamin status and provision with micro- and macroelements of limited groups of people at different time periods since the accident at Chornobyl nuclear power plant]. Ukrainskii biokhimicheskii zhurnal. 2006;78(2):5-26.
22. Stepanova EI, Davidenko OA, Vdovenko V. [Peculiarities of superficial architectonics of peripheral blood erythrocytes in children irradiated in utero and living in the radioactive-contaminated areas]. Tsitologiya i genetika. 2006;40(6):40-3.
23. WHO. Health Effects of the Chernobyl Accident and Special Health Care Programmes. Geneva: 2006.

24. Svendsen ER, Kolpakov IE, Stepanova YI, Vdovenko VY, Naboka MV, Mousseau TA, et al. <sup>137</sup>Cesium exposure and spirometry measures in Ukrainian children affected by the Chernobyl nuclear incident. *Environmental health perspectives*. 2010;118(5):720-5.

25. The Ukrainian Government. Standards of nutrition and partial compensation of nutritional expenses of children affected by Chernobyl accident [Internet]. Available at: <http://zakon4.rada.gov.ua/laws/show/258-92-%D0%BF> Accessed 05/31/2013.

26. Perevoznikov O.M., Vasylenko V.V., Litvinetz, L.A., Yakovleva, G.N. 2009. Mass many years WBC-monitoring of Ukrainian inhabitants involved Chernobyl NPP accident. *Radiation Hygiene* Volume 2, Issue 2, 2009, pp.40-47. (Abstract in English);

Available at:

[http://radhyg.ru/index.php?option=com\\_content&view=article&id=127&catid=47&Itemid=59&lang=ru](http://radhyg.ru/index.php?option=com_content&view=article&id=127&catid=47&Itemid=59&lang=ru) Accessed 5/31/2013.

27. The Institute of Radiation Safety "BELRAD". [web page]. Available at <http://www.belrad-institute.org/> Accessed 06/01/2013.

28. IAEA - USSR Whole Body Counter Intercomparison/R. V. Griffith, R. Ouvrard, V. Bad'in, S. Ulanovsky, A. Kovtun, L. Litvinets, F. Steger// *Proceedings of a sympos., Monreal, 20 - 25 May, 1992*. Available at: [http://www.irpa.net/irpa8/cdrom/VOL.1/M1\\_254.PDF](http://www.irpa.net/irpa8/cdrom/VOL.1/M1_254.PDF) Accessed 06/01/2013.



29. SCRM AMS of Ukraine. Recommendations on assessment of internal dose of irradiation from radiocesium using whole body counters. Kiev, 1994, 24p.[in Russian]

30. SCRM AMS of Ukraine. Recommendations on measurements with whole body counters during the dosimetric pasportization of residential areas of Ukraine. Kiev, 1996, 73p.[in Russian].

31. Recommendations for assessment of dose of internal irradiation from radioactive cesium in residents using whole body counters. Scientific Center of Radiation Medicine Academy of Medical Sciences of Ukraine. Kiev, 1994, p.24 [in Russian].

32. Hoshi M, Konstantinov YO, Evdeeva TY, Kovalev AI, Aksenov AS, Koulikova NV et al. Radiocesium in children residing in the western districts of the Bryansk Oblast from 1991-1996. Health physics. 2000;79:182-6.

33. Hayashida N, Sekitani Y, Kozlovsky A, Rafalsky R, Gutevich A, Daniliuk V, et al. Screening for  $^{137}\text{Cs}$  body burden due to the Chernobyl accident in Korosten City, Zhitomir, Ukraine: 1996-2008. J Radiat Res. 2011;52(5):629-33.

34. ICRP. Age-dependent Doses to the Members of the Public from Intake of Radionuclides Part 5, Compilation of Ingestion and Inhalation Coefficients. ICRP publication 72. Annals of the ICRP. 1995; 26(1).

35. Thöml H, Diem H, Haferlach T. Color Atlas of Hematology. Practical Microscopic and Clinical Diagnosis. Thieme New York: New York, USA, 2004.
36. Shabalov NP. Diseases of children. 4th ed. Peter: Saint-Petersburg, Russia, 2001.
37. Nicholson JF, Pesce M. Reference Ranges for Laboratory Tests and Procedures. In: Behrman RE, Kliegman RM, Jenson HB (eds). Nelson textbook of pediatrics 16th ed. W.B. Saunders Company : Philadelphia, USA, 2000.
38. Hatcher L. A Step-by-Step Approach to Using SAS for Factor Analysis and Structural Equation Modeling. SAS Institute Inc: Cary, NC, USA, 1994.
39. Chernobyl Ministry of Ukraine of Emergencies and Affairs of Population Protection from the consequences of Chernobyl Catastrophe. Atlas. Ukraine Radioactive contamination. Kyiv, Ukraine, 2008.
40. Fliedner TM, Graessle D, Paulsen C, Reimers K. Structure and function of bone marrow hemopoiesis: mechanisms of response to ionizing radiation exposure. Cancer biotherapy & radiopharmaceuticals. 2002;17(4):405-26.
41. Stepanova Ye A, Korobova, Ye M, Orlov DS, Petrovskaya, I V. Lead in soils of the remote zone of the Chernobyl APS accident effect. Pochvovedenie. 1990;10:61-9.
42. Mamajakyp uuly C, Kamchibeckov, A.B, Magazova C.M., Nazarbeckov, D.K., Nartaeva, A.K., Mamatov, C.M. 2006. Changes in the periphery blood cells at people living in a region

of burial places of radioactive waste. [In Russian, Abstract in English]. News of Russian Scientific Center of Rentgenradiology. Issue 6. Available at:

[http://vestnik.rncrr.ru/vestnik/v6/papers/mam\\_v6.htm](http://vestnik.rncrr.ru/vestnik/v6/papers/mam_v6.htm) Accessed 05/31/2013.

43. Kamchibeckov, A.B, Magazova C.M., Nazarbeckov, D.K., Mamajakyp uuly C, Nartaeva, A.K., Mamatov, C.M. 2006. Study of hematopoiesis and immune status of children residing in and outside of the area of tailing pits. News of Russian Scientific Center of Rentgenradiology. Issue 6. [in Russian, abstract in English]. Available at:

[http://vestnik.rncrr.ru/vestnik/v6/papers/kamch\\_v6.htm](http://vestnik.rncrr.ru/vestnik/v6/papers/kamch_v6.htm) Accessed 05/31/2013.

44. ICRP. The 2007 Recommendations of the International Commission on Radiological Protection. ICRP Publication 103. Annals of the ICRP. 2007; 37 (2-4).




Fig 1. Exposure distributions of a)  $^{137}\text{Cs}$  Whole body concentration (WBC, Bq/kg), b)  $^{137}\text{Cs}$  in Soil(kBq/m<sup>2</sup>), and c) the correlation between  $^{137}\text{Cs}$  in soil (x-axis) and  $^{137}\text{Cs}$  WBC (y-axis). (n=566).

Figure 2.




Fig 2. Whole body concentration (WBC) by season. Boxplots displaying the exposure distributions of  $^{137}\text{Cs}$  whole body concentration (Bq/kg), depending on season of the measurement (n=566). The cross refers to the mean WBC.

Figure 3.




Fig 3. Individual Whole Body Concentration (WBC) of  $^{137}\text{Cs}$  (Bq/kg) and  $\log(\text{WBC})$ , plotted against individual blood counts: hemoglobin (Hb) g/L, erythrocyte count (RBC)  $\times 10^{12}$  cells/L, and thrombocyte count (PLT)  $\times 10^9$  platelets/L, (n=566).

Figure 4.




Fig 4. Residential activity of <sup>137</sup>Cs in soil (kBq/m<sup>2</sup>), plotted against individual blood counts: a) hemoglobin (Hb) g/L, b) erythrocyte count (RBC) x10<sup>12</sup> cells/L, and c) thrombocyte count (PLT)x10<sup>9</sup> platelets/L ( n=566 ).