



LUND UNIVERSITY

The Development of Severe Neonatal Alloimmune Thrombocytopenia due to Anti-HPA-1a Antibodies Is Correlated to Maternal ABO Genotypes

Ahlen, Maria Therese; Husebekk, Anne; Killie, Mette Kjaer; Kjeldsen-Kragh, Jens; Olsson, Martin L; Skogen, Bjorn

Published in:
Clinical & Developmental Immunology

DOI:
[10.1155/2012/156867](https://doi.org/10.1155/2012/156867)

2012

[Link to publication](#)

Citation for published version (APA):

Ahlen, M. T., Husebekk, A., Killie, M. K., Kjeldsen-Kragh, J., Olsson, M. L., & Skogen, B. (2012). The Development of Severe Neonatal Alloimmune Thrombocytopenia due to Anti-HPA-1a Antibodies Is Correlated to Maternal ABO Genotypes. *Clinical & Developmental Immunology*, Article 156867. <https://doi.org/10.1155/2012/156867>

Total number of authors:
6

General rights

Unless other specific re-use rights are stated the following general rights apply:
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

Research Article

The Development of Severe Neonatal Alloimmune Thrombocytopenia due to Anti-HPA-1a Antibodies Is Correlated to Maternal ABO Genotypes

Maria Therese Ahlen,^{1,2} Anne Husebekk,^{1,2} Mette Kjær Killie,¹
Jens Kjeldsen-Kragh,^{3,4} Martin L. Olsson,⁵ and Bjørn Skogen^{1,2}

¹ Department of Laboratory Medicine, University Hospital of North Norway, 9038 Tromsø, Norway

² Department of Immunology, Institute of Medical Biology, University of Tromsø, 9037 Tromsø, Norway

³ Department of Immunology and Transfusion Medicine, Oslo University Hospital, Ullevål, 0407 Oslo, Norway

⁴ Faculty Division Ullevål University Hospital, University of Oslo, 0407 Oslo, Norway

⁵ Division of Hematology and Transfusion Medicine, Department of Laboratory Medicine, Lund University, SE-221 00 Lund, Sweden

Correspondence should be addressed to Maria Therese Ahlen, therese.ahlen@unn.no

Received 28 June 2011; Revised 16 August 2011; Accepted 16 September 2011

Academic Editor: Raivo Uibo

Copyright © 2012 Maria Therese Ahlen et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Background. Maternal alloantibodies against HPA-1a can cross placenta, opsonize foetal platelets, and induce neonatal alloimmune thrombocytopenia (NAIT). In a study of 100, 448 pregnant women in Norway during 1995–2004, 10.6% of HPA-1a negative women had detectable anti-HPA-1a antibodies. **Design and Methods.** A possible correlation between the maternal ABO blood group phenotype, or underlying genotype, and severe thrombocytopenia in the newborn was investigated. **Results.** We observed that immunized women with blood group O had a lower risk of having a child with severe NAIT than women with group A; 20% with blood group O gave birth to children with severe NAIT, compared to 47% among the blood group A mothers (relative risk 0.43; 95% CI 0.25–0.75). **Conclusion.** The risk of severe neonatal alloimmune thrombocytopenia due to anti-HPA-1a antibodies is correlated to maternal ABO types, and this study indicates that the observation is due to genetic properties on the maternal side.

1. Introduction

Foetal-maternal incompatibility in the human platelet antigen (HPA)-1 alloantigen system is the most common underlying cause of neonatal alloimmune thrombocytopenia (NAIT), a condition where maternal alloantibodies opsonize foetal platelets during pregnancy and reduce their survival in circulation. The incompatibility is based on a single-nucleotide polymorphism (SNP) which results in a leucine/proline substitution at residue 33 in the $\beta 3$ integrin that constitutes membrane glycoprotein $\beta 3$ [GPIIIa] present on platelets in complex with α Ib integrin [GPIIb] [1]. On platelets, the α Ib $\beta 3$ [GPIIb/IIIa] is also the major carrier of blood group A antigen [2].

About 10% of HPA-1a negative women who have been pregnant with an HPA-1-incompatible child have detectable

HPA-1a antibodies [3]. In several studies, a correlation between maternal antibody level and the severity of thrombocytopenia in the newborn has been shown [4–6]. The alloimmunization is strongly associated with the *HLA-DRB3*01:01* allele [3, 7, 8]; however, only about 30% of the women with this HLA antigen are immunized. Except for the incompatibility in platelet antigen and the association to HLA, other factors which may influence the immune response to HPA-1a have not been identified.

In the present study, we have examined the maternal ABO blood groups and frequency of HPA-1a-immunization of the women identified in the large prospective screening and intervention study carried out in Norway from 1995 to 2004. We included 152 HPA-1a-immunized women, 146 of whom had altogether 158 HPA-1-incompatible pregnancies in the screening study. The ABO distribution among

immunized women was investigated, and the maternal ABO phenotype and *ABO* genotype was correlated to the severity of thrombocytopenia of the newborn.

2. Materials and Methods

2.1. Patients. Pregnant women were recruited for HPA-1 allotyping from three regions in Norway between December, 1995 and March, 2004 [3]. Samples for routine Rh(D) typing were also used for determining HPA-1 allotype by flow cytometry (anti-CD61 mAb), enzyme-linked immunosorbent assay (ELISA), or polymerase chain reaction (PCR) as previously described [9]. A total of 100,448 pregnant women were typed for the platelet antigen HPA-1a, and 2,111 of those were HPA-1a negative (2.1%). Of these, 1,990 were further tested, and anti-HPA-1a antibodies were detected in 154 women during the pregnancy. In total, 146 of these immunized women underwent 158 HPA-1a-incompatible pregnancies. ABO blood group typing was performed by conventional technique. Genomic typing of HPA-1 (*ITGB3*; rs5918 in dbSNP) and *ABO* in the neonates was performed in samples from cord blood or buccal swabs. For the newborns, the *ABO* genotype was used to predict the ABO blood group. In this context, we have defined ABO incompatibility only as an A₁ phenotype in the newborn, in blood group O mothers, because individuals with A₂, and the majority of individuals with B phenotype, express only low levels of corresponding antigens on the surface of platelets [2, 10–12]. Thrombocytopenia was defined as a platelet count $\leq 150 \times 10^9/L$, and severe thrombocytopenia less than $50 \times 10^9/L$ measured in cord blood and/or capillary blood at birth. Detection of anti-HPA-1a IgG antibodies was performed by flow cytometry and quantified with monoclonal antibody immobilization of platelet antigen assay (MAIPA) [3], by using the anti-CD61 monoclonal antibody clone Y2/51 (Dako, Glostrup, Denmark) for immobilisation of platelet glycoproteins. Women were tested at several time points during the pregnancy, and those with a positive antibody test at any time during the pregnancy were characterized as immunized. Nineteen women were primary immunized during the studied pregnancy, 13 of these were primigravida. All others may have been immunized in connection with a prior pregnancy. Prior affected pregnancies were not excluded as a cause of severe NAIT. The NAIT diagnosis was based on maternal anti-HPA-1a antibodies and HPA-1a antigen incompatibility. Other possible reasons for thrombocytopenia (infection, maternal ITP, etc.) were not registered. Informed consent was provided in accordance with the declaration of Helsinki. The study was approved by the Regional Committee for Medical Research Ethics, North Norway (approval no. P-REK V 13/1995).

2.2. ABO Genotyping. ABO genotyping was performed by PCR-RFLP analysis to detect six major alleles, A¹, A², B, O¹/O^{1v}, and O² (also known as A101/A201/B101/O01/O02/O03) according to the nomenclature used by the Blood Group Antigen Gene Mutation Database, dbRBC [13], and further discrimination between the common O¹ and O^{1v} alleles [O01/O02] was performed using primers and reaction

conditions as described by Olsson and Chester [14, 15] with some modifications: HotStarTaq polymerase 5 U/ μ L (QIAGEN, Hilden, Germany) was used with the following cycling programs for both analyses: 95°C 15 min, 10 cycles of 94°C for 10 seconds, 63°C for 30 seconds and 72°C for 30 seconds followed by 25 cycles of 94°C for 10 seconds, 61°C for 30 seconds and 72°C for 30 seconds for samples with 100 ng DNA template. For samples with ~ 25 ng template and < 10 ng DNA template, 1 and 2 extra cycles at each of the annealing temperatures was performed, respectively. Subsequently, digestion with endonucleases and qualitative analyses of product were performed [14, 15]. Ambiguous results were confirmed with selected primer sets from a recently published PCR-ASP method for ABO genotyping [16]. For simplicity, only one terminology, dbRBC, will be used throughout this paper. Genotypes are written as X/X.

2.3. Statistical Analysis. Standard statistical calculations as mean, relative risk, Chi-square test, analysis of variance with Bonferroni's test as post hoc analysis, and plots were performed with computer software SPSS for windows (Statistical Package for the Social Sciences, Version 16.0 SPSS Inc., Chicago, Ill, USA). $P < 0.05$ was considered significant.

3. Results

3.1. The ABO Phenotype Distribution of the Immunized Mothers. The ABO phenotype distribution among 154 HPA-1a immunized women was similar to the distribution of the general Norwegian population [17] adjusted to statistics for 2005 (data not shown), indicating that the maternal ABO type does not influence the risk of HPA-1a immunization.

3.2. Maternal ABO Blood Group and Risk of Severe NAIT. In 158 HPA-1-incompatible pregnancies with 83 cases of NAIT, there were 54 cases of severe NAIT. The maternal ABO phenotypes were compared to the platelet count in their neonates (Table 1); 46.6% of the immunized women with blood group A gave birth to children with severe NAIT, compared to 20.0% among the immunized mothers with blood group O. The relative risk of NAIT (platelet count $\leq 150 \times 10^9/L$) in the neonates of HPA-1a immunized women with blood group O as compared to blood group A was 0.67 (95% CI 0.48–0.94), whereas the relative risk of severe NAIT (platelet count $< 50 \times 10^9/L$) in the neonates of HPA-1a immunized women with blood group O was 0.43 (95% CI 0.25–0.75) as compared to the neonates of women with blood group A. However, the frequency of moderate NAIT (platelet count $50\text{--}150 \times 10^9/L$) was not lower among the blood group O mothers. 91% of the immunized women carried the *HLA-DRB3*01:01* allele. There were no cases of severe NAIT among *HLA-DRB3*01:01*-negative mothers.

3.3. ABO Incompatibilities between Mothers and Newborns. To investigate whether the ABO incompatibility between mother and foetus could explain the difference in the frequency of severe NAIT, newborns were ABO genotyped as basis for the prediction of their ABO phenotype. One hundred and thirty out of 146 newborns were ABO genotyped

TABLE 1: The maternal ABO type distribution in the pregnancies compared to the severity of NAIT.

Maternal ABO type	Numbers of newborns with platelet count <math>< 50 \times 10^9/L</math> (% within ABO type; 95% CI)	Numbers of newborns with platelet count 50–150 $\times 10^9/L$	Numbers of newborns with platelet count >150 $\times 10^9/L$	P value [†]
A	34 (46.6; 0.36–0.58)	13	26	0.005
O	12 (20.0; 0.12–0.32)	14	34	
B	7 (38.9; 0.20–0.61)	1	10	
AB	1 (14.3; 0.03–0.51)	1	5	
Total	54	29	75	

Relative risk of NAIT was 0.67 (95% CI 0.48–0.94) in neonates born of women with blood group O versus blood group A. Relative risk of severe NAIT was 0.43 (95% CI 0.25–0.75) in neonates born of women with blood group O versus blood group A. [†]Chi-square test (two-sided) for frequencies of NAIT and severe NAIT in blood group O compared to blood group A.

(restricted by lack of material). For ABO-incompatibility studies, thus only 52 of 60 blood group O mother-child pairings could be included. The fifty-two mothers with blood group O gave birth to 16 A-incompatible (blood group A₁) and 36 compatible (sixteen blood group O, one B, and five A₂) children. Four of the 16 A-incompatible pregnancies resulted in a newborn with severe thrombocytopenia compared to 6 of the 36 ABO-compatible pregnancies. This indicates that ABO incompatibility is not the underlying cause of the observed phenomenon reported in the present study.

3.4. The ABO Genotype of the Mothers and Platelet Counts in the Newborn. The ABO genotype of 143 HPA-1a-immunized women who gave birth to 155 HPA-1a-positive neonates was determined (data not shown). The overall O allele frequencies among the immunized women were O01 0.56, O02 0.42, and O03 0.02. Individuals with blood groups A, B, and O were further subgrouped based on genotyping, and thus, the frequencies of newborns with severe NAIT within each subgroup were compared. The cases with maternal blood group AB were excluded for further analysis due to the low number of individuals. Analysis of the platelet counts in newborns of mothers with different ABO genotypes revealed that the frequency of newborns with severe NAIT differed (Pearson Chi-square $P = 0.0036$) among the maternal ABO genotype groups.

Among blood group A mothers, the frequency of newborns with severe NAIT was 42% in pregnancies where the mother carried only one A allele (A101 or A201), compared to 69% where mothers carried two A alleles (relative risk 0.61; 95% CI 0.38–0.98). In pregnancies where the mother had blood group O, the frequency of newborns with severe NAIT was 9%, where the mother did not carry any O02 allele, compared to 27% where the mother carried one or two O02 alleles; however, this did not reach statistical significance (relative risk 0.33 NS $P = 0.13$).

Platelet counts in newborns of mothers with blood group A and O are plotted in Figure 1. The mean antibody levels between these groups were not significantly different: 11.6 IU/mL for blood group A mothers, 1.8 IU/mL for O02-negative blood group O mothers, and 11.1 IU/mL for O02-positive blood group O mothers ($P = 0.18$ one-way ANOVA). However, the correlation between the maternal




FIGURE 1: The platelet count at delivery in HPA-1a-positive newborns was grouped according to maternal ABO type (A or O). Platelet count $\leq 150 \times 10^9/L$ (dashed line) is defined as NAIT and $< 50 \times 10^9/L$ (dotted line) as severe NAIT. The mean platelet counts in the three groups are indicated, with error bars representing 95% CI. Mean platelet counts are significantly different ($P < 0.019$, one-way ANOVA). The mean antibody levels between these groups are not significantly different: 11.6 IU/mL for blood group A mothers, 11.1 IU/mL for O02-positive blood group O mothers, and 1.8 IU/mL O02-negative blood group O mothers.

antibody level and platelet count in the newborn for these cases was described in Killie et al. [6].

Among the NAIT cases, defined as platelet count $\leq 150 \times 10^9/L$, the mean platelet count in newborns of homozygous O01/O01 mothers was higher ($83.2 \times 10^9/L$) than in the newborns with O02-positive mothers with blood group O ($43.7 \times 10^9/L$) or in newborns of women with blood group A ($46.1 \times 10^9/L$) (Table 2). Together, these data support

TABLE 2: Platelet counts in the newborns with NAIT.

Maternal ABO types		N*	Median platelet count	Mean platelet count (95% CI)	P value with Bonferroni correction
Blood group O	Genotype <i>O01/O01</i>	9	86	83.2 (56.9–109.6)	0.043 [†]
	Genotypes <i>O01/O02</i> and <i>O02/O02</i>	17	21	43.7 (22.7–64.7)	
Blood group A		47	31	46.1 (35.0–57.2)	0.028 [†]

* Total number of pregnancies resulting in a newborn with NAIT (platelet count ≤ 150) was 83. DNA for genotyping was available for 68 women with blood group O or A, and they had altogether 73 HPA-1-incompatible pregnancies.

[†] Comparison with platelet counts in children born of women with the genotype *O01/O01*.

our hypothesis that there are genetic properties among the immunized women influencing the risk of severe NAIT in the newborn.

4. Discussion

The ABO phenotype distribution among the HPA-1a immunized women is similar to the distribution in the Norwegian population, indicating that the generation of an immune response with antibody synthesis is independent of the ABO blood group of the mother. However, we observed that whereas only 20% of pregnancies among the immunized women with blood group O resulted in severe NAIT in the newborn, 47% of the immunized women with blood group A had newborns with severe thrombocytopenia. A recent retrospective study by Bertrand et al. did not find any significant correlation between the severity of the thrombocytopenia and the ABO genotype [18]. As these authors propose, the discrepancy between Bertrand's and our study may be due to the retrospective/prospective nature of the studies. We found no indications that the low frequency of severe NAIT in the children of women with blood group O was due to ABO incompatibility between mother and foetus. Additional measurements of maternal anti-RBC IgG antibody in the women with blood group O could have given further information of any influence of potential antibodies directed against the A antigen carried by α IIb on platelets. Another hypothesis that could explain the lower frequency of newborns with severe NAIT among the immunized mothers with blood group O, compared to blood group A, is that the ABO gene is located close to a gene encoding an immunoregulatory factor with polymorphic variants. In order to approach this question, we compared NAIT to ABO genotypes. The allelic differences in the gene encoding the A/B glycosyltransferases are defined by SNPs that changes the amino acid sequence of the enzyme and thereby its glycosylating properties. The *A101*, *A201*, *B101*, *O01*, *O02*, and *O03* alleles all produce transcripts (although A transcripts are virtually undetectable in peripheral blood) [19, 20], but the *O01* and *O02* transcripts both contain a shift in the reading frames that will severely truncate any resulting protein and leave it without enzymatic activity [21]. It is still unclear if these short nonfunctional proteins are expressed at all although it has been suggested [22].

The ABO genotype frequencies in the Norwegian population are not known, but the O allele frequencies observed in immunized women are similar to the frequencies reported

for a Swedish population [15], where the *O02* constitutes about 40% of all O alleles and *O03* allele is infrequent. This further shows that the generation of an immune response to HPA-1a is independent of ABO blood groups. However, when it comes to development of NAIT, the different risks of severe thrombocytopenia observed in genetic subgroups of blood group A support the hypothesis that a genetic linkage may be involved, rather than the ABO phenotype itself even though the mechanism is still not understood. Although the differences in the *O02*-positive and *O02*-negative subgroups of blood group O do not reach statistical significance, an interesting trend is observed.

Phylogenetic analyses of the ABO locus have shown that the *O02* probably is an ancient allelic lineage at the ABO locus, separate from the *A101* and *O01* alleles [23]. Therefore, it is interesting to subdivide the blood group O women according to their genotype. The 9q34 chromosomal region, where the ABO gene is located [24], contains several loci encoding immune response regulating genes. There is obviously no genetic linkage between the ABO [9q34] and *ITGB3* [17q21] loci. The association of the ABO type to the development of severe NAIT could be due to a potential linkage to one or more gene(s) encoding regulatory factors. Further investigation has to be conducted to find out whether such factors are linked to the ABO locus in a way that can explain our observation.

5. Conclusions

The development of severe NAIT in newborns is caused by transfer of platelet-reactive antibodies during pregnancy; however, several biological factors likely play a role in the immune response mechanism. In the present study, with data from a prospective NAIT study, we showed that the risk of severe NAIT due to anti-HPA-1a antibodies is correlated to maternal ABO types. The results indicate that there are genetic properties related to the maternal ABO genotype that influence the immune response that cause severe thrombocytopenia in the newborn of anti-HPA-1a immunized mothers.

Authors' Contribution

B. Skogen was responsible for conception of the study. M. T. Ahlen contributed to study design, performed the experiments, collected data, and performed statistical analyses. A. Husebekk supervised the research; M. L. Olsson

contributed with study design and interpretation of ABO analyses; J. Kjeldsen-Kragh and M. K. Killie contributed to study design and interpretation of data, the paper was written by M. T. Ahlen and B. Skogen, with contributions from A. Husebekk, J. Kjeldsen-Kragh, M. K. Killie, and M. L. Olsson. All authors critically reviewed the paper and approved the paper for publication.

Conflicts of Interests

The authors reported no potential conflicts of interest.

Acknowledgments

The authors would like to thank Dr. Åsa Hellberg for help with technical advice regarding ABO genotyping. This work was supported by a Grant from the North Norway Regional Health Authority (Bodø, Norway).

References

- [1] P. J. Newman, R. S. Derbes, and R. H. Aster, "The human platelet alloantigens, PI(A1) and PI(A2), are associated with a leucine33/proline33 amino acid polymorphism in membrane glycoprotein IIIa, and are distinguishable by DNA typing," *Journal of Clinical Investigation*, vol. 83, no. 5, pp. 1778–1781, 1989.
- [2] L. L. W. Cooling, K. Kelly, J. Barton, D. Hwang, T. A. W. Koerner, and J. D. Olson, "Determinants of ABH expression on human blood platelets," *Blood*, vol. 105, no. 8, pp. 3356–3364, 2005.
- [3] J. Kjeldsen-Kragh, M. K. Killie, G. Tomter et al., "A screening and intervention program aimed to reduce mortality and serious morbidity associated with severe neonatal alloimmune thrombocytopenia," *Blood*, vol. 110, no. 3, pp. 833–839, 2007.
- [4] G. Bertrand, C. Martageix, V. Jallu, F. Vitry, and C. Kaplan, "Predictive value of sequential maternal anti-HPA-1a antibody concentrations for the severity of fetal alloimmune thrombocytopenia," *Journal of Thrombosis and Haemostasis*, vol. 4, no. 3, pp. 628–637, 2006.
- [5] S. Jægtvik, A. Husebekk, B. Aune, P. Øian, L. B. Dahl, and B. Skogen, "Neonatal alloimmune thrombocytopenia due to anti-HPA 1a antibodies; the level of maternal antibodies predicts the severity of thrombocytopenia in the newborn," *British Journal of Obstetrics and Gynaecology*, vol. 107, no. 5, pp. 691–694, 2000.
- [6] M. K. Killie, A. Husebekk, J. Kjeldsen-Kragh, and B. Skogen, "A prospective study of maternal anti-HPA 1a antibody level as a potential predictor of alloimmune thrombocytopenia in the newborn," *Haematologica*, vol. 93, no. 6, pp. 870–877, 2008.
- [7] D. L'Abbe, L. Tremblay, M. Filion et al., "Alloimmunization to platelet antigen HPA-1a (PIA1) is strongly associated with both HLA-DRB3*0101 and HLA-DQB1*0201," *Human Immunology*, vol. 34, no. 2, pp. 107–114, 1992.
- [8] L. M. Williamson, G. Hackett, J. Rennie et al., "The natural history of fetomaternal alloimmunization to the platelet-specific antigen HPA-1a (PIA1, Zwa) as determined by antenatal screening," *Blood*, vol. 92, no. 7, pp. 2280–2287, 1998.
- [9] M. K. Killie, J. Kjeldsen-Kragh, I. Randen, B. Skogen, and A. Husebekk, "Evaluation of a new flow cytometric HPA 1a screening method: a rapid and reliable tool for HPA 1a screening of blood donors and pregnant women," *Transfusion and Apheresis Science*, vol. 30, no. 2, pp. 89–92, 2004.
- [10] B. Skogen, B. R. Hansen, A. Husebekk, T. Havnes, and K. Hannestad, "Minimal expression of blood group A antigen on thrombocytes from A2 individuals," *Transfusion*, vol. 28, no. 5, pp. 456–459, 1988.
- [11] B. R. Curtis, A. Fick, A. J. Lochowicz et al., "Neonatal alloimmune thrombocytopenia associated with maternal-fetal incompatibility for blood group B," *Transfusion*, vol. 48, no. 2, pp. 358–364, 2008.
- [12] K. Ogasawara, J. Ueki, M. Takenaka, and K. Furihata, "Study on the expression of ABH antigens on platelets," *Blood*, vol. 82, no. 3, pp. 993–999, 1993.
- [13] O. O. Blumenfeld and S. K. Patnaik, "Allelic genes of blood group antigens: a source of human mutations and cSNPs documented in the blood group antigen gene mutation database," *Human Mutation*, vol. 23, no. 1, pp. 8–16, 2004.
- [14] M. L. Olsson and M. A. Chester, "A rapid and simple ABO genotype screening method using a novel B/O2 versus A/O2 discriminating nucleotide substitution at the ABO locus," *Vox Sanguinis*, vol. 69, no. 3, pp. 242–247, 1995.
- [15] M. L. Olsson and M. A. Chester, "Frequent occurrence of a variant O1 gene at the blood group ABO locus," *Vox Sanguinis*, vol. 70, no. 1, pp. 26–30, 1996.
- [16] B. Hosseini-Maaf, A. Hellberg, M. A. Chester, and M. L. Olsson, "An extensive polymerase chain reaction-allele-specific polymorphism strategy for clinical ABO blood group genotyping that avoids potential errors caused by null, subgroup, and hybrid alleles," *Transfusion*, vol. 47, no. 11, pp. 2110–2125, 2007.
- [17] O. Hartmann and P. Stavem, "ABO blood-groups and cancer," *The Lancet*, vol. 283, no. 7346, pp. 1305–1306, 1964.
- [18] G. Bertrand, M. Drame, C. Martageix, and C. Kaplan, "Prediction of the fetal status in noninvasive management of alloimmune thrombocytopenia," *Blood*, vol. 117, no. 11, pp. 3209–3213, 2011.
- [19] B. Thuresson, M. A. Chester, J. R. Storry, and M. L. Olsson, "ABO transcript levels in peripheral blood and erythropoietic culture show different allele-related patterns independent of the CBF/NF-Y enhancer motif and multiple novel allele-specific variations in the 5'- and 3'-noncoding regions," *Transfusion*, vol. 48, no. 3, pp. 493–504, 2008.
- [20] Y.-C. Twu, C.-Y. Hsieh, and L.-C. Yu, "Expression of the histo-blood group B gene predominates in AB-genotype cells," *Transfusion*, vol. 46, no. 11, pp. 1988–1996, 2006.
- [21] F. Yamamoto, H. Clausen, T. White, J. Marken, and S. Hakomori, "Molecular genetic basis of the histo-blood group ABO system," *Nature*, vol. 345, no. 6272, pp. 229–233, 1990.
- [22] B. Eiz-Vesper, A. Seltsam, and R. Blasczyk, "ABO glycosyltransferases as potential source of minor histocompatibility antigens in allogeneic peripheral blood progenitor cell transplantation," *Transfusion*, vol. 45, no. 6, pp. 960–968, 2005.
- [23] F. Roubinet, S. Despiau, F. Calafell et al., "Evolution of the O alleles of the human ABO blood group gene," *Transfusion*, vol. 44, no. 5, pp. 707–715, 2004.
- [24] E. P. Bennett, R. Steffensen, H. Clausen, D. O. Weghuis, and A. G. van Kessel, "Genomic cloning of the human histo-blood group ABO locus," *Biochemical and Biophysical Research Communications*, vol. 206, no. 1, pp. 318–325, 1995.