



LUND UNIVERSITY

Endoscopic sinus surgery improve quality of life and decrease absenteeism in patients with chronic rhinosinusitis - a multi-centre study

Sahlstrand Johnson, Pernilla; Ohlsson, Bodil; Ahlner Elmqvist, Marianne

Published in:
Clinical and Translational Allergy

DOI:
[10.1186/2045-7022-5-S4-O6](https://doi.org/10.1186/2045-7022-5-S4-O6)

2015

Document Version:
Publisher's PDF, also known as Version of record

[Link to publication](#)

Citation for published version (APA):
Sahlstrand Johnson, P., Ohlsson, B., & Ahlner Elmqvist, M. (2015). Endoscopic sinus surgery improve quality of life and decrease absenteeism in patients with chronic rhinosinusitis - a multi-centre study. *Clinical and Translational Allergy*, 5(suppl 4), 2-3. Article 06. <https://doi.org/10.1186/2045-7022-5-S4-O6>

Total number of authors:
3

General rights

Unless other specific re-use rights are stated the following general rights apply:
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00



ORAL PRESENTATION

Open Access

Endoscopic sinus surgery improve quality of life and decrease absenteeism in patients with chronic rhinosinusitis - a multi-centre study

Pernilla Sahlstrand-Johnson^{1*}, Bodil Ohlsson², Marianne Ahlner-Elmqvist³

From The 10th Symposium of Experimental Rhinology and Immunology of the Nose (SERIN 2015) Stockholm, Sweden. 19-21 February 2015

Background

We have previously reported that chronic rhinosinusitis (CRS) has a significant impact on health-related quality of life (HRQOL). In this study we aimed to analyse whether sino-nasal symptom scoring can be used to predict the postoperative outcome measured by HRQOL. Secondly, we wanted to investigate whether absenteeism caused by sinus symptoms decrease after endoscopic sinus surgery (ESS). We also wanted to assess whether the duration of sinus disease had an effect on the outcome of ESS.

Methods

Two hundred and seven patients with CRS with and without nasal polyps (NP) admitted for ESS from ten hospitals were enrolled. EPOS2012 definitions of CRS and NP were used. The patients completed the short-form 36-item questionnaire (SF-36), the 22 Sinonasal Outcome Test (SNOT-22) and a total Visual Analogue Scale (VAS) regarding rhinosinusitis symptoms preoperatively, 6 and 12 months after surgery.

Results

Of the 207 patients, 135 were diagnosed with CRS+NP and 72 CRS-NP. Sixty-nine percent of the population had anterior ethmoidectomy, 36% posterior ethmoidectomy and 24% surgery of the frontal sinus recess. The quality of life scores measured by SF-36 improved in all eight domains but one after surgery. The total SNOT-22 score decreased significantly from 51.8 to 33.0 at 6 months postop ($p < 0.0001$) and stayed at this level 12 months after surgery. There was no statistically significant difference in total SNOT-22 scoring between patients with and without

NP, but in four of the SNOT-22 questions there were differences. The total VAS scores diminished from 67 to 33 postoperatively. Sick-leave due to rhinosinusitis dropped from 8-14 days to 1-7 days 12 months after ESS. The questions on "Lack of a good night's sleep" and "Frustrated/restless/irritable" in SNOT-22 were predictive ($p = 0.001$) for the postoperative total SNOT-22 score, when performing a backwards stepwise regression analysis. Those patients who had had a history of less than 12 months of sinus disease scored best postoperatively measured by SNOT-22.

Conclusion

Quality of life improved significantly after ESS measured by SF-36, SNOT-22 and VAS. Scoring of two of the questions in SNOT-22 proved to be significantly predictive for the postoperative outcome. Patients with CRS±NP reported less absenteeism due to sino-nasal disease 12 months postoperatively. The patients with shorter sinus disease duration seem to gain more by sinus surgery, which is something we need to look further at.

Authors' details

¹Skåne University Hospital, Dep of Otorhinolaryngology, 20502, Malmö, Sweden. ²Skåne University Hospital, Department of Clinical Sciences, Malmö, Sweden. ³Lund University, Department of Health Sciences, Lund, Sweden.

Published: 26 June 2015

doi:10.1186/2045-7022-5-S4-O6

Cite this article as: Sahlstrand-Johnson et al.: Endoscopic sinus surgery improve quality of life and decrease absenteeism in patients with chronic rhinosinusitis - a multi-centre study. *Clinical and Translational Allergy* 2015 **5**(Suppl 4):O6.

¹Skåne University Hospital, Dep of Otorhinolaryngology, 20502, Malmö, Sweden

Full list of author information is available at the end of the article