



# LUND UNIVERSITY

## Co-segregation of the PROS1 locus and protein S deficiency in families having no detectable mutations in PROS1.

Lanke, Elsa; Johansson, Anna; Hillarp, Andreas; Lethagen, Stefan; Zöller, Bengt; Dahlbäck, Björn; Halldén, Christer

*Published in:*

Journal of Thrombosis and Haemostasis

*DOI:*

[10.1111/j.1538-7836.2004.00950.x](https://doi.org/10.1111/j.1538-7836.2004.00950.x)

2004

[Link to publication](#)

*Citation for published version (APA):*

Lanke, E., Johansson, A., Hillarp, A., Lethagen, S., Zöller, B., Dahlbäck, B., & Halldén, C. (2004). Co-segregation of the PROS1 locus and protein S deficiency in families having no detectable mutations in PROS1. *Journal of Thrombosis and Haemostasis*, 2(11), 1918-1923. <https://doi.org/10.1111/j.1538-7836.2004.00950.x>

*Total number of authors:*

7

### General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117  
221 00 Lund  
+46 46-222 00 00

## Co-segregation of the *PROS1* locus and protein S deficiency in families having no detectable mutations in *PROS1*

E. LANKE,\* A. M. JOHANSSON,† A. HILLARP,‡ S. LETHAGEN,\* B. ZÖLLER,§ B. DAHLBÄCK‡ and C. HALLDÉN‡

\*Department for Coagulation Disorders, University Hospital, Malmö; †Department of Cell and Organism Biology, Lund University, Sweden;

‡Department of Clinical Chemistry, University Hospital, Malmö; and §Department of Internal Medicine, Ystad Hospital, Sweden

**To cite this article:** Lanke E, Johansson AM, Hillarp A, Lethagen S, Zöller B, Dahlbäck B, Halldén C. Co-segregation of the *PROS1* locus and protein S deficiency in families having no detectable mutations in *PROS1*. *J Thromb Haemost* 2004; 2: 1918–23.

**Summary.** Inherited deficiency of protein S constitutes an important risk factor of venous thrombosis. Many reports have demonstrated that causative mutations in the protein S gene are found only in approximately 50% of the cases with protein S deficiency. It is uncertain whether the protein S gene is causative in all cases of protein S deficiency or if other genes are involved in cases where no mutation is identified. The aim of the current study was to determine whether haplotypes of the protein S gene cosegregate with the disease phenotype in cases where no mutations have been found. Eight protein S-deficient families comprising 115 individuals where previous DNA sequencing had failed to detect any causative mutations were analyzed using four microsatellite markers in the protein S gene region. Co-segregation between microsatellite haplotypes and protein S deficiency was found in seven of the investigated families, one family being uninformative. This suggests that the causative genetic defects are located in or close to the protein S gene in a majority of such cases where no mutations have been found.

**Keywords:** coagulation, *PROS1*, protein C, protein S, vitamin K, thrombosis.

### Introduction

Protein S (PS) is a vitamin K-dependent single-chain plasma protein that regulates blood coagulation. Its main function is to serve as a nonenzymatic cofactor to activated protein C (APC) in the proteolytic degradation of procoagulant factors Va and VIIIa [1]. Inherited deficiency of PS is an autosomal dominant disorder that is associated with venous thrombosis. Homozygous or compound heterozygous deficiency of PS is extremely rare. Heterozygous PS deficiency is present in approximately

2–5% of thrombosis patients [2,3], whereas the prevalence in the general population has been estimated to be between 0.03 and 0.13% [4]. In plasma, PS circulates both as free protein and in complex with C4b-binding protein. Approximately 30% of PS is found in the free form and it is only the free form that can function as a cofactor to APC [5]. Due to the different forms of PS, three types of PS deficiency have been proposed. Type I is characterized by low levels of both total and free PS. Type II entails diminished activity of PS but normal levels of free and total PS. Type III deficiency is characterized by a low level of free PS and a normal level of total PS. Type I and III deficiency have been shown to coexist in many families [6–12], indicating that these two types of deficiencies are phenotypic variants of the same genetic disease. There are a number of factors that influence the plasma level of PS, including the genetic [13] and physiologic status, which lead to an overlap in PS levels between healthy individuals and PS deficient patients. PS increases with age [8] and decreases during pregnancy and oral contraception [14]. Furthermore, different assays for PS are influenced by preanalytical variables and suffer from limitations in analytical performance making a correct laboratory diagnosis a difficult task [15–17]. Of the assays, a low free PS level has been found to be the most reliable way of diagnosing PS deficiency [6,18].

The gene for protein S (*PROS1*) is located near the centromere on chromosome 3p11.1-q11.2 [19]. It is approximately 80 kb long and contains 15 exons [19–22]. The large size of the gene, in addition to the presence of a pseudogene (*PROS2*), complicates genetic analysis of the *PROS1* gene. Nevertheless, screening for *PROS1* gene mutations has been performed in a number of studies. The proportion of cases where no mutations are detected varies widely between studies [10,17,23–33]. When all cases are pooled together, *PROS1* gene mutations are not detected in 47% of PS deficient families. The reason for the low prevalence of *PROS1* mutations in well-characterized PS deficient patients is unknown.

The present study investigates whether microsatellite marker haplotypes cosegregate with PS deficiency in type III or mixed type I and type III PS deficient families, where mutations in the *PROS1* gene despite sequencing have not been found.

Correspondence: Björn Dahlbäck, Department of Clinical Chemistry, Malmö University Hospital, SE-205 02 Malmö, Sweden.

Tel.: +46 40 331501; fax: +46 40 337044; e-mail: bjorn.dahlback@klkemi.mas.lu.se

Received 24 February 2004, accepted 23 June 2004

## Materials and methods

### Patients

In total, eight unrelated type III or mixed type I and type III PS deficient families, comprising 115 individuals, with a family history of venous thrombosis were investigated (Table 1). The characteristics and the diagnostic criteria of the phenotypic validation have been described previously [6,33]. Of the 56 individuals with PS deficiency, 21 had experienced venous thrombosis. The current study uses the same family identification codes as was used in the previous studies [6,33]. A PS deficient member of each family was previously investigated by DNA-sequencing of all exons and exon-intron boundaries for the presence of mutations that could explain the PS deficiency [33]. That study included a total of 17 families. In eight of these families (47%) no mutation could be identified [33]. All of these eight families are analyzed in the current study. The ethical committee approved the study and appropriate consent was obtained from the participating individuals.

### Microsatellite marker analysis

Genomic DNA was extracted from blood collected in EDTA using QIAamp DNA Blood Maxi Kit (Qiagen, Germany). DNA concentrations were determined by fluorometry using Pico Green. Four microsatellite markers, *D3S2388*, *D3S1595*, *D3S1552* and *D3S3634* were used for segregation analysis. The map positions of the four markers have been established by using a combination of the deCode, Marshfield and Genethon maps since neither of them includes all four markers. The four microsatellite markers were positioned in the above-mentioned order with the *PROS1* gene located between markers *D3S1552* and *D3S3634*. The distances from *D3S2388* to *D3S1595*, to *D3S1552* and to *D3S3634* were 1.58, 3.20 and 4.30 cM, respectively. The microsatellite markers were amplified using PCR conditions as recommended by the manufacturer of the Taq polymerase (Applied Biosystems, USA) and an annealing temperature of 58 °C for marker *D3S1552* and 62 °C for the

remaining three markers. PCR products were resolved using capillary electrophoresis run on an ABI PRISM™ 310 sequencer employing GeneScan software (Applied Biosystems, USA). The allele sizes of the microsatellite markers were determined in relation to a size marker, GeneScan-350™ ROX (Applied Biosystems, USA). All participating individuals were successfully genotyped for all markers; hence there were no missing values.

### Genetic analysis

The pedigrees were constructed using Progeny (Progeny Software LLC, USA). Multi-point LOD-scores were calculated using Genehunter [34] and Simwalk2 [35]. Genehunter was used for families 11, 20, 26 and 31, whereas Simwalk2 was used for families 8, 13, 15 and 17, as these families were too extensive for Genehunter. Simwalk2 calculates location-score, which is fully comparable to LOD-score. The penetrance was assumed to be 100% and the frequency of the disease allele 0.001 [18]. In family number 26, the cosegregation analysis suggested incomplete penetrance and the disease status of the two healthy siblings in this family was therefore given unknown status. In family number 15 the affected family members had a null allele for *D3S3634*, and in family number 31 the affected family members had a null allele for both *D3S1595* and *D3S3634*. For this reason, these markers were not used in the calculations for these families. Microsatellite marker allele frequencies were calculated from 48 unrelated individuals not affected by PS deficiency. These individuals are sampled from the same population as the investigated families. Genotype inconsistencies were identified manually and using PedCheck [36]. Repeated genotyping made it possible to correct all inconsistencies. Haplotypes for all individuals were reconstructed using Simwalk2, which also searched for recombination events.

## Results

A total of 115 individuals from eight protein S deficient families were genotyped using four microsatellite markers from the

**Table 1** Description of the investigated PS deficient families


Family number	Number of individuals	Number of PS-deficient individuals*	Disease-associated haplotype	Multi-point LOD-scores			
				<i>D3S2388</i>	<i>D3S1595</i>	<i>D3S1552</i>	<i>D3S3634</i>
8	16	9 (4)	4,4,4,3	3.64	3.69	3.75	3.70
11	6	4 (2)	3,4,4,5	0.79	0.80	0.81	0.82
13†	31	10 (1)	3,2,4,2 and 4,4,4,2	3.43	3.38	3.78	3.81
15‡	22	12 (3)	3,8,4,-	3.24	3.26	3.20	-
17	20	10 (6)	3,4,3,3	3.83	4.46	4.72	4.68
20	8	4 (1)	4,5,1,2	0.78	0.82	0.88	0.88
26§	5	2 (2)					
31‡	7	5 (1)	6,-,4,-	1.18	-	1.09	-

\*All families, except family 26, consist of mixed type I and type III phenotypes. The number of type III deficient individuals is given in parenthesis and is defined by low free protein S concentration and a total protein S concentration within the normal reference range [33]. †Two disease-associated haplotypes identified in two branches of this family. ‡Null alleles present. Markers indicated by - were not used in linkage analysis. §This type III family is uninformative due to the small size of the family.

*PROS1* gene region. There was cosegregation between PS deficiency and certain microsatellite haplotypes in seven informative families, family 26 being uninformative. Each family had its own unique disease-associated haplotype (Table 1). As an example, the pedigree of family number 8 is shown in Fig. 1(A). Haplotype 4, 4, 4, 3 is present in all PS deficient individuals and is absent in all healthy individuals in this pedigree. In family number 13, there were two clear divisions of the family with a certain microsatellite haplotype cosegregating with the disease for each division (Fig. 1B). The two haplotypes shared alleles for markers *D3S1552* and *D3S3634* (flanking the *PROS1* gene), indicating that one of the haplotypes most likely has emerged from the other by recombination. All PS deficient individuals within a family shared the same haplotype and there was no case of a nonaffected member with the disease-associated haplotype, except for two possible cases: one healthy individual in family number 13 (individual IV:6, Fig. 1B) could not be excluded from having the disease-associated haplotype; and family number 26 where all four children shared the same haplotypes, yet only two of them were clearly affected by PS deficiency. However, as the father in this family was deceased, his protein S status was unknown and his DNA unavailable for analysis, the family was uninformative. Apart from these possible exceptions, no healthy individuals had a disease-associated haplotype. Thus, the penetrance was 100% in the informative families.

The pattern of cosegregation observed for all informative families is compatible with an explanation where the *PROS1* gene is causative of the protein S deficiency. The strength of this conclusion at the family level was determined by performing multipoint linkage analysis. LOD scores for linkage (recombination fraction = 0) between PS deficiency and the four microsatellite markers are shown in Table 1. Four of the families showed very high LOD-scores between 3.2 and 4.8. Three of the remaining four families produced lower but positive LOD-scores between 0.78 and 1.2. The four families with high LOD-scores contained between 16 and 31 individuals, whereas the families with slightly lower LOD-scores contained between six and eight individuals. The lower LOD scores in three of the families are a direct result of their small sizes and the low number of affected individuals. In addition, family number 26 was uninformative. Family number 20 most likely contains a recombination event, which contributes to the lower LOD-score observed in this family. Thus, the positive LOD-scores detected in all informative families support cosegregation between the markers flanking the *PROS1* gene and the PS deficiency phenotype.

Simwalk2 suggested three recombinant haplotypes in total, one of which is the above-mentioned in family number 13. This recombination event has occurred between *D3S1595* and *D3S1552*, which correspond to the two divisions within the family. In family number 17, recombination has occurred between *D3S2388* and *D3S1595* or between *D3S1595* and *D3S1552*. In family number 20, Simwalk2 suggested a recombination between *D3S1595* and *D3S1552*. According to the



**Fig. 1.** Pedigrees demonstrating cosegregation between PS deficiency and microsatellite haplotypes of the *PROS1* gene region. Arrows indicate index cases, which have previously been sequenced with no causative mutations found. Filled symbols indicate diagnosed PS deficiency, and the capital letter N denotes healthy individuals, whereas empty symbols indicate unknown status. We have had access to blood samples of individuals with haplotypes indicated below. Microsatellite markers are shown in the order as suggested by the available mapping data. Haplotypes was suggested by Simwalk2. The disease-associated haplotype (marked in bold) is detected in all genotyped individuals diagnosed with PS deficiency but in no healthy individuals, which clearly states cosegregation between the disease and the *PROS1* gene locus. (A) Pedigree showing cosegregation between PS deficiency and the haplotype 4, 4, 4, 3 in family number 8. (B) Pedigree of family number 13 illustrating recombination. PS deficiency in the two branches cosegregates with the haplotypes 3, 2, 4, 2 and 4, 4, 4, 2, respectively, of the *PROS1* gene region. The only individual whose haplotypes cannot be ascertained is individual IV:6, which is indicated by slashes. (C) Pedigree of family 31 showing cosegregation between PS deficiency and the haplotype 6, -4, -. The existence of a deletion is indicated by reanalysis of marker *D3S3634* using another pair of primers.

order in which the markers and the *PROS1* gene appear, none of the recombination events separated the gene from its two flanking markers.

In families 15 and 31, null-alleles appeared for some markers in the disease-associated haplotypes. As null-alleles are usually the result of polymorphisms occurring in the primer annealing sites or, more seldom, the result of deletions, the marker *D3S3634* was re-analyzed using a different set of primers. Also in these cases, the disease-associated haplotypes carried a null-allele in both families. This indicates that deletions exist in both of these families. This pattern of inheritance is illustrated in Fig. 1(C) where the hemizygosity for marker *D3S3634* result in several cases of apparent non-Mendelian inheritance in family 31.

## Discussion

The identification of *PROS1* mutations in PS deficiency has been reported in several studies, including more than 300 unrelated PS deficient families [10,17,23–33]. In 53% of all these cases a mutation, likely to cause the decreased level of PS, was found, leaving the remaining 47% unexplained. Different screening procedures for mutation detection have been used in these studies including sequence analysis via RT-PCR, single-strand conformation polymorphism (SSCP) analysis, heteroduplex analysis, analysis by denaturing gradient gel electrophoresis (DGGE) and PCR amplification of *PROS1* exons followed by direct sequence analysis. The mRNA approach is fairly rapid but mutation detection is limited by the occurrence of allelic exclusion [7,25,38]. SSCP and DGGE analysis have been successfully applied as mutation screening procedures in many genes. However, the sensitivity of these techniques is not 100%. Thus, the preferred way to screen for mutations is by direct sequencing of the *PROS1* coding sequences, although it is more expensive than other screening techniques. The direct sequencing approach has been used in the majority of studies, including 147 families with PS deficiency [10,17,23,25,26,29,32,33]. Despite using this approach, 46% of the cases did not reveal any *PROS1* mutations. The sequencing approach was also used for our PS families with similar results, i.e. no mutation could be found in eight out of 17 PS deficient families [33]. This situation thus implicates either the existence of other loci involved in PS deficiency, or the existence of common types of genetic defects in the *PROS1* gene region that are not easily detected by sequencing of the coding regions.

The aim of the current study was to determine whether there was cosegregation between certain microsatellite haplotypes of the *PROS1* gene region and the disease phenotype in families in whom earlier sequencing had not revealed any causative mutations in the *PROS1* gene. We found PS deficiency to cosegregate with the *PROS1* haplotypes in seven of the eight families lacking an identified mutation, one family remaining uninformative [33]. The four large families (families 8, 13, 15 and 17) showed very high LOD-scores clearly demonstrating linkage. The smaller families (families 11, 20, and 31) showed

lower but positive LOD-scores. As we only test one candidate locus for cosegregation, the existence of positive LOD-scores in all informative families is an argument for the involvement of *PROS1* in a majority of protein S deficiency cases. In addition, in families 15 and 31 the cosegregating disease haplotypes carried deletions. Whether these deletions are causally related to the protein S deficiency phenotype remains to be demonstrated. Taken together, these results clearly indicate that in the informative families, the mutations causing PS deficiency are located in or within close proximity of the *PROS1* gene locus.

Our results are not in accordance with the results of two other studies, where it has been suggested that the molecular basis of type III PS deficiency is not a single defect in the *PROS1* gene that is inherited in a Mendelian way [10,39]. However, in this context it is noteworthy that type III PS deficiency is difficult to diagnose accurately due to a considerable overlap in PS values between individuals with and without the genetic defect. In addition, the number and sizes of the previously investigated type III PS deficient families were smaller than the ones now investigated. Moreover, we believe that the use of a panel of four informative microsatellite markers increase the validity of the present study. Based on our results, we propose that PS deficiency is a single-gene disorder with Mendelian inheritance and close to complete penetrance. This most likely includes most or all PS deficiency cases for which no causative mutations have been identified despite sequencing of the entire coding region. One explanation for the absence of identified mutations in previous studies might be that sequencing of the coding regions fails to detect mutations in the promoter or in the introns. As the promoter region of *PROS1* is largely uncharacterized and the introns are rather large, the screening for mutations in these regions is an extensive and uncertain task, which is even further complicated by the existence of the homologous *PROS2* pseudo gene. Other possible causes of protein S deficiency, which are difficult to identify with a sequencing approach, are larger deletions or duplications covering the whole or part of the *PROS1* gene. Given the results obtained in the present study for families 15 and 31, a dedicated screen for deletions in all exons of the gene seems warranted.

An implication of our finding is that a PS deficiency diagnosis can now be confirmed genetically, provided an entire family with several affected family members is analyzed. By using informative microsatellite markers, the inheritance of a cosegregating haplotype can be identified. As the penetrance of PS deficiency is close to complete, expressing a disease-associated haplotype is a very strong indication for having the disease. Considering the difficulty with which PS deficiency is diagnosed phenotypically, this genotypic approach will facilitate diagnosing familial PS deficiency.

In conclusion, our data contribute to a clearer understanding of the genetic background of inherited PS deficiency in terms of cosegregation between the disease and the *PROS1* gene locus. This new knowledge will help unravel the molecular genetics involved in PS deficiency.

## Acknowledgements

This work was supported by grants from the Swedish Science Council (07143) and by grants from the Malmö University Hospital funds and from Region Skåne.

## References

- Dahlbäck B. Blood coagulation. *Lancet* 2000; **355**: 1627–32.
- Lane DA, Mannucci PM, Bauer KA, Bertina RM, Bochkov NP, Boulyjenkov V, Chandry M, Dahlbäck B, Ginter EK, Miletich JP, Rosendaal FR, Seligsohn U. Inherited thrombophilia: part 1. *Thromb Haemost* 1996; **76**: 651–62.
- Seligsohn U, Lubetsky A. Genetic susceptibility to venous thrombosis. *N Eng J Med* 2001; **344**: 1222–31.
- Dykes AC, Walker ID, McMahon AD, Islam SIAM, Tait RC. A study of protein S antigen levels in 3788 healthy volunteers: influence of age, sex and hormone use, and estimate for prevalence of deficiency state. *Br J Haematol* 2001; **113**: 636–41.
- Dahlbäck B. Inhibition of protein C cofactor function of human and bovine protein S by C4b-binding protein. *J Biol Chem* 1986; **261**: 12022–7.
- Zöller B, Garcia de Frutos P, Dahlbäck B. Evaluation of the relationship between protein S and C4b-binding protein isoforms in hereditary protein S deficiency demonstrating type I and type III deficiencies to be phenotypic variants of the same genetic disease. *Blood* 1995; **85**: 3524–31.
- Mustafa S, Pabinger I, Varadi K, Halbmayer WM, Lechner K, Schwarz HP, Fischer M, Mannhalter C. A hitherto unknown splice site defect in the protein S gene (PROS1): the mutation results in allelic exclusion and causes type I and type III protein S deficiency. *Br J Haematol* 1997; **99**: 298–300.
- Simmonds RE, Zoller B, Ireland H, Thompson E, de Frutos PG, Dahlbäck B, Lane DA. Genetic and phenotypic analysis of a large (122-member) protein S-deficient kindred provides an explanation for the familial coexistence of type I and type III plasma phenotypes. *Blood* 1997; **89**: 4364–70.
- Hermida J, Faioni EM, Mannucci PM. Poor relationship between phenotypes of protein S deficiency and mutations in the protein S alpha gene. *Thromb Haemost* 1999; **82**: 1634–8.
- Espinosa-Parrilla Y, Morell M, Souto JC, Tirado I, Fontcuberta J, Estivill X, Sala N. Protein S gene analysis reveals the presence of a cosegregating mutation in most pedigrees with type I but not type III PS deficiency. *Hum Mutat* 1999; **14**: 30–9.
- Gandrille S, Borgel D, Sala N, Espinosa-Parrilla Y, Simmonds R, Rezende S, Lind B, Mannhalter C, Pabinger I, Reitsma PH, Formstone C, Cooper DN, Saito H, Suzuki K, Bernardi F, Aiach M. Protein S deficiency: a database of mutations – summary of the first update. *Thromb Haemost* 2000; **84**: 918.
- Andersen BD, Bisgaard ML, Lind B, Philips M, Villoutreix BO, Thorsen S. Characterization and structural impact of five novel PROS1 mutations in eleven protein S-deficient families. *Thromb Haemost* 2001; **86**: 1392–9.
- Leroy-Matheron C, Duchemin J, Levent M, Gouault-Heilmann M. Genetic modulation of plasma protein S levels by two frequent dimorphisms in the PROS1 gene. *Thromb Haemost* 1999; **82**: 1088–92.
- Malm J, Laurell M, Dahlbäck B. Changes in the plasma levels of vitamin K-dependent proteins C and S and of C4b-binding protein during pregnancy and oral contraception. *Br J Haematol* 1988; **68**: 437–43.
- Persson KEM, Hillarp A, Dahlbäck B. Analytical considerations for free protein S assays in protein S deficiency. *Thromb Haemost* 2001; **86**: 1144–7.
- Persson KEM, Dahlbäck B, Hillarp A. Diagnosing protein S deficiency: analytical considerations. *Clin Lab* 2003; **49**: 103–10.
- Labrousse S, Reboul M-P, Guérin V, Vergnes C, Freyburger G. Protein C and protein S assessment in hospital laboratories: which strategy and what role for DNA sequencing? *Blood Coagul Fibrinolysis* 2003; **14**: 531–8.
- Makris M, Leach M, Beauchamp NJ, Daly ME, Cooper PC, Hampton KK, Bayliss P, Peake IR, Miller GJ, Preston FE. Genetic analysis, phenotypic diagnosis, and risk of venous thrombosis in families with inherited deficiencies of protein S. *Blood* 2000; **95**: 1935–41.
- Watkins PC, Eddy R, Fukushima Y, Byers MG, Cohen EH, Dackowski WR, Wydro RM, Shows TB. The gene for protein S maps near the centromere of human chromosome 3. *Blood* 1988; **71**: 238–41.
- Edenbrandt CM, Lundwall A, Wydro R, Stenflo J. Molecular analysis of the gene for vitamin K dependent protein S and its pseudogene: cloning and partial gene organization. *Biochemistry* 1990; **29**: 7861–8.
- Ploos van Amstel HK, Reitsma PH, van der Logt CPE, Bertina RM. Intron-exon organization of the active human protein S gene PS alpha and its pseudogene PS beta: duplication and silencing during primate evolution. *Biochemistry* 1990; **29**: 7853–61.
- Schmidel DK, Tatro AV, Phelps LG, Tomczak JA, Long GL. Organization of the human protein S genes. *Biochemistry* 1990; **29**: 7845–52.
- Reitsma PH, Ploos van Amstel HK, Bertina RM. Three novel mutations in five unrelated subjects with hereditary protein S deficiency type I. *J Clin Invest* 1994; **93**: 486–92.
- Formstone CJ, Wacey AI, Berg LP, Rahman S, Bevan D, Rowley M, Voke J, Bernardi F, Legnani C, Simioni P, Girolami A, Tuddenham EGD, Kakkar VV, Cooper DN. Detection and characterization of seven novel protein S (PROS) gene lesions: evaluation of reverse transcriptase-polymerase chain reaction as a mutation screening strategy. *Blood* 1995; **86**: 2632–41.
- Gomez E, Poort SR, Bertina RM, Reitsma PH. Identification of eight point mutations in protein S deficiency type I – analysis of 15 pedigrees. *Thromb Haemost* 1995; **73**: 750–5.
- Mustafa S, Pabinger I, Mannhalter C. Protein S deficiency type I. identification of point mutations in 9 of 10 families. *Blood* 1995; **86**: 3444–51.
- Borgel D, Duchemin J, Alhenc-Gelas M, Matheron C, Aiach M, Gandrille S and the French network on molecular abnormalities responsible for protein C and protein S deficiencies. Molecular basis for protein S hereditary deficiency: genetic defects observed in 118 patients with type I and type IIa deficiencies. *J Lab Clin Med* 1996; **128**: 218–27.
- Duchemin J, Borg JY, Borgel D, Vasse M, Leveque H, Aiach M, Gandrille S. Five novel mutations of the protein S active gene (PROS 1) in 8 Norman families. *Thromb Haemost* 1996; **75**: 437–44.
- Simmonds RE, Ireland H, Kunz G, Lane DA and the protein S Study Group. Identification of 19 protein S gene mutations in patients with phenotypic protein S deficiency and thrombosis. *Blood* 1996; **88**: 4195–204.
- Bustorff TC, Freire I, Gago T, Crespo F, David D. Identification of three novel mutations in hereditary protein S deficiency. *Thromb Haemost* 1997; **77**: 21–5.
- Espinosa-Parrilla Y, Morell M, Borrell M, Souto JC, Fontcuberta J, Estivill X, Sala N. Optimization of a simple and rapid single-strand conformation analysis for detection of mutations in the PROS1 gene: identification of seven novel mutations and three novel, apparently neutral, variants. *Hum Mutat* 2000; **15**: 463–73.
- Iwaki T, Mastushita T, Kobayashi T, Yamamoto Y, Nomura Y, Kagami K, Nakayama T, Sugiura I, Kojima T, Takamatsu J, Kanayama N, Saito H. DNA sequence analysis of protein S deficiency – identification of four point mutations in twelve Japanese subjects. *Semin Thromb Hemost* 2001; **27**: 155–60.
- Rezende SM, Lane DA, Zoller B, Mille-Baker B, Laffan M, Dahlbäck B, Simmonds RE. Genetic and phenotypic variability between

- families with hereditary protein S deficiency. *Thromb Haemost* 2002; **87**: 258–65.
- 34 Kruglyak L, Daly MJ, Reeve-Daly MP, Lander ES. Parametric and nonparametric linkage analysis: a unified multipoint approach. *Am J Hum Genet* 1996; **58**: 1347–63.
- 35 Sobel E, Lange K. Descent graphs in pedigree analysis: applications to haplotyping, location scores, and marker sharing statistics. *Am J Hum Genet* 1996; **58**: 1323–37.
- 36 O'Connell JR, Weeks DE. PedCheck: a program for identification of genotype incompatibilities in linkage analysis. *Am J Hum Genet* 1998; **63**: 259–66.
- 37 Huang QY, Xu FH, Shen H, Deng HY, Liu YJ, Liu YZ, Li JL, Recker RR, Deng HW. Mutation patterns at dinucleotide microsatellite loci in humans. *Am J Hum Genet* 2002; **70**: 625–34.
- 38 Sacchi E, Pinotti M, Marchetti G, Merati G, Tagliabue L, Mannucci PM. Protein S mRNA in patients with protein S deficiency. *Thromb Haemost* 1995; **73**: 746–9.
- 39 Espinosa-Parrilla Y, Morell M, Souto JC, Borrell M, Heine-Suner D, Tirado I, Volpini V, Estivill X, Sala N. Absence of linkage between type III protein S deficiency and the *PROS1* and *C4 BP* genes in families carrying the protein S Heerlen allele. *Blood* 1997; **89**: 2799–806.