



LUND UNIVERSITY

Myocardial infarction in an urban population: worse long term prognosis for patients from less affluent residential areas.

Tydén, Patrik; Hansen, Ole; Engström, Gunnar; Hedblad, Bo; Janzon, Lars

Published in:
Journal of Epidemiology and Community Health

DOI:
[10.1136/jech.56.10.785](https://doi.org/10.1136/jech.56.10.785)

2002

[Link to publication](#)

Citation for published version (APA):
Tydén, P., Hansen, O., Engström, G., Hedblad, B., & Janzon, L. (2002). Myocardial infarction in an urban population: worse long term prognosis for patients from less affluent residential areas. *Journal of Epidemiology and Community Health*, 56(10), 785-790. <https://doi.org/10.1136/jech.56.10.785>

Total number of authors:
5

General rights

Unless other specific re-use rights are stated the following general rights apply:
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00



Myocardial infarction in an urban population: worse long term prognosis for patients from less affluent residential areas

P Tydén, O Hansen, G Engström, B Hedblad and L Janzon

J. Epidemiol. Community Health 2002;56:785-790
doi:10.1136/jech.56.10.785

Updated information and services can be found at:
<http://jech.bmjjournals.com/cgi/content/full/56/10/785>

These include:

References

This article cites 44 articles, 27 of which can be accessed free at:
<http://jech.bmjjournals.com/cgi/content/full/56/10/785#BIBL>

Rapid responses

You can respond to this article at:
<http://jech.bmjjournals.com/cgi/eletter-submit/56/10/785>

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article

Topic collections

Articles on similar topics can be found in the following collections

[Other Cardiovascular Medicine](#) (1718 articles)
[Socioeconomic Determinants of Health](#) (694 articles)

Notes

To order reprints of this article go to:
<http://www.bmjjournals.com/cgi/reprintform>

To subscribe to *Journal of Epidemiology and Community Health* go to:
<http://www.bmjjournals.com/subscriptions/>

RESEARCH REPORT

Myocardial infarction in an urban population: worse long term prognosis for patients from less affluent residential areas

P Tydén, O Hansen, G Engström, B Hedblad, L Janzon

J Epidemiol Community Health 2002;**56**:785–790

Study objective: The objective in this follow up study from the Malmö myocardial infarction register has been to assess whether long term survival following discharge after first myocardial infarction has any relation with the socioeconomic environment and to assess to what extent intra-urban differences in mortality from ischaemic heart disease can be accounted for by covariance with long term survival following discharge after acute myocardial infarction.

Design: Register based surveillance study.

Setting: Seventeen residential areas in the city of Malmö, Sweden.

Participants: The cohort contains all 2931 male and 2083 female patients with myocardial infarction who were discharged for the first time between 1986–95 from Malmö University Hospital.

Main results: During the on average 4.9 years of follow up 55% of the patients died. The sex adjusted and age adjusted all cause mortality rate/1000 patient years ranged between residential areas from 85.5 to 163.6. The area specific relative risk of death after discharge was associated with a low socioeconomic score, $r=-0.56$, $p=0.018$. Major risk factors for cardiovascular disease were more prevalent in areas with low socioeconomic score and low rates of survival. Of the intra-urban differences in mortality from ischaemic heart disease, 41% could be accounted for by differences with regard to the survival rate after discharge.

Conclusions: The results are compatible with the hypothesis that the socioeconomic environment plays an important part in the survival rate of patients with myocardial infarction. To assess the preventive potential, the extent to which socioeconomic circumstances covary with severity of disease, respectively with the use and compliance with secondary preventive measures, needs to be evaluated.

Correspondence to:
Dr P Tydén, Department of
Community Medicine, Unit
of Epidemiology Malmö
University Hospital, 205
02 Malmö, Sweden;
patrik.tyden@smi.mas.lu.se

Accepted for publication
14 January 2002

Between and within countries there are marked differences in mortality in ischaemic heart disease.^{1–3} In a study from the city of Malmö in Sweden it was shown that between residential areas the age adjusted and sex adjusted annual mortality rate in ischaemic heart disease ranged from 286 to 446/100 000.⁴ These differences could only partly be accounted for by covariance with incidence of disease and short-term case fatality rate.^{4–5} Areas with high mortality rates deviated in terms of their socioeconomic circumstances unfavourably from the city average. This association remained statistically significant after adjustment for differences between areas with regard to the prevalence of other major risk factors associated with cardiovascular disease.⁶

The severity of coronary atherosclerosis, the degree of damage to the myocardium, and the use and compliance with secondary preventive measures are some of the factors known to modify the prognosis after myocardial infarction.^{7–16} It has been our hypothesis that many of these prognostic markers covary with the patients' socioeconomic circumstances and that hence the prognosis would be poorer for patients from less affluent areas.

The objective in this follow up study from the Malmö myocardial infarction register¹⁷ has been to assess whether, in an urban population, long term survival following discharge after first myocardial infarction has any relation with the socioeconomic environment and to assess to what extent intra-urban differences in mortality from ischaemic heart disease can be accounted for by covariance with long term survival following discharge after acute myocardial infarction.

METHODS

Retrieval of cases

The city of Malmö, located in southern Sweden, has approximately 250 000 inhabitants. Malmö University Hospital is the single referral unit for patients with acute myocardial infarction. After discharge the patients are taken care of by general practitioners and private physicians. From the hospital patient administrative register it is possible to retrieve, for each patient, name, 10 digit personal identification number, diagnosis, and dates of admittance and discharge. For patients with ICD codes 410.00–410.99 respectively 410A–410X according to the 8th and 9th versions, this information has been transferred to the Malmö myocardial infarction register since 1972. The percentage of patients classified correctly has been estimated at 90%–95%.¹⁸ Residential area at the time of the treatment occasion is available for cases from 1986 and onwards. During the time period 1986 to 1995 there were in all 2931 men, mean (SD) age 68.5 (11.5), and 2083 women, mean age 75.3 (10.7) who for the first time were discharged from Malmö University Hospital after treatment because of myocardial infarction. The 42% female patients can be accounted for by differences with regard to the population at risk—that is, an overrepresentation of women in age groups with the highest incidence of myocardial infarction.¹⁹ At least two of the following three criteria were required for diagnosis: (1) Central chest pain, lung oedema or shock. (2) ECG changes indicating acute myocardial infarction. (3) Increased serum activities of cardiac enzymes.¹⁷

Vital status on 31 December 1998 was updated by record linkage with the National Cause of Death Register. The average time of follow up was 4.9 years (ranging from one day to 13 years). Vital status was updated on all patients.

Information regarding the underlying cause of death was at the time of this study only available for cases who died before 1997. Ischaemic heart disease was in 60% of the cases the underlying cause of death. The National Cause of Death Register supplied the information needed to calculate, for each of the residential areas, the average annual rate of mortality from ischaemic heart disease (ICD codes 410–414 according to the 8th and 9th version) during the time period 1986–1995. These estimates are based on 7922 deaths.

Socioeconomic circumstances of residential areas

Within the city there are 18 residential areas that in terms of morbidity and socioeconomic circumstances are not only very different from each other but have remained so over time.^{4 20 21} The harbour area, because of the small number of people living there, has not been included in this study. A comprehensive socioeconomic score, SES, was used for the comparison of socioeconomic circumstances. For the computation of this score, which is based on information supplied by Malmö City Council and Statistics Sweden,²² we used data on the rate of migration, the percentage of residents with foreign citizenship among the ones with foreign background, the percentage of people receiving social welfare support (negative signs), and the rate of employment (positive sign). These parameters were selected in order to cover different aspects of the concept of socioeconomic deprivation in Sweden today. The four variables included in the SES were highly intercorrelated (all $r \geq 0.76$, all $p < 0.01$), indicating that they all reflect some aspect of socioeconomic deprivation. The area value for each of these four variables was standardised by subtracting from it the mean value for the city and dividing the difference by the standard deviation for all the 17 areas.²³ The sum of these standardised values is the socioeconomic score of the residential area. This score, which has been used in several other studies,^{6 20 24} correlates with other well known measures of socioeconomic conditions, for example, mean income ($r = 0.83$, $p < 0.01$) and percentage of blue collar workers ($r = -0.70$, $p < 0.01$).

Values on the percentage of people, who in each area were receiving social welfare support, are from 1991. The average value that year was 11%, ranging from 0.8% to 28.6% between areas. Area specific rates of migration, which is the percentage that per year moves within or in/out from a residential area, is similarly based on values from that year. Between areas the percentages range from 5.9% to 22.6% with an average of 15% in Malmö. Foreign background was defined as foreign citizens, Swedish citizens who were born as foreign citizens, or children under 18 years of age with one or two foreign born parents. The percentage of residents with foreign citizenship as a proportion of all citizens with foreign background was used as a measure of the integration of immigrants. It should be emphasised, however, that this measure is influenced by a number of both social and economic circumstances. In 1992 there were 48% who met these criteria, with a range of 22.6% to 63.2% between areas. In 1991 the rate of employment, which is the percentage of all inhabitants between 20 and 64 years employed in the free labour market, was 79% in Malmö, ranging from 63.1% to 94.4% in residential areas.

Area of living remained unchanged for more than 90% of the patients who died during the follow up period.

Prevalence of cardiovascular risk factors in residential areas

The Malmö Diet and Cancer cohort²⁵ was used to assess the area specific prevalence of smokers and of patients with hypertension, diabetes, and hypercholesterolaemia. This cohort consists of 28 466 residents (11 206 men and 17 260 women) aged 45–73 years who were examined at Malmö University Hospital between March 1991 and December 1996.




Figure 1 Kaplan-Meier curves of survival after discharge following treatment for first myocardial infarction. Men and women below 65, 65–74, and above 74 years of age.

Mean (SD) age was 59.2 (7.0) years for men and 57.4 (7.9) for women. The number of participants per residential area ranged from 462 (area no 8) to 3285 (area no 1) and the participants/residents ratio from 27% (area no 16) to 54% (area no 8).

Smoking habits were assessed by a questionnaire. Those who confirmed regular or occasional smoking were counted as smokers.

Blood pressure was measured twice in the right arm after five minutes rest. Subjects who confirmed the use of blood pressure lowering medication, together with subjects whose systolic or diastolic blood pressure was equal to or exceeded 160 and/or 95 mm Hg, were classified as hypertensive.

The prevalence of hypercholesterolaemia is based on a subsample of 5362 subjects (3148 women and 2214 men) who entered the cohort between October 1991 and February 1994. Subjects who reported the use of cholesterol lowering medication and subjects whose cholesterol value exceeded 6.5 mmol/l were considered to have hypercholesterolaemia.

Subjects who reported treatment for diabetes and subjects whose blood glucose level was ≥ 6.7 mmol/l were considered to have diabetes.

The cardiovascular risk factor score

These four major risk factors were used to compute an area specific cardiovascular risk factor score. This was achieved by first calculating for each area the sex adjusted and age adjusted prevalences using the equivalent average method.²⁶ Each of these values was standardised by subtracting from it the average value for Malmö and then dividing the difference by the standard deviation for the city.²³ The area score is the sum of these standardised values.

Statistical methods

Area specific rates of all cause mortality/1000 patients years after discharge have been adjusted for differences with regard to age distribution and sex distribution. The entire cohort of patients between 1986–95 was used as the standard population. Three age groups—that is, below 65, 65–74, and 75 and above—were used in these computations.

Area specific rates of mortality in ischaemic heart disease have similarly been adjusted for differences with regard to age and sex. The population of Malmö above 20 years of age 1986–95 was used as the standard population. The interval used for age stratification was five years.

The Kaplan-Meier method was used to illustrate survival in relation to age and sex. Cox proportional hazards analysis was

Table 1 Age and sex adjusted all cause mortality rates after discharge in relation to the socioeconomic and the cardiovascular risk factor scores

Residential area	Socioeconomic score	Cardiovascular risk factor score	Number of residents	Number of patients	Dead during follow up (n (%))	*Mortality rates per 1000 patient years
Low SES						
16	-7.18	0.13	18289	273	144 (52.7)	140.1
10	-5.41	-2.09	7691	110	44 (40.0)	124.4
2	-4.42	0.78	34341	651	406 (62.4)	159.8
3	-3.11	5.01	22734	600	362 (60.3)	129.1
15	-2.17	5.21	5002	130	80 (61.5)	163.6
4	-2.12	2.44	10597	233	119 (51.1)	123.8
Median SES						
13	-1.52	4.52	9557	176	89 (50.6)	152.1
12	-1.46	2.09	26212	554	281 (50.7)	125.8
9	-0.22	0.07	12174	408	243 (59.6)	138.2
1	1.54	-1.67	24297	715	432 (60.4)	125.5
5	2.43	-1.76	11266	260	141 (54.2)	159.6
High SES						
14	2.59	-3.90	10612	110	35 (31.8)	103.9
11	2.87	1.24	10527	260	130 (50.0)	128.4
6	3.89	-3.83	11078	257	137 (53.3)	128.0
17	4.44	-3.09	9321	180	74 (41.1)	99.0
7	4.84	-1.97	4582	63	24 (38.1)	129.9
8	5.01	-3.13	2839	34	14 (41.2)	85.5


*Age and sex adjusted.

used to illustrate survival rates for patients from residential areas with low, median, and high socioeconomic score after adjustment for differences with regard to age and sex. Age adjusted and sex adjusted relative risks of death after discharge in each of the 17 residential areas were calculated in a similar fashion by Cox proportional hazards analysis. Linear regression was used to illustrate to what extent these area specific relative risks of death after discharge covaried with the socioeconomic and the cardiovascular risk factor scores. The same method was used in the analysis of area specific rates of mortality in ischaemic heart disease in relation to rates of long term survival after discharge.

RESULTS

Rate of mortality after discharge in relation to age and sex

At the end of follow up 2755 of the 5014 patients were dead (54.9%). The relative risk of dying after discharge increased by 7% per year of age (RR 1.07; 95% CI 1.068 to 1.077). The lower mortality rate for women remained statistically significant after age adjustment (RR 0.9; 95% CI 0.84 to 0.99) (fig 1).

**Figure 2** Relative risk of death after discharge in relation to the socioeconomic score of residential areas. Regression line with 95% confidence intervals.

Rates of mortality after discharge in relation to the socioeconomic score

Table 1 illustrates dissimilarities between residential areas in terms of socioeconomic circumstances, the prevalence of major risk factors for cardiovascular disease, and the rates of mortality after discharge.

The age adjusted and sex adjusted all cause mortality rate following discharge after first myocardial infarction ranged between areas from 85.5 to 163.6/1000 patient years of follow up. The area specific relative risk of death after discharge covaried in a linear fashion with the socioeconomic area score, $r = -0.56$, $p = 0.018$, fig 2. Table 2 illustrates how of each of the included socioeconomic variables covaries with the risk of death after discharge. When stratified for age (<65 ($n = 1275$), $65-74$ ($n = 1516$) and >74 ($n = 2223$) years of age) survival differences after discharge among patients living in areas with high, median, and low socioeconomic scores showed statistical significance only in the age group below 65. The mortality rates in this age group were 22% and 40% lower for patients who came from areas with median (RR 0.78; 95% CI = 0.60 to 0.99, $p = 0.046$) and high socioeconomic score (RR 0.60; 95% CI = 0.44 to 0.81, $p < 0.001$), fig 3. The average follow up



**Figure 3** Survival after discharge among patients below 65 years of age from residential areas with high, median, and low socioeconomic score. Cox regression curves with adjustment for age and sex.

Table 2 Correlation coefficients (*r*) for the relation between socioeconomic circumstances in residential areas and the age and sex adjusted relative risks of death after discharge

	<i>r</i>	<i>P</i>
Migration rate (%)	0.46	0.06
Foreign citizens among foreign background (%)	0.56	<0.05
Social welfare support (%)	0.45	0.07
Employment rate (%)	-0.68	<0.01
Socioeconomic score (SES)	-0.56	<0.05

**Figure 4** Relative risk of death after discharge in relation to the cardiovascular risk factor score of residential areas. Regression line with 95% confidence intervals.


time for patients from areas with high, median, and low socioeconomic scores was very similar. Only minor differences occurred when the calculations were done separately for men and women.

Rates of mortality after discharge in relation to the cardiovascular risk factor score

The age adjusted and sex adjusted area specific relative risks of death after discharge covaried in a linear fashion with the cardiovascular risk factor scores, $r=0.64$, $p=0.005$, fig 4. According to this association, 41% of the intra-urban variance of mortality after discharge can be accounted for by covariance with the cardiovascular risk factor score. The corresponding value for the socioeconomic score was 32% and together the two scores accounted for 48 % of the variance of mortality following discharge, table 3.

Relation between area specific rate of mortality in ischaemic heart disease and survival following discharge after first myocardial infarction

The area specific relative risks of death following discharge after first myocardial infarction covaried in a statistically sig-

**Figure 5** Age and sex adjusted rates of mortality from ischaemic heart disease per 100 000 in relation to the relative risk of death after discharge. Regression line with 95% confidence intervals.

nificant fashion with the rates of mortality in ischaemic heart disease, $r=0.64$, $p=0.006$, fig 5. Forty one per cent of the intra-urban variance in mortality from ischaemic heart disease can be accounted for by the variance in survival following discharge after first myocardial infarction according to this association.

DISCUSSION

Five years after discharge less than half of the patients were alive. Our results are compatible with the hypothesis that the socioeconomic environment may play an important part in the survival rate of patients with myocardial infarction and that the prognosis is poorer for patients from less affluent areas. Considering the ecological design it is however not appropriate to make any causal inferences.

Further studies are needed to explore whether the association with socioeconomic circumstances could have been confounded by covariance with other risk factors of importance for the long term prognosis, for example, the severity of coronary atherosclerosis, the degree of damage to the myocardium, the use and compliance with secondary preventive measures. Major risk factors for cardiovascular disease were more prevalent in areas with low socioeconomic score and low rates of survival. The important issue from a preventive perspective, however, is that, between groups defined in terms of residential area, there were marked differences in the age adjusted and sex adjusted rate of mortality after discharge and that these areas can be described in terms of socioeconomic circumstances and prevalence of major risk factors for cardiovascular disease. Differences in ischaemic heart disease mortality may be related to incidence,⁴ short-term case fatality rate,⁵ and survival after hospitalisation. It is our conclusion that in this city differences, in terms of long term survival after discharge, significantly contribute to the pattern of ischaemic heart disease mortality. This is an important observation in evaluating the preventive potential. An inverse relation

Table 3 Percentage of the intra-urban variance in survival after discharge accounted for by the socioeconomic and the cardiovascular risk factor scores

	<i>r</i>	<i>r</i> ² × 100 (%)	<i>P</i>
Cardiovascular risk factor score	0.64	41	0.005
Socioeconomic score (SES)	-0.56	32	0.018
Cardiovascular risk factor score + SES*	0.69	48	0.011

*Correlation between cardiovascular risk factor score and socioeconomic score: $r=-0.55$, $r^2=0.31$, $p=0.021$.

Key points

- Geographical differences in mortality from ischaemic heart disease imply a preventive potential.
- Long term survival after first myocardial infarction is in this urban population significantly worse for patients from less affluent residential areas.
- In this city differences in long term survival after first myocardial infarction significantly contribute to the geographical pattern of mortality from ischaemic heart disease.

between age and rate of survival did, as expected, dilute the influence of the socioeconomic score on the rate of survival in older age groups. The higher survival rate for patients from areas with median and high score was statistically significant only for those who were below 65 years of age. This is consistent with earlier studies showing that the relative influence of socioeconomic^{27, 28} as well as biological^{29, 30} risk factors on incidence and survival decreases with age.^{29, 31, 32}

After discharge the patients were taken care of by primary healthcare physicians. A number of trials have reported the benefits of secondary preventive measures—that is, treatment of hypertension and hyperlipidaemia, help to quit smoking, and antithrombotic treatment. Yet it seems that far from all patients receive appropriate care.³³ Studies from Italy have reported a lack of standardisation with regard to risk stratification strategies after infarction.³⁴ In studies from Sweden it has been shown that approximately 50% of patients treated for hypertension still have diastolic blood pressure exceeding the national treatment goal of <90 mm Hg.³⁵ Whether this proportion differs between groups defined in terms of their socioeconomic circumstances remains to be evaluated. The ability to quit and remain free from smoking seems on the other hand to be related to level of education and annual income.³⁶

Differences with regard to the severity of coronary atherosclerosis and damage to the myocardium could explain why the survival curves for patients from areas with high, median, and low socioeconomic scores started to deviate already during the first year after discharge. However, as the curves of survival continued to deviate during the entire follow up period it seems reasonable to assume that rates of survival in groups defined in terms of residential area may be attributable to some extent to differences with regard to the use and compliance with secondary preventive measures.

Many of the sociodemographic circumstances associated with the rate of survival have been identified as independent risk factors for the incidence of myocardial infarction in prospective cohort studies.^{37, 38} According to migrant studies it seems that the sociodemographic environment in itself may influence a person's susceptibility.³⁹ Whether these associations are applicable to patients being discharged after myocardial infarction, remains to be evaluated.

Results from both observational studies and clinical trials suggest that giving myocardial infarction patients emotional and social support can improve the survival rate.^{40–44} The true nature of this relation is not known. According to a mailed questionnaire survey it has been shown that, between residential areas in the city of Malmö, there are marked differences with regard to the availability of emotional support and degree of social anchorage.^{45, 46}

Some methodological issues should be considered. The follow up of patients with regard to residential area before and after infarction did not indicate that the pattern of survival could have been confounded by misclassification with regard to residential area. Residential area remained unchanged for more than 90% of the patients who died during the follow up period. All patients were treated at Malmö University Hospital, which is the only hospital for myocardial infarction and other

somatic disorders in the city. The National Board of Health and Welfare has shown that the diagnosis of myocardial infarction collected via the patient administrative register is correct in 90%–95% of the cases.¹⁸ Previous long term follow up studies from the register have shown that the validity regarding vital status is close to 100%.²⁴ It thus seems unlikely that the results could be explained by geographical differences with regard to case retrieval and validation of cases and deaths.

The survival rate in our patient cohort may, compared with survival rates reported from Minnesota,⁴⁷ the SPRINT registry,⁴⁸ and randomised mega trials of thrombolytic therapy^{49, 50} seem lower than expected. However, because of differences in regard to age distribution and criteria used for inclusion and exclusion of patients, results are not readily comparable.

In this follow up of survival after myocardial infarction, 55% of the patients were no longer alive five years later. Rate of mortality after discharge was significantly higher for patients from residential areas with inferior socioeconomic circumstances. Differences in terms of long term survival contributed to the geographical pattern of mortality from ischaemic heart disease in a significant fashion. In order to exploit the preventive potential, the extent to which socioeconomic circumstances covary with severity of disease, respectively with the use and compliance with secondary preventive measures, needs to be evaluated.

Authors' affiliations

P Tydén, G Engström, B Hedblad, L Janzon, Department of Community Medicine, Malmö University Hospital, Lund University, Malmö, Sweden

O Hansen, Department of Cardiology, Malmö University Hospital

Funding: this study was supported by grants from the Swedish Council for Social Research.

Conflicts of interest: none.

REFERENCES

- 1 Sans S, Kesteloot H, Kromhout D. The burden of cardiovascular diseases mortality in Europe. *Eur Heart J* 1997;**18**:1213–48.
- 2 Nerbrand C, Svärdsudd K, Hörte LG, et al. Geographical variation of mortality from cardiovascular diseases. The project "Myocardial Infarction in mid-Sweden". *Eur Heart J* 1991;**12**:4–9.
- 3 Hammar N, Ahlbom A, Theorell T. Geographical differences in myocardial infarction incidence in eight Swedish counties, 1976–1981. *Epidemiology* 1992;**3**:348–55.
- 4 Tydén P, Hansen O, Janzon L. Intra-urban variations in incidence and mortality in myocardial infarction. A study from the myocardial infarction register in the city of Malmö, Sweden. *Eur Heart J* 1998;**19**:1795–801.
- 5 Tydén P, Engström G, Hansen O, et al. Geographical pattern of female deaths from myocardial infarction in an urban population: fatal outcome out-of-hospital related to socio-economic deprivation. *J Intern Med* 2001;**250**:201–7.
- 6 Engström G, Berglund G, Göransson M, et al. Distribution and determinants of ischaemic heart disease in an urban population. A study from the myocardial infarction register in Malmö, Sweden. *J Intern Med* 2000;**247**:588–96.
- 7 Volpi A, De Vita C, Franzosi MG, et al. Determinants of 6-month mortality in survivors of myocardial infarction after thrombolysis. Results of the GISSI-2 database. *Circulation* 1993;**88**:416–29.
- 8 Fibrinolytic Therapy Trialists (FTT) Collaborative Group: Indications for fibrinolytic therapy in suspected acute myocardial infarction: collaborative overview of early mortality and major morbidity results from all randomised trials of more than 1000 patients. *Lancet* 1994;**343**:311–22.
- 9 Michels KB, Yusuf S. Does PTCA in acute myocardial infarction affect mortality and reinfarction rates? A quantitative overview (meta-analysis) of the randomised clinical trials. *Circulation* 1995;**91**:476–85.
- 10 ISIS-2 (Second International Study of Infarct Survival) Collaborative Group. Randomised trial of intravenous streptokinase, oral aspirin, both, or neither among 17 187 cases of suspected myocardial infarction: ISIS-2. *Lancet* 1988;ii:349–60.
- 11 Ades PA, Pashkow FJ, Nestor JR. Cost-effectiveness of cardiac rehabilitation after myocardial infarction. *J Cardiopulm Rehabil* 1997;**17**:222–31.
- 12 Yusuf S, Peto R, Lewis J, et al. Beta-blockade during and after myocardial infarction: an overview of the randomised trials. *Progr Cardiovasc Dis* 1985;**17**:335–71.
- 13 Pfeffer MA, Braunwald E, Moyé LA, et al. Effect of captopril on mortality and morbidity in patients with left ventricular dysfunction after myocardial infarction. *N Engl J Med* 1992;**327**:669–77.

- 14 **Alistair SH**, Murray GD, Ball SG on behalf of the Airex Study Investigators: Follow-up study of patients randomly allocated ramipril or placebo for heart failure after myocardial infarction: AIRE Extension (AIREX) Study. *Lancet* 1997;**349**:1493–7.
- 15 **Scandinavian Simvastatin Survival Study Group**. Randomised trial of cholesterol lowering in 4444 patients with coronary heart disease: the Scandinavian Simvastatin Survival Study (4S). *Lancet* 1994;**344**:1383–9.
- 16 **Sacks FM**, Pfeffer MA, Moye LA, *et al*. The effect of pravastatin on coronary events after myocardial infarction in patients with average cholesterol levels. *N Engl J Med* 1996;**335**:1001–9.
- 17 **Hansen O**, Johansson BW. Epidemiologic aspects of coronary heart disease in Malmö 1935–1988. *Am J Epidemiol* 1991;**133**:721–33.
- 18 **The National Board of Health and Welfare**. *Värdering av diagnoskvaliteten för akut hjärtinfarkt i tre svenska län 1995*. Artikelnr 1997–84–8. Stockholm: 1997.
- 19 **Mosca L**, Manson JE, Sutherland SE, *et al*. Cardiovascular disease in women. A statement for healthcare professionals from the American Heart Association. *Circulation* 1997;**96**:2468–82.
- 20 **Engström G**, Jerntorp I, Pessah-Rasmussen H, *et al*. Geographic distribution of stroke incidence within an urban population: relations to socioeconomic circumstances and prevalence of cardiovascular risk factors. *Stroke* 2001;**32**:1098–103.
- 21 **Manjer J**, Berglund G, Bondesson L, *et al*. Intra-urban differences in breast cancer mortality: a study from the city of Malmö in Sweden. *J Epidemiol Community Health* 2000;**54**:279–85.
- 22 **The Unit of Planning and Statistics**. *Områdesfakta för Malmö 1991–1996*. Malmö: Malmö City Council, 1991–1996.
- 23 **Woodward M**. Small area statistics as markers for personal social status in the Scottish heart health study. *J Epidemiol Community Health* 1996;**50**:570–6.
- 24 **Engström G**, Göransson M, Hansen O, *et al*. Trends in long-term survival after myocardial infarction: less favourable patterns for patients from deprived areas. *J Intern Med* 2000;**248**:425–34.
- 25 **Berglund G**, Elmstahl S, Janzon, *et al*. Design and feasibility. *J Intern Med* 1993;**233**:45–51.
- 26 **Merlo J**, Ranstam J, Råstam L, *et al*. Age standardisation of drug utilisation: comparisons of different methods using cardiovascular drug data from Sweden and Spain. *Eur J Clin Pharmacol* 1994;**46**:393–8.
- 27 **Peltonen M**, Rosén M, Lundberg V, *et al*. Social patterning of myocardial infarction and stroke in Sweden: incidence and survival. *Am J Epidemiol* 2000;**151**:283–92.
- 28 **Kunst AE**, Mackenbach JP. The size of mortality differences associated with educational level in nine industrialized countries. *Am J Public Health* 1994;**84**:932–7.
- 29 **Anderson KM**, Wilson P, Odell PM, *et al*. An updated coronary risk profile. A statement for health professionals. *Circulation* 1991;**83**:356–62.
- 30 **Dawber TR**. *The Framingham study: the epidemiology of atherosclerotic disease*. Cambridge, MA: Harvard University Press, 1980.
- 31 **Wannamethee G**, Whincup P, Shaper AG, *et al*. Factors determining case fatality in myocardial infarction “Who dies in a heart attack?” *Br Heart J* 1995;**74**:324–31.
- 32 **Njölstad I**, Arnesen E. Preinfarction blood pressure and smoking are determinants for fatal outcome of myocardial infarction. *Arch Intern Med* 1998;**158**:1326–32.
- 33 **Euroaspire Study Group**. Euroaspire. A European Society of Cardiology survey of secondary prevention of coronary heart disease: Principal results. *Eur Heart J* 1997;**18**:1569–82.
- 34 **Maggioni AP**, Tavazzi L, Fabbri G, *et al*. Epidemiology of post-infarction risk stratification strategies in a country with low volume of revascularisation procedures. The GISSI Prognosis Registry. *Eur Heart J* 1998;**19**:1784–94.
- 35 **Nilsson P**, Andersson DKG, Andersson P-E, *et al*. Cardiovascular risk factors in treated hypertensives – a nationwide, cross-sectional study in Sweden. *J Intern Med* 1993;**233**:239–45.
- 36 **Engström G**, Tydén P, Berglund G, *et al*. Incidence of myocardial infarction in women. A cohort study of risk factors and modifiers of effect. *J Epidemiol Community Health* 2000;**54**:104–7.
- 37 **Suadicani P**, Hein HO, Gyntelberg F. Serum validated tobacco use and social inequalities in risk of ischemic heart disease. *Int J Epidemiol* 1994;**23**:293–300.
- 38 **Holme I**, Helgeland A, Hjermann I, *et al*. Four year mortality by some socioeconomic indicators: the Oslo study. *J Epidemiol Community Health* 1980;**34**:48–52.
- 39 **Marmot MG**, Syme SL, Kagan A, *et al*. Epidemiologic studies of coronary heart disease and stroke in Japanese men living in Japan, Hawaii, and California: prevalence of coronary and hypertensive heart disease and associated risk factors. *Am J Epidemiol* 1975;**102**:514–25.
- 40 **Carinci F**, Nicolucci A, Ciampi A, *et al*. Role of interactions between psychological and clinical factors in determining 6-month mortality among patients with acute myocardial infarction. *Eur Heart J* 1997;**18**:835–45.
- 41 **Thomas SA**, Friedmann E, Wimbush F, *et al*. Psychological factors and survival in the cardiac arrhythmia trial (CAST): a reexamination. *Am J Crit Care* 1997;**6**:116–26.
- 42 **Kawachi I**, Colditz GA, Ascherio A, *et al*. A prospective study of social networks in relation to total mortality and cardiovascular disease in men in the USA. *J Epidemiol Community Health* 1996;**50**:245–51.
- 43 **Friedmann E**, Thomas SA. Pet ownership, social support, and one-year survival after acute myocardial infarction in the Cardiac Arrhythmia Suppression Trial (CAST). *Am J Cardiol* 1995;**76**:1213–17.
- 44 **Berkman LF**, Leo-Summers L, Horwitz RI. Emotional support and survival after myocardial infarction. A prospective, population-based study of the elderly. *Ann Intern Med* 1992;**117**:1003–9.
- 45 **Nilsson Å**, Hanson BS, Isacson S-O. *Public Health Report – Malmö County 1991 (Folkhälsorapport – Malmö kommun 1991)*. Malmö: Department of Community Medicine, Malmö University Hospital, 1991.
- 46 **Lindström M**, Bexell A, Hanson BS, *et al*. *Public Health Report – Malmö County 1994 (Folkhälsorapport – Malmö kommun 1994)*. Malmö: Department of Community Medicine, Malmö University Hospital, 1994.
- 47 **Orencia A**, Bailey K, Yawn BP, *et al*. Effect of gender on long-term outcome of angina pectoris and myocardial infarction/sudden unexpected death. *JAMA* 1993;**269**:2392–7.
- 48 **Benderly M**, Behar S, Reicher-Reiss H, *et al* for the SPRINT study Group. Long-term prognosis of women after myocardial infarction. *Am J Epidemiol* 1997;**146**:153–60.
- 49 **Baigent C**, Collins R, Appleby P, *et al*. ISIS-2: 10 year survival among patients with suspected acute myocardial infarction in randomised comparison of intravenous streptokinase, oral aspirin, both, or neither. The ISIS-2 (Second International Study of Infarct Survival) Collaborative Group. *BMJ* 1998;**316**:1337–43.
- 50 **Franzosi MG**, Santoro E, De Vita C, *et al*. Ten-year follow up of the first megatrial testing thrombolytic therapy in patients with acute myocardial infarction. *Circulation* 1998;**98**:2659–65.