



# LUND UNIVERSITY

## Neonatal ACE inhibition in rats interferes with lung development.

Lasaitiene, Daina; Chen, Yun; Nannmark, Ulf; Wollmer, Per; Friberg, Peter

*Published in:*  
Clinical Physiology and Functional Imaging

*DOI:*  
[10.1046/j.1475-0961.2003.00530.x](https://doi.org/10.1046/j.1475-0961.2003.00530.x)

2004

[Link to publication](#)

*Citation for published version (APA):*  
Lasaitiene, D., Chen, Y., Nannmark, U., Wollmer, P., & Friberg, P. (2004). Neonatal ACE inhibition in rats interferes with lung development. *Clinical Physiology and Functional Imaging*, 24(1), 65-68.  
<https://doi.org/10.1046/j.1475-0961.2003.00530.x>

*Total number of authors:*  
5

### General rights

Unless other specific re-use rights are stated the following general rights apply:  
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117  
221 00 Lund  
+46 46-222 00 00

# Neonatal ACE inhibition in rats interferes with lung development

Daina Lasaitiene<sup>1</sup>, Yun Chen<sup>1</sup>, Ulf Nannmark<sup>2</sup>, Per Wollmer<sup>3</sup> and Peter Friberg<sup>1,4</sup>

<sup>1</sup>Department of Physiology and <sup>2</sup>Department of Anatomy and Cell Biology, University of Gothenburg, Gothenburg, Sweden, <sup>3</sup>Department of Clinical Physiology, Malmo University Hospital, Malmo, Sweden and <sup>4</sup>Department of Clinical Physiology, Sahlgrenska University Hospital, Gothenburg, Sweden

## Summary

### Correspondence

Daina Lasaitiene, Department of Physiology,  
Institute of Physiology and Pharmacology,  
University of Gothenburg, Box 432, SE-405 30,  
Gothenburg, Sweden  
E-mail: daina.lasaitiene@fysiologi.gu.se

### Accepted for publication

Received 11 April 2003;  
accepted 15 October 2003

### Key words

angiotensin-converting enzyme; lung development;  
renin–angiotensin system; surface tension; Wilhelmy  
balance

The renin–angiotensin system (RAS) is developmentally up-regulated and it is essential for kidney development in several species. Given the fact that the rat lung undergoes postnatal development, the mammalian lung possesses the highest angiotensin-converting enzyme (ACE) levels and ACE activity increases during the first weeks postpartum, we tested the hypothesis that ACE inhibition influences postnatal lung development. Rats were given the ACE inhibitor enalapril ( $10 \text{ mg kg}^{-1}$ ) from 0 to 9 days of age and their lungs were examined at day 4 and 9. Lung structure was evaluated by means of light microscopy, and surface tension of bronchoalveolar lavage fluid was measured by means of a Wilhelmy balance. Neonatal ACE inhibition lowered the surface tension of bronchoalveolar lavage fluid and caused widening of respiratory airspaces and thinning of alveolar septa. Our results suggest that early postnatal ACE inhibition in rats interferes with lung development.

## Introduction

The activity and distribution of the renin–angiotensin system (RAS) are developmentally regulated in a tissue-specific manner, and angiotensin II (Ang II) can act as a modulator of growth in a variety of cells and tissues (Gomez, 1990; Touyz & Schiffrin, 2000). The RAS has also been shown to be critically important in development. There is ample evidence for its mandatory role in kidney development (Friberg *et al.*, 1994; Tufro-McReddie *et al.*, 1995; Guron & Friberg, 2000). Different components of the RAS, including angiotensin-converting enzyme (ACE) and both angiotensin type 1 ( $\text{AT}_1$ ) and type 2 ( $\text{AT}_2$ ) receptors, are expressed in lung endothelial and type II epithelial cells, respectively (Morrell *et al.*, 1996; Wang *et al.*, 1999). The mammalian lung possesses the highest ACE levels (Esther *et al.*, 1996) and ACE activity increases during the first weeks postpartum (Wallace *et al.*, 1978). Rats are born with immature lungs and, within the first two postnatal weeks, the alveolization (Scheuermann *et al.*, 1988), together with subcellular reorganization in type II epithelial cells (Young *et al.*, 1991) and changes in surfactant homeostasis (Griese *et al.*, 1999), occurs in rats. The aim of this study was to test the hypothesis that ACE inhibition (ACEI) influences lung development. Hence, rats were given ACEI during 9 days starting at day 0 after birth and

their lungs were examined at day 4 and 9. Structural changes in ACEI lung were evaluated at light microscopical level, and surface tension of bronchoalveolar lavage fluid (BALF), a reflector of quantitative and/or qualitative abnormalities of alveolar surfactant, was measured by means of a Wilhelmy balance (Schmekel *et al.*, 1992).

## Methods

### General procedures

Time-mated, female Wistar rats (B & K Universal AB, Sollentuna, Sweden) were transported to our facility on the 14th day of pregnancy. They were observed carefully for determination of the day of delivery. Both male and female pups were included in the study. Weight matched pups were divided into two groups receiving daily intraperitoneal injections of either enalapril maleate ( $10 \text{ mg kg}^{-1}$ ; Sigma-Aldrich Sweden AB) or isotonic saline vehicle in equivalent volumes of  $10 \text{ ml kg}^{-1}$  from day 0 (within 12 h after birth) to day 9. We chose  $10 \text{ mg kg}^{-1}$  dose of enalapril maleate, which was demonstrated to inhibit nearly completely ACE activity (Guron *et al.*, 1999). Throughout the study rats had free access to standard rat chow and tap water and were kept in rooms with a controlled temperature of  $24^\circ\text{C}$  and a

12 h dark (6 p.m.–6 a.m.)/light cycle. The regional animal ethics committee in Gothenburg approved all experiments.

### Light microscopy and semiquantitative assessment

For microscopical evaluation, seven to nine pups from enalapril maleate and saline treated group were anaesthetized with sodium pentobarbital ( $60 \text{ mg kg}^{-1}$ , intraperitoneally; Apoteks-bolaget, Umeå, Sweden) at day 4 and 9 after birth. After thoracotomy, the lungs were excised and immersion-fixed in 4% paraformaldehyde. The upper, middle and lower part of the left lung was cross-dissected into  $\approx 4 \text{ mm}$  thick planes and their fixation was continued for 24 h. After fixation planes were dehydrated through an ethanol series and all three planes from each lung were embedded in paraffin en bloc. Keeping the distance of  $50 \mu\text{m}$  in between, three  $3\text{-}\mu\text{m}$ -thick serial sections were cut and stained with haematoxylin and eosin. Nine sections from each lung were used for the semiquantitative assessment of lung structural changes. Assessments were made by an investigator blind to treatment group. The presence of wider respiratory airspaces and thinner alveolar septa was scored semiquantitatively with the use of arbitrary scale: 1, mild; 2, moderate; and 3, severe changes.

### Measurement of surface tension of bronchoalveolar lavage fluid

For measurements of surface tension of BALF, seven to nine pups from enalapril maleate and saline treated groups were anaesthetized with sodium pentobarbital ( $60 \text{ mg kg}^{-1}$ , i.p.) at day 4 and 9 after birth. After thoracotomy the lungs of each animal were washed gently three times with  $37^\circ\text{C}$  saline ( $30 \text{ ml kg}^{-1}$ ) and BALF aliquots were pooled, yielding in  $2\text{--}2.5 \text{ ml}$  of lavage fluid. The lavage fluid was centrifuged at  $150 \text{ g}$  for 10 min to remove any cells and frozen at  $-20^\circ\text{C}$ . Surface tension of BALF was measured with a Wilhelmy balance (Biegler Electronic, Mauerbach, Austria). A teflon trough ( $115 \times 18 \text{ mm}$ , depth  $17 \text{ mm}$ ) was filled with  $35 \text{ ml}$  of saline. The measurements were made at room temperature. A platinum plate, partly immersed in the liquid, is suspended in a balance. The trough contains a movable barrier, which changes the area into which the platinum plate is immersed. The duration of a cyclic area change to 20% of the initial area is 3 min. A baseline recording of the surface tension of saline was first established. The surface tension of saline is  $73 \text{ mN m}^{-1}$  and is independent of area change. A  $200 \mu\text{l}$  of lavage fluid was added to the trough. A stable recording of surface tension versus area was rapidly obtained, and the maximum ( $\gamma_{\text{max}}$ ) and minimum ( $\gamma_{\text{min}}$ ) surface tension of the third cycle were recorded, thus, yielding in total two measurements on each sample of lavage fluid. We measured the surface tension of alveolar surfactant rather than its amount or composition since the surface tension reflects quantitative and/or qualitative abnormalities of alveolar surfactant.

At the end of experiments rats were killed by an overdose of sodium pentobarbital.

### Statistical analysis

Values are expressed as mean values  $\pm$  SD. All data were analysed by ANOVA followed by Fisher's *post hoc* least significant difference (LSD) test, using the computer software SYSTAT (version 5.02; Evanston, IL, USA).  $P < 0.05$  was considered statistically significant.

## Results

### Morphological and semiquantitative observations

Enalapril treatment induced structural alterations in the lungs on postnatal days 4 and 9. Wider respiratory airspaces and thinner alveolar septa were observed in enalapril-treated rats (Figs 1b, d and 2), but it remains open how much of it reflects a real widening of the airspaces and how much is caused by the thinning of the septa.


### Surface tension of bronchoalveolar lavage fluid

Enalapril treatment lowered minimum ( $\gamma_{\text{min}}$ ) surface tension of BALF, as measured with  $200 \mu\text{l}$  quantity (Fig. 3). The maximum ( $\gamma_{\text{max}}$ ) surface tension of BALF did not differ between groups either on postnatal day 4 ( $70.3 \pm 1.7$  versus  $68.4 \pm 2.6 \text{ mN m}^{-1}$ , in control and enalapril treated rats, respectively) or day 9 ( $69.4 \pm 3.2$  versus  $69.9 \pm 2.1 \text{ mN m}^{-1}$ , in control and enalapril treated rats, respectively).

## Discussion

The main finding of this study was that ACEI decreased surface tension of BALF, reflecting higher amount of alveolar surfactant, in neonatal rats on day 4 and 9. This was paralleled by structural alterations in the lungs. Thus, early postnatal ACE inhibition in rats interferes with lung development.

Surface tension of BALF, as measured in this study, mirrors the functional integrity of alveolar surfactant system. Pulmonary surfactant is a surface-active material composed of phospholipid-rich lipoproteins that reduce surface tension at the alveolar surface, promoting lung expansion during inspiration and preventing lung collapse during expiration (De Paepe et al., 1998). The lower minimum surface tension of BALF in enalapril treated rats suggests that there is more surface-active surfactant in alveolar space, which, in turn, could result in lung structural alterations, characterized by thinner alveolar septa and wider respiratory airspaces, as evident from our study. Pulmonary surfactant is synthesized, stored, secreted, taken up and catabolized by type II pneumocytes (Griese et al., 1999). The lamellar body within type II pneumocyte is the final secretory structure of pulmonary surfactant (Voorhout et al., 1993). In a pilot study, we evaluated the morphology of type II pneumocytes at ultrastructural level and observed no alterations in enalapril-subjected lungs, implying preserved intracellular stores of



**Figure 1** (a)–(d) Lung sections from 4 days-old (a, b) and 9 days-old (c, d) rats treated with saline vehicle (a, c) or enalapril (b, d). Enalapril-treated rat shows thinner alveolar septa and wider respiratory airspaces. Magnification 10X.


surfactant (data not shown). Collectively, it appears that early postnatal ACE inhibition increases the alveolar surfactant pool.

At the present time, we could only speculate about the mechanism of the changes we observed because we have not established whether the alveolar surfactant pool is increased due to enalapril-stimulated secretion or enalapril-inhibited uptake or both. Our finding of the disturbance in alveolar surfactant system in enalapril-treated rats suggests a role for ACE in the regulation of surfactant homeostasis in developing rat lungs. In this context, there is evidence to suggest that ACE inhibition results in increased levels of bradykinin and/or prostaglandins (PGs) (Buttar, 1997). Prostaglandins, in turn, enhance


production and secretion of surfactant by type II pneumocytes (Oyarzun & Clements, 1978; Nagai *et al.*, 1995). Therefore, it is possible that the disturbance in surfactant homeostasis following neonatal ACEI is due to increased PGs, such as  $\text{PGE}_2$  (Marin-Castano *et al.*, 2002). Indeed, rat  $\text{PGE}_2$  receptors are expressed in lung tissue (Nemoto *et al.*, 1997), and  $\text{PGE}_2$  is a potent local mediator of cell growth and differentiation in various tissues, and it plays a role in early postnatal lung development (Nagai *et al.*, 1995).

In summary, ACE regulates surfactant homeostasis in immature rat lung and, therefore, may play an important role in lung development.





**Figure 2** Semiquantitative assessment of lung structural changes, expressed as a number of points, in enalapril and saline vehicle treated neonatal rats at 4 and 9 days of age. The presence of wider respiratory airspaces and thinner alveolar septa was scored semiquantitatively with the use of arbitrary scale: 1, mild; 2, moderate; 3, severe changes. Nine sections per each lung were evaluated. \* $P < 0.05$ , indicating a statistically significant overall effect of enalapril versus saline vehicle. Values are mean values  $\pm$  SD ( $n = 7-9$  per group and time point).



**Figure 3** Minimum surface tension of bronchoalveolar lavage fluid in enalapril and saline vehicle treated neonatal rats at 4 and 9 days of age. \* $P < 0.05$ , indicating a statistically significant overall effect of enalapril versus saline vehicle. Values are mean values  $\pm$  SD ( $n = 7-9$  per group and time point).

## Acknowledgments

This study was supported by the Swedish Medical Research Council (Grants 9047 and 10841) and IngaBritt and Arne Lundberg Research Foundation. The technical assistance of Yvonne Josefsson and Gunnel Bokhede is gratefully acknowledged.

## References

- Buttar HS. An overview of the influence of ACE inhibitors on fetal-placental circulation and perinatal development. *Mol Cell Biochem* (1997); **176**: 61–71.
- De Paep ME, Papadakis K, Johnson BD, Luks FI. Fate of the type II pneumocyte following tracheal occlusion in utero: a time-course study in fetal sheep. *Virchows Arch* (1998); **432**: 7–16.

- Esther CR Jr, Howard TE, Zhou Y, Capecchi MR, Marrero MB, Bernstein KE. Lessons from angiotensin-converting enzyme-deficient mice. *Curr Opin Nephrol Hypertens* (1996); **5**: 463–467.
- Friberg P, Sundelin B, Bohman SO et al. Renin-angiotensin system in neonatal rats: induction of a renal abnormality in response to ACE inhibition or angiotensin II antagonism. *Kidney Int* (1994); **45**: 485–492.
- Gomez RA. Molecular biology of components of the renin-angiotensin system during development. *Pediatr Nephrol* (1990); **4**: 421–423.
- Griese M, Beck J, Feuerhake F. Surfactant lipid uptake and metabolism by neonatal and adult type II pneumocytes. *Am J Physiol* (1999); **277**: L901–L909.
- Guron G, Friberg P. An intact renin-angiotensin system is a prerequisite for normal renal development. *J Hypertens* (2000); **18**: 123–137.
- Guron G, Marcussen N, Nilsson A, Sundelin B, Friberg P. Postnatal time frame for renal vulnerability to enalapril in rats. *J Am Soc Nephrol* (1999); **10**: 1550–1560.
- Marin-Castano ME, Schanstra JP, Neau E et al. Induction of functional bradykinin b(1)-receptors in normotensive rats and mice under chronic Angiotensin-converting enzyme inhibitor treatment. *Circulation* (2002); **105**: 627–632.
- Morrell NW, Grieshaber SS, Danilov SM, Majack RA, Stenmark KR. Developmental regulation of angiotensin converting enzyme and angiotensin type 1 receptor in the rat pulmonary circulation. *Am J Respir Cell Mol Biol* (1996); **14**: 526–537.
- Nagai A, Katayama M, Thurlbeck WM, Matsui R, Yasui S, Konno K. Effect of indomethacin on lung development in postnatal rats: possible role of prostaglandin in alveolar formation. *Am J Physiol* (1995); **268**: L56–L62.
- Nemoto K, Pilbeam CC, Bilak SR, Raisz LG. Molecular cloning and expression of a rat prostaglandin E2 receptor of the EP2 subtype. *Prostaglandins* (1997); **54**: 713–725.
- Oyarzun MJ, Clements JA. Control of lung surfactant by ventilation, adrenergic mediators, and prostaglandins in the rabbit. *Am Rev Respir Dis* (1978); **117**: 879–891.
- Scheuermann DW, Van Meir F, Adriaensen D, Timmermans JP, De Groodt-Lasseel MH. Development of alveolar septa and formation of alveolar pores during the early postnatal period in the rat lung. *Acta Anat (Basel)* (1988); **131**: 249–261.
- Schmekel B, Bos JA, Khan AR, Wohlfart B, Lachmann B, Wollmer P. Integrity of the alveolar-capillary barrier and alveolar surfactant system in smokers. *Thorax* (1992); **47**: 603–608.
- Touyz RM, Schiffrin EL. Signal transduction mechanisms mediating the physiological and pathophysiological actions of angiotensin II in vascular smooth muscle cells. *Pharmacol Rev* (2000); **52**: 639–672.
- Tufro-McReddie A, Romano LM, Harris JM, Ferder L, Gomez RA. Angiotensin II regulates nephrogenesis and renal vascular development. *Am J Physiol* (1995); **269**: F110–F115.
- Voorhout WF, Weaver TE, Haagsman HP, Geuze HJ, Van Golde LM. Biosynthetic routing of pulmonary surfactant proteins in alveolar type II cells. *Microsc Res Tech* (1993); **26**: 366–373.
- Wallace KB, Bailie MD, Hook JB. Angiotensin-converting enzyme in developing lung and kidney. *Am J Physiol* (1978); **234**: R141–R145.
- Wang R, Zagariya A, Ibarra-Sunga O et al. Angiotensin II induces apoptosis in human and rat alveolar epithelial cells. *Am J Physiol* (1999); **276**: L885–L889.
- Young SL, Fram EK, Spain CL, Larson EW. Development of type II pneumocytes in rat lung. *Am J Physiol* (1991); **260**: L113–L122.