



# LUND UNIVERSITY

## Mechanisms of exocytosis in insulin-secreting B-cells and glucagon-secreting A-cells.

Barg, Sebastian

*Published in:*  
Pharmacology and Toxicology

*DOI:*  
[10.1034/j.1600-0773.2003.920102.x](https://doi.org/10.1034/j.1600-0773.2003.920102.x)

2003

[Link to publication](#)

*Citation for published version (APA):*  
Barg, S. (2003). Mechanisms of exocytosis in insulin-secreting B-cells and glucagon-secreting A-cells. *Pharmacology and Toxicology*, 92(1), 3-13. <https://doi.org/10.1034/j.1600-0773.2003.920102.x>

*Total number of authors:*  
1

### General rights

Unless other specific re-use rights are stated the following general rights apply:  
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117  
221 00 Lund  
+46 46-222 00 00

# Mechanisms of Exocytosis in Insulin-Secreting B-Cells and Glucagon-Secreting A-Cells

Sebastian Barg

Department of Physiological Sciences, Molecular and Cellular Physiology, Lund University, Sölvegatan 19, BMC F11, S-221 84 Lund, Sweden

(Received November 28, 2001; Accepted June 10, 2002)

**Abstract:** In pancreatic B- and A-cells, metabolic stimuli regulate biochemical and electrical processes that culminate in  $\text{Ca}^{2+}$ -influx and release of insulin or glucagon, respectively. Like in other (neuro)endocrine cells,  $\text{Ca}^{2+}$ -influx triggers the rapid exocytosis of hormone-containing secretory granules. Only a small fraction of granules (<1% in insulin-secreting B-cells) can be released immediately, while the remainder requires translocation to the plasma membrane and further “priming” for release by several ATP- and  $\text{Ca}^{2+}$ -dependent reactions. Such functional organization may account for systemic features such as the biphasic time course of glucose-stimulated insulin secretion. Since this release pattern is altered in type-2 diabetes mellitus, it is conceivable that disturbances in the exocytotic machinery underlie the disease. Here I will review recent data from our laboratory relevant for the understanding of these processes in insulin-secreting B-cells and glucagon-secreting A-cells and for the identification of novel targets for antidiabetic drug action. Two aspects are discussed in detail: 1) The importance of a tight interaction between L-type  $\text{Ca}^{2+}$ -channels and the exocytotic machinery for efficient secretion; and 2) the role of intragranular acidification for the priming of secretory granules and its regulation by a granular 65-kDa sulfonyleurea-binding protein.

## The importance of insulin

The introduction of a novel medical treatment has rarely had such a dramatic impact on the patient's life as the introduction and use of the blood glucose-lowering hormone insulin. Within two years of its discovery in 1921, insulin was available in large quantities and used routinely in the treatment of diabetes mellitus. It took only a few weeks of insulin treatment to revitalize severely starved diabetic patients, illustrating the fundamental role the hormone plays in the metabolic regulation of the body. Diabetes mellitus is characterized by chronic hyperglycaemia, which results from a failure of the body to release adequate amounts of the blood glucose-lowering hormone insulin, the inability of the target organs to respond to insulin with increased uptake of glucose, or a combination of both. The two main classes, “insulin-dependent diabetes” (IDDM or type-1) and non-insulin-dependent diabetes” (NIDDM or type-2) are easily distinguished by the severity of insulin deficiency, which is more profound in IDDM (Pickup & Williams 1997). However, NIDDM is becoming an ever greater public health problem, currently affecting an estimated 150 million people worldwide (Amos *et al.* 1997). NIDDM is character-

ized by insufficient insulin secretion relative to the blood glucose level. Additionally, already at early stages of the disease the normal pattern of insulin release is disturbed so that the rapid but transient peak of secretion in response to a glucose challenge (1st phase) is absent, while a slow but sustained component remains (2nd phase; 1992). Because of the secretional defects associated with diabetes, it is important to understand the molecular mechanisms underlying normal and pathological insulin release.

## Insulin secretion from pancreatic B-cells

Insulin is produced in the B-cells of the islets of Langerhans, where it is stored in ~10000 secretory vesicles or “granules” waiting to be released into the blood stream. The latter occurs by fusion of the granule with the plasma membrane, a process referred to as exocytosis and used in most cells of the body to release substances or to insert newly synthesized proteins into the plasma membrane. Like in other endocrine cells and in neurones, exocytosis in B-cells is regulated and the amount of insulin circulating in the blood depends more on the rate of exocytosis than on the rate of insulin biosynthesis.

B-cells are electrically active and respond to elevated blood glucose by generating action potentials (fig. 1). Electrophysiological experiments during the past 15 years have thrown light on the way that release of insulin is coupled to stimulation. In this model, glucose is taken up by the cell

Author for correspondence: Sebastian Barg, Department of Physiological Sciences, Molecular and Cellular Physiology, Lund University, Sölvegatan 19, BMC F11, S-221 84 Lund, Sweden (fax +46 222 7763, e-mail sebastian.barg@mphy.lu.se).




Fig. 1. Stimulus secretion coupling in pancreatic B-cells. Glucose enters the cell via GLUT-2 transporters, and is metabolised to produce ATP. The resulting increase in the cytosolic ATP/ADP-ratio closes K<sub>ATP</sub>-channels in the plasma membrane, which leads to depolarization of the cell and influx of Ca<sup>2+</sup> through voltage dependent L-type Ca<sup>2+</sup>-channels (VDCC). The increase in Ca<sup>2+</sup> at the granule then triggers exocytosis and thus insulin secretion. Sulfonylurea close the K<sub>ATP</sub>-channel by binding to its SUR-subunits, and thereby cause membrane depolarization independent of the metabolic state of the cell.

and metabolized to ATP. The increase in the ATP/ADP ratio then leads to closure of ATP-sensitive K<sup>+</sup>-channels (K<sub>ATP</sub>-channels) in the plasma membrane (fig 1). Since these channels are responsible for maintaining a resting potential of about  $-70$  mV, i.e., close to the K<sup>+</sup>-equilibrium potential, this leads to a gradual depolarization of the cell. Eventually, voltage-dependent Ca<sup>2+</sup>-channels become activated and initiate action potentials. Influx of Ca<sup>2+</sup> through these channels and the resulting elevation of the cytosolic Ca<sup>2+</sup>-concentration then triggers exocytosis of the insulin-containing granules (Ashcroft & Rorsman 1989).

### The exocytotic machinery

B-cells are fairly typical examples of peptide-secreting endocrine cells. Proinsulin, the precursor of insulin, is synthesized in the endoplasmic reticulum and undergoes a series of maturation steps, starting already in the Golgi. The product is then packaged into secretory granules that gradually acidify, allowing further processing to insulin (Hutton 1994). These granules are found throughout the cytosol and eventually translocated to the plasma membrane. The ultimate fusion of the granule with the plasma membrane is triggered by Ca<sup>2+</sup> and controlled by a complex network of protein-protein and protein-lipid interactions that are similar in all cellular membrane fusion events, and largely conserved from yeast to man. A vast body of molecular and physiological data accumulated over the past years has led to a unifying model for membrane fusion (Calakos & Scheller 1996). It is therefore not surprising that many of the proteins involved in the regulation of neurotransmitter release have also been identified in the pancreatic B-cell and demonstrated to participate in insulin secretion (Lang 1999). These proteins include the SNAREs (sol-

uble *N*-ethylmaleimide-sensitive fusion protein [NSF] attachment protein [SNAP] receptors) synaptobrevin-2, syntaxin-1A, and SNAP-25 (25-kDa synaptosomal-associated protein). It is now clear that a ternary complex consisting of these proteins plays a central role in exocytosis (fig 4; Söller *et al.* 1993), by tethering the granules to the plasma membrane (Sutton *et al.* 1998). Synaptotagmin is the most likely candidate for the Ca<sup>2+</sup>-sensor of the exocytotic machinery (Fernandez-Chacon *et al.* 2001), and essential for Ca<sup>2+</sup>-dependent neurotransmission (Littleton *et al.* 1993) and exocytosis of insulin-containing granules (Lang *et al.* 1997). It is a protein of 65 kDa that spans the vesicle membrane once and contains a large cytoplasmic domain with two Ca<sup>2+</sup>-binding C2 domains (protein kinase C-homology domains), C2A and C2B (Sutton *et al.* 1995). In addition to binding Ca<sup>2+</sup>, the C2 domains are involved in the association of syt with phospholipids, SNAREs, and a variety of other proteins. Several isoforms of the protein have been identified, of which synaptotagminIII and synaptotagminVII are present in B-cells (Lang *et al.* 1997; Gao *et al.* 2000).

### Docking, priming, ATP-dependence, and kinetics of exocytosis

Fusion occurs rapidly upon Ca<sup>2+</sup>-influx but obviously requires the granule to be situated in the immediate vicinity of the plasma membrane. Electron microscopy on various neuronal and endocrine cells (von Gersdorff *et al.* 1996; Plattner *et al.* 1997), including B-cells (Olofsson *et al.* 2002), has revealed a subset of secretory granules that is "docked" beneath the plasma membrane. These findings correlate with data obtained with capacitance measurements (fig. 2) demonstrating a burst of exocytosis in response to a step elevation of Ca<sup>2+</sup>, followed by a slow but sustained phase (fig. 3). In B-cells, the size of the burst that can be released immediately upon Ca<sup>2+</sup>-influx corresponds to less than 1% of the total number of granules (Barg *et al.* 2001a, 2002a & b). Refilling of this functionally defined "readily releasable pool" is ATP-, Ca<sup>2+</sup>- and temperature-dependent (Bittner & Holz 1992; Parsons *et al.* 1995, Renström *et al.* 1996a & b, Eliasson *et al.* 1997; Gromada *et al.* 1999), and commonly referred to as "priming" (Hay & Martin 1992). In the B-cells, the "readily releasable pool" can be depleted in less than a second (Barg *et al.* 2001a & 2002a), whereas its refilling takes up to a minute (Gromada *et al.* 1999; Barg *et al.* 2002a). Under conditions of prolonged stimulation, the priming of vesicles for exocytosis therefore becomes rate-limiting, which may account for the biphasic time course in glucose-stimulated insulin secretion. In this scenario, the early rapid component (1st phase secretion) corresponds to the Ca<sup>2+</sup>-dependent release of granules already belonging to the "readily releasable pool", while the sustained component (2nd phase) reflects the relatively slow priming of granules and to some extent probably their translocation to the release site (docking) (Rorsman *et al.* 2000; Barg *et al.* 2002a). Interestingly, priming occurs after




Fig. 2. Exocytosis measured as capacitance increase. In this method, a high resistance seal is formed between a patch of the cell surface membrane and a recording electrode made of glass. Electrical contact with the cytosol is then established by rupturing the membrane within the patch. Capacitance can be measured by circuit analysis, and since the cell capacitance is proportional to its surface area ( $10 \text{ fF}/\mu\text{m}^2$ ), any membrane that is added by exocytosis is detected as capacitance increase. Exocytosis can be measured in single cells at millisecond time resolution, making the study of release kinetics feasible. A. Schematic of a cell undergoing exocytosis; granule membrane is added to the plasma membrane and thereby increases the cell's capacitance. B. Typical capacitance recording of a cell stimulated with  $\text{Ca}^{2+}$  in the recording electrode. C. Means of eliciting exocytosis in patch-clamped cells; 1. infusion of an EGTA/ $\text{Ca}^{2+}$ -mixture leads to slow, sustained exocytosis (see B); 2. voltage-clamp depolarization evokes the opening of voltage-gated  $\text{Ca}^{2+}$ -channels (VDCC),  $\text{Ca}^{2+}$ -influx, and triggering of exocytosis in a near physiological manner and results in steep spatial  $\text{Ca}^{2+}$ -gradients; 3. flash photolysis of "caged"  $\text{Ca}^{2+}$ ;  $\text{Ca}^{2+}$  is chelated by the NP-EGTA cage and introduced into the cell via the recording electrode. A flash ( $<1 \text{ msec}$ ) is then directed at the cell to photolyse the cage, increasing  $\text{Ca}^{2+}$  uniformly within a few hundreds of microseconds. Note that in all cases access to the cytosol is conveniently provided through the recording electrode, and test substances can be applied intracellularly simply by including them in the electrode solution.

docking of the granule at the release site and the ATP-dependence is therefore not due to the energy required for translocation. Several sequential steps are necessary to render the granule release-competent, and numerous ideas have been put forward to explain the ATP-dependence of priming. These include the ATP-dependent synthesis of phosphoinositides and protein kinase-mediated protein phosphorylation (Klenchin & Martin 2000). More recently, the presynaptic proteins munc-13 and Rim (Rab3-interacting molecule) have also been implicated in priming, mainly because they act after docking but before fusion (Ashery *et al.* 2000; Koushika *et al.* 2001).

### Coupling of $\text{Ca}^{2+}$ -influx and exocytosis

Despite many similarities of the exocytotic machinery in (neuro-) endocrine cells and that in neurotransmitter-releasing fast synapses, there are important differences: 1) synaptic vesicles are small (diameter  $<30 \text{ nm}$ ; von Gersdorff 1996) compared with peptide containing vesicles (hun-

dreds of nm; Plattner *et al.* 1997; Olofsson *et al.* 2002); 2) synaptic exocytosis occurs without delay after the influx of  $\text{Ca}^{2+}$  and is at least one order-of magnitude faster than peptide secretion (Ämmälä *et al.* 1993; Mennerick & Matthews 1996); 3) synaptic vesicles may stay largely intact during exocytosis and are refilled with new transmitter molecules immediately after endocytosis, whereas peptide-containing granules are assembled and filled with secretory peptides in the Golgi apparatus (Betz & Bewick 1992; Hutton 1994); 4) synaptic exocytosis generally requires higher concentrations of  $\text{Ca}^{2+}$  (Ämmälä *et al.* 1993; Mennerick & Matthews 1996, Chow *et al.* 1996; Bollmann *et al.* 2000); and 5) although most secretory cells contain more than one type of voltage-gated  $\text{Ca}^{2+}$ -channel, synaptic transmitter exocytosis depends on the faster N- or P-type (Hirning *et al.* 1988; Uchitel *et al.* 1992), whereas endocrine cells preferentially use the L-type (Lopez *et al.* 1994; Barg *et al.* 2001a). Taken together, the findings in neurones suggest that the site of  $\text{Ca}^{2+}$ -influx is within minimal distance of the  $\text{Ca}^{2+}$ -sensor at the synaptic vesicle and functionally associated within release sites. This "coupling" appears to be crucial for fast synapses in order to avoid the diffusional delay between  $\text{Ca}^{2+}$ -influx and exocytosis. Notably, the high concentrations of  $\text{Ca}^{2+}$  required for exocytosis in certain neurones ( $\sim 100 \mu\text{M}$ ; Heidelberger *et al.* 1994) are not observed throughout the cytosol, and can in fact be expected only at the pore of the  $\text{Ca}^{2+}$ -channel. Such an organization can easily be envisaged to be energetically advantageous as it restricts the  $\text{Ca}^{2+}$ -signal to a small part of the cell where it can quickly trigger exocytosis, because less  $\text{Ca}^{2+}$  has to be cleared from the cytosol after the stimulation. In addition,  $\text{Ca}^{2+}$  may still be used for other signal transduction purposes like the control of gene transcription or synaptic plasticity (Berridge 1998).

In endocrine cells such functional coupling between  $\text{Ca}^{2+}$ -channel and exocytotic machinery seems less critical. In chromaffin cells for example, exocytosis persists even after  $\text{Ca}^{2+}$ -entry has stopped, due to the relatively low  $\text{Ca}^{2+}$ -dependence of exocytosis (Augustine & Neher 1992) and the time required to remove  $\text{Ca}^{2+}$  from the cytosol (Chow *et al.* 1996). However, recent estimates of the  $K_D$




Fig. 3. Kinetics of exocytosis. A. Three distinct phases of exocytosis are readily distinguished in mouse pancreatic B-cells: (1) exocytosis of readily releasable pool (2–3) priming/retrieval of new granules; data combined from Barg *et al.* (2001). B. Schematic of regulated exocytosis. Granules are translocated to the plasma membrane, where they dock and are primed for release by an ATP-dependent reaction.  $\text{Ca}^{2+}$ -influx triggers release of the primed granules.



of  $\text{Ca}^{2+}$ -dependent exocytosis in B-cells are around  $20 \mu\text{M}$  (Takahashi *et al.* 1997; Barg *et al.* 2001a), and in fact similar to that of some neuronal terminals (Bollmann *et al.* 2000; Beutner *et al.* 2001). This value is surprisingly high because the average cytosolic (global)  $\text{Ca}^{2+}$  in the B-cell does not exceed  $\sim 2 \mu\text{M}$  during stimulation (Ämmälä *et al.* 1993; Bokvist *et al.* 1995; Dryselius *et al.* 1999). Obviously, steep spatial  $\text{Ca}^{2+}$ -gradients must exist and exocytosis is probably triggered in these zones where  $[\text{Ca}^{2+}]_i$  may be 10 times higher than in the remainder of the cell (Bokvist *et al.* 1995; Qian & Kennedy 2001).

### $\text{Ca}^{2+}$ -channels are part of the fusion machinery

The N-type  $\text{Ca}^{2+}$ -channel binds to the synaptic core complex via syntaxin, SNAP-25, and the C2B domain of synaptotagminI (fig. 4), suggesting that in neurones the  $\text{Ca}^{2+}$ -channel is an integral part of the exocytotic machinery (Mochida *et al.* 1996). The interacting site of the channel is located between transmembrane regions II and III of its pore-forming  $\alpha 1$  subunit (Sheng *et al.* 1996; Wiser *et al.* 1997; Atlas 2001), and is often referred to as the “synprint”-region. Recombinant peptides containing the synprint region dissociate the channel from the exocytotic complex and, when injected into presynaptic neurones, reversibly inhibit synaptic transmission without affecting  $\text{Ca}^{2+}$ -influx (Mochida *et al.* 1996). Syntaxin appears to act as a scaffold for the exocytotic machinery, keeping the proteins of the vesicle docking site in place (Wu *et al.* 1999). Interestingly, syntaxin alone (i.e., in the absence of other synaptic proteins) binds to the N-type channel and *decreases* its activity (Bezprozvanny *et al.* 1995). This could serve to prevent  $\text{Ca}^{2+}$ -influx at sites that are not occupied with docked secretory vesicles. Once the site is occupied with a vesicle, it would bind to the  $\text{Ca}^{2+}$ -channels in the vicinity with its synaptotagmin, and thus promote  $\text{Ca}^{2+}$ -channel activity.




Fig. 4. Proteins involved in exocytosis. A, an insulin-containing granule that approaches the plasma membrane. B, a docked vesicle in the moment just before fusion. The SNARE proteins have formed a parallel four-helix bundle, which pulls the granule down the plasma membrane C, after fusion, the SNARE protein-complex is disassembled by the ATPase NSF.

Syntaxin, SNAP-25, and synaptotagmin also bind to the synprint site of the L-type  $\text{Ca}^{2+}$ -channel, which mediates  $\text{Ca}^{2+}$ -influx in endocrine cells (Wiser *et al.* 1997 & 1999; Atlas 2001). Like in neurones, interfering with the interaction between the  $\text{Ca}^{2+}$ -channel and the core complex by introducing the corresponding L-type synprint peptide (Lc<sub>753–893</sub>) completely abolishes depolarization-evoked insulin exocytosis (Wiser *et al.* 1999; Barg *et al.* 2001a). Importantly, neither  $\text{Ca}^{2+}$ -dependent exocytosis itself nor  $\text{Ca}^{2+}$ -currents are affected, suggesting that the exogenous synprint peptide dissociates  $\text{Ca}^{2+}$ -channel/SNARE/synaptotagmin complexes, thereby increasing the average distance between channels and docked granules. This interpretation is supported by the finding that synprint-blocked exocytosis can be rescued by strong, prolonged stimulation, which increases the total  $\text{Ca}^{2+}$ -influx and leaves more time for its diffusion away from the channel pore (Barg *et al.* 2001a). The latter should increase the  $\text{Ca}^{2+}$ -concentration sufficiently to trigger exocytosis even at sites that are not in the immediate vicinity of  $\text{Ca}^{2+}$ -influx.

### How many channels are needed to make a granule fuse?

In B-cells, the magnitude of the whole-cell  $\text{Ca}^{2+}$ -current is very small, and non-stationary fluctuation analysis shows that mouse B-cells are equipped with fewer than 500  $\alpha 1\text{C}$  L-type  $\text{Ca}^{2+}$ -channels (Barg *et al.* 2001a). This corresponds to a  $\text{Ca}^{2+}$ -channel density of  $\sim 0.9$  channel per  $\mu\text{m}^2$  or only 5–10% of that reported for other endocrine cells ( $9\text{--}20/\mu\text{m}^2$ ; Fenwick *et al.* 1982). Nevertheless, B-cells are capable of fast exocytosis. Analysis of the kinetics of exocytosis in these cells during voltage-clamp depolarizations reveals an early, very rapid component of about 60 granules. Exocytosis of this pool takes place at a peak rate of  $\sim 700$  granules per second (Barg *et al.* 2001a), suggesting very high levels of  $\text{Ca}^{2+}$  (probably  $>100 \mu\text{M}$ ) at the site of exocytosis. Assuming that the 60 granules that can be released at maximum rate are associated with about 500  $\text{Ca}^{2+}$ -channels, it seems likely that more than one channel binds to each exocytotic site. Indeed, cell-attached patches on B-cells ( $\sim 1\%$  of the cell surface) predominantly contain either no channels at all or three active  $\text{Ca}^{2+}$ -channels, indicating non-random distribution of the channels on the plasma membrane (Barg *et al.* 2001a). Further support for  $\text{Ca}^{2+}$ -channel clustering comes from experiments with the L-type  $\text{Ca}^{2+}$ -channel agonist BayK8644, which acts by increasing the open lifetimes. Prolonged openings of single  $\text{Ca}^{2+}$ -channel should make exocytosis more likely, because of the increased probability that the  $\text{Ca}^{2+}$ -concentration around the granule lasts sufficiently long to trigger exocytosis. However, the efficiency of exocytosis was not altered in presence of BayK8644, despite 6 times longer  $\text{Ca}^{2+}$ -channel openings (Barg *et al.* 2001a). The reason might be that the openings of several clustered  $\text{Ca}^{2+}$ -channels overlap in time and thereby achieve a similar effect as BayK8644 would if the channels were not clustered.

### Effects of $\text{Ca}^{2+}$ on the recruitment of granules into the readily releasable pool

In addition to triggering exocytosis,  $\text{Ca}^{2+}$  stimulates the recruitment of release-competent granules (Ämmälä *et al.* 1993; Heidelberger *et al.* 1994). Thus, the size of readily releasable pool depends on the concentration of cytosolic  $\text{Ca}^{2+}$  that is present during the minute or so *before* the initiation of exocytosis (Påhlman & Barg, unpublished results). The previous finding that exogenous  $\text{Ca}^{2+}$ -chelators inhibit exocytosis in endocrine cells (Ämmälä *et al.* 1993; Chow *et al.* 1996) may therefore be due to reduced rate of supply of granules ready for release due to artificially low basal  $[\text{Ca}^{2+}]_i$ . However, even at elevated  $[\text{Ca}^{2+}]_i$  recruitment of granules in the B-cell is a slow process ( $\sim 5$  granules/sec., compared with the exocytotic burst of 700 granules/sec.). During sustained stimulation the rate of supply of new granules quickly becomes rate-limiting for the process of exocytosis. A number of  $\text{Ca}^{2+}$ -binding proteins are known to affect exocytosis, and may underlie the  $\text{Ca}^{2+}$ -dependence of granule recruitment. These include calmodulin (Chen *et al.* 1999), the  $\text{Ca}^{2+}$ -dependent phosphatase calcineurin (Renström *et al.* 1996b), CAPS ( $\text{Ca}^{2+}$ -dependent activator protein for secretion; Ann *et al.* 1997), and  $\text{Ca}^{2+}$ /calmodulin kinase II (Ämmälä *et al.* 1993; Gromada *et al.* 1999). Blockage of the latter inhibited the slow phase, and suggested a myosin-actin-dependent step in the granule recruitment (Bi *et al.* 1997). The basal concentration of cytosolic  $\text{Ca}^{2+}$  depends to a great extent on intracellular  $\text{Ca}^{2+}$ -handling, and there is evidence for  $\text{Ca}^{2+}$ -release from the ER (Tengholm *et al.* 1998; Holz *et al.* 1999), for  $\text{Ca}^{2+}$ -uptake by the mitochondria (Rutter *et al.* 1993), and for accumulation of  $\text{Ca}^{2+}$  in the granules (Scheenen *et al.* 1998; Mitchell *et al.* 2001).

### Comparison between insulin-secreting B-cells and glucagon-secreting A-cells

Glucagon is a major catabolic and hyperglycaemic 29 a.a. hormone, which is secreted from the A-cells. Its main biological effect is to enhance synthesis and mobilization of glucose in the liver. Secretion of glucagon is stimulated by low blood glucose (Gorus *et al.* 1984), amino acids (Pipeleers *et al.* 1985), and a variety of hormones and neurotransmitters, such as adrenaline, glucose-dependent insulinotropic polypeptide (GIP), and glucagon-like peptide-1 (Ding *et al.* 1997; Gromada *et al.* 1997). Insulin, somatostatin, and fatty acids reduce glucagon secretion (Gerich *et al.* 1976; Ding *et al.* 1997). Whereas insulin levels are inadequately low in hyperglycaemic diabetic patients, glucagon levels are elevated, and this aggravates the disease (Unger 1971). The reason for this abnormality of glucagon secretion is not known. Electrical activity has been observed in A-cells (Rorsman & Hellman 1988; Barg *et al.* 2000; Göpel *et al.* 2000) and is, at least in part, attributable to voltage-gated  $\text{Ca}^{2+}$ -channels. A- and B-cells in the mouse can be distinguished electrophysiologically because A-cells

1) are small in size (capacitance of 3 pF compared with 6–7 pF in B-cells, corresponding to diameters of 10 and 15  $\mu\text{m}$ ; Barg *et al.* 2000), 2) are equipped with a large tetrodotoxin-sensitive and transient  $\text{Na}^+$ -current (Barg *et al.* 2000; Göpel *et al.* 2000), 3) have almost no  $\text{K}_{\text{ATP}}$ -conductance (Barg *et al.* 2000; Göpel *et al.* 2000), and 4) are, contrary to B-cells, electrically active at low glucose levels (Barg *et al.* 2000; Göpel *et al.* 1999 & 2000). The latter observation is in agreement with the tentative physiological role of A-cells and has been explained by a combination of a small (hyperpolarizing)  $\text{K}_{\text{ATP}}$ -current and the fact that regenerative electrical activity depends on  $\text{Na}^+$ , rather than  $\text{Ca}^{2+}$  in these cells (Göpel *et al.* 2000). In this scenario, elevation of glucose results in an increased cytoplasmic ATP/ADP-ratio, which closes the few  $\text{K}_{\text{ATP}}$ -channels that are present in A-cells (Bokvist *et al.* 1999; Barg *et al.* 2000; Göpel *et al.* 2000). This is associated with membrane depolarization, which in turn prevents reactivation of the  $\text{Na}^+$ -channels after an action potential.

### Kinetics of glucagon exocytosis

Exocytosis of glucagon (A-cells) is  $\text{Ca}^{2+}$ -dependent and slightly faster than exocytosis of insulin (B-cells; fig. 5). The difference reflects the number of granules belonging to the RRP ( $\sim 120$  granules in A-cells and  $\sim 60$  granules in B-cells), and the rate of refilling after depletion of the readily releasable pool ( $\sim 20$  granules/s in A-cells and 5–10 granules/sec. in B-cells). In addition, there is a strong secondary acceleration of the exocytotic rate in A-cells after depletion of the readily releasable pool (fig. 5), which may reflect fast  $\text{Ca}^{2+}$ -dependent translocation of granules from the interior of the cell towards the release sites. Interestingly, the *maximum* rate of 750 granules/s during a depolarization is achieved within about 20 msec. and fairly similar to that observed in B-cells ( $\sim 700$  granules/sec. after 15 msec.). Taken together, the data therefore suggest a fairly similar organization of  $\text{Ca}^{2+}$ -dependent exocytosis in both cell types, but a higher number of release sites in A-cells. It seems further likely that the large  $\text{Ca}^{2+}$ -currents of the relatively small A-cells, leading to larger and less localized  $\text{Ca}^{2+}$  transients, may explain part of the differences,




Fig. 5. Comparison of exocytosis in B- and A-cells. A single B-cell (left) or A-cell (right) was stimulated with a train of depolarizations (upper trace). While the exocytotic responses in the B-cell become progressively smaller throughout the train, there is a secondary acceleration in A-cells.

e.g., faster  $\text{Ca}^{2+}$ -dependent recruitment of granules and the secondary acceleration. It can be speculated that these features enable the A-cell to respond with a bout of glucagon release during short periods of glucose demand, as required during intense physical activity and in certain life-threatening situations such as shock (van der Meer *et al.* 1981) and trauma (Efendic *et al.* 1974).

### Sulfonylureas stimulate insulin release

Sulfonylureas have been used in the treatment of NIDDM for almost 45 years, but it is only recently that the molecular mechanisms of their action have been elucidated. The principal effect of the sulfonylureas is to stimulate insulin secretion from the B-cells (Ashcroft & Ashcroft 1992). This is a consequence of their ability to selectively inhibit the  $\text{K}_{\text{ATP}}$ -channels in the B-cell plasma membrane (Trube *et al.* 1986) by binding to its sulfonylurea receptor (Aguilar-Bryan *et al.* 1995) subunit. This short-circuits the normal physiological regulation of the channel by the metabolic state of the cell (via the ATP/ADP-ratio) and results in membrane depolarization,  $\text{Ca}^{2+}$ -influx and insulin secretion (Ashcroft & Rorsman 1989).

In addition to the stimulation induced by the inhibition of the  $\text{K}_{\text{ATP}}$ -channel, therapeutic concentrations of sulfonylureas potentiate insulin secretion by a mechanism that is exerted downstream of the depolarization, i.e., *directly* on the exocytotic machinery (Eliasson *et al.* 1996; Tian *et al.* 1998; Barg *et al.* 1999 & 2001b). Thus, sulfonylureas remain stimulatory in conditions that preclude the effects of sulfonylureas on the membrane potential, i.e. after permeabilization (Tian *et al.* 1998) or voltage-clamping of the cells (Eliasson *et al.* 1996; Barg *et al.* 1999 & 2001b). However, since sulfonylureas normally depolarize the B-cell and thereby indirectly stimulate insulin release, any *direct* stimulation of the exocytotic machinery may easily be overlooked. Our laboratory has used patch-clamp capacitance measurements to elucidate the mechanisms underlying sulfonylurea-dependent stimulation of exocytosis (Barg *et al.* 1999 & 2001b). In these experiments, sulfonylureas (tolbutamide) failed to stimulate exocytosis in the absence of  $\text{Ca}^{2+}$  and stimulated at concentrations that were only slightly higher than resting  $[\text{Ca}^{2+}]_i$ . This suggests that sulfonylureas do not induce, but rather stimulate ( $\text{Ca}^{2+}$ -dependent) exocytosis (Barg *et al.* 1999). Similar effects of sulfonylureas have been reported in pancreatic A-cells (Høy *et al.* 2000) and clonal neuroendocrine PC12-cells (Taylor *et al.* 1999), suggesting that the underlying mechanism is not restricted to B-cells and insulin secretion.

### Regulation of exocytosis by a granular 65-kDa sulfonylurea-binding protein

Pharmacologically, the mechanism that mediates the *direct* stimulation of the exocytotic machinery is very similar to the B-cell variety of the  $\text{K}_{\text{ATP}}$ -channel, which consists of an octameric complex of four type 1 sulphonylurea receptors

(SUR1) and four  $\text{K}^+$ -channel (Kir6.2) subunits. Thus, exocytosis is inhibited by intracellular application of diazoxide, an activator of B-cell  $\text{K}_{\text{ATP}}$ -channels, while pinacidil, an activator of vascular  $\text{K}_{\text{ATP}}$ -channels that contain SUR2A (Shindo *et al.* 1998), has no effect (Barg *et al.* 1999).  $\text{K}_{\text{ATP}}$ -channels are physiologically activated with reduced ATP/ADP ratio, and a similar regulation leads to inhibition of the exocytotic machinery. Interestingly, this inhibition of  $\text{Ca}^{2+}$ -dependent exocytosis is fully rescued by addition of tolbutamide (Barg *et al.* 2001b). The latter suggests that a direct stimulation of the exocytotic machinery could contribute to the insulinotropic action of sulfonylureas in diabetic patients, where the ATP/ADP ratio may be reduced as a consequence of impaired glucose metabolism.

ATP- and sulfonylurea-sensitive  $\text{K}^+$  and  $\text{Cl}^-$ -conductances have been found in isolated zymogen granules of the exocrine pancreas (Thevenod *et al.* 1992 & 1994). In these granules, a 65-kDa sulfonylurea-binding protein is involved in the modulation of the activity of the granular ion channels. This protein cross-reacts with antibodies originally raised against a SUR-related protein, the 170-kDa mdr1 (multi-drug-resistance P-glycoprotein; Thevenod *et al.* 1994; Braun *et al.* 1997). Interestingly, the majority (90%) of the high-affinity sulfonylurea binding sites in the B-cell are intracellular and associated with the insulin-containing secretory granules (Ozanne *et al.* 1995). It seems therefore possible that such channels are also present in B-cell granules, and there are several arguments in favor of this idea. First, two different antibodies (JSB-1 and C219) originally raised against mdr1 detect a protein of 65 kDa in B-cell granules. Second, intracellular application of one of these antibodies (JSB-1) abolishes the stimulatory action of sulfonylureas on exocytosis, yet does not interfere with their capacity to block the  $\text{K}_{\text{ATP}}$ -channels (Barg *et al.* 1999). Third, the sulfonylurea glibenclamide binds to a 65 kDa protein in insulin-containing granules (Renström *et al.* 2002). Finally, both SUR and mdr1 belong to the family of ATP-binding cassette (ABC)-proteins (Schwiebert *et al.* 1999), and tamoxifen, a blocker of ABC-protein function in a variety of contexts, prevents the stimulation of exocytosis by tolbutamide (Barg *et al.* 1999). Collectively, the data argue that sulfonylureas bind to a mdr1-related granular protein of 65 kDa, similar to the protein found in the zymogen granules of the exocrine pancreas. The binding of sulfonylureas to the 65 kDa protein lead to an augmentation of the exocytotic machinery, possibly via activation of ion channels in the granule membrane.

### Granular $\text{Cl}^-$ -flux through CIC-3 channels is required for insulin exocytosis

By analogy with other ABC-proteins, especially SUR, it is conceivable that the 65-kDa granular sulfonylurea binding protein acts as a regulatory subunit of a  $\text{K}^+$ -channel. This may indeed be the case in A-cells, where intracellular application of certain  $\text{K}^+$ -channel blockers interferes with exo-

cytosis (Høy *et al.* 2000). However, exocytosis in the B-cell is unaffected by replacing intracellular  $K^+$  with  $Cs^+$ . It is instead strongly inhibited by the  $Cl^-$ -channel blocker DIDS (4,4'-di-isothiocyanato-stilbene-2,2'-disulfonic acid) or by replacing intracellular  $Cl^-$  with glutamate (Barg *et al.* 2001b). This suggests that intracellular  $Cl^-$ , rather than  $K^+$ -fluxes, participate in the control of exocytosis in the B-cell. Indeed, antibodies against the  $Cl^-$ -channel CIC-3 label B-cell granules and detect a protein with a molecular weight of  $\sim 90$  kDa in B-cell homogenates and in a granule-enriched subcellular fraction of clonal insulin-containing Ins-1 cells. One of the antibodies, antiCIC-3-C, selectively blocks CIC-3 channel function, which can be exploited to test the role of this channel in the process of exocytosis. Notably, the antibody abolishes exocytosis when it is dialyzed into the cell via the patch pipette, while the non-functional antibody antiCIC-3-N has no effect (Barg *et al.* 2001b). These data raise the interesting possibility that granular  $Cl^-$ -flux carried by CIC-3 channels is required for  $Ca^{2+}$ -dependent exocytosis in the B-cell. This phenomenon does not seem to be limited to the B-cell, as  $Cl^-$  is required for hormone release in a variety of other cells, including pancreatic acinar cells (Fuller *et al.* 1989) and pituitary melanotrophs (Rupnik & Zorec 1992).

#### Chloride influx enables granule acidification

How does influx of  $Cl^-$  into the granule stimulate exocytosis? One consequence of  $Cl^-$ -influx would be net accumulation of electrolytes, water uptake, and increased hydrostatic pressure inside the granules. The latter has been suggested to promote membrane fusion (Woodbury 1995), but there is evidence to the contrary (Breckenridge & Almers 1987). Osmotic effects are unlikely to be involved in the B-cell because reversing osmotic gradients with strongly hyperosmotic intracellular solutions has no effect on exocytosis (Barg *et al.* 2001b). Alternatively, electrogenic  $H^+$ -pumping, as required for granule acidification, might require a shunt-conductance to prevent the development of a large electrical potential (positive inside). Indeed, regulation of vesicle acidification by chloride counter-ion flux seems to be a common phenomenon (al Awqati *et al.* 1992). For example, CIC-5 was found on endocytic vesicles of renal cells (Günther *et al.* 1998), and disruption of this channel impaired endocytosis and vesicle acidification (Piwon *et al.* 2000).

In the B-cell, interfering with either the granular v-type  $H^+$ -ATPase with bafilomycin, or with intracellular  $H^+$ -gradients by introducing the protonophore carbonyl cyanide m-chlorophenylhydrazone (CCCP) suppresses exocytosis (Barg *et al.* 2001b). This implicates granule acidification as an important step during the preparation of granules for exocytosis.  $H^+$ -pumping itself may be controlled by sulfonyleurea-sensitive mechanism, as direct measurements of granular pH with the fluorescent probe LysoSensor show that tolbutamide acidifies granules. Importantly, CCCP-induced equilibration of the granule pH can be prevented by

blocking CIC-3 channel function with DIDS or antiCIC-3-C. Why is this so? Apparently, protons leave the granule only when the  $Cl^-$ -channel is open, because the (electrogenic) proton efflux requires charge neutralization. The electrical potential (negative inside) across the granule membrane would otherwise balance the  $H^+$ -gradient and prevent equilibration. This feature can be used to study the pharmacology of the granular  $Cl^-$ -channel. It turns out that the CCCP-dependent  $H^+$ -leak is prevented by the very same maneuvers that also block exocytosis. ADP or diazoxide inhibit the leak, while tolbutamide enhances it (Barg *et al.* 2001b). Thus, exocytosis in the B-cell depends, like CCCP-dependent proton leakage from the granule, on open CIC-3 channels.

#### Priming for exocytosis by granular acidification and $Cl^-$ -fluxes

Which step during the life of the insulin granule is modulated by granular ion fluxes, tolbutamide, and the ATP/ADP-ratio? Although capacitance measurements monitor only membrane fusion, kinetic analysis of the capacitance




Fig. 6. Model for the regulation of granular pH and exocytosis. A. Metabolically regulated  $Cl^-$ -influx into the granule through an ion channel complex comprised of CIC-3 and an mdr1-like 65-kDa protein, determines the rate of granular acidification by the V-type  $H^+$ -ATPase. Tolbutamide, diazoxide and ADP are likely to interact with the regulatory 65-kDa protein, whereas DIDS and anti-hCIC-3-C bind to and inhibit CIC-3 directly. B. Replenishment of readily releasable pool depends on intragranular acidification. See text for details.



data allows limited predictions of processes that precede exocytosis, e.g., the rate at which granules become readily releasable (Parsons *et al.* 1995). During trains of depolarizations, exocytosis proceeds throughout the train, with the fastest rate during the first depolarization and progressively smaller responses during subsequent depolarizations (fig. 5). This is usually interpreted to mean that the first depolarizations rapidly deplete the readily releasable pool, and that the remainder of the train is associated with the slow recruitment of new granules. Such experiments show that ADP selectively blocks the slower second phase, and that the effect of ADP is counteracted by tolbutamide (Barg *et al.* 2001b). Similarly, intracellular application of the  $\text{Cl}^-$ -channel blocker DIDS or the antibody anti-hCIC-3-C remove the slow phase, but leave the initial burst intact. These data agree with a model in which ADP, tolbutamide, and granular  $\text{Cl}^-$ -flux control the refilling of the readily releasable pool, without affecting exocytosis of granules that have already progressed into this pool. This is in contrast to the effects of the protonophore CCCP, which abolishes not only the late component of the release, but eliminates the readily releasable pool as well. The latter may indicate that low intragranular pH is required not only for the priming of the granules but also for maintaining their release competence. Importantly, granule acidification depends on a bafilomycin-sensitive ATP-driven protonpump. It is therefore conceivable, that acidification of the insulin-granule is not required merely for processing of insulin (Hutton 1994), but also represents one of the ATP-dependent priming reactions.

A possible model that accounts for the available data is presented in fig 6. Secretory granules become release-competent by intragranular acidification, which depends on the simultaneous activity of the v-type  $\text{H}^+$ -ATPase and the CIC-3  $\text{Cl}^-$ -channel. By analogy to the regulation of the  $\text{K}_{\text{ATP}}$ -channel in the plasma membrane (Gribble *et al.* 1997), it seems possible that CIC-3 together with a regulatory mdr1-like protein of 65-kDa (Barg *et al.* 1999; Renström *et al.* 2002) form an ion channel complex in the granular membrane. In this scenario, ATP enhances granular  $\text{Cl}^-$  uptake, acidification and priming by a direct interaction with CIC-3, whereas ADP attenuates the same chain of events by binding to the regulatory 65-kDa protein. Tolbutamide interacts with the 65-kDa protein and prevents ADP binding, thereby counteracting the inhibitory effects of ADP on priming. How does granular acidification affect the exocytotic machinery? It can only be speculated that low intragranular pH may induce conformational changes in some of the proteins constituting the exocytotic machinery, thereby perhaps rendering them more fusogenic (Sutton *et al.* 1998). Indeed, vacuole acidification has recently been demonstrated to be required for the pairing of the SNARE-proteins in yeast (Ungermann *et al.* 1999).

Finally, the model implies that the priming of granules for release is controlled by the metabolic state of the cell, which may be of significance for the coupling between the level of blood glucose and insulin release. Data from B-cells

of diabetic individuals are scarce, but it appears possible that slow priming would produce secretional defects similar to those seen in NIDDM. Since the granular sulfonylurea receptor is involved in the modulation of priming, it may represent an exciting target for the development of new antidiabetic drugs.

#### Acknowledgements

I thank Dr. Patrik Rorsman for his support and for critically reading the manuscript. Funding was provided by the Medical Faculty at the University of Lund, the Dr. P Håkansson's Foundation, the Swedish Research Council (no. 8647), the Juvenile Diabetes Research Foundation (JDRF), the Swedish Diabetes Association, the NovoNordisk Foundation, the Crafoord foundation, the Swedish Society for Medical Research (SSMF), and the Royal Physiographic Society Lund, and Thuring's stiftelse.

#### References

- Aguilar-Bryan, L., C. G. Nichols, S. W. Wechsler, J. P. Clement, A. E. Boyd, G. Gonzalez, H. Herrera-Sosa, K. Nguy, J. Bryan & D. A. Nelson: Cloning of the beta cell high-affinity sulfonylurea receptor: a regulator of insulin secretion. *Science* 1995, **268**, 423–426.
- al Awqati, Q., J. Barasch & D. A. Landry: Chloride channels of intracellular organelles and their potential role in cystic fibrosis. *J. Exp. Biol.* 1997, **172**, 245–266.
- Amos, A., D. McCarty & P. Zimmet: The rising global burden of diabetes and its complications: estimates and projections. *Diabetes Care* 1997, **21**, S1–S85.
- Ann, K., J. A. Kowalchuk, K. M. Loyet & T. F. Martin: Novel  $\text{Ca}^{2+}$ -binding protein (CAPS) related to UNC-31 required for  $\text{Ca}^{2+}$ -activated exocytosis. *J. Biol. Chem.* 1997, **272**, 19637–19640.
- Ashcroft, F. M. & P. Rorsman: Electrophysiology of the pancreatic beta-cell. *Prog. Biophys. Mol. Biol.* 1989, **54**, 87–143.
- Ashcroft, S. J. & F. M. Ashcroft: The sulfonylurea receptor. *Biochim. Biophys. Acta* 1992, **1175**, 45–59.
- Ashery, U., F. Varoqueaux, T. Voets, A. Betz, P. Thakur, H. Koch, E. Neher, N. Brose & J. Rettig: Munc13-1 acts as a priming factor for large dense-core vesicles in bovine chromaffin cells. *EMBO J.* 2000, **19**, 3586–3596.
- Atlas, D.: Functional and physical coupling of voltage-sensitive calcium channels with exocytotic proteins: ramifications for the secretion mechanism. *J. Neurochem.* 2001, **77**, 972–985.
- Augustine, G. J. & E. Neher: Calcium requirements for secretion in bovine chromaffin cells. *J. Physiol. (Lond.)* 1992, **450**, 247–271.
- Barg, S., E. Renström, P. O. Berggren, A. Bertorello, K. Bokvist, M. Braun, L. Eliasson, W. E. Holmes, M. Köhler, P. Rorsman & F. Thevenod: The stimulatory action of tolbutamide on  $\text{Ca}^{2+}$ -dependent exocytosis in pancreatic beta cells is mediated by a 65-kDa mdr-like P-glycoprotein. *Proc. Natl. Acad. Sci. USA* 1999, **96**, 5539–5544.
- Barg, S., J. Galvanovskis, S. O. Göpel, P. Rorsman & L. Eliasson: Tight coupling between electrical activity and exocytosis in mouse glucagon-secreting alpha-cells. *Diabetes* 2000, **49**, 1500–1510.
- Barg, S., X. Ma, L. Eliasson, J. Galvanovskis, S. O. Göpel, S. Obermüller, J. Platzer, E. Renström, M. Trus, D. Atlas, J. Striessnig & P. Rorsman: Fast exocytosis with few  $\text{Ca}^{2+}$ -channels in insulin-secreting mouse pancreatic B-cells. *Biophys. J.* 2001a, **81**, 3308–3323.
- Barg, S., P. Huang, L. Eliasson, D. J. Nelson, S. Obermüller, P. Rorsman, F. Thevenod & E. Renström: Priming of insulin gran-

- ules for exocytosis by granular  $\text{Cl}^-$  uptake and acidification. *J. Cell Sci.* 2001b, **114**, 2145–2154.
- Barg, S., C. S. Olofsson, J. Schriever-Abeln, A. Wendt, S. Gebre-Medhin, E. Renström & P. Rorsman: Delay between fusion pore opening and peptide release from large dense-core vesicles in neuroendocrine cells. *Neuron* 2002a, **33**, 287–299.
- Barg, S., L. Eliasson, E. Renström & P. Rorsman: A subset of 50 secretory granules in close contact with L-type  $\text{Ca}^{2+}$ -channels accounts for first-phase insulin secretion in mouse  $\beta$ -cells. *Diabetes* 2002b, **51**, S74–S82.
- Berridge, M. J.: Neuronal calcium signaling. *Neuron* 1998, **21**, 13–26.
- Betz, W. J. & G. S. Bewick: Optical analysis of synaptic vesicle recycling at the frog neuromuscular junction. *Science* 1992, **255**, 200–203.
- Beutner, D., T. Voets, E. Neher & T. Moser: Calcium dependence of exocytosis and endocytosis at the cochlear inner hair cell afferent synapse. *Neuron* 2001, **29**, 681–690.
- Bezprozvanny, I., R. H. Scheller & R. W. Tsien: Functional impact of syntaxin on gating of N-type and Q-type calcium channels. *Nature* 1995, **378**, 623–626.
- Bi, G. Q., R. L. Morris, G. Liao, J. M. Alderton, J. M. Scholey & R. A. Steinhardt: Kinesin- and myosin-driven steps of vesicle recruitment for  $\text{Ca}^{2+}$ -regulated exocytosis. *J. Cell. Biol.* 1997, **138**, 999–1008.
- Bittner, M. A. & R. W. Holz: A temperature-sensitive step in exocytosis. *J. Biol. Chem.* 1992, **267**, 16226–16229.
- Bokvist, K., L. Eliasson, C. Åmmälä, E. Renström & P. Rorsman: Co-localization of L-type  $\text{Ca}^{2+}$  channels and insulin-containing secretory granules and its significance for the initiation of exocytosis in mouse pancreatic B-cells. *EMBO J.* 1995, **14**, 50–57.
- Bokvist, K., H. L. Olsen, M. Hoy, C. F. Gotfredsen, W. F. Holmes, K. Buschard, P. Rorsman & J. Gromada: Characterisation of sulphonylurea and ATP-regulated  $\text{K}^+$  channels in rat pancreatic A-cells. *Pflügers Arch.* 1999, **438**, 428–436.
- Bollmann, J. H., B. Sakmann & J. G. Borst: Calcium sensitivity of glutamate release in a calyx-type terminal. *Science* 2000, **289**, 953–957.
- Braun, M., I. Anderie & F. Thevenod: Evidence for a 65 kDa sulphonylurea receptor in rat pancreatic zymogen granule membranes. *FEBS Lett.* 1997, **411**, 255–259.
- Breckenridge, L. J. & W. Almers: Final steps in exocytosis observed in a cell with giant secretory granules. *Proc. Natl. Acad. Sci. USA* 1987, **84**, 1945–1949.
- Calakos, N. & R. H. Scheller: Synaptic vesicle biogenesis, docking, and fusion: a molecular description. *Physiol. Rev.* 1996, **76**, 1–29.
- Chen, Y. A., V. Duvvuri, H. Schulman & R. H. Scheller: Calmodulin and protein kinase C increase  $\text{Ca}^{2+}$ -stimulated secretion by modulating membrane-attached exocytic machinery. *J. Biol. Chem.* 1999, **274**, 26469–26476.
- Chow, R. H., J. Klingauf, C. Heinemann, R. S. Zucker & E. Neher: Mechanisms determining the time course of secretion in neuroendocrine cells. *Neuron* 1996, **16**, 369–376.
- Ding, W. G., E. Renström, P. Rorsman, K. Buschard & J. Gromada: Glucagon-like peptide I and glucose-dependent insulinotropic polypeptide stimulate  $\text{Ca}^{2+}$ -induced secretion in rat  $\alpha$ -cells by a protein kinase A-mediated mechanism. *Diabetes* 1997, **46**, 792–800.
- Dryselius, S., E. Grapengiesser, B. Hellman & E. Gylfe: Voltage-dependent entry and generation of slow  $\text{Ca}^{2+}$  oscillations in glucose-stimulated pancreatic  $\beta$ -cells. *Amer. J. Physiol.* 1999, **276**, E512–E518.
- Efendic, S., E. Cerasi & R. Luft: Trauma: hormonal factors with special reference to diabetes mellitus. *Acta anaesthesiol. scand.* 1974, **55**, Suppl. 107–119.
- Eliasson, L., E. Renström, C. Åmmälä, P. O. Berggren, A. M. Bertorello, K. Bokvist, A. Chibalin, J. T. Deeney, P. R. Flatt, J. Gabel, J. Gromada, L. O. Larsson, P. Lindstrom, C. J. Rhodes & P. Rorsman: PKC-dependent stimulation of exocytosis by sulphonylureas in pancreatic  $\beta$  cells. *Science* 1996, **271**, 813–815.
- Eliasson, L., E. Renström, W. G. Ding, P. Proks & P. Rorsman: Rapid ATP-dependent priming of secretory granules precedes  $\text{Ca}^{2+}$ -induced exocytosis in mouse pancreatic B-cells. *J. Physiol.* 1997, **503**, 399–412.
- Fenwick, E. M., A. Marty & E. Neher: Sodium and calcium channels in bovine chromaffin cells. *J. Physiol. (Lond.)* 1982, **331**, 599–635.
- Fernandez-Chacon, R., A. Königstorfer, S. H. Gerber, J. Garcia, M. F. Matos, C. F. Stevens, N. Brose, J. Rizo, C. Rosenmund & T. C. Südhof: Synaptotagmin I functions as a calcium regulator of release probability. *Nature* 2001, **410**, 41–49.
- Fuller, C. M., H. H. Deetjen, A. Piiper & I. Schulz: Secretagogue and second messenger-activated  $\text{Cl}^-$  permeabilities in isolated pancreatic zymogen granules. *Pflügers Arch.* 1989, **415**, 29–36.
- Gao, Z., J. Reavey-Cantwell, R. A. Young, P. Jegier & B. A. Wolf: Synaptotagmin III/VII isoforms mediate  $\text{Ca}^{2+}$ -induced insulin secretion in pancreatic islet  $\beta$ -cells. *J. Biol. Chem.* 2000, **275**, 36079–36085.
- Gerich, J. E., M. A. Charles & G. M. Grodsky: Regulation of pancreatic insulin and glucagon secretion. *Annu. Rev. Physiol.* 1976, **38**, 353–388.
- Gorus, F. K., W. J. Malaisse & D. G. Pipeleers: Differences in glucose handling by pancreatic A- and B-cells. *J. Biol. Chem.* 1984, **259**, 1196–1200.
- Gribble, F. M., S. J. Tucker & F. M. Ashcroft: The interaction of nucleotides with the tolbutamide block of cloned ATP-sensitive  $\text{K}^+$  channel currents expressed in *Xenopus* oocytes: a reinterpretation. *J. Physiol. (Lond.)* 1997, **504**, 35–45.
- Gromada, J., K. Bokvist, W. G. Ding, S. Barg, K. Buschard, E. Renström & P. Rorsman: Adrenaline stimulates glucagon secretion in pancreatic A-cells by increasing the  $\text{Ca}^{2+}$  current and the number of granules close to the L-type  $\text{Ca}^{2+}$  channels. *J. Gen. Physiol.* 1997, **110**, 217–228.
- Gromada, J., M. Hoy, E. Renström, K. Bokvist, L. Eliasson, S. Göpel & P. Rorsman: CaM kinase II-dependent mobilization of secretory granules underlies acetylcholine-induced stimulation of exocytosis in mouse pancreatic B-cells. *J. Physiol. (Lond.)* 1999, **518**, 745–759.
- Günther, W., A. Luchow, F. Cluzeaud, A. Vandewalle & T. J. Jentsch: CIC-5, the chloride channel mutated in Dent's disease, colocalizes with the proton pump in endocytotically active kidney cells. *Proc. Natl. Acad. Sci. USA* 1998, **95**, 8075–8080.
- Göpel, S., T. Kanno, S. Barg, J. Galvanovskis & P. Rorsman: Voltage-gated and resting membrane currents recorded from B-cells in intact mouse pancreatic islets. *J. Physiol. (Lond.)* 1999, **521**, 717–728.
- Göpel, S. O., T. Kanno, S. Barg, X. Weng, J. Gromada & P. Rorsman: Regulation of glucagon release in mouse  $\alpha$ -cells by  $\text{K}_{\text{ATP}}$  channels and inactivation of TTX-sensitive  $\text{Na}^+$  channels. *J. Physiol. (Lond.)* 2000, **528**, 509–520.
- Hay, J. C. & T. F. Martin: Resolution of regulated secretion into sequential MgATP-dependent and calcium-dependent stages mediated by distinct cytosolic proteins. *J. Cell. Biol.* 1992, **119**, 139–151.
- Heidelberger, R., C. Heinemann, E. Neher & G. Matthews: Calcium dependence of the rate of exocytosis in a synaptic terminal. *Nature* 1994, **371**, 513–515.
- Hirning, L. D., A. P. Fox, E. W. McCleskey, B. M. Olivera, S. A. Thayer, R. J. Miller & R. W. Tsien: Dominant role of N-type  $\text{Ca}^{2+}$  channels in evoked release of norepinephrine from sympathetic neurons. *Science* 1988, **239**, 57–61.
- Holz, G. G., C. A. Leech, R. S. Heller, M. Castonguay & J. F. Habener: cAMP-dependent mobilization of intracellular  $\text{Ca}^{2+}$  stores by activation of ryanodine receptors in pancreatic  $\beta$ -cells. A  $\text{Ca}^{2+}$  signaling system stimulated by the insulinotropic hormone glucagon-like peptide-1(7–37). *J. Biol. Chem.* 1999, **274**, 14147–14156.

- Hutton, J. C.: Insulin secretory granule biogenesis and the proinsulin-processing endopeptidases. *Diabetologia* 1994, **37**, S48–S56.
- Høy, M., H. L. Olsen, K. Bokvist, K. Buschard, S. Barg, P. Rorsman & J. Gromada: Tolbutamide stimulates exocytosis of glucagon by inhibition of a mitochondrial-like ATP-sensitive  $K^+$  ( $K_{ATP}$ ) conductance in rat pancreatic A-cells. *J. Physiol. (Lond.)* 2000, **527**, 109–120.
- Klenchin, V. A. & T. F. Martin: Priming in exocytosis: attaining fusion-competence after vesicle docking. *Biochimie* 2000, **82**, 399–407.
- Koushika, S. P., J. E. Richmond, G. Hadwiger, R. M. Weimer, E. M. Jorgensen & M. L. Nonet: A post-docking role for active zone protein Rim. *Nat. Neurosci.* 2001, **4**, 997–1005.
- Lang, J., M. Fukuda, H. Zhang, K. Mikoshiba & C. B. Wollheim: The first C2 domain of synaptotagmin is required for exocytosis of insulin from pancreatic  $\beta$ -cells: action of synaptotagmin at low micromolar calcium. *EMBO J.* 1997, **16**, 5837–5846.
- Lang, J.: Molecular mechanisms and regulation of insulin exocytosis as a paradigm of endocrine secretion. *Eur. J. Biochem.* 1999, **259**, 3–17 (review).
- Littleton, J. T., M. Stern, K. Schulze, M. Perin & H. J. Bellen: Mutational analysis of *Drosophila* synaptotagmin demonstrates its essential role in  $Ca^{2+}$ -activated neurotransmitter release. *Cell* 1993, **74**, 1125–1134.
- Lopez, M. G., A. Albillos, M. T. de la Fuente, R. Borges, L. Gandia, E. Carbone, A. G. Garcia & A. R. Artalejo: Localized L-type calcium channels control exocytosis in cat chromaffin cells. *Pflügers Arch.* 1994, **427**, 348–354.
- Mennerick, S. & G. Matthews: Ultrafast exocytosis elicited by calcium current in synaptic terminals of retinal bipolar neurons. *Neuron* 1996, **17**, 1241–1249.
- Mitchell, K. J., P. Pinton, A. Varadi, C. Tacchetti, E. K. Ainscow, T. Pozzan, R. Rizzuto & G. A. Rutter: Dense core secretory vesicles revealed as a dynamic  $Ca^{2+}$  store in neuroendocrine cells with a vesicle-associated membrane protein aequorin chimera. *J. Cell. Biol.* 2001, **155**, 41–52.
- Mochida, S., Z. H. Sheng, C. Baker, H. Kobayashi & W. A. Catterall: Inhibition of neurotransmission by peptides containing the synaptic protein interaction site of N-type  $Ca^{2+}$  channels. *Neuron* 1996, **17**, 781–788.
- Olofsson, C. S., S. O. Göpel, S. Barg, J. Galvanovskis, X. Ma, A. Salehi, P. Rorsman & L. Eliasson: Fast insulin secretion reflects exocytosis of docked granules in mouse pancreatic B-cells. *Pflügers Arch.* 2002, **444**, 43–51.
- Ozanne, S. E., P. C. Guest, J. C. Hutton & C. N. Hales: Intracellular localization and molecular heterogeneity of the sulphonylurea receptor in insulin-secreting cells. *Diabetologia* 1995, **38**, 277–282.
- Parsons, T. D., J. R. Coorsen, H. Horstmann & W. Almers: Docked granules, the exocytic burst, and the need for ATP hydrolysis in endocrine cells. *Neuron* 1995, **15**, 1085–1096.
- Pickup, J. & G. Williams: *Textbook of diabetes*. Blackwell Science, Malden, Massachusetts, USA, 1997.
- Pipeleers, D. G., F. C. Schuit, C. F. Van Schravendijk & M. Van de Winkel: Interplay of nutrients and hormones in the regulation of glucagon release. *Endocrinology* 1985, **117**, 817–823.
- Piwon, N., W. Günther, M. Schwake, M. R. Bosl & T. J. Jentsch: CIC-5  $Cl^-$  channel disruption impairs endocytosis in a mouse model for Dent's disease. *Nature* 2000, **408**, 369–373.
- Plattner, H., A. R. Artalejo & E. Neher: Ultrastructural organization of bovine chromaffin cell cortex-analysis by cryofixation and morphometry of aspects pertinent to exocytosis. *J. Cell. Biol.* 1997, **139**, 1709–1717.
- Porte, D.: Banting lecture 1990. Beta-cells in type II diabetes mellitus. *Diabetes* 1991, **40**, 166–180 (review).
- Qian, W. J. & R. T. Kennedy: Spatial organization of  $Ca^{2+}$ -entry and exocytosis in mouse pancreatic  $\beta$ -cells. *Biochem. Biophys. Res. Commun.* 2001, **286**, 315–321.
- Renström, E., W. G. Ding, K. Bokvist & P. Rorsman: Neurotransmitter-induced inhibition of exocytosis in insulin-secreting  $\beta$ -cells by activation of calcineurin. *Neuron* 1996a, **17**, 513–522.
- Renström, E., L. Eliasson, K. Bokvist & P. Rorsman: Cooling inhibits exocytosis in single mouse pancreatic B-cells by suppression of granule mobilization. *J. Physiol. (Lond.)* 1996, **494**, 41–52.
- Renström, E., S. Barg, F. Thevenod & P. Rorsman: Sulfonylurea-mediated stimulation of insulin exocytosis via an ATP-sensitive  $K^+$ -channel-independent action. *Diabetes* 2002, **51**, S33–S36.
- Rorsman, P. & B. Hellman: Voltage-activated currents in guinea pig pancreatic  $\alpha 2$  cells. Evidence for  $Ca^{2+}$ -dependent action potentials. *J. Gen. Physiol.* 1988, **91**, 223–242.
- Rorsman, P., L. Eliasson, E. Renström, J. Gromada, S. Barg & S. Göpel: The cell physiology of biphasic insulin secretion. *News Physiol. Sci.* 2000, **15**, 72–77.
- Rupnik, M. & R. Zorec: Cytosolic chloride ions stimulate  $Ca^{2+}$ -induced exocytosis in melanotrophs. *FEBS Lett.* 1992, **303**, 221–223.
- Rutter, G. A., J. M. Theler, M. Murgia, C. B. Wollheim, T. Pozzan & R. Rizzuto: Stimulated  $Ca^{2+}$  influx raises mitochondrial free  $Ca^{2+}$  to supramicromolar levels in a pancreatic  $\beta$ -cell line. Possible role in glucose and agonist-induced insulin secretion. *J. Biol. Chem.* 1993, **268**, 22385–22390.
- Scheenen, W. J., C. B. Wollheim, T. Pozzan & C. Fasolato:  $Ca^{2+}$  depletion from granules inhibits exocytosis. A study with insulin-secreting cells. *J. Biol. Chem.* 1998, **273**, 19002–19008.
- Schwiebert, E. M., D. J. Benos, M. E. Egan, M. J. Stutts & W. B. Guggino: CFTR is a conductance regulator as well as a chloride channel. *Physiol. Rev.* 1999, **79**, S145–S166.
- Sheng, Z. H., J. Rettig, T. Cook & W. A. Catterall: Calcium-dependent interaction of N-type calcium channels with the synaptic core complex. *Nature* 1996, **379**, 451–454.
- Shindo, T., M. Yamada, S. Isomoto, Y. Horio & Y. Kurachi: SUR2 subtype (A and B)-dependent differential activation of the cloned ATP-sensitive  $K^+$  channels by pinacidil and nicorandil. *Brit. J. Pharmacol.* 1998, **124**, 985–991.
- Sutton, R. B., B. A. Davletov, A. M. Berghuis, T. C. Südhof & S. R. Sprang: Structure of the first C2 domain of synaptotagmin I: a novel  $Ca^{2+}$ /phospholipid-binding fold. *Cell* 1995, **80**, 929–938.
- Sutton, R. B., D. Fasshauer, R. Jahn & A. T. Brünger: Crystal structure of a SNARE complex involved in synaptic exocytosis at 2.4 Å resolution. *Nature* 1998, **395**, 347–353.
- Söllner, T., S. W. Whiteheart, M. Brunner, H. Erdjument-Bromage, S. Geromanos, P. Tempst & J. E. Rothman: SNAP receptors implicated in vesicle targeting and fusion. *Nature* 1993, **362**, 318–324.
- Takahashi, N., T. Kadowaki, Y. Yazaki, Y. Miyashita & H. Kasai: Multiple exocytotic pathways in pancreatic beta cells. *J. Cell. Biol.* 1997, **138**, 55–64.
- Taylor, S. C., E. Carpenter, M. L. Roberts & C. Peers: Potentiation of quantal catecholamine secretion by glibenclamide: evidence for a novel role of sulphonylurea receptors in regulating the  $Ca^{2+}$ -sensitivity of exocytosis. *J. Neurosci.* 1999, **19**, 5741–5749.
- Tengholm, A., C. Hagman, E. Gylfe & B. Hellman: *In situ* characterization of nonmitochondrial  $Ca^{2+}$  stores in individual pancreatic  $\beta$ -cells. *Diabetes* 1998, **47**, 1224–1230.
- Thevenod, F., K. V. Chathadi, B. Jiang & U. Hopfer: ATP-sensitive  $K^+$  conductance in pancreatic zymogen granules: block by glyburide and activation by diazoxide. *J. Membr. Biol.* 1992, **129**, 253–266.
- Thevenod, F., I. Anderie & I. Schulz: Monoclonal antibodies against MDR1 P-glycoprotein inhibit chloride conductance and label a 65-kDa protein in pancreatic zymogen granule membranes. *J. Biol. Chem.* 1994, **269**, 24410–24417.
- Tian, Y. A., G. Johnson & S. J. Ashcroft: Sulfonylureas enhance exocytosis from pancreatic beta-cells by a mechanism that does not involve direct activation of protein kinase C. *Diabetes* 1998, **47**, 1722–1726.
- Trube, G., P. Rorsman & T. Ohno-Shosaku: Opposite effects of tol-

- butamide and diazoxide on the ATP-dependent  $K^+$  channel in mouse pancreatic  $\beta$ -cells. *Pflügers Arch.* 1986, **407**, 493–499.
- Uchitel, O. D., D. A. Protti, V. Sanchez, B. D. Cherksey, M. Sugimori & R. Llinas: P-type voltage-dependent calcium channel mediates presynaptic calcium influx and transmitter release in mammalian synapses. *Proc. Natl. Acad. Sci. U S A* 1992, **89**, 3330–3333.
- Unger, R. H.: Glucagon physiology and pathophysiology. *New Engl. J. Med.* 1971, **285**, 443–449.
- Ungermann, C., W. Wickner & Z. Xu: Vacuole acidification is required for trans-SNARE pairing, LMA1 release, and homotypic fusion. *Proc. Natl. Acad. Sci. U S A* 1999, **96**, 11194–11199.
- van der Meer, C., P. W. Valkenburg & J. A. Versluys-Broers: Disturbances in the glucose metabolism in intestinal ischemia shock. *Circ. Shock* 1981, **8**, 375–392.
- von Gersdorff, H., E. Vardi, G. Matthews & P. Sterling: Evidence that vesicles on the synaptic ribbon of retinal bipolar neurons can be rapidly released. *Neuron* 1996, **16**, 1221–1227.
- Wiser, O., D. Tobi, M. Trus & D. Atlas: Synaptotagmin restores kinetic properties of a syntaxin-associated N-type voltage sensitive calcium channel. *FEBS Lett.* 1997, **404**, 203–207.
- Wiser, O., M. Trus, A. Hernandez, E. Renström, S. Barg, P. Rorsman & D. Atlas: The voltage sensitive Lc-type  $Ca^{2+}$ -channel is functionally coupled to the exocytotic machinery. *Proc. Natl. Acad. Sci. USA*, 1999, **96**, 248–253.
- Woodbury, D. J.: Evaluation of the evidence for ion channels in synaptic vesicles. *Mol. Membr. Biol.* 1995, **12**, 165–171.
- Wu, M. N., T. Fergestad, T. E. Lloyd, Y. He, K. Broadie & H. J. Bellen: Syntaxin 1A interacts with multiple exocytic proteins to regulate neurotransmitter release in vivo. *Neuron* 1999, **23**, 593–605.
- Ämmälä, C., L. Eliasson, K. Bokvist, O. Larsson, F.M. Ashcroft & P. Rorsman: Exocytosis elicited by action potentials and voltage-clamp calcium currents in individual mouse pancreatic B-cells. *J. Physiol (Lond)*, 1993, **472**, 665–688.