

# LUND UNIVERSITY

## Optimal pumping of orbital entanglement with single-particle emitters

Sherkunov, Y.; d'Ambrumenil, N.; Samuelsson, Peter; Buettiker, M.

Published in: Physical Review B (Condensed Matter and Materials Physics)

DOI: 10.1103/PhysRevB.85.081108

2012

Link to publication

Citation for published version (APA): Sherkunov, Y., d'Ambrumenil, N., Samuelsson, P., & Buettiker, M. (2012). Optimal pumping of orbital entanglement with single-particle emitters. Physical Review B (Condensed Matter and Materials Physics), 85(8), Article 081108. https://doi.org/10.1103/PhysRevB.85.081108

Total number of authors: 4

General rights

Unless other specific re-use rights are stated the following general rights apply: Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

· Users may download and print one copy of any publication from the public portal for the purpose of private study

or research.
You may not further distribute the material or use it for any profit-making activity or commercial gain

· You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: https://creativecommons.org/licenses/

#### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

#### LUND UNIVERSITY

**PO Box 117** 221 00 Lund +46 46-222 00 00

### **Optimal pumping of orbital entanglement with single-particle emitters**

Y. Sherkunov,<sup>1,2</sup> N. d'Ambrumenil,<sup>2</sup> P. Samuelsson,<sup>3</sup> and M. Büttiker<sup>4</sup>

<sup>1</sup>Physics Department, Lancaster University, Lancaster LA1 4YB, United Kingdom

<sup>2</sup>Department of Physics, University of Warwick, Coventry CV4 7AL, United Kingdom

<sup>3</sup>Division of Mathematical Physics, Lund University, Box 118, SE-221 00 Lund, Sweden

<sup>4</sup>Département de Physique Théorique, Université de Genève, CH-1211 Genève 4, Switzerland

(Received 6 January 2012; published 27 February 2012)

We propose a method for the optimal time-controlled generation of entangled itinerant particles, using ondemand sources in a conductor in the quantum Hall regime. This entanglement pump is realized by applying periodic, tailored voltage pulses to pairs of quantum dots or quantum point contacts. We show that the pump can produce orbital Bell pairs of both electrons and holes at the optimal rate of half a pair per pumping cycle. The entanglement can be detected by a violation of a Bell inequality formulated in terms of low-frequency current cross correlations.

DOI: 10.1103/PhysRevB.85.081108

PACS number(s): 73.23.-b, 42.50.Lc, 03.67.Mn

Entanglement of itinerant electrons in mesoscopic and nanoscale systems continues to attract great attention. Despite intense effort, a clear demonstration of the generation, spatial separation and detection of entangled pairs of electrons is still missing. Of particular interest would be an on-demand source for entanglement,<sup>1-4</sup> while a key component in quantum information processing is the time-controlled production of entangled flying quantum bits. A scheme for the dynamical generation, or pumping, of orbitally entangled electron-hole pairs was proposed by two of us in Ref. 1. However, as the proposed pump operated in the weak amplitude regime, it produced on average much less than one Bell pair per cycle. Subsequently it was shown that, for strong amplitude pumping of noninteracting particles, the optimal production rate is half a Bell pair per cycle.<sup>2</sup> Optimal electron-hole pair entanglement pumps have also been proposed, but without any scheme for entanglement detection.<sup>2,4</sup>

An important step toward entanglement pumping was taken experimentally by Fève *et al.*,<sup>5</sup> who realized a singleparticle on-demand source in a conductor in the quantum Hall regime.<sup>6–11</sup> Under ideal conditions, the source produces exactly one electron and one hole per cycle. A scheme for the production of pairs of orbitally entangled particles in different time bins based on two on-demand sources in a double electronic Mach-Zehnder interferometer was proposed in Refs. 12 and 13. The origin of the entanglement was two-particle interference,<sup>14–16</sup> manifested by a nonlocal two-particle Aharonov-Bohm effect. The maximum production was 1/4 Bell pair per cycle, i.e., half the optimal rate.

Here we propose an entanglement pump, aiming for the simplest scheme for detection and optimal production of orbital entanglement in the same system (see Fig. 1). A conductor in the quantum Hall regime with two on-demand sources, C and D, is connected to four terminals via electronic beam splitters at A and B. Using a single spin-polarized edge state and the interferometer of Ref. 1, the sources operate in the strong amplitude regime and generate pairs of entangled particles at an optimal rate.

The two types of sources, driven by voltage pulses  $V_C(t)$ and  $V_D(t)$ , are shown in Figs. 1(b) and 1(c). Figure 1(b) shows the level of a quantum dot (QD) driven up (down) through Fermi energy, generating a single electron (hole), as in the experiment of Ref. 5. However, with the QD coupled to separate quantum Hall edge states, the particle is emitted into a superposition of states at the two edges. In Fig. 1(c), a quantum point contact (QPC) is opened and closed to generate exactly one particle and one hole, which are independently emitted into edge state superpositions. We show below that, for QD as well as QPC sources, driving C and D periodically generates orbitally entangled pairs of both electron and hole wave packets, with one particle propagating toward A and one toward B. For synchronized and spatially symmetric sources the entanglement production rate is optimal, with half a Bell pair per cycle. Using earlier results for many-body states of emitted particles of on-demand sources,<sup>4,6,9</sup> we derive explicit expressions for the entangled wave function. The entanglement, arising from two-particle interference,<sup>14–16</sup> can be detected via low-frequency cross correlations of currents at the four terminals 1-4. Throughout the Rapid Communication we consider zero temperature and put  $\hbar, e = 1$ .

*QD sources*. We first consider QD sources. The aim is to obtain the many-body wave function of the entangled particles emitted from the sources toward *A* and *B*. We consider the scattering properties of source *C* for a single-electron-emission event. The QD has a single localized level at energy E(t) and tunnel couplings to the edges toward *A* and *B*. If the energy of the level is varied slowly on the scale of the Wigner delay time (adiabatic approximation), the time development of the scattering matrix  $S_C$ . When the energy of the level is varied at constant speed,  $E(t) = v(t - t_C)$ ,  $S_C$  takes the Breit-Wigner form

$$S_C = \frac{1}{t - t_C - i\tau_C} \begin{pmatrix} t - t_C + i\overline{\tau}_C & -2i\sqrt{\tau_{CA}\tau_{CB}} \\ -2i\sqrt{\tau_{CA}\tau_{CB}} & t - t_C - i\overline{\tau}_C \end{pmatrix}, \quad (1)$$

with  $\tau_C = \tau_{CA} + \tau_{CB}$  and  $\bar{\tau}_C = \tau_{CA} - \tau_{CB}$ . Here  $\nu \tau_{CA}$  and  $\nu \tau_{CB}$  are tunnel rates through the barriers into states propagating toward *A* and *B*.



FIG. 1. (Color online) (a) Optimal orbital entanglement pump with two single-particle emitters, C and D, connected to conductors in the quantum Hall regime. The emitted electrons and holes propagate along edge states to controllable electronic beam splitters at A and Band are detected at terminals 1–4. The sources C and D can be either (b) quantum dots or (c) quantum point contacts. Tailored voltage profiles  $V_C(t)$  and  $V_D(t)$  are applied to a gate in case (b) to move a localized level with energy E(t) up (down) through the Fermi energy releasing an electron (hole), or in case (c) to cycle the transmission probability, T(t), from zero through unity and back to generate a particle-hole excitation.

The scattering matrix  $S_C$  can be diagonalized in a time-independent basis  $\tilde{S}_C = V S_C V^{\dagger} = \begin{pmatrix} e^{i\phi_C} & 0 \\ 0 & 1 \end{pmatrix}$ , with V =

$$\frac{1}{\sqrt{\tau_C}} \left( \frac{-\sqrt{\tau_{CA}}}{\sqrt{\tau_{CB}}} \frac{\sqrt{\tau_{CB}}}{\sqrt{\tau_{CA}}} \right) \text{ and}$$

$$e^{i\phi_C(t)} = \frac{t - t_C + i\tau_C}{t - t_C - i\tau_C}.$$
(2)

The low-temperature many-body state incident on *C* is a filled Fermi sea of electrons originating from terminals 2 and 3. The incident many-body state is  $|\psi_{in}\rangle = \prod_{\epsilon < 0} a^{\dagger}_{C+}(\epsilon)a^{\dagger}_{C-}(\epsilon)|\rangle$ , where the  $a^{\dagger}_{C\pm}(\epsilon)$  are creation operators in the diagonal basis for particles with energy  $\epsilon$  in the incoming channels + or -, and  $|\rangle$  is the vacuum. Denoting the operators for particles in the corresponding outgoing states by  $b^{\dagger}_{C\pm}(\epsilon)$ , the many-body state after impinging on *C* is

$$|\psi_C\rangle = \prod_{\epsilon < 0} b^{\dagger}_{C+}(\epsilon) \sum_{\epsilon'} (e^{i\phi_C})_{\epsilon,\epsilon'} b^{\dagger}_{C-}(\epsilon')|\rangle.$$
(3)

Here  $(e^{i\phi_C})_{\epsilon,\epsilon'}$  is the Fourier transform of  $e^{i\phi_C(t)}$  with respect to the energy difference  $(\epsilon - \epsilon')$ .<sup>17–19</sup>

With the phase profile of (2) and taking  $\nu > 0$ , the many-body state becomes  $|\psi_C^e\rangle = \sqrt{2\tau_C} \sum_{\epsilon>0} \exp[\epsilon(it_C - \tau_C)]b_{C-}^{\dagger}(\epsilon)|0\rangle$ . It describes a single electron above the unperturbed Fermi sea  $|0\rangle$ .<sup>6,9</sup> The wave function of the single-particle excitation is  $\psi(x,t) \sim 1/(x - v_{dr}[t - t_C - i\tau_C])$ , where *x* is the distance from *C* and  $v_{dr}$  the drift velocity along the edge. Using  $[b_{CA}^{\dagger}(\epsilon), b_{CB}^{\dagger}(\epsilon)] = V[b_{C-}^{\dagger}(\epsilon), b_{C+}^{\dagger}(\epsilon)]$  to write  $|\psi_C^e\rangle$  in terms of operators  $b_{CA}^{\dagger}(\epsilon)$  and  $b_{CB}^{\dagger}(\epsilon)$ , which

PHYSICAL REVIEW B 85, 081108(R) (2012)

create particles propagating along the edges toward A and B, we obtain

$$\left|\psi_{C}^{e}\right\rangle = \frac{1}{\sqrt{\tau_{C}}} \left[-\sqrt{\tau_{CA}} p_{CA}^{\dagger} + \sqrt{\tau_{CB}} p_{CB}^{\dagger}\right] \left|0\right\rangle. \tag{4}$$

Here the normalized wave-packet operator  $p_{CA}^{\dagger} = \sqrt{2\tau_C} \sum_{\epsilon>0} \exp[\epsilon(it_C - \tau_C)] b_{CA}^{\dagger}(\epsilon)$  and similarly for  $p_{CB}^{\dagger}$ . Equation (4) shows that driving the QD level at *C* up through the Fermi level injects a single electron into a linear superposition of states propagating toward *A* and *B*.

The corresponding result to (4), when driving the QD level at *D* up through the Fermi level, is  $|\psi_D^e\rangle = (1/\sqrt{\tau_D})[-\sqrt{\tau_{DA}}e^{-i\eta}p_{DA}^{\dagger} + \sqrt{\tau_{DB}}e^{i\eta}p_{DB}^{\dagger}]|0\rangle$ . Here  $p_{DA}^{\dagger}$  and  $p_{DB}^{\dagger}$  are obtained from  $p_{CA}^{\dagger}$  and  $p_{CB}^{\dagger}$  by changing indices  $C \leftrightarrow D$ . We have also included a phase factor  $e^{i\eta}$  to account for a possible Aharonov-Bohm phase. The total wave function for the electrons emitted from *C* and *D* is then the product  $|\psi_{out}^{ee}\rangle = |\psi_C^e\rangle|\psi_D^e\rangle$ . We are interested in the nonlocal properties of  $|\psi_{out}^{ee}\rangle$ , with the particles emitted toward different beam splitters. Projecting  $|\psi_{out}^{ee}\rangle$  onto the subspace with one particle at *A* and one at *B* gives the normalized wave function<sup>20</sup>

$$\begin{split} \left|\psi_{AB}^{ee}\right\rangle &= \frac{1}{\sqrt{N}} [\sqrt{\tau_{CA} \tau_{DB}} p_{CA}^{\dagger} p_{DB}^{\dagger} e^{i\eta} \\ &+ \sqrt{\tau_{CB} \tau_{DA}} p_{CB}^{\dagger} p_{DA}^{\dagger} e^{-i\eta} ] \left|0\right\rangle, \end{split}$$
(5)

where  $N = \tau_{CA}\tau_{DB} + \tau_{CB}\tau_{DA}$ . The weight, or probability that  $|\psi_{AB}^{ee}\rangle$  is generated, is  $w_{AB} = N/(\tau_C \tau_D)$ . The wave function  $|\psi_{AB}^{ee}\rangle$  describes two particles above

The wave function  $|\psi_{AB}^{ee}\rangle$  describes two particles above the filled Fermi sea. It is entangled in the orbital (i.e., source *C* and *D*) degree of freedom—a result of two-particle interference.<sup>14</sup> The superposition in  $|\psi_{AB}^{ee}\rangle$  results from the indistinguishability of the two emission processes leading to a particle at *A* and one at *B*. In one process a particle moves from *C* to *A* and another from *D* to *B*. In the second process particles move from *D* to *A* and from *C* to *B*. To quantify the orbital entanglement we consider the reduced  $4 \times 4$  orbital density matrix  $\rho_{AB}$ . This is obtained by tracing  $|\psi_{AB}^{ee}\rangle\langle\psi_{AB}^{ee}|$ over energy:<sup>21</sup>

$$\rho_{AB} = N^{-1} (\tau_{CA} \tau_{DB} | CD \rangle \langle CD | + \tau_{CB} \tau_{DA} | DC \rangle \langle DC | + \chi \sqrt{\tau_{CA} \tau_{CB} \tau_{DA} \tau_{DB}} [e^{i\eta} | CD \rangle \langle DC | + \text{H.c.}]), \quad (6)$$

where  $|CD\rangle \equiv |C\rangle_A |D\rangle_B$  and  $|C\rangle_A$  denotes an electron at *A* emitted from *C*, etc. Here

$$\chi = \frac{4\tau_C \tau_D}{(t_C - t_D)^2 + (\tau_C + \tau_D)^2}$$
(7)

quantifies the overlap  $(0 \le \chi \le 1)$  between the two wave packets emitted from *C* and *D*. We note that a reduced overlap  $\chi < 1$  plays the same role as a nonzero dephasing for two-particle interference.<sup>16</sup>

The entanglement of  $\rho_{AB}$  is conveniently quantified via the concurrence C, which ranges from 0 for a separable nonentangled state to 1 for a maximally entangled state.<sup>22</sup> For  $\rho_{AB}$  we find

$$\mathcal{C} = 2(\chi/N)\sqrt{\tau_{CA}\tau_{CB}\tau_{DA}\tau_{DB}},\tag{8}$$

which is nonzero for arbitrary small overlap  $\chi$  and arbitrary tunnel couplings.<sup>12</sup> For QDs synchronized in time,  $t_C = t_D$ ,

with symmetric couplings  $\tau_{CA} = \tau_{CB} = \tau_C/2$  and  $\tau_{DA} = \tau_{DB} = \tau_D/2$ , the density matrix  $\rho_{AB} = |\varphi_{AB}^{ee}\rangle\langle\varphi_{AB}^{ee}|$  with  $|\varphi_{AB}^{ee}\rangle = 1/\sqrt{2}[e^{i\eta}|CD\rangle + e^{-i\eta}|DC\rangle]$ . This state is an orbital Bell pair, i.e., it is maximally entangled (C = 1). Moreover, the weight  $w_{AB} = 1/2$  and hence the concurrence production per cycle is  $w_{AB}C = 1/2$ , the theoretical maximum.<sup>20</sup>

If the QD level at *C* is driven down through the Fermi level at time  $t'_C$ , a hole is generated in a linear superposition of states in the edges toward *A* and *B*, with a wave function  $|\psi^h_C\rangle =$  $(1/\sqrt{\tau_C})[-\sqrt{\tau_{CA}}h^{\dagger}_{CA} + \sqrt{\tau_{CB}}h^{\dagger}_{CB}]|0\rangle$ . Here the hole wavepacket creation operator is  $h^{\dagger}_{CA} = \sqrt{2\tau_C} \sum_{\epsilon < 0} \exp[\epsilon(it'_C + \tau_C)]b_{CA}(\epsilon)$ . A similar relation holds for  $|\psi^h_D\rangle$ , giving a total hole wave function  $|\psi^{hh}_{out}\rangle = |\psi^h_C\rangle|\psi^h_D\rangle$ .

For a large amplitude driving up and subsequently down through the Fermi energy, with  $t'_C - t_C \gg \tau_C$  and  $t'_D - t_D \gg \tau_D$ , the electron and hole emissions are well separated in time and can be treated as independent.<sup>8,9</sup> The total wave function for the emitted particles is then  $|\psi_{out}^{QD}\rangle = |\psi_{out}^{ee}\rangle|\psi_{out}^{hh}\rangle$ . In the experimentally relevant situation,<sup>5</sup> with a cycling of the QD level with period  $T \gg \tau_C, \tau_D$ , the pump produces pairs of both entangled electrons and holes which, for symmetric and syncronized sources, reaches the optimal production rate of 1/2 a Bell pair per period.

*QPC sources*. For QPC sources, the emitted state is created by applying tailored voltage pulses  $V_{C,D}(t)$  to the gates *C* and *D*.<sup>4</sup> The instantaneous scattering matrix of gate *C* can be written

$$S_C = \begin{pmatrix} \lambda_C(t) & \kappa_C(t) \\ -\kappa_C^*(t) & \lambda_C^*(t) \end{pmatrix}, \tag{9}$$

where  $|\kappa_C(t)|^2$  and  $|\lambda_C(t)|^2$  are the transmission and reflection probabilities through the QPC, respectively. Provided no voltage is applied across the device,  $S_C$  can be diagonalized<sup>4</sup> in a time-independent basis  $\tilde{S}_C = V S_C V^{\dagger} = \begin{pmatrix} e^{i\phi_C} & 0\\ 0 & e^{-i\phi_C} \end{pmatrix}$ , with  $V = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & -i\\ 1 & i \end{pmatrix}$  and  $e^{i\phi_C(t)} = \lambda_C(t) + i\kappa_C(t)$ . The outgoing manybody wave function can then be written in the diagonal basis

$$\bar{\psi}_C \rangle = \frac{1}{2} \prod_{\epsilon < 0} \sum_{\epsilon', \epsilon''} (e^{i\phi_C})_{\epsilon, \epsilon'} (e^{-i\phi_C})_{\epsilon, \epsilon''} b^{\dagger}_{-}(\epsilon') b^{\dagger}_{+}(\epsilon'') |\rangle.$$
(10)

By varying  $V_C(t)$  so that the transmission amplitude  $\kappa_C(t) = 2\tau_C(t - t_C)/[(t - t_C)^2 + \tau_C^2]$ , the phase factor  $e^{i\phi_C(t)}$  is again given by (2). In this case we can write the outgoing many-body wave function<sup>6</sup>

$$\left|\psi_{C}^{eh}\right\rangle = \frac{1}{2}(p_{CA}^{\dagger} + ip_{CB}^{\dagger})(h_{CA}^{\dagger} - ih_{CB}^{\dagger})|0\rangle \tag{11}$$

in the unrotated basis with a corresponding result for  $|\psi_D^{eh}\rangle$ . The wave functions  $|\psi_{C,D}^{eh}\rangle$  describe one electron and one hole wave packet, independently emitted on top of the unperturbed Fermi sea, in superpositions which describe excitations propagating toward *A* and *B*. Consequently, the total state  $|\psi_{out}^{QPC}\rangle = |\psi_C^{eh}\rangle |\psi_D^{eh}\rangle$ , as for the QD, is the direct product of independently emitted pairs of electrons and holes. The difference is that for the QPC the electrons and holes at *C*(*D*) are emitted at the same time,  $t'_C = t_C$  and  $t'_D = t_D$ . The reduced density matrix and the concurrence of the electron and hole states at *A* and *B* are given by Eqs. (6) and (8), respectively,

#### PHYSICAL REVIEW B 85, 081108(R) (2012)

after taking  $\tau_{CA} = \tau_{CB} = \tau_C/2$  and  $\tau_{DA} = \tau_{DB} = \tau_D/2$ . For perfectly synchronized driving  $t_C = t_D$  the QPC also works as an optimal entanglement pump, producing independent pairs of entangled electrons and holes at *A* and *B* with a rate 1/2 per cycle.

*Bell inequality.* The existence of entanglement in the system illustrated in Fig. 1 can be verified experimentally by demonstrating violation of a Bell inequality (BI). A BI can conveniently be formulated in terms of the joint, or coincident, probabilities to detect quasiparticle excitations,<sup>14</sup> as  $M \leq 2$  for the Bell parameter

$$M = |E(\alpha, \beta) - E(\alpha, \beta') + E(\alpha', \beta) + E(\alpha', \beta')|.$$
(12)

Here the correlation function

$$E(\alpha,\beta) = \frac{P_{24} + P_{13} - P_{14} - P_{23}}{P_{24} + P_{13} + P_{14} + P_{23}},$$
(13)

where  $P_{ij}$  (=  $P_{ij}^{ee} + P_{ij}^{hh}$ ) is the sum of the probabilities to jointly detect an electron or hole at A (i = 1,2) and B (i = 3,4) during a pumping cycle,<sup>1</sup> and is given by

$$P_{ij}^{ee} = \int_{0}^{\infty} d\epsilon \int_{0}^{\infty} d\epsilon' \langle b_{i}^{\dagger}(\epsilon) b_{j}^{\dagger}(\epsilon') b_{j}(\epsilon') b_{i}(\epsilon) \rangle,$$

$$P_{ij}^{hh} = \int_{-\infty}^{0} d\epsilon \int_{-\infty}^{0} d\epsilon' \langle b_{i}(\epsilon) b_{j}(\epsilon') b_{j}^{\dagger}(\epsilon') b_{i}^{\dagger}(\epsilon) \rangle.$$
(14)

The annihilation operators for the states entering the terminals i, j at A are

$$\binom{b_2}{b_1} = \binom{\cos\alpha & \sin\alpha}{-\sin\alpha & \cos\alpha} \binom{b_{CA}}{b_{DA}}.$$
 (15)

The operators at *B* are obtained after setting  $\alpha \rightarrow \beta$ ,  $1 \rightarrow 3$ , and  $2 \rightarrow 4$ . From the emitted many-body wave functions for the QD and the QPC, together with the operator relations at the beam splitters (15), we obtain the joint detection probability

$$P_{13}^{ee} = (2\tau_{CA}\tau_{DB}\sin^2\alpha\cos^2\beta + 2\tau_{CB}\tau_{DA}\sin^2\beta\cos^2\alpha - \chi\cos\eta\sqrt{\tau_{CA}\tau_{DB}\tau_{CB}\tau_{DA}}\sin2\alpha\sin2\beta)/(2\tau_C\tau_D),$$
(16)

with corresponding results for the other  $P_{ij}^{ee}$   $(P_{ij}^{hh} = P_{ij}^{ee})^{.23}$ For the QPC source we put  $\tau_{CA} = \tau_{CB} = \tau_C/2$  and  $\tau_{DA} = \tau_{DB} = \tau_D/2$ . Inserting the  $P_{ij}$  into (13) gives

$$E(\alpha,\beta) = -(\cos 2\alpha \cos 2\beta + C \cos \eta \sin 2\alpha \sin 2\beta), \quad (17)$$

where C is the concurrence in Eq. (8). By optimizing the settings  $\alpha$ ,  $\alpha'$ ,  $\beta$ , and  $\beta'$  of the beam splitters *A* and *B*,<sup>24</sup> the BI reduces to  $2\sqrt{1 + C^2 \cos^2 \eta} \leq 2$ , which can be violated for arbitrary overlap  $\chi$  and phase  $\eta$ .

The joint detection probabilities in Eq. (14) are presently not directly accessible in mesoscopic conductors, as they require time-resolved correlation measurements on the time scale of the period *T*. It is, however, possible to express  $P_{ij}$  in terms of experimentally available currents and low-frequency current cross correlators. The period-averaged, zero-frequency cross correlations  $S_{ij}$  (Ref. 25) are found to be

$$S_{ij} = T^{-1} \left( P_{ij}^{ee} - P_i^e P_j^e + P_{ij}^{hh} - P_i^h P_j^h \right).$$
(18)

Note that  $S_{ij}$  does not contain any electron-hole correlations, as a consequence of the independent emission of electrons and

holes, discussed above. In Eq. (18) the (single-particle) probability to detect an electron in lead *i* is  $P_i^e = \int_0^\infty d\epsilon \langle b_i^{\dagger}(\epsilon) b_i(\epsilon) \rangle$ , while  $P_i^h$  is the probability to detect a hole. For the QD,  $P_i^e$ and  $P_i^h$  are available from  $I_i(t)$ , which is the experimentally accessible time-resolved current.<sup>5</sup> By integrating over half cycles, driving the QD-level up (down) during the first (second) half, we have  $(1/e) \int_0^{T/2} dt I_i(t) = Q_i^e/e = P_i^e$  and  $(1/e) \int_{T/2}^T dt I_i(t) = Q_i^h/e = P_i^h$ . Here  $Q_i^e$  and  $Q_i^h$  are the electron and hole charge flowing into contact *i* during the cycle. For the QPC,  $P_i^e$  and  $P_i^h$  can instead be obtained in a more indirect way via the low-frequency current autocorrelations, which we do not discuss here. Taken together, this allows PHYSICAL REVIEW B 85, 081108(R) (2012)

us to express the correlation function  $E(\alpha,\beta)$ , and hence the Bell inequality, directly in terms of currents and current correlations.

In conclusion, we have proposed an optimal entanglement on-demand source in a noninteracting mesoscopic conductor in the quantum Hall regime. Pairs of entangled, spatially separated electrons and holes are generated by applying tailored, time-dependent voltage pulses to quantum dots or quantum point contacts. The entanglement can be investigated by current and current cross-correlation measurements.

We acknowledge discussions with B. Muzykantskii.

- <sup>1</sup>P. Samuelsson and M. Büttiker, Phys. Rev. B **71**, 245317 (2005).
- <sup>2</sup>C. W. J. Beenakker, M. Titov, and B. Trauzettel, Phys. Rev. Lett. **94**, 186804 (2005).
- <sup>3</sup>K. K. Das, S. Kim, and A. Mizel, Phys. Rev. Lett. **97**, 096602 (2006).
- <sup>4</sup>Y. Sherkunov, J. Zhang, N. d'Ambrumenil, and B. Muzykantskii, Phys. Rev. B **80**, 041313 (2009).
- <sup>5</sup>G. Fève, A. Mahé, J.-M. Berroir, T. T. Kontos, B. Plaçais, D. C. Glattli, A. Cavanna, B. Etienne, and Y. Jin, Science **316**, 1169 (2007).
- <sup>6</sup>J. Keeling, I. Klich, and L. S. Levitov, Phys. Rev. Lett. **97**, 116403 (2006).
- <sup>7</sup>M. Vanević, Y. V. Nazarov, and W. Belzig, Phys. Rev. Lett. **99**, 076601 (2007).
- <sup>8</sup>M. Moskalets, P. Samuelsson, and M. Büttiker, Phys. Rev. Lett. **100**, 086601 (2008).
- <sup>9</sup>J. Keeling, A. V. Shytov, and L. S. Levitov, Phys. Rev. Lett. **101**, 196404 (2008).
- <sup>10</sup>J. Zhang, Y. Sherkunov, N. d'Ambrumenil, and B. Muzykantskii, Phys. Rev. B **80**, 245308 (2009).
- <sup>11</sup>F. Battista and P. Samuelsson, Phys. Rev. B 83, 125324 (2011).
- <sup>12</sup>S. Ol'khovskaya, J. Splettstoesser, M. Moskalets, and M. Büttiker, Phys. Rev. Lett. **101**, 166802 (2008).
- <sup>13</sup>J. Splettstoesser, M. Moskalets, and M. Büttiker, Phys. Rev. Lett. 103, 076804 (2009).

- <sup>14</sup>P. Samuelsson, E. V. Sukhorukov, and M. Büttiker, Phys. Rev. Lett. 92, 026805 (2004).
- <sup>15</sup>I. Neder, N. Ofek, Y. Chung, M. Heiblum, D. Mahalu, and V. Umansky, Nature (London) **448**, 333 (2007).
- <sup>16</sup>P. Samuelsson, I. Neder, and M. Büttiker, Phys. Rev. Lett. **102**, 106804 (2009).
- <sup>17</sup>Y. Adamov and B. Muzykantskii, Phys. Rev. B **64**, 245318 (2001).
- <sup>18</sup>M. Moskalets and M. Büttiker, Phys. Rev. B 66, 205320 (2002).
- <sup>19</sup>N. d'Ambrumenil and B. Muzykantskii, Phys. Rev. B 71, 045326 (2005).
- <sup>20</sup>C. W. J. Beenakker, in *Quantum Computers, Algorithms and Chaos* Proceedings of the International School of Physics "Enrico Fermi," Course CLXII, Varenna, 2005, edited by G. Casati, D. L. Shepelyansky, P. Zoller, and G. Benenti (IOS Press, Amsterdam, 2006), pp. 307–347, arxiv:cond-mat/0508488.
- <sup>21</sup>P. Samuelsson, E. V. Sukhorukov, and M. Büttiker, New. J. Phys. 7, 176 (2005).
- <sup>22</sup>W. K. Wootters, Phys. Rev. Lett. 80, 2245 (1998).
- <sup>23</sup>We point out that the same result is obtained by employing a Floquet scattering approach (Ref. 18), where no use is made of the wave functions  $|\Psi_{out}^{QD}\rangle$  or  $|\Psi_{out}^{QPC}\rangle$ .
- <sup>24</sup>P. Samuelsson, E. V. Sukhorukov, and M. Büttiker, Phys. Rev. Lett. 91, 157002 (2003).
- <sup>25</sup>Y. M. Blanter and M. Büttiker, Phys. Rep. **336**, 1 (2000).