

## LUND UNIVERSITY

## Identification of Reaction Products in <sup>50</sup>Ti+<sup>249</sup>Cf Reactions at TASCA

Di Nitto, A.; Khuyagbaatar, J.; Ackermann, D.; Adamczewski-Musch, J.; Andersson, Lise-Lotte; Badura, E.; Block, M; Brand, H.; Cox, D.M.; Düllmann, Ch.E.; Dvorak, J.; Eberhardt, K.; Ellison, P. A.; Esker, N. E.; Even, J.; Fahlander, Claes; Forsberg, Ulrika; Gates, J.M.; Golubev, Pavel; Gothe, O.; Gregorich, K.E.; Hartmann, W.; Herzberg, R.-D.; Heßberger, F.P.; Hoffmann, J.; Hollinger, R.; Hübner, A.; Jäger, E.; Jeppsson, J.; Kindler, B.; Klein, S.; Kojouharov, I.; Kratz, J.V.; Krier, J.; Kurz, N.; Lahiri, S.; Linev, S.; Lommel, B.; Maiti, M.; Mändl, R.; Merchán, E.; Minami, S.; Mistry, A.K.; Mokry, Ch.; Nitsche, H.; Omtvedt, J.P.; Pang, G.; Pysmenetska, I.; Renisch, D.; Rudolph, Dirk

Published in: GSI Scientific Report 2016

DOI:

10.15120/GR-2017-1

2017

Document Version:
Publisher's PDF, also known as Version of record

Link to publication

Citation for published version (APA):

Di Nitto, A., Khuyagbaatar, J., Ackermann, D., Adamczewski-Musch, J., Andersson, L.-L., Badura, E., Block, M., Brand, H., Cox, D. M., Düllmann, C. E., Dvorak, J., Eberhardt, K., Ellison, P. A., Esker, N. E., Even, J., Fahlander, C., Forsberg, U., Gates, J. M., Golubev, P., ... Yakusheva, V. (2017). Identification of Reaction Products in Ti+<sup>249</sup>Cf Reactions at TASCA. In K. Große (Ed.), *GSI Scientific Report 2016* (Vol. 2017-1, pp. 203-203). Article RESEARCH-NUSTAR-SHEC-3 (GSI Scientific Report; Vol. 2017-1).. https://doi.org/10.15120/GR-2017-1

Total number of authors:

Creative Commons License: CC BY

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- or research.

   You may not further distribute the material or use it for any profit-making activity or commercial gain

· You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: https://creativecommons.org/licenses/

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Download date: 23. Aug. 2025

**LUND UNIVERSITY** 

PO Box 117 221 00 Lund +46 46-222 00 00

## Identification of reaction products in $^{50}$ Ti + $^{249}$ Cf reactions at TASCA $^*$

A. Di Nitto †1,2, J. Khuyagbaatar<sup>2,3</sup>, D. Ackermann ‡2, J.Adamczewski-Musch², L.-L. Andersson³,4, E. Badura², M. Block², H. Brand², D.M. Cox⁴, Ch.E. Düllmann¹,2,3, J. Dvorak³, K. Eberhardt¹, P.A. Ellison⁵, N.E. Esker⁵, J. Even¹,3, C. Fahlander⁶, U. Forsberg⁶, J.M. Gates⁵, P. Golubev⁶, O. Gothe⁵, K.E. Gregorich⁵, W. Hartmann², R.D. Herzberg⁴, F.P. Heβberger²,3, J. Hoffmann², R. Hollinger², A. Hübner², E. Jäger², J. Jeppsson⁶, B. Kindler², S. Klein¹, I. Kojouharov², J.V. Kratz¹, J. Krier², N. Kurz², S. Lahiri³, S.Linev², B. Lommel², M. Maiti¬, R. Mändl², E. Merchan², S. Minami², A. Mistry⁴, Ch. Mokry¹, H. Nitsche⁵, J.P. Omtvedt⁶, G.K. Pang⁵, I. Pysmenetska², D. Renisch¹,³, D. Rudolph⁶, J. Runke², L.G. Sarmiento⁶,9, M. Schädel²,¹¹⁰, H. Schaffner², B. Schausten², A. Semchenkov⁶, J. Steiner², P. Thörle-Pospiech¹, T. Torres De Heidenreich², N. Trautmann¹, A. Türler¹¹, J. Uusitalo¹², D. Ward⁶, M. Wegrzecki¹³, P. Wieczorek², N. Wiehl¹, A. Yakushev², and V. Yakusheva³

<sup>1</sup>U. Mainz, Germany; <sup>2</sup>GSI, Darmstadt, Germany; <sup>3</sup>HIM, Mainz, Germany; <sup>4</sup>U. Liverpool, UK; <sup>5</sup>LBNL+UC Berkeley, CA, USA; <sup>6</sup>Lund U., Sweden; <sup>7</sup>SINP, Kolkata, India; <sup>8</sup>U. Oslo, Norway; <sup>9</sup>UNAL Bogotá, Colombia; <sup>10</sup>JAEA Tokai, Japan; <sup>11</sup>U. Bern+PSI Villigen, Switzerland; <sup>12</sup>U. Jyväskylä, Finland; <sup>13</sup>ITE, Warsaw, Poland

During the last decades heavy ion induced reactions were applied to explore the chart of nuclei up to the superheavy elements (SHE), which has resulted in the discovery of the SHE up to Og (Z=118) in complete fusion reactions [1]. Alternatively, recent model calculations suggest the possibility to produce exotic nuclei including SHE in non-fusion channels of heavy ion induced reactions [2]. Pioneering studies on the possible production of such exotic nuclei in non-fusion reactions were performed in the late 1970s by applying chemical separation techniques [3], which are suitable for longer-lived nuclei ( $\gtrsim 1\,\mathrm{h}$ ). Many properties of the multi-nucleon transfer reactions have been established, but still detailed information on the reaction mechanism/kinematics is missing [5, 6, 4].




Figure 1: Energy spectrum measured with the focal plane detector during beam-off periods for the  $^{50}$ Ti +  $^{249}$ Cf reaction. Isotopic identification of some lines is given.

Recently, at the velocity filter SHIP new short-lived (down to  $10^{-6}$  h) neutron-deficient nuclei of heavy ele-

ments have been synthesized in the <sup>48</sup>Ca + <sup>248</sup>Cm reaction [7]. These and other results at SHIP (see [8]), benefiting of the kinematic separation, demonstrate the relevance of the forward angle measurements for the investigation of the reaction dynamics.

At the gas-filled recoil separator TASCA, non-fusion products of the  $^{50}{\rm Ti}$   $+^{249}$  Cf reaction have been investigated. They were produced during the experiment for searching the SHE with Z=120 [9]. The magnetic settings of TASCA were tuned to collect the products of fusion-evaporation reactions, but even under these conditions some amount of non-fusion products were passing through TASCA and were implanted into the focal plane detector. Here their subsequent radioactive decays were measured. A typical energy spectrum containing lines from the  $\alpha$  decay of the implanted nuclei is shown in Figure 1. By exploiting the  $\alpha$  decay properties, the identification of nuclei was performed employing a position and time correlation analyses between implantation and/or  $\alpha$ -like events. In total, 57 isotopes with Z=83-90 were identified.

Experimental details and the final analysis will be given in a forthcoming publication [10].

## References

- Y. T. Oganessian and V. K. Utyonkov, Rep. Prog. Phys. 78, 036301 (2015).
- [2] V. I. Zagrebaev and W. Greiner PRC 83, 044618 (2011).
- [3] M. Schädel et al., PRL 41, 469 (1978), and 48, 852 (1982).

203

- [4] L. Corradi et al., Jour. of Phys. G 36, 113101 (2009).
- [5] J.V. Kratz et al., NPA **944**, 117 (2015).
- [6] M. Götz et al., NPA 961, 1 (2017).
- [7] S. Heinz et al., EPJA **52**, 278 (2016).
- [8] S. Heinz et al., EPJA **51**, 140 (2015).
- [9] Ch. E. Düllmann et al. to be published.
- [10] A. Di Nitto et al., to be published.



 $<sup>^{\</sup>ast}$  We thank the ion-source and accelerator staff at GSI. Work supported by BMBF contract-No. 06MZ7164.

<sup>†</sup> a.dinitto@gsi.de

<sup>&</sup>lt;sup>‡</sup> Present address: GANIL, Caen, France