



LUND UNIVERSITY

Regulation of Insulin Secretion in Relation to Nitric Oxide, Carbon Monoxide and Acid alpha-Glucoside Hydrolase Activities

Mosén, Henrik

2005

[Link to publication](#)

Citation for published version (APA):

Mosén, H. (2005). *Regulation of Insulin Secretion in Relation to Nitric Oxide, Carbon Monoxide and Acid alpha-Glucoside Hydrolase Activities*. [Doctoral Thesis (compilation), Clinical Physiology (Lund)]. Department of Experimental Medical Science, Lund University.

Total number of authors:

1

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

Insulin release transduction mechanism through acid glucan 1,4- α -glucosidase activation is Ca^{2+} regulated

ALBERT SALEHI, HENRIK MOSÉN, AND INGMAR LUNDQUIST

Department of Pharmacology, University of Lund, S-223 62 Lund, Sweden

Salehi, Albert, Henrik Mosén, and Ingmar Lundquist.

Insulin release transduction mechanism through acid glucan 1,4- α -glucosidase activation is Ca^{2+} regulated. *Am. J. Physiol.* 274 (*Endocrinol. Metab.* 37): E459–E468, 1998.—An important signal involved in glucose-stimulated insulin secretion is transduced through the action of a lysosomal acid, glucan 1,4- α -glucosidase. We investigated the Ca^{2+} dependency of this enzyme activity in relation to insulin release. In isolated islets, increased levels of extracellular Ca^{2+} induced a large increase in acid glucan 1,4- α -glucosidase activity accompanied by a similar increase in insulin release at both substimulatory and stimulatory concentrations of glucose. At low glucose the Ca^{2+} “inflow” blocker nifedipine unexpectedly stimulated enzyme activity without affecting insulin release. However, nifedipine suppressed $^{45}\text{Ca}^{2+}$ outflow from perfused islets at low glucose and at Ca^{2+} deficiency when intracellular Ca^{2+} was mobilized by carbachol. This nifedipine-induced retention of Ca^{2+} was reflected in increased acid glucan 1,4- α -glucosidase activity. Adding different physiological Ca^{2+} concentrations or nifedipine to islet homogenates did not increase enzyme activity. Neither selective glucan 1,4- α -glucosidase inhibition nor the ensuing suppression of glucose-induced insulin release was overcome by a maximal Ca^{2+} concentration. Hence, Ca^{2+} -induced changes in acid glucan 1,4- α -glucosidase activity were intimately coupled to similar changes in Ca^{2+} -glucose-induced insulin release. Ca^{2+} did not affect the enzyme itself but presumably activated either glucan 1,4- α -glucosidase-containing organelles or closely interconnected messengers.

pancreatic islets; lysosomal enzymes; nifedipine; emiglitate; carbachol; calcium ion

IT IS A WELL-KNOWN FACT (8, 25, 34, 35) that Ca^{2+} plays an essential role in the stimulus-secretion coupling for insulin release. The cytosolic accumulation and intracellular distribution of Ca^{2+} are important for a growing number of different events in the secretory process, which is affected not only by the major secretagogue glucose but also by a variety of cholinergic, adrenergic, and peptidergic influences (2, 3, 8–10, 12, 22, 25, 33–35). Moreover, glucose-induced insulin release itself is a complex cascade of events, the details of which are far from elucidated (2, 25, 35). We have previously proposed (14–20, 28–32) that one of these glucose-induced multiple signals is transduced via the vacuolar system, involving the activation of a lysosome-acid glucan 1,4- α -glucosidase system. The acid glucan 1,4- α -glucosidase (EC 3.2.1.3) and the acid α -glucosidase (EC 3.2.1.20) are members of the α -glucosidehydrolase family. The acid glucan 1,4- α -glucosidase is known to preferentially attack α -1,4-linked polymers such as glycogen (24) and thereby to have the ability to produce high local concentrations of nonphosphorylated glucose within the vacuolar system. This glucose production, in turn, could act as a transducer (e.g., metabolic, osmotic,

or cybernetic) in the multifactorial process of insulin release. Glycogen is a normal constituent of islet tissue (11, 21), and it is known to display a surprisingly constant concentration at a wide range of blood glucose levels (21). Hence an important part of islet glycogen is probably not integrated in the metabolic pool of glucose phosphorylation processes in the cytoplasm but rather is restricted to a compartmentalized vacuolar pool of signal glycogen available to the acid glucan 1,4- α -glucosidase. It is worth noting that the phosphorolytic breakdown of glycogen *in vitro* from mouse islets is reportedly very slow (21).

In a recent report (28) we showed that the defective glucose-induced insulin release from isolated mouse islets in a Ca^{2+} -deficient medium was accompanied by markedly reduced activities of islet lysosomal α -glucosidehydrolases. In contrast, the activity of another lysosomal glycosidase, *N*-acetyl- β -D-glucosaminidase, was completely unaffected by Ca^{2+} deficiency. Likewise, islet activities of the lysosomal enzyme acid phosphatase and the neutral α -glucosidase (endoplasmic reticulum) were not influenced in a Ca^{2+} -deficient medium. A similar pattern of a greatly suppressed glucose-induced insulin release in parallel with a reduced acid α -glucosidehydrolase activity was accomplished by different selective α -glucosidehydrolase inhibitors such as miglitol, emiglitate, and acarbose (19, 20, 28–32).

Hence, because the activity of the lysosomal acid α -glucosidehydrolases (but not the neutral α -glucosidase) seemed to be one of several important intracellular key factors in glucose-induced insulin release, we found it mandatory to study the Ca^{2+} dependency of these enzyme activities in more detail. In the present investigation we tested 1) high concentrations of extracellular Ca^{2+} , 2) the Ca^{2+} channel blocker nifedipine and the intracellular Ca^{2+} mobilizer carbachol, as well as 3) the selective α -glucosidehydrolase inhibitor emiglitate (26) to further elucidate the role of Ca^{2+} in regulating islet acid α -glucosidehydrolase activities and insulin release.

MATERIALS AND METHODS

Animals

Adult female mice of the NMRI strain (B & K Universal, Sollentuna, Sweden), 3–4 mo old and weighing 25–30 g, fed *ad libitum*, and with free access to water, were used throughout the study. The experiments were approved by the Ethical Committee for Animal Research at the University of Lund.

Drugs and Chemicals

Collagenase (CLS 4) was obtained from Worthington Biochemicals (Freehold, NJ). Nifedipine, ethylene glycol-bis(β -

aminoethyl ether)-*N,N,N',N'*-tetraacetic acid (EGTA), and carbachol as well as methylumbelliferyl-coupled substrates were purchased from Sigma Chemical (St. Louis, MO). Emiglitate, *N*-[β -(4-ethoxycarbonylphenoxy) ethyl]-1-deoxyojirimycin (Bay 0 1248), was kindly supplied by Bayer (Leverkusen, Germany). Bovine serum albumin was from ICN Biomedicals (High Wycombe, UK). ⁴⁵CaCl₂ was from Radiochemical Centre (Amersham). All other chemicals were from British Drug Houses (Poole, UK) or Merck (Darmstadt, Germany). The radioimmunoassay kits for insulin determination were obtained from Novo Nordisk (Bagsvaerd, Denmark).

Experimental Procedure

Isolation of pancreatic islets from freely fed mice was accomplished by retrograde injection of a collagenase solution via the bile-pancreatic duct (5). The animals were killed from 8 to 9 AM by elongation of the neck and were immediately injected with collagenase.

Batch incubation of isolated islets. The freshly isolated islets were preincubated for 30 min at 37°C in Krebs-Ringer bicarbonate (KRB) buffer, pH 7.4, supplemented with 10 mmol/l *N*-2-hydroxyethylpiperazine-*N'*-2-ethanesulfonic acid (HEPES), 0.1% bovine serum albumin, and 1 mmol/l glucose. Each incubation vial contained 30 islets in 1.5 ml buffer solution and was gassed with 95% O₂-5% CO₂ to obtain constant pH and oxygenation. After preincubation, the buffer was changed to a medium containing 1 or 16.7 mmol/l glucose \pm test substances, and the islets were incubated for 120 min. Emiglitate, when used, was included also during the preincubation period (20). All incubations were performed at 37°C in an incubation box (30 cycles/min). It should be noted that, in the experiments with high concentrations of Ca²⁺, phosphate and sulfate in the KRB-HEPES buffer were replaced with equimolar amounts of chloride (9). The change in osmotic pressure with high Ca²⁺ was compensated for by reduction of NaCl, as described by Hellman (9). Immediately after incubation, an aliquot of the medium was removed for assay of insulin. Insulin was determined radioimmunologically (7).

Lysosomal enzyme activities. If not otherwise stated, the islets were then thoroughly washed in glucose-free KRB buffer and collected and stored in 200 μ l acetate-EDTA buffer (1.1 mmol/l EDTA and 5 mmol/l acetate, pH 5.0) at -20°C (14). Ancillary experiments showed that the collected islets could be kept frozen for several months without loss of enzyme activity. After sonication, islet homogenates were analyzed for lysosomal enzyme activities. In experiments in which we studied the direct influence of different Ca²⁺ concentrations added to islet homogenates on the lysosomal enzyme activities, the islets were washed in a glucose- and Ca²⁺-free KRB buffer and collected and stored in 5 mmol/l acetate in the absence of EDTA. The procedures for determination of acid phosphatase (pH 4.5), acid α -glucosidase (pH 4.0 and 5.0), *N*-acetyl- β -D-glucosaminidase (pH 5.0), and neutral α -glucosidase (pH 6.5) with methylumbelliferyl-coupled substrates have previously been described (16). Islet glucan 1,4- α -glucosidase activity with glycogen as substrate was determined at pH 4.0, as described in detail elsewhere (14, 19). The acid α -glucosidase activity was assayed at both pH 4.0 and pH 5.0, because previous studies (23) have shown that inhibition of α -glucosidase by the lysosomotropic drug suramin was dependent on the prevailing pH value. Furthermore, it should be recalled that lysosomal enzyme activities are subjected to circadian and seasonal variations (see Ref. 6). Therefore, all experiments were always performed with both control groups and experimental groups at each occasion. Protein was analyzed according to the method of Lowry et al. (13).

Islet perfusion experiments. In the perfusion experiments, islets (150–200) were first incubated for 90 min in 800 μ l of KRB medium containing 20 mmol/l glucose and 50 μ l ⁴⁵CaCl₂ (50–100 μ Ci), which was added from a stock solution with a specific activity of 10–40 mCi/mg Ca²⁺. The islets were then washed three times with nonradioactive medium, divided into two or three groups with 75–100 islets per group, and transferred to perfusion columns. The islets were thereby sandwiched between two layers of gel (Bio-gel P-4, 200–400 mesh; Bio-Rad Laboratory, Richmond, CA) and perfused at a rate of 0.1 ml/min with the KRB buffer supplemented with 1 mmol/l glucose. Test substances were introduced according to the protocols. A Ca²⁺-deficient medium was obtained by omitting calcium chloride and adding 0.5 mmol/l EGTA. The radioactivity lost by the islets was measured in effluent fractions collected every 2 min (50 μ l of the sample were added to 5 ml of scintillation fluid) and counted in a scintillation counter (Packard Instrument, Downers Grove, IL). The fractional efflux rate was calculated for each period (radioactivity lost by islets during the time interval/radioactivity present in the islets during the same time interval), and the mean value calculated for *minutes* 40 and 42 was then normalized to 100%. Insulin was determined with a radioimmunoassay (7).

Statistics

Probability levels of random differences were determined by Student's unpaired *t*-test or analysis of variance followed by Tukey-Kramer's multiple comparisons test where applicable.

RESULTS

Effect of a Maximal Concentration of Extracellular Ca²⁺ on Basal Insulin Release and Islet Lysosomal Enzyme Activities

Taking advantage of a previous dose-response study by Hellman (9) showing that increasing extracellular Ca²⁺ concentrations up to 30 mmol/l in the presence of a substimulatory glucose level could increase insulin release from isolated *ob/ob* islets, we performed a series of experiments at 1 mmol/l glucose with either a normal Ca²⁺ (2.5 mmol/l) or a high maximal (9) concentration of Ca²⁺ (30 mmol/l) in the extracellular medium. Figure 1 shows that increasing the Ca²⁺ concentration from 2.5 to 30 mmol/l induced an almost threefold increase in basal insulin release. This enhanced insulin secretion was accompanied by a marked increase in islet lysosomal α -glucosidase activities, i.e., acid glucan 1,4- α -glucosidase (3-fold increase), acid α -glucosidase pH 4.0 (+40%), and pH 5.0 (+95%), whereas other lysosomal enzyme activities such as acid phosphatase and *N*-acetyl- β -D-glucosaminidase were totally unaffected. In contrast, the activity of the neutral α -glucosidase, an enzyme attributed to the endoplasmic reticulum, was modestly reduced (-30%) by 30 mmol/l Ca²⁺ in the incubation medium.

Effects of Direct Addition of Different Concentrations of Ca²⁺ on Lysosomal Enzyme Activities in Islet Homogenates

Because a high concentration of extracellular Ca²⁺ greatly increased the acid α -glucosidase activity




Fig. 1. Effect of Ca²⁺ on islet activities of different lysosomal enzymes and neutral α -glucosidase, as well as insulin secretion, at a low substimulatory concentration of glucose (1 mmol/l): acid α -glucosidase, pH 4.0 (A); acid α -glucosidase, pH 5.0 (B); acid glucan 1,4- α -glucosidase (C); acid phosphatase (D); *N*-acetyl- β -D-glucosaminidase (E); neutral α -glucosidase (F); insulin release (G). Islets were incubated for 2 h at 2.5 mmol/l Ca²⁺ (open bars) or at 30 mmol/l Ca²⁺ (solid bars). Enzyme activities are expressed as μ mol glucose (acid glucan 1,4- α -glucosidase) or 4-methylumbelliferone liberated per g protein per min. Insulin secretion is expressed as ng insulin released per islet per 2 h. Values are means \pm SE for 9–10 batches of islets in each group obtained from 3 independent experiments. * P < 0.05, ** P < 0.01, *** P < 0.001.

ties in intact islets (Fig. 1), we studied whether addition of Ca²⁺ to islet homogenates could influence these enzyme activities directly. The effects of direct addition to islet homogenates of a wide range of Ca²⁺ concentrations (0.05 μ mol/l–30 mmol/l) on the different enzyme activities are illustrated in the absence (Fig. 2, A and B) and presence (Fig. 2, C and D) of calmodulin. No appreciable influence of Ca²⁺ was seen within known intracellular fluctuations of the cation. However, there was a large decrease in acid α -glucosidase activities (about –80%) and a marked increase in acid glucan 1,4- α -glucosidase activity (about +50%) at very high “unphysiological” intracellular concentrations of Ca²⁺ (2, 10, and 30 mmol/l).

Influence of the Ca²⁺ Channel Blocker Nifedipine on Islet Lysosomal Enzyme Activities and Insulin Release at Low and High Glucose Concentrations

To further investigate, in intact islets, the effect of Ca²⁺ perturbations on insulin release in relation to the activities of the acid α -glucosidase, we studied the influence of nifedipine on basal and glucose-induced insulin secretion and islet lysosomal enzyme activities. Figure 3A shows the effect of nifedipine on insulin secretion from incubated islets at low or high glucose. As expected, glucose-induced insulin secretion was greatly suppressed in the presence of nifedipine. Moreover, we found, unexpectedly, that the islet lysosomal acid α -glucosidase activities were significantly increased by nifedipine at basal glucose (Fig. 4), i.e., acid glucan 1,4- α -glucosidase (+35%) and acid

α -glucosidase pH 4.0 (+45%) and pH 5.0 (+40%). However, nifedipine had no effects on these enzymes in the presence of high glucose (16.7 mmol/l), a glucose concentration which by itself markedly enhanced the acid α -glucosidase activities (Fig. 4).

Effects of Direct Addition of Nifedipine on Lysosomal Enzyme Activities in Islet Homogenates

To elucidate whether nifedipine could directly activate the islet α -glucosidase in islet homogenates, a dose-response study was performed. Table 1 shows the effect of nifedipine on the activities of the different lysosomal enzymes and the neutral α -glucosidase in islet homogenates. Nifedipine at 1 μ mol/l induced a marked increase in *N*-acetyl- β -D-glucosaminidase activity. Furthermore, nifedipine at 30 μ mol/l induced a modest suppression of the activities of the acid α -glucosidase (about –10 to –15%) and a pronounced decrease of *N*-acetyl- β -D-glucosaminidase activity (about –55%). Acid phosphatase activity was not influenced, whereas the activity of the neutral α -glucosidase was moderately reduced at 10 and 30 μ mol/l nifedipine.

Effect of Nifedipine on ⁴⁵Ca²⁺ Efflux from Isolated Islets Either at Low (1 mmol/l) Glucose and Normal Ca²⁺ or in a Ca²⁺-Deficient Medium in Which the Islets Were Stimulated by the Intracellular Ca²⁺ Mobilizer Carbachol at Low Glucose

Because high extracellular Ca²⁺ (Fig. 1) as well as nifedipine (Fig. 4) could increase the acid α -glucosidase





Fig. 2. Influence of different concentrations of Ca²⁺ added directly to enzyme assay mixtures containing aliquots of Ca²⁺-free islet homogenates in the absence (A and B) or presence (C and D) of 25 U/ml calmodulin. Enzyme activities are expressed as percentage of control incubations. Values are means \pm SE for 5–9 observations in each group obtained from 3 independent experiments.

drolase activities in intact islets, we searched for an effect of nifedipine on intracellular Ca²⁺ that was independent of its Ca²⁺ channel-blocking effects. Therefore, to study whether nifedipine could influence the efflux of Ca²⁺ from the β -cell, we performed a series of perfusion experiments with ⁴⁵Ca²⁺-loaded islets. The first experiment was conducted at substimulatory (1 mmol/l) glucose and normal Ca²⁺ (Fig. 5). It is seen that nifedipine induced a marked decrease of ⁴⁵Ca²⁺ efflux at this low glucose concentration, which indeed does not open the nifedipine-sensitive Ca²⁺ channels. The basal insulin release was not influenced by nifedipine (see also Fig. 3A). Because the cholinergic muscarinic receptor agonist carbachol is known to mobilize intracellularly stored Ca²⁺ (12), we performed another experiment at 1 mmol/l glucose with carbachol as a stimulus in a Ca²⁺-deficient perfusion medium supplemented with 0.5 mmol/l EGTA, to preclude any influx of Ca²⁺ into the β -cells. Figure 6 shows the effect of nifedipine on ⁴⁵Ca²⁺ efflux and insulin release during stimulation with carbachol (50 μ mol/l). In the absence of nifedipine, carbachol induced a clear biphasic increase of ⁴⁵Ca²⁺ efflux. Addition of nifedipine converted the initial carbachol-induced increase of ⁴⁵Ca²⁺ into a marked but transient decrease followed by a modest increase, which, however, was strongly suppressed compared with the carbachol controls. Thus nifedipine induced a powerful inhibition of carbachol-stimulated ⁴⁵Ca²⁺ efflux in a Ca²⁺-deficient medium. As expected, carbachol did not

notably influence insulin release from either controls or nifedipine-treated islets in the absence of extracellular Ca²⁺ (Fig. 6, *bottom*).

Effect of Nifedipine on Islet Lysosomal Enzyme Activities and Insulin Release in a Ca²⁺-Deficient Medium in the Presence and Absence of the Intracellular Ca²⁺ Mobilizer Carbachol

The next experiment was performed to test whether the carbachol-mobilized Ca²⁺, which was retained within the β -cell by the action of nifedipine (see Fig. 6), might be redistributed to affect the acid α -glucosidase activity. To avoid any influence of extracellular Ca²⁺, the experiment was performed in a Ca²⁺-deficient medium. We therefore incubated isolated islets for 60 min in the absence and presence of nifedipine and carbachol after a preincubation period of 40 min, i.e., largely mimicking the perfusion experiments. Figure 7 shows that nifedipine induced a large increase (almost 2-fold) in islet lysosomal acid α -glucosidase activities at basal glucose (1 mmol/l) and Ca²⁺ deficiency, i.e., a much higher increase than in the presence of a normal concentration of extracellular Ca²⁺ (see Fig. 4). The activities of other lysosomal enzymes and the neutral α -glucosidase were not influenced by nifedipine (Fig. 7). Carbachol had no notable influence on the different enzyme activities in the basal state and did not induce a greater increase in enzyme activities



together with nifedipine than that brought about by nifedipine itself, except for inducing a modest but significant increase in the acid glucan 1,4- α -glucosidase activity (+22%) in the presence of the Ca²⁺ channel blocker (Fig. 7). Hence, nifedipine stimulated acid glucan 1,4- α -glucosidase activity in both the absence and the presence of carbachol. As expected, no effect of either carbachol or nifedipine on insulin release in the Ca²⁺-deficient medium was observed (Fig. 3B).

Effect of a Maximal Concentration of Extracellular Ca²⁺ on Glucose-Stimulated Insulin Release and Islet Lysosomal Enzyme Activities in the Absence and Presence of the Selective α -Glucosidase Inhibitor Emiglitate

In a further attempt to elucidate whether the Ca²⁺ dependency of islet acid α -glucosidase and insulin release in intact islets was intimately coupled in the secretory process, we performed a series of experiments with the selective α -glucosidase inhibitor emiglitate (20, 26, 29, 31) in the presence of high glucose and using either a normal (2.5 mmol/l) or a maximal concentration of Ca²⁺ (10 mmol/l). At this high glucose concentration 10 mmol/l Ca²⁺ is maximal, because higher Ca²⁺ concentrations are even inhibitory to insulin release (8). Figures 3C and 8 show that raising the extracellular Ca²⁺ from 2.5 to 10 mmol/l at 16.7 mmol/l glucose (open bars) did increase both insulin release (+55%; Fig. 3C) and acid glucan 1,4- α -glucosidase activity (+50%) as well as acid α -glucosidase pH 4.0 (+55%) and pH 5.0 (+40%; Fig. 8). The activities of acid phosphatase and *N*-acetyl- β -D-glucosaminidase, however, were significantly suppressed by high Ca²⁺ (-25%; Fig. 8). Moreover, addition of emiglitate almost totally suppressed both glucose-induced and Ca²⁺-induced increase of acid α -glucosidase activities (Fig. 8) as well as the release of insulin (Fig. 3C).

DISCUSSION

The present results lend further support to our previous hypothesis (14–20, 28–32) that one of the multiple signals involved in glucose-induced insulin release is transduced through the activation of a glyco-gen-hydrolyzing acid, glucan 1,4- α -glucosidase.

Fig. 3. *A*: effect of 30 μ mol/l nifedipine (solid bars) on basal (1 mmol/l glucose) and glucose-stimulated (16.7 mmol/l glucose) insulin secretion from isolated pancreatic islets at normal extracellular Ca²⁺ (2.5 mmol/l). Controls are shown by open bars. Values are means \pm SE for 6–9 batches of islets in each group obtained from 4 independent experiments. ****P* < 0.001 vs. controls. *B*: insulin release in a Ca²⁺-deficient medium at 1 mmol/l glucose in the presence and absence of 50 μ mol/l carbachol and 30 μ mol/l nifedipine (solid bars). Controls are shown by open bars. Values are means \pm SE for 8–10 batches of islets in each group obtained from 4 independent experiments. *C*: effect of Ca²⁺ (10 mmol/l) on insulin release at a high stimulatory concentration of glucose (16.7 mmol/l) in the absence (open bars) or presence (solid bars) of the selective α -glucosidase inhibitor emiglitate (1 mmol/l). Values are means \pm SE for 10–12 batches of islets in each group obtained from 4 independent experiments. ****P* < 0.001 vs. controls.




Fig. 4. Effect of 30 $\mu\text{mol/l}$ nifedipine (solid bars) on activities of different lysosomal enzymes and neutral α -glucosidase in islets exposed to 1 or 16.7 mmol/l glucose. Controls are shown by open bars. Enzyme activities are expressed as micromoles glucose (acid glucan 1,4- α -glucosidase) or 4-methylumbelliferone liberated per g protein per minute. Values are means \pm SE for 6–9 batches of islets in each group obtained from 4 independent experiments. * $P < 0.05$, *** $P < 0.001$ vs. controls.

Effects of Ca²⁺ Deficiency and High Extracellular Ca²⁺

In a recent study we reported (28) that the activities of the acid α -glucosidases were highly depressed in islets incubated in a Ca²⁺-deficient medium. In contrast, classical lysosomal enzyme activities, such as acid phosphatase and *N*-acetyl- β -D-glucosaminidase, as well as the neutral α -glucosidase (endoplasmic reticulum), were unaffected by the absence of extracellular Ca²⁺. In addition, we observed (28) that the presence of a high concentration of glucose in the incubate induced a significant increase of islet acid

α -glucosidase activities also in a Ca²⁺-deficient medium. The measured increase, however, never achieved the level of enzyme activity recorded in the presence of a normal physiological Ca²⁺ concentration (28). These observations suggested to us that both glucose and Ca²⁺ were needed for the full expression of islet acid α -glucosidase activity. We show here that a high concentration of Ca²⁺ itself, in the presence of a substimulatory concentration of glucose (1 mmol/l), greatly enhanced the islet acid α -glucosidase activities in isolated islets in parallel with an increased insulin release. These effects are most likely a result of increased intracellular Ca²⁺ concentration ([Ca²⁺]_i), because recent data suggest that high extracellular Ca²⁺ can induce a rise in [Ca²⁺]_i, which primarily is accounted for by Ca²⁺ influx through dihydropyridine- and voltage-insensitive, nonselective cation channels (33). Other lysosomal enzyme activities, such as acid phosphatase and *N*-acetyl- β -D-glucosaminidase, were unaffected by the high extracellular Ca²⁺ level. Moreover, a maximal (8) extracellular Ca²⁺ concentration at high glucose did further increase both insulin release and the acid α -glucosidase activities, supporting the idea that Ca²⁺ is a key regulatory factor for islet acid α -glucosidase activities.

Effects of Blockade of Ca²⁺ Influx by Nifedipine and of Intracellular Ca²⁺ Mobilization by Carbachol

Because high extracellular Ca²⁺ enhanced the acid α -glucosidase activity at low substimulatory glucose (Fig. 1, present study) and because high glucose could partially enhance the enzyme activity in the absence of extracellular Ca²⁺ (28), the question arose whether activation of the enzyme(s) was dependent on both the influx of extracellular Ca²⁺ and the redistribution/sequestration of intracellular Ca²⁺. It is well established that glucose-induced insulin release is accompanied by closure of ATP-sensitive K⁺ channels, followed by membrane depolarization (in normal mouse islets at glucose concentrations >7 mmol/l) (4), opening of the voltage-dependent Ca²⁺ channels, and influx of Ca²⁺. The subsequent rise in intracellular Ca²⁺ elicits multiple signals that induce the recruitment and extrusion of secretory granules (2, 25, 35). The present data show that nifedipine, a well-known blocker of voltage-dependent Ca²⁺ channels, unexpectedly did induce a marked increase in acid α -glucosidase activity in the presence of a very low nondepolarizing concentration of glucose (1 mmol/l). Hence, nifedipine might have changed the intracellular distribution of Ca²⁺ and/or inhibited Ca²⁺ outflow, because its classical Ca²⁺ channel-inhibiting effects could not be operating at 1 mmol/l glucose. Our ⁴⁵Ca²⁺ efflux experiments showed that such an assumption was justified. Nifedipine induced a marked suppression of the basal efflux of ⁴⁵Ca²⁺ in the presence of 1 mmol/l glucose and a normal extracellular Ca²⁺ concentration. This effect was in all probability not solely, if at all, a consequence of an inhibitory effect of nifedipine on the influx of Ca²⁺ through nonvoltage-dependent Ca²⁺ channels, because ⁴⁵Ca²⁺ efflux in a Ca²⁺-deficient medium with very low glucose (1 mmol/l)

Table 1. Influence of nifedipine on different lysosomal enzyme activities and neutral α -glucosidase in islet homogenates

	Control	Nifedipine, $\mu\text{mol/l}$				
		0.1	1	10	30	
Acid phosphatase	100 \pm 1.0	99 \pm 1.1	100 \pm 1.1	93 \pm 1.0	90 \pm 3.5	
N-acetyl- β -D-glucosaminidase	100 \pm 2.4	99 \pm 2.0	224 \pm 7.5 \ddagger	114 \pm 2.5*	44 \pm 8.6 \ddagger	
Acid glucan 1,4- α -glucosidase	100 \pm 2.1	105 \pm 3.1	99 \pm 2.2	95 \pm 1.5	84 \pm 6.2*	
Acid α -glucosidase						
pH 4.0	100 \pm 1.6	101 \pm 2.8	105 \pm 1.5	94 \pm 1.5	84 \pm 1.9 \ddagger	
pH 5.0	100 \pm 1.4	101 \pm 1.8	102 \pm 1.6	98 \pm 4.0	90 \pm 1.3 \ddagger	
Neutral α -glucosidase	100 \pm 2.7	94 \pm 1.0	97 \pm 2.3	76 \pm 4.1 \ddagger	84 \pm 6.0 \ddagger	

Aliquots of pancreatic islet homogenates were assayed with nifedipine in a concentration range from 0.1 $\mu\text{mol/l}$ to 30 $\mu\text{mol/l}$. Enzyme activities are expressed as percentage of control incubations. Values are means \pm SE for 4–9 observations. * $P < 0.05$; $\ddagger P < 0.01$; $\ddagger P < 0.001$.

was strongly inhibited by nifedipine also during stimulation by the intracellular Ca²⁺ mobilizer carbachol. Carbachol has previously been shown to be a very efficient mobilizer of intracellular Ca²⁺ in isolated islets (12). Hence, in addition to its blocking effect on voltage-dependent Ca²⁺ channels, nifedipine, at least under our experimental conditions, also inhibited the outflow of Ca²⁺ across the plasma membrane and/or caused a redistribution of intracellular Ca²⁺, leading to accumulation of Ca²⁺ in acid α -glucosidase-containing organelles in the vacuolar system. Such an assumption is in accordance with a very recent finding (1), showing that a high concentration (30 $\mu\text{mol/l}$) of the nifedipine

analog nifedipine did enhance the cytoplasmic Ca²⁺ concentration in mouse thymocytes in the absence of extracellular Ca²⁺. Indeed, a Ca²⁺ redistribution effect of nifedipine (or a nifedipine-induced messenger) rather than, or in addition to, an inhibition of ⁴⁵Ca²⁺ efflux across the plasma membrane, is suggested by the observation that the initial acute increase in ⁴⁵Ca²⁺ efflux in the biphasic response to carbachol stimulation was converted into a marked initial decrease (negative peak; see Fig. 6) by nifedipine followed by a modest and highly suppressed second phase. Such a pattern is in




Fig. 5. Effect of nifedipine on ⁴⁵Ca²⁺ efflux (top) and insulin release (bottom) from perfused islets at a low substimulatory concentration of glucose (1 mmol/l) and normal Ca²⁺ (2.5 mmol/l). Nifedipine (○, 30 $\mu\text{mol/l}$) or solvent (controls, ●) was introduced at minute 42, as shown by dotted vertical line. Fractional efflux rate was normalized, as described in MATERIALS AND METHODS. Values are means \pm SE for 6–7 perfusions in each group obtained from 3 independent experiments.




Fig. 6. Effect of carbachol on ⁴⁵Ca²⁺ efflux (top) and insulin release (bottom) in the presence (○) and absence (●) of nifedipine (30 $\mu\text{mol/l}$) at a low substimulatory concentration of glucose (1 mmol/l) in a Ca²⁺-deficient medium supplemented with 0.5 mmol/l EGTA. Carbachol (50 $\mu\text{mol/l}$) was introduced at minute 42, as indicated by dotted vertical line. Basal controls in the absence of nifedipine and carbachol are shown by dashed line. Fractional efflux rate was normalized, as described in MATERIALS AND METHODS. Values are means \pm SE for 5 perfusions in each group obtained from 4 independent experiments.




Fig. 7. Effect of 30 μ mol/l nifedipine (solid bars) on activities of different lysosomal enzymes and neutral α -glucosidase in islets incubated in the absence and presence of 50 μ mol/l carbachol at 1 mM glucose in a Ca²⁺-deficient medium. Controls are shown by open bars. Enzyme activities are expressed as micromoles glucose (acid glucan 1,4- α -glucosidase) or 4-methylumbelliferone liberated per g protein per minute. Values are means \pm SE for 8–10 batches of islets in each group obtained from 4 independent experiments. * P < 0.05, ** P < 0.01, *** P < 0.001.

fact much like the ⁴⁵Ca²⁺ efflux curve obtained by glucose stimulation (3) in the presence of extracellular Ca²⁺ (and in the absence of nifedipine).

The putative Ca²⁺ redistribution leading to activation of the acid α -glucosidase seemed in no way obligatory to the inflow of extracellular Ca²⁺ through the voltage-dependent Ca²⁺ channels, because enzyme activity could be induced by nifedipine in a Ca²⁺-deficient medium. From these experiments it is also obvious that the nifedipine-stimulated Ca²⁺ redistribu-

tion and the subsequent increase in acid glucan 1,4- α -glucosidase activity are not sufficient by themselves to induce an insulin secretory response, because nifedipine at the same time blocks the voltage-dependent Ca²⁺ channels. Furthermore, it seems very unlikely that Ca²⁺ itself is directly modulating the enzyme(s), because physiological [Ca²⁺]_i (10, 25, 27, 33) in both the absence and presence of calmodulin did not display any notable effect on the enzyme activities after addition to islet homogenates. It should be noted, however, that very high concentrations of Ca²⁺ (2 mmol/l–30 mmol/l) did activate considerably the acid glucan 1,4- α -glucosi-




Fig. 8. Effect of Ca²⁺ on islet activities of different lysosomal enzymes and neutral α -glucosidase at a high stimulatory concentration of glucose (G; 16.7 mmol/l) in the absence (open bars) or presence (solid bars) of the selective α -glucosidase inhibitor emiglitate (1 mmol/l). Islets were incubated for 2 h at either 2.5 or 10 mmol/l Ca²⁺. Enzyme activities are expressed as μ mol glucose (acid glucan 1,4- α -glucosidase) or 4-methylumbelliferone liberated per g protein per min. Values are means \pm SE for 10–12 batches of islets in each group obtained from 4 independent experiments. * P < 0.05; ** P < 0.01; *** P < 0.001.

dase but inhibited the acid α -glucosidases (Fig. 2). Such high Ca²⁺ concentrations are not likely to occur intracellularly, although theoretically they cannot be completely ruled out in certain Ca²⁺-rich subcellular compartments and organelles. Furthermore, this highly differential action of 2 and 10 mmol/l Ca²⁺ on the acid glucan 1,4- α -glucosidase, compared with the acid α -glucosidases in islet homogenates, was indeed not reflected in the various experiments with isolated intact islets, where these activities were always increased by the supraphysiological extracellular Ca²⁺ concentrations used in our studies (see Figs. 1 and 8).

With regard to glucose-stimulated insulin release, it is conceivable that the initial glucose-induced decrease in ⁴⁵Ca²⁺ efflux (3), which was recently shown to be the result of the ability of glucose to induce sequestration of cytoplasmic Ca²⁺ in a slowly exchangeable "organelle pool" (10), may be a key event in this context. Interestingly, the glucose-induced redistribution and sequestration of intracellular Ca²⁺ are manifested earlier and at lower glucose concentrations than those required to open the voltage-dependent Ca²⁺ channels (10). It is not inconceivable that one of these sequestration targets is the acid glucan 1,4- α -glucosidase-containing organelles. Such an assumption is in accordance with previous data showing, in a Ca²⁺-deficient medium, that an increase in glucose concentration from 1 to 4 mmol/l also increased the acid α -glucosidase activities (28). Hence, in addition to other factors, glucose-stimulated insulin release is apparently dependent on both a redistribution and a sequestration of intracellular Ca²⁺, which in turn activate the lysosomal-acid glucan 1,4- α -glucosidase system as well as the inflow of extracellular Ca²⁺ through voltage-dependent Ca²⁺ channels at depolarizing glucose concentrations. This redistribution/sequestration hypothesis also conforms with recent observations (29) showing that the acid α -glucosidase activities were profoundly suppressed in islets incubated in a Ca²⁺-deficient medium in the presence of the phosphodiesterase inhibitor 3-isobutyl-1-methylxanthine (IBMX), which is known to induce perturbations of intracellular organelle-bound Ca²⁺ in the β -cell (2).

Effects of the Selective α -Glucosidase Inhibitor Emiglitate

Finally, our experiments with emiglitate strongly suggest that the activating effect of Ca²⁺ on the lysosome-acid glucan 1,4- α -glucosidase system and glucose-induced insulin release are closely interconnected. Emiglitate almost totally suppressed insulin release stimulated both by high glucose alone and by high extracellular Ca²⁺ in the presence of high glucose. This powerful inhibition of insulin release was accompanied by greatly suppressed activities of the islet acid α -glucosidase. Hence, it appears that a most important Ca²⁺ effect in the stimulus-secretion coupling of glucose-stimulated insulin release is exerted closely proximal to the action of the acid glucan 1,4- α -glucosidase, the inhibition of which greatly impairs the signal transduction. This inhibition of enzyme activity

and subsequent insulin release apparently cannot be overcome by greatly increasing the Ca²⁺ concentration (see Fig. 8). Thus it is not inconceivable that this particular effect of Ca²⁺ is exerted on certain membrane components of acidic organelles and/or key factor(s) assisting the acid α -glucosidase in their *in vivo* catalytic function. It should be noted that emiglitate is reportedly (26) a selective α -glucosidase inhibitor. Hence, our results suggest a direct cause-effect relationship between islet acid glucan 1,4- α -glucosidase activity on the one hand and glucose-Ca²⁺-induced insulin release on the other. The present data are thus in accordance with previous observations in our laboratory showing that nutrient-induced insulin release is greatly suppressed by different selective α -glucosidase inhibitors, such as the pseudotetraraccharide acarbose or the deoxynojirimycin derivatives miglitol and emiglitate (20, 28–32), whereas Ca²⁺-independent insulin secretion induced by IBMX is not (29). Moreover, receptor-activated insulin release induced by carbachol is unaffected by selective α -glucosidase inhibition (30, 32). These data also conform with the present results showing that carbachol itself had no influence on islet acid glucan 1,4- α -glucosidase activity in a Ca²⁺-deficient medium (see Fig. 7). In contrast, glucose has previously been shown to greatly enhance the enzyme activity during Ca²⁺ deficiency (28).

In summary, in intact islets, high supraphysiological concentrations of extracellular Ca²⁺ brought about a marked enhancement of the islet acid α -glucosidase activities, accompanied by a large insulin release. The Ca²⁺ channel blocker nifedipine unexpectedly brought about an increase in acid α -glucosidase activity at low glucose. This increase was explained by showing that nifedipine suppressed ⁴⁵Ca²⁺ outflow from perfused islets at substimulatory glucose and normal Ca²⁺, as well as after intracellular mobilization of ⁴⁵Ca²⁺ by carbachol in a Ca²⁺-deficient medium. The inhibition of ⁴⁵Ca²⁺ efflux was probably accomplished through increased intracellular sequestration and impaired outflow of Ca²⁺ across the plasma membrane. The Ca²⁺-induced effects were shown not to be exerted by a direct action of either nifedipine or Ca²⁺ on the acid α -glucosidases. Instead we suggest that this signal function of Ca²⁺ is exerted on a step closely proximal to enzyme activation, e.g., on certain membrane constituents of acidic organelles and/or key factor(s) modulating the acid α -glucosidase in their *in vivo* catalytic function. This was further emphasized by the finding that selective inhibition of the acid α -glucosidases by emiglitate almost abolished glucose-induced insulin release, an effect which could not be overcome by increased Ca²⁺. Taken together with data on islet acid α -glucosidase activities obtained from previous experiments in Ca²⁺-deficient media (28), a redistribution of Ca²⁺ induced by glucose (or by pharmacological agents such as nifedipine) that is directed to acid α -glucosidase-containing organelles appears an attractive mechanism in this context. The intimate details of Ca²⁺ redistribution, seques-

tration, and induction of acid glucan 1,4- α -glucosidase activity in nutrient-induced insulin release will await further investigations.

The skillful assistance of Elsy Ling and Britt-Marie Nilsson and the secretarial help of Eva Björkbo and Björn Otterlin are gratefully acknowledged.

This study was supported by the Swedish Medical Research Council (14X-4286), the Crafoord Foundation, the Swedish Diabetes Association, the Albert Pålsson Foundation, and the Åke Wiberg Foundation.

Address for reprint requests: A. Salehi, Dept. of Pharmacology, Sölvegatan 10, S-223 62 Lund, Sweden.

Received 12 June 1997; accepted in final form 24 November 1997.

REFERENCES

- Chen, W. R., F. Liu, Y. J. Feng, and J. T. Zhang. Nicardipine enhances cytoplasmic calcium concentration in mouse thymocytes in vitro. *Acta Pharmacol. Sinica* 16: 489–492, 1995.
- Draznin, B., J. P. Steinberg, M. Goodman, J. W. Leitner, and K. E. Sussman. Control of secretion vesicle margination and lysis by glucose, IBMX, and glyburide. *Am. J. Physiol.* 248 (*Endocrinol. Metab.* 11): E375–E380, 1985.
- Frankel, B. J., W. T. Imagawa, M. D. L. O'Connor, I. Lundquist, J. A. Kromhout, R. E. Fanska, and G. M. Grodsky. Glucose-stimulated ⁴⁵calcium efflux from isolated rat pancreatic islets. *J. Clin. Invest.* 62: 525–531, 1978.
- Gao, Z.-Y., M. Gerard, and J.-C. Henquin. Glucose- and concentration-dependence of vasopressin-induced hormone release by mouse pancreatic islets. *Regul. Pept.* 38: 89–98, 1992.
- Gotoh, M., T. Maki, T. Kiyozumi, S. Satomi, and A. P. Monaco. An improved method for isolation of mouse pancreatic islets. *Transplantation* 40: 437–438, 1985.
- Groh, V., and H. von Mayersbach. Enzymatic and functional heterogeneity of lysosomes. *Cell Tissue Res.* 214: 613–621, 1981.
- Heding, L. G. A simplified insulin radioimmunoassay method. In: *Labelled Proteins in Tracer Studies*, edited by L. Donato, G. Milhaud, and J. Sirchis. Brussels: Euratom, 1966, p. 345–350.
- Hellman, B. The significance of calcium for glucose stimulation of insulin release. *Endocrinology* 97: 392–398, 1975.
- Hellman, B. Stimulation of insulin release after raising extracellular calcium. *FEBS Lett.* 63: 125–128, 1976.
- Hellman, B., E. Gylfe, P. Bergsten, E. Grapengiesser, P. E. Lund, A. Berts, S. Dryselius, A. Tengholm, Y. J. Liu, M. Eberhardson, and R. H. Chow. The role of Ca²⁺ in the release of pancreatic islet hormones. *Diabetes Metab.* 20: 123–131, 1994.
- Hellman, B., and L.-Å. Idahl. On the functional significance of the pancreatic β -cell glycogen. In: *The Structure and Metabolism of the Pancreatic Islets*, edited by S. Falkmer, B. Hellman, and I.-B. Täljedal. Oxford, UK: Pergamon, 1970, p. 253–262.
- Henquin, J.-C., M. C. Garcia, G. M. Bozem, M. P. Hermans, and M. Nenquin. Muscarinic control of pancreatic B cell function involves sodium-dependent depolarization and calcium influx. *Endocrinology* 122: 2134–2142, 1988.
- Lowry, O. H., N. J. Rosebrough, A. L. Farr, and R. J. Randall. Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* 193: 265–275, 1951.
- Lundquist, I. Method for determination of acid amyloglucosidase in isolated islets of the pancreas. *Enzyme* 12: 647–657, 1971.
- Lundquist, I. Carbohydrate content and regulation following injection of different glycogenolytic enzymes. *Enzyme* 20: 234–247, 1975.
- Lundquist, I. Lysosomal enzyme activities in pancreatic islets from normal and obese hyperglycemic mice. *Metab. Clin. Exp.* 34: 1–9, 1985.
- Lundquist, I. Islet amyloglucosidase activity: some characteristics, and its relation to insulin secretion stimulated by various secretagogues. *Diabetes Res.* 3: 31–41, 1986.
- Lundquist, I., and R. Lövdahl. Effect of fasting on islet lysosomal enzyme activities and the in vivo insulin response to different secretagogues. *Horm. Metab. Res.* 15: 11–14, 1983.
- Lundquist, I., and G. Panagiotidis. The relationship of islet amyloglucosidase activity and glucose-induced insulin secretion. *Pancreas* 7: 352–357, 1992.
- Lundquist, I., G. Panagiotidis, and A. Salehi. Islet acid glucan-1,4- α -glucosidase: a putative key enzyme in nutrient-stimulated insulin secretion. *Endocrinology* 137: 1219–1225, 1996.
- Matschinsky, F. M., and J. E. Ellerman. Metabolism of glucose in the islets of Langerhans. *J. Biol. Chem.* 243: 2730–2736, 1967.
- McDermott, A. M., and G. W. G. Sharp. Inhibition of insulin secretion: a fail-safe system. *Cell. Signalling* 5: 229–234, 1993.
- Panagiotidis, G., A. Salehi, and I. Lundquist. Effect of the lysosomotropic drug suramin on islet lysosomal enzyme activities and insulin-secretory response induced by various secretagogues. *Pharmacology* 43: 163–168, 1991.
- Pazur, J. H., and K. Kleppe. The hydrolysis of α -D-glucosides by amyloglucosidase from *Aspergillus niger*. *J. Biol. Chem.* 237: 1002–1006, 1962.
- Prentki, M., and F. M. Matschinsky. Ca²⁺, cAMP, and phospholipid-derived messengers in coupling mechanisms of insulin secretion. *Physiol. Rev.* 67: 1185–1248, 1987.
- Puls, W., H. P. Krause, L. Muller, H. Schutt, R. Sill, and G. Thomas. Inhibitors of the rate of carbohydrate and lipid absorption by the intestine. *Int. J. Obesity* 8, Suppl. 1: 181–190, 1984.
- Rorsman, P., K. Bokvist, C. Åmmälä, L. Eliasson, E. Renström, and J. Gäbel. Ion channels, electrical activity and insulin secretion. *Diabetes Metab.* 20: 138–145, 1994.
- Salehi, A., and I. Lundquist. Ca²⁺ deficiency, selective α -glucosidase inhibition, and insulin secretion. *Am. J. Physiol.* 265 (*Endocrinol. Metab.* 28): E1–E9, 1993.
- Salehi, A., and I. Lundquist. Islet glucan-1,4- α -glucosidase: differential influence on insulin secretion induced by glucose and isobutylmethylxanthine in mice. *J. Endocrinol.* 138: 391–400, 1993.
- Salehi, A., and I. Lundquist. Changes in islet glucan-1,4- α -glucosidase activity modulate sulphonylurea-induced but not cholinergic insulin secretion. *Eur. J. Pharmacol.* 243: 185–191, 1993.
- Salehi, A., and I. Lundquist. Modulation of islet G-proteins, α -glucosidase inhibition and insulin release stimulated by various secretagogues. *Biosci. Rep.* 16: 23–34, 1996.
- Salehi, A., G. Panagiotidis, L. A. H. Borg, and I. Lundquist. The pseudotetrasaccharide acarbose inhibits pancreatic islet glucan-1,4- α -glucosidase activity in parallel with a suppressive action on glucose-induced insulin release. *Diabetes* 44: 830–836, 1995.
- Silva, A. M., L. M. Rosario, and M. Santos. Background Ca²⁺ influx mediated by a dihydropyridine- and voltage-insensitive channel in pancreatic β -cells. *J. Biol. Chem.* 269: 17095–17103, 1994.
- Wolf, B. A., J. R. Colca, J. Turk, J. Florholmen, and M. L. McDaniel. Regulation of Ca²⁺ homeostasis by islet endoplasmic reticulum and its role in insulin secretion. *Am. J. Physiol.* 254 (*Endocrinol. Metab.* 17): E121–E136, 1988.
- Zawalich, W. S., and H. Rasmussen. Control of insulin secretion: a model involving Ca²⁺, cAMP and diacylglycerol. *Mol. Cell. Endocrinol.* 70: 119–137, 1990.