

LUND UNIVERSITY

Journey from the centre of the Earth

Scherstén, Anders; Elliott, Tim; Hawkesworth, Chris

Published in: Planet Earth

2005

Link to publication

Citation for published version (APA): Scherstén, A., Elliott, T., & Hawkesworth, C. (2005). Journey from the centre of the Earth. Planet Earth, (Spring issue), 22-22. http://www.nerc.ac.uk/publications/planetearth/2005/spring/spr05-journey.pdf

Total number of authors: 3

General rights

Unless other specific re-use rights are stated the following general rights apply: Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

• Users may download and print one copy of any publication from the public portal for the purpose of private study

or research. • You may not further distribute the material or use it for any profit-making activity or commercial gain

· You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: https://creativecommons.org/licenses/

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117 221 00 Lund +46 46-222 00 00

Journey from the centre of the Earth?

More that make up the Earth's surface. But some sit in the middle – the volcanoes of Hawaii are the best example – fuelled by unusually hot mantle below. Lavas from these volcanoes should tell us about the inside of our planet, but it is difficult to establish exactly where they come from. Some researchers argue the hot plumes in the mantle that drive these volcanoes contain material recycled from the Earth's crust in a process called subduction (see box). Others think the mantle plumes come from much deeper, perhaps even incorporating small amounts of the Earth's core from a staggering 2,900 km below the crust.

Recently, researchers found telltale signatures in lava from Hawaii suggesting it came from deep within the planet. The Earth's outer core may have unusually high levels of the metals platinum and rhenium compared to another metal, osmium. Isotopes of platinum and rhenium undergo radioactive decay, creating forms of osmium with different molecular weights to the

The central iron core of our planet has a solid inner and a liquid outer layer. A silicate mantle surrounds the core, and on top of that is the Earth's solid crust. The mantle gets heat both from the core, and by the decay of radioactive isotopes. Hot material is buoyant, and this sets up a slow convective circulation that dissipates the heat. As mantle material moves upwards and outwards it melts, and some forces through the Earth's crust, forming active volcanoes. New oceanic crust is made this way, along ridges in the middle of the oceans. But since the Earth is not expanding, this constant growth is balanced. Elsewhere, the plates in the Earth's crust are pushed together, driving one down into the mantle. This subduction takes place beneath arcs of volcanoes, such as those in the Western Pacific, Western North America and the Caribbean.

most abundant form (isotopes). So researchers finding high levels of these osmium isotopes in the lava inferred at least some of it must come from the core.

But there's a problem. Since platinum and rhenium still exist in the silicate portion of the Earth (i.e. all but the metallic core), and continue to decay into isotopes of osmium, the ratios of isotopes will change over geological time in the silicate Earth too. So our team turned to a more robust test, using an isotope whose 'parent' isotope hasn't existed on Earth for more than four billion years.

The isotope hafnium 182 was present at the birth of our solar system. It survived for only around 50 million years before it had all decayed to form an isotope of tungsten. We know the proportions of this tungsten isotope in the Earth's mantle, but intriguingly, there are about two parts per 10,000 more in the Earth's mantle than in meteorites made up of the same material from which the Earth formed. It might not sound much, but it's significant. If you accept that the overall proportions in the Anders Scherstén, Tim Elliott and Chris Hawkesworth explain how an extinct isotope can tell us about our planet's inner workings.



meteorites and newly-formed Earth would have been the same, the tungsten isotope on Earth must have become slightly concentrated in the mantle, almost certainly because of core formation. In turn, this means that any material from the core that eventually reaches the surface as lava should have lower levels of the tungsten isotope than mantle material. Crucially, since all hafnium 182 decayed over four billion years ago, there will have been no changes in the

ratio of the tungsten isotope to its more usual form since then.

Our team reanalysed the Hawaiian rocks. None had tungsten isotope ratios suggesting they came from the core. But we know metals like osmium are scarce in the Earth's mantle, so why do the unusual osmium ratios occur? We now think they support the idea that the Earth's crust is recycled when it's forced downwards where plates collide. Sediments on ocean floors are often rich in metals, such as manganese, which can even form nodules or crusts on the seafloor. These crusts could have similar osmium ratios to those seen in the Hawaiian lavas because they too have relatively high ratios of platinum to osmium, and so with time develop distinctive osmium isotopes.

Chris Hawkesworth and Tim Elliott can be contacted through the Department of Earth Sciences, University of Bristol, Wills Memorial Building, Queen's Road, Bristol, BS8 1RJ. Tel: 0117 9545400, email: c.j.hawkesworth@bris.ac.uk or Tim.Elliott@bris.ac.uk. Anders Scherstén is now at Faculteit der Aard- en Levenswetenschappen. Tel: +31 (0)20 598 7335, email: Anders.Schersten@falw.vu.nl.