



# LUND UNIVERSITY

## Experiences from treatment-predictive KRAS testing; high mutation frequency in rectal cancers from females and concurrent mutations in the same tumor

Jönsson, Mats; Isinger Ekstrand, Anna; Edekling, Thomas; Eberhard, Jakob; Grabau, Dorthe; Borg, David; Nilbert, Mef

*Published in:*  
BMC Clinical Pathology

*DOI:*  
[10.1186/1472-6890-9-8](https://doi.org/10.1186/1472-6890-9-8)

2009

[Link to publication](#)

### *Citation for published version (APA):*

Jönsson, M., Isinger Ekstrand, A., Edekling, T., Eberhard, J., Grabau, D., Borg, D., & Nilbert, M. (2009). Experiences from treatment-predictive KRAS testing; high mutation frequency in rectal cancers from females and concurrent mutations in the same tumor. *BMC Clinical Pathology*, 9(Oct 15), 8. Article 8. <https://doi.org/10.1186/1472-6890-9-8>

*Total number of authors:*  
7

### **General rights**

Unless other specific re-use rights are stated the following general rights apply: Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117  
221 00 Lund  
+46 46-222 00 00

Research article

Open Access

## Experiences from treatment-predictive *KRAS* testing; high mutation frequency in rectal cancers from females and concurrent mutations in the same tumor

Mats Jönsson<sup>1</sup>, Anna Ekstrand<sup>1</sup>, Thomas Edekling<sup>2</sup>, Jakob Eberhard<sup>1</sup>,  
Dorthe Grabau<sup>3</sup>, David Borg<sup>1</sup> and Mef Nilbert\*<sup>1,4</sup>

Address: <sup>1</sup>Department of Oncology, Institute of Clinical Sciences, Lund university, Lund, Sweden, <sup>2</sup>Department of Oncology, Växjö Hospital, Växjö, Sweden, <sup>3</sup>Department of Pathology, Institute of Clinical Sciences, Lund university, Lund, Sweden and <sup>4</sup>Clinical Research Centre, Hvidovre Hospital, Copenhagen University, Hvidovre, Denmark

Email: Mats Jönsson - mats.jonsson@med.lu.se; Anna Ekstrand - anna.ekstrand@med.lu.se;  
Thomas Edekling - thomas.edekling@ltkronoberg.se; Jakob Eberhard - jakob.eberhard@med.lu.se; Dorthe Grabau - dorthe.grabau@skane.se;  
David Borg - david.borg@skane.se; Mef Nilbert\* - mef.nilbert@med.lu.se

\* Corresponding author

Published: 15 October 2009

Received: 15 June 2009

*BMC Clinical Pathology* 2009, **9**:8 doi:10.1186/1472-6890-9-8

Accepted: 15 October 2009

This article is available from: <http://www.biomedcentral.com/1472-6890/9/8>

© 2009 Jönsson et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Abstract

**Background:** *KRAS* mutations represent key alterations in colorectal cancer development and lead to constitutive EGFR signaling. Since EGFR inhibition represents a therapeutic strategy in advanced colorectal cancer, *KRAS* mutation analysis has quickly been introduced as a treatment-predictive test.

**Methods:** We used a real-time PCR based method to determine *KRAS* mutations in 136 colorectal cancers with mutations identified in 53 (39%) tumors.

**Results:** *KRAS* mutations were significantly more often found in rectal cancer (21/38, 55%) than in colon cancer (32/98, 33%) ( $P = 0.02$ ). This finding was explained by marked differences mutation rates in female patients who showed mutations in 33% of the colon cancers and in 67% of the rectal cancers ( $P = 0.01$ ). Concurrent *KRAS* mutations were identified in three tumors; two colorectal cancers harbored Gly12Asp/Gly13Asp and Gly12Cys/Gly13Asp and a third tumor carried Gly12Cys/Gly12Asp in an adenomatous component and additionally acquired Gly12Val in the invasive component.

**Conclusion:** The demonstration of a particularly high *KRAS* mutation frequency among female rectal cancer patients suggests that this subset is the least likely to respond to anti-EGFR therapies, whereas the observation of concurrent *KRAS* mutations imply that repeated *KRAS* targeting may occur during tumor progression in a subset of colorectal cancers.

### Background

Inhibition of the epidermal growth factor receptor (EGFR) signaling pathway represents a therapeutic option in advanced colorectal cancer. Improved response rates and

prolonged time to metastasis/survival has been demonstrated with the currently registered EGFR blocking antibodies cetuximab and panitumumab, and additional EGFR inhibitors are in various stages of clinical trials.

Mutations in the *KRAS* oncogene typically occur already in the late adenoma stage and have since long been recognized as a key event in colorectal cancer development [1]. Overall, *KRAS* mutations are found in about 40% of the tumors and are predominantly located in codons 12 (82% of the mutations reported) and 13 (17%) <http://www.sanger.ac.uk/genetics/CGP/cosmic>. Activating mutations lead to permanently GTP bound *KRAS* and constitutive downstream pathway signaling, also in the absence of upstream EGFR stimulation. Presence of *KRAS* mutations therefore represents a negative predictor of response to EGFR therapy and *KRAS* mutation testing has rapidly moved into the molecular diagnostic work-up of colorectal cancers considered for EGFR treatment [2,3]. Quality assurance programs for *KRAS* mutation testing and practice guidelines related to e.g. optimal testing material, methodological considerations and recommendations for the reporting of the results are currently being developed [4]. PCR-based assays constitute the cornerstone for clinical *KRAS* testing since these analyses allow high-throughput testing and have a favorable sensitivity, also in samples with low tumor cell content. We report the experiences from our first 136 treatment-predictive *KRAS* tests and herein report significant differences in mutation frequencies in colon cancer and rectal cancer and coexisting *KRAS* mutations in a subset of the tumors.


## Methods

*KRAS* mutation testing was performed in 136 adenocarcinomas of the colon ( $n = 98$ ) and the rectum ( $n = 38$ ). The mean age was 56 (21-81) years and the series included 64 (47%) females. Representative tumor blocks were selected and were in the majority of the cases derived from the primary tumor, in 13 cases from metastatic tissue and in four cases from a local recurrence. Presence of at least 20% tumor cells in the tissue was verified by a pathologist. DNA was extracted from serial sections of formalin-fixed, paraffin-embedded tumor tissue using the Qiamp DNA FFPE tissue Kit (Qiagen, Hilsen, Germany) according to the manufacturer's recommendations. Standard clinical analysis applied the DxS real-time PCR based kit (Roche Diagnostics, Basel, Switzerland), which identifies 7 different *KRAS* mutations in exons 12 and 13 with high sensitivity. In order to confirm presence of double or triple coexisting mutations, samples with such unusual patterns were also subjected to pyrosequencing (PyroMark™ Q96 *KRAS* v2.0, Qiagen, Hilsen, Germany) according to the manufacturer's recommendations on PSQ™HS96A [5]. Mutations were quantified using the machine's software. Statistical analysis used a Chi2 test and the level of significance was set at 5%. *KRAS* mutation testing was carried out as part of standard care, all patients provided informed consent for testing, and the study was conducted according to the Helsinki declaration.

## Results

*KRAS* mutations were identified in 53/136 (39%) colorectal cancers. Overall, mutation status did not correlate with sex when analyzed in the whole cohort with mutations in 28/64 (44%) women and in 5/72 (35%) men ( $P = 0.28$ ) or age (mean age 55 years in the mutant group and 57 years in the wild-type group). *KRAS* mutations, however, significantly correlated with tumor location with mutations found in 32/98 (33%) colon cancers and 21/38 (55%) rectal cancers ( $P = 0.02$ ; Chi2-test). This difference was related to a high mutation rate in female rectal cancer patients (14/21, 67%) compared to females with colon cancer (14/43, 33%) ( $p = 0.01$ ), whereas no significant difference related to tumor location was identified in men (41% in rectal cancer and 33% in colon cancer,  $P = 0.52$ ). Though the materials in subsets are small, women with *KRAS* mutant rectal cancers were diagnosed mean 10 years earlier (mean age 48, range 25-71, years) than those with *KRAS* wild-type rectal cancers (mean age 58, range 52-70, years).

The mutation spectrum revealed the expected codon 12 and 13 mutations with the p.Gly12Asp, p.Gly13Asp, and p.Gly12Val being the most commonly found (table 1) [6]. Three tumors showed two-three different *KRAS* mutations (figure 1), which were identified using real-time PCR and verified in a separate tumor block and using pyrosequencing as a complementary method. The first case was a colon cancer from a 65-year old man with hereditary nonpolyposis colorectal cancer and a germline mutation in the *MLH1* gene. The tumor was found to harbor two *KRAS* mutations, Gly12Asp ( $\Delta$ CT value 6.9, cutoff 8) and Gly13Asp ( $\Delta$ CT value 5.6, cutoff 9). The patient had 10 years earlier developed two synchronous colon cancers, both of which were retrospectively analyzed, but neither of these contained any *KRAS* mutation (data not shown). The second case was a 68-year old man with a rectal cancer metastatic to the liver. The primary tumor revealed coexisting coexistence of the *KRAS* mutations Gly12Cys ( $\Delta$ CT value 5.4, cutoff 7.0) and Gly13Asp ( $\Delta$ CT 4.8, cutoff 9.0). The third case was a 47-year old woman who presented with a rectal cancer and concurrent liver metastases. The initial biopsy from the rectal tumor revealed only adenomatous components, whereas a second biopsy contained infiltrative adenocarcinoma. Unfortunately, no histologic sample from the liver lesions was available. The tubulovillous adenoma harbored the mutations Gly12Asp ( $\Delta$ CT value 6.2, cutoff 8.0) and Gly12Cys ( $\Delta$ CT value 3.9, cutoff 7.0), whereas the sample with also an invasive component contained both these mutations and in addition (contained) a Gly12Val mutation ( $\Delta$ CT value 4.1, cutoff 6.5). All mutations were identified by real-time PCR and verified using pyrosequencing.



**Figure 1**  
**Real-time PCR analysis identified coexisting KRAS mutations in three cases.** Case 1) colon cancer with the Gly12Asp and Gly13Asp mutations, Case 2) colon cancer with coexisting Gly12Cys and Gly13Asp mutations, and Case 3) a rectal tumor that revealed Gly12Asp and Gly12Cys in adenomatous components and an additional Gly12Val mutation in the invasive tumor component. All mutations were verified using pyrosequencing. Case 1 contained 12% Gly12Asp and 24% Gly13Asp, and case 2 4% Gly12Cys and 34% Gly13Asp. Case 3 harboured 17% Gly12Asp and 22% Gly12Cys in the adenomatous component, and 10% Gly12Asp, 7% Gly12Cys, and 12% Gly12Val in the invasive component (data not shown).

**Discussion**

The overall KRAS mutation frequency of 39% is in line with the results from previous studies <http://www.sanger.ac.uk/genetics/CGP/cosmic>, though the different mutation rates - 33% in colon cancer and 55% in rectal cancer (p = 0.02) - has not been reported in conjunction with treatment-predictive testing. Distinct molecular pathways are preferentially involved in the adenoma-carcinoma sequence in the proximal and the distal colon with TP53 and KRAS mutations being favored in rectal cancer, whereas mismatch repair defects are more common in proximal colon tumors [7]. A higher muta-

tion frequency in rectal cancers compared to colon cancer has previously been described, although the magnitude has been lower than identified herein [8]. The effect seems to be limited to females with rectal cancer, 67% of whom harbored KRAS mutations. This observation is in line with the findings in a large population-based series from the Netherlands in which Brink et al. reported a higher KRAS mutation rate in female rectal cancer patients [8]. Furthermore, in the subset of females with rectal cancer the KRAS mutant tumors were diagnosed mean 10 years earlier (mean age 48 years) behaviour than the wild-type tumors (mean age 58). Though this observation is new and needs

**Table 1: Summary of KRAS mutations identified.**

KRAS mutation	Amino acid change	*Number (%) mutations	Distribution colon/rectum
c.35G>A	p.Gly12 Asp	19 (33)	13/6
c.38G>A	p.Gly13Asp	12 (21)	8/5
c.35G>T	p.Gly12Val	10 (18)	4/6
c.34G>T	p.Gly12Cys	7 (12)	6/1
c.34G>A	p.Gly12Ser	5 (9)	1/4
c.35G>C	p.Gly12Ala	4 (7)	3/1
c.34G>C	p.Gly12Arg	-	-

\*Represents 53 tumors, including 3 with coexisting mutations

confirmation, a recent study has demonstrated that early-onset colorectal cancers frequently show pathologic features associated with aggressive clinical features [9]. The tumors here investigated were selected based on potential EGFR treatment, which may bias the results toward tumors prone to metastatic development, though the overall *KRAS* mutation frequency of 39% and the relation between colon cancer and rectal cancer are as expected. The differences in *KRAS* mutation rates identified suggest that sex and tumor location influence the chance of clinical usefulness of EGFR inhibitors with a particularly low likelihood of benefit among female rectal cancer patients.

A stepwise accumulation of mutations in different pathways constitutes a hallmark of colorectal cancer development. The observation of coexisting *KRAS* mutations in a small number of colorectal cancers, however, suggests that some tumors benefit from repeated targeting of the same pathway. This observation is in line with the association between *KRAS* mutations and 12p gain in colorectal adenomas [10]. Recently, the "two-hit-hypothesis" was challenged in a study of inactivating *APC* gene mutations in colorectal cancer, in which three hits - including mutations, deletions, and copy number gain - were demonstrated in a subset of the tumors [11]. *APC* gene mutations, as well as *KRAS* mutations, are recognized as early events in colorectal cancer development. Based on our results and previous observations, concurrent *KRAS* mutations seem to occur in about 3% of colorectal cancers. The majority of these cases show coexisting mutations in codon 12, whereas a smaller number have mutations in different codons (12 and 13 or 12 and another codon) [8,12-15]. Different *KRAS* mutations have shown variable efficacy in downstream signaling transduction and also seem to involve different effector molecules [16,17]. The specific *KRAS* mutation may thereby influence different aspects of tumor progression, which could explain why repeated *KRAS* targeting is observed in a subset of colorectal cancer. Interestingly the p.Gly12Val mutation was found in 10 rectal cancers and 4 colon cancers, suggesting an overrepresentation in the former tumor type. This mutation has also been linked to a worse prognosis [16]. The addition of a p.Gly12Val mutation in the invasive component of a tumor already containing dual *KRAS* mutations (figure 1, case 3) suggests a role for repeated *KRAS* targeting during tumor progression and serves as a reminder to consider coexisting mutations when unexpected patterns occur in treatment predictive testing.

## Conclusion

In conclusion, our experience from predictive testing for *KRAS* mutations reveal a high mutation frequency (67%) in rectal cancers from females and thus implicates that this subset is the least likely to respond to anti-EGFR ther-

apies. Concurrent *KRAS* mutations were observed in 3/136 tumors and although it remains a rare finding suggests that repeated *KRAS* targeting may occur during colorectal cancer progression.

## Competing interests

The investigators have received an unrestricted research grant and the principal investigator (M.N.) has in the past five years received lecture honorary from Merck A/S. No other competing interests apply.

## Authors' contributions

MJ carried out mutation analysis and drafted the manuscript. AE performed mutation analysis. TE, JE and DB identified and clinically managed patients and herein contributed with clinical data, DG selected tumor tissues, MN planned the study and was responsible for the clinical testing. All authors have contributed to and approved of the manuscript.

## Acknowledgements

This study was supported by the Swedish Cancer Fund, the Medical Research Council and an unrestricted research grant from Merck A/S.

## References

1. Cho KR, Vogelstein B: **Genetic alterations in the adenoma-carcinoma sequence.** *Cancer* 1992, **70**:1727-31.
2. Lindarou H, Dahabreh J, Kanaloipiti D, Siannis F, Bafaloukos D, Kosmidis P, Papadimitriou CA, Murray S: **Assessment of somatic *KRAS* mutations as a mechanism associated with resistance to EGFR-targeted agents: a systematic review and meta-analysis of studies in advanced non-small-cell lung cancer and metastatic colorectal cancer.** *Lancet Oncol* 2008, **9**:962-72.
3. Allegra CJ, Jessup JM, Somerfield MR, Hamilton SR, Hammond EH, Hayes DF, McAllister PK, Morton RF, Schilsky RL: **American Society of Clinical Oncology provisional clinical opinion: testing for *KRAS* gene mutations in patients with metastatic colorectal carcinoma to predict response to anti-epidermal growth factor receptor monoclonal antibody therapy.** *J Clin Oncol* 2009, **27**:2091-6.
4. Van Krieken JHJM, Jung A, Kirchner T, et al.: ***KRAS* mutation testing for predicting response to anti-EGFR therapy for colorectal carcinoma: proposal for an European quality assurance programme.** *Virchow Arch* 2008, **453**:417-31.
5. Ogino S, Kawasaki T, Brahmandam M, Ogino S, Kawasaki T, Brahmandam M, Yan L, Cantor M, Namgyal C, Mino-Kenudson M, Lauwers GY, Loda M, Fuchs CS: **Sensitive sequencing method for *KRAS* mutation detection by Pyrosequencing.** *J Mol Diagn* 2005, **7**:413-21.
6. Andreyev HJN, Norman AR, Cunningham D, et al.: **Kirsten ras mutation in patients with colorectal cancer: the "RASCAL II" study.** *Br J Cancer* 2001, **85**:692-6.
7. Iacopetta B: **Are there two sides to colorectal cancer?** *Int J Cancer* 2002, **101**:403-8.
8. Brink M, de Goeij FPM, Weijenberg MP, Brink M, Weijenberg MP, De Goeij AF, Schouten LJ, Koedijk FD, Roemen GM, Lentjes MH, De Bruine AP, Goldbohm RA, Brandt PA Van Den: **K-ras oncogene mutations in sporadic colorectal cancer in the Netherlands cohort study.** *Carcinogenesis* 2003, **24**:703-10.
9. Yantiss R, Goodarzi M, Zhou X, Rennert H, Pirog EC, Banner BF, Chen YT: **Clinical, pathologic, and molecular features of early-onset colorectal carcinoma.** *Am J Surg Pathol* 2009, **33**:572-82.
10. Gattenlohner S, Germer C, Muller-Hermelink H-K: **Letter to the editor.** *N Engl J Med* 2009, **380**:835.

11. Leslie A, Stewart A, Baty DU, et al: **Chromosomal changes in colorectal adenomas: relationship to gene mutations and potential for clinical utility.** *Genes Chrom Cancer* 2006, **45**:126-35.
12. Segdits S, Rowan AJ, Howarth K, et al: **APC and the three-hit hypothesis.** *Oncogene* 2009, **28**:146-55.
13. Miyakura Y, Sugano K, Fukayama N, Konishi F, Nagai H: **Concurrent mutations of K-ras oncogene at codons 12 and 22 in colon cancer.** *Jpn J Clin Oncol* 2002, **32**:219-21.
14. Martinez-Garza SG, Nunez-Salazar AJ, Calderon-Garciduenas AL, Bosques-Padilla FJ, Niderhauser-García A, Barrera-Saldaña HA: **Frequency and clinicopathology associations of K-ras mutations in colorectal cancer in a northeast Mexican population.** *Dig Dis* 1999, **17**:225-9.
15. Moerkerk P, Arends JW, van Driel M, de Bruine A, de Goeij A, ten Kate J: **Type and number of Ki-ras point mutations relate to stage of human colorectal cancer.** *Cancer Res* 1994, **54**:3376-8.
16. Al-Mulla F, Milner-White EJ, Going J, Birnie G: **Structural differences between Valine-12 and Aspartate-12 Ras proteins may modify carcinoma aggression.** *J Pathology* 1999, **187**:433-38.
17. Monticone M, Biollo E, Maffei M, Donadini A, Romeo F, Storlazzi CT, Giaretti W, Castagnola P: **Gene expression deregulation by KRAS G12D and G12V in a BRAF V600E context.** *Mol Cancer* 2008, **7**:92.

### Pre-publication history

The pre-publication history for this paper can be accessed here:

<http://www.biomedcentral.com/1472-6890/9/8/prepub>

Publish with **BioMed Central** and every scientist can read your work free of charge

*"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."*

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

