



# LUND UNIVERSITY

## Clinical and Microbiological Characteristics of Severe Streptococcus pyogenes Disease in Europe.

Luca, Bogdan; Darenberg, Jessica; Neal, Shona; Siljander, Tuula; Strakova, Lenka; Tanna, Asha; Creti, Roberta; Ekelund, Kim; Koliou, Maria; Tassios, Panayotis T; van der Linden, Mark; Straut, Monica; Vuopio-Varkila, Jaana; Bouvet, Anne; Efstratiou, Androulla; Schalén, Claës; Henriques-Normark, Birgitta; Jasir, Aftab

*Published in:*  
Journal of Clinical Microbiology

*DOI:*  
[10.1128/JCM.02155-08](https://doi.org/10.1128/JCM.02155-08)

2009

[Link to publication](#)

### *Citation for published version (APA):*

Luca, B., Darenberg, J., Neal, S., Siljander, T., Strakova, L., Tanna, A., Creti, R., Ekelund, K., Koliou, M., Tassios, P. T., van der Linden, M., Straut, M., Vuopio-Varkila, J., Bouvet, A., Efstratiou, A., Schalén, C., Henriques-Normark, B., & Jasir, A. (2009). Clinical and Microbiological Characteristics of Severe Streptococcus pyogenes Disease in Europe. *Journal of Clinical Microbiology*, 47, 1155-1165.  
<https://doi.org/10.1128/JCM.02155-08>

*Total number of authors:*  
18

### General rights

Unless other specific re-use rights are stated the following general rights apply:  
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117  
221 00 Lund  
+46 46-222 00 00



LUND UNIVERSITY  
Faculty of Medicine

---

LUP

*Lund University Publications*

Institutional Repository of Lund University

---

This is an author produced version of a paper published in Journal of clinical microbiology. This paper has been peer-reviewed but does not include the final publisher proof-corrections or journal pagination.

Citation for the published paper:

Bogdan Luca, Jessica Darenberg, Shona Neal, Tuula Siljander, Lenka Strakova, Asha Tanna, Roberta Creti, Kim Ekelund, Maria Koliou, Panayotis T Tassios, Mark van der Linden, Monica Straut, Jaana Vuopio-Varkila, Anne Bouvet, Androulla Efstratiou, Claës Schalén, Birgitta Henriques-Normark, Aftab Jasir

"Clinical and Microbiological Characteristics of Severe Streptococcus pyogenes Disease in Europe."

Journal of clinical microbiology,  
2009 Jan 21. [Epub ahead of print]

<http://dx.doi.org/10.1128/JCM.02155-08>

Access to the published version may  
require journal subscription.

Published with permission from:  
American Society for Microbiology

**Clinical and Microbiological Characteristics of Severe *Streptococcus pyogenes* Disease in Europe**

**Bogdan Luca-Harari<sup>1\*</sup>, Jessica Darenberg<sup>2\*</sup>, Shona Neal<sup>3</sup>, Tuula Siljander<sup>4</sup>, Lenka Strakova<sup>5</sup>, Asha Tanna<sup>3</sup>, Roberta Creti<sup>6</sup>, Kim Ekelund<sup>7</sup>, Maria Koliou<sup>8</sup>, Panayotis T. Tassios<sup>9</sup>, Mark van der Linden<sup>10</sup>, Monica Straut<sup>11</sup>, Jaana Vuopio-Varkila<sup>4</sup>, Anne Bouvet<sup>12</sup>, Androulla Efstratiou<sup>3</sup>, Claes Schalén<sup>1</sup>, Birgitta Henriques-Normark<sup>2</sup>, the Strep-EURO study group<sup>†</sup>, and Aftab Jasir<sup>1,13§</sup>**

<sup>1</sup>Department of Laboratory Medicine, Division of Medical Microbiology, Lund University, Lund, Sweden; <sup>2</sup>Swedish Institute for Infectious Disease Control, Solna, Sweden; <sup>3</sup>Respiratory and Systemic Infections Laboratory, Health Protection Agency, London, UK; <sup>4</sup>National Public Health Institute, Helsinki, Finland; <sup>5</sup>National Institute of Public Health, Prague, Czech Republic; <sup>6</sup>Department of Infectious, Parasitic and Immune-mediated Diseases, Istituto Superiore di Sanità, Rome, Italy; <sup>7</sup>Statens Serum Institut, Copenhagen, Denmark; <sup>8</sup>Archbishop Makarios Hospital, Nicosia, Cyprus; <sup>9</sup>University of Athens, Athens, Greece; <sup>10</sup>German National Reference Center for Streptococci, Department of Medical Microbiology, University Hospital RWTH Aachen, Germany; <sup>11</sup>Molecular Epidemiology Laboratory, Cantacuzino Institute, Bucharest, Romania; <sup>12</sup>National Reference Center for Streptococci, Associated Laboratory for group A streptococci, Department of Microbiology, Hotel Dieu AP-HP, Paris Descartes University, France and <sup>13</sup> Clinical Microbiology and Immunology, Lund University Hospital (USIL), Lund, Sweden

1 † **The Strep-EURO study group (except above already listed authors):** Cyprus (Nasia  
2 Hannidou), Czech Republic (Paula Kriz, Jitka Motlova), Denmark (Margit S. Kaltoft), Finland  
3 (Joonas Iivonen, Jari Jalava), France (Julien Loubinoux, Liliana Mihaila), Germany (Rudolf  
4 Lütticken, Ralf René Reinert), Greece (Joseph Papaparaskevas, Levantia Zacharidou, Nicholas J.  
5 Legakis), Italy (Lucilla Baldassarri, Monica Imperi, Graziella Orefici), Romania (Vasilica  
6 Ungureanu), Sweden (Anna Norrby-Teglund, Lars Björck), UK (Neelam Alhaddad, Michaela  
7 Emery, Catherine Keshishian, Theresa Lamagni)

8  
9 \*The first two authors contributed equally to this study

10  
11 §Corresponding author. Mailing address: Department of Clinical Microbiology and Immunology,  
12 Lund University hospital (USIL), Sölvegatan 23, 22362, Lund, Sweden.  
13 Telephone: +46 46 173286. Fax: +46 46 135936. E-mail: [aftab.jasir@med.lu.se](mailto:aftab.jasir@med.lu.se)

14  
15 **Key words:** *Streptococcus pyogenes*; Virulence factors; *emm*-type; Superantigens; Streptococcal  
16 toxic shock syndrome; necrotizing fasciitis; Puerperal sepsis

## 1    **ABSTRACT**

2    In an attempt to compare the epidemiology of severe *S. pyogenes* infection within Europe,  
3    prospective data were collected through the Strep-EURO programme. Surveillance of severe *S.*  
4    *pyogenes* infection diagnosed during 2003 and 2004 was undertaken in eleven countries across  
5    Europe using a standardised case definition and questionnaire. Patient data as well as bacterial  
6    isolates were collected and characterized by T- and, M/*emm*-typing and selected strains were  
7    analysed for presence of superantigen genes. Data were analysed to compare the clinical and  
8    microbiological patterns of infections across participating countries.

9    Totally 4353 isolates were collected from 5521 cases with severe *S. pyogenes* infection identified.  
10    It was wide diversity of M/*emm*-types (104) found among the *S. pyogenes* clinical isolates but  
11    M/*emm*-type distribution varied broadly between participating countries. The ten most  
12    predominant M/*emm*-types were 1, 28, 3, 89, 87, 12, 4, 83, 81, and 5 in descending order. A  
13    correlation was found between some specific disease manifestation, age of patients and *emm*-types.  
14    Streptococcal toxic shock syndrome and necrotizing fasciitis, although caused by a large number  
15    of types, were particularly associated with M/*emm*-types 1 and 3.

16    The *emm*-types included in the 26-valent vaccine under development, were generally well  
17    represented in the present material; 16 of the vaccine types accounted for 69% of isolates. The  
18    Strep-EURO collaborative programme has contributed to enhance the knowledge on the spread of  
19    invasive disease caused by *S. pyogenes* within Europe and encourage future surveillance with  
20    notification of cases and characterisation of strains, important for vaccine strategies and other  
21    health care issues.

## 1 INTRODUCTION

2 *Streptococcus pyogenes* (group A streptococcus, GAS), a major human pathogen (9) studied for  
3 decades may give rise to common throat and skin infections, but also to invasive diseases, such as  
4 arthritis, septicaemia, cellulitis, puerperal fever, necrotising fasciitis (NF) and streptococcal toxic  
5 shock syndrome (STSS) (14). Since the mid 1980's there are increasing numbers of reports  
6 describing severe GAS manifestations, however the underlying factors of this pathogens  
7 worldwide resurgence remaining unknown (20).

8 The M-protein, encoded by the *emm*-gene, is an important virulence factor, and also an  
9 epidemiological marker that are used throughout the world to characterize GAS (5, 21-23). The  
10 type specificity of the M-protein, of which there are more than 100 different types known, is  
11 largely determined by the epitope located in 40 to 50 amino acid residues at the amino-terminal (4,  
12 16, 27). These regions of M-proteins have been shown to evoke antibodies with great bactericidal  
13 activity, not likely cross-reactive with human tissues (3, 16). Hence,, an approach in the  
14 development of a GAS vaccine has been to combine small amino-terminal M-protein peptides to  
15 make multivalent vaccines that would elicit opsonic antibodies against epidemiologically  
16 important GAS serotypes (15). Also other surface proteins, like the serum opacity factor (SOF)  
17 and the T-protein are used to characterize different GAS types. In addition to the known linkage  
18 between T-serotype, SOF production, and *emm*-type (25, 26), several studies also indicated  
19 correlations between *emm*-types, disease manifestations, and also other virulence factors,  
20 especially the superantigens (SAg) (7, 10, 40, 42).

21 Epidemiological studies, providing the type distributions in the communities, are of basic  
22 importance for identification and control of streptococcal infections. Furthermore, by tracing  
23 selected virulence features of isolates causing disease, the understanding of pathogenic

mechanisms of the various disease manifestations would be enhanced. In order to improve knowledge on severe GAS infections, the Strep-EURO programme was implemented during 2003-2004. Overall epidemiological findings of the programme were reported recently (29). In the present paper, type characteristics and SAg repertoire of the streptococcal isolates are described and also possible associations with clinical findings.

## **MATERIAL AND METHODS**

**Clinical data and isolates.** Through collaboration between eleven European countries (Cyprus, Czech Republic, Denmark, Finland, France, Germany, Greece, Italy, Romania, Sweden and the UK) on the epidemiology of invasive GAS disease, enhanced surveillance was undertaken between January 1<sup>st</sup> 2003 and December 31<sup>st</sup> 2004. Methods employed to identify cases varied by country, but mostly implied invited submission of isolates from local microbiology laboratories to the national streptococcal reference centre. Demographic and clinical data, as well as risk factor information were collected through a standardized questionnaire, with the exception of Denmark and Sweden, where surveillance with earlier designed questionnaires was already operational.

**Case definition and isolate identification.** For invasive GAS disease and STSS the consensus definition proposed by the Working Group on Severe Streptococcal Infections in 1993 was used (45). Identification of GAS isolates was confirmed using morphological and growth characteristics, bacitracin susceptibility, or pyrolidonyl-arylamidase testing and latex agglutination with group A antisera commercially available.

**Typing of isolates.** Isolates were T-typed using commercial poly- and monospecific T-antisera, according to the manufacturer's recommendation (Sevapharma, Prague, Czech Republic) (Moody *et al.*, 1965). M/*emm*-typing was performed using somewhat different methods between countries,

thus evaluated and further described in an external quality assurance (EQA) study (33). Although both serological and/or genotypical methods were used to determine the M/*emm*-types, the results are hereafter referred to as *emm*-types. The *emm*-sequences obtained by sequence based methods were identified by comparisons to available sequences in the CDC database ([ftp://ftp.cdc.gov/pub/infectious\\_diseases/biotech/tsemml/](ftp://ftp.cdc.gov/pub/infectious_diseases/biotech/tsemml/)). Unusual type-combinations between T- and *emm*-types (rare or previously not reported) were verified blindly by another participating reference center.

**SAg gene detection.** Presence of SAg genes was tested in Lund, Sweden for 1127 isolates from five countries (Czech Republic, Denmark, Finland, France, and Romania) and included *speA*, *speB*, *speC*, *speF*, *speG*, *speH*, *speI*, *speJ*, *ssa*, and *smeZ* (30). Isolates from remaining countries were tested in the respective national centres: Swedish isolates were tested for all the above mentioned SAg genes but *speI* (18), Greek and Italian isolates for *speA*, *speB*, and *speC* (13), and German isolates for *speA*, *speC*, and *ssa* (47). A fraction consisting of 256 UK isolates (18% of the total *emm*-typed strains) were tested locally for the presence of *speA*, *speB*, and *speC*. In addition, 193 isolates (covering 38 out of the 74 different *emm*-types identified in the UK) were tested in Lund as described above.

**Statistical analysis.** Data were analyzed using GraphPad Prism, version 4 (GraphPad Software) and SAS, version 9.1.3, proc logistic (SAS Institute). For nominal data,  $\chi^2$  test or Fisher's exact test were used when appropriate. Logistic regression was performed using *emm*-type as outcome and clinical conditions or risk factors as predictors. The analyses were performed separately for each of the 10 most prevalent *emm*-types, and compared to the group consisting of cases caused by all other types (i.e. except the 10 most prevalent ones). Each model was reduced by backward

- 1 elimination where the significant level was set at 5%. In the logistic regression analyses, only
- 2 cases with age, gender and clinical/ risk factor information available were included.

## RESULTS

From a total of 5521 patients with invasive streptococcal disease, 4354 (79%) bacterial isolates were submitted to the reference centers in the participating countries. Clinical information was available for 3404 isolates (62% of all cases).

**T-types.** In total 4171 isolates were subjected to T-typing, 408 (10%) of these being non typeable (NT). Fifty different T-types, or type profiles, were recognized, the most prevalent were T1 (19%), T28 (18%), T3/13/B3264 (23%), T12 (8%), T4 (5%), T5 (3%), T6, T11, and T8/25/Imp19 (2% each) (table 1).

**emm-types.** Among 4353 *emm*-typed isolates one hundred and four different types were identified, of which the most prevalent ( $\geq 2\%$ ) were *emm1* (19%), 28 (12%), 3 (10%), 89 (8%), 87 (6%), 12 (5%), 4 (5%), 83 (3%), 81 (3%) and 5, 77, 6, 22, and 18 (2% each) (table 1). The type distribution varied significantly between the eleven countries, but the overall prevalence was strongly influenced by the large proportion of isolates originating from the UK (figure 1), and also from Sweden. Although *emm87* and *emm83* were the fifth and eighth overall most common types, majority of these isolates were from the UK (93% and 90% of isolates respectively). In total 34 different *emm*-types encompassed the ten most prevalent types in the eleven countries. Importantly, *emm1* was the most abundant type in the majority of countries, with a proportion ranging between 15% and 33% of isolates. In contrast, within Denmark, Finland, and Sweden *emm28* was the most prevalent type, ranging from 16% to 45% of isolates. As shown in figure1B, certain types among the overall ten most prevalent *emm*-types were absent in some of the countries; *e.g* in Romania only three of the overall ten most prevalent types were found. Type *emm3* was infrequent in the Czech Republic, Finland, Greece, and Sweden with prevalence ranging from 1% to 5%, and absent in Romania. Type *emm43* was found exclusively in the UK.

Other types almost confined to the UK were *emm82* (93% from the UK), *emm5* (91%), *emm83* (90%), and *emm68* (81%). Type *emm53* was found only in the Czech Republic, Greece and the UK. All the *emm118* isolates (n=34) originated from either Denmark or Sweden.

**T/*emm*-type combinations.** As shown in table 1, the number of T/*emm*- type combinations was high (N=314), some of these were unfrequented, other previously not reported (underlined in the table). The most prevalent T-type was 3/13/B3264 (or combinations thereof *e.g.* 3/13, 13/B3264, 3/B3264) and associated to no less than 40 different *emm*-types. In general, *emm1* was limited to T1 (98%) but a small number of these isolates expressed T-types 3, 3/13/B3264 or 4.

**Correlation between age, gender and *emm*-types.** Among 600 isolates collected from children (0-17 years), the most frequent *emm*-types were, in descending order 1 (26%), 12 (11%), 4, 3 (10% each), and 28 (7%). In patients aged 18 and older, the most prevalent type was also *emm1* (19%), but followed by *emm28* (13%), *emm3* (10%) and *emm89* (9%).

A significant female predominance for *emm87* and *emm28* (58%,  $p<0.001$  for both) was found. Type *emm28* was also more prevalent in age groups 30-39 years (17%) and 70-79 years (19%), in the younger group strongly associated to females (80%,  $p<0.001$ ). Types *emm81* and *emm83* were significantly overrepresented among males (62%,  $p<0.05$ , and 68%,  $p<0.001$ , respectively).

**Seasonal fluctuations.** During the study period, several *emm*-types presented a steady seasonal prevalence, whereas other showed fluctuations (figure 2). Overall, 59% of cases were reported in the 6 winter months (January to April + November and December) in both years. In contrast, tendencies towards higher frequencies of *emm12* was noted during the warmer months (May-August;  $p<0.05$ ).

## **Disease manifestations, risk factors and *emm*-types.**

The most severe manifestations, STSS and NF, were caused by 45 different types, of which *emm1* was the most prevalent, accounting for 37% and 31% of cases respectively (table 2); in addition, a considerable proportion were caused by *emm3* isolates (17% and 14%, respectively). In the statistical regression model, when comparing each of the 10 most prevalent types versus the other types combined, STSS was statistically more often caused by *emm1* or *emm3* ( $p < 0.001$  for each).

Patients without focal symptoms were less often infected by *emm1* (17%,  $p < 0.05$ ), in contrast to types *emm81* (45%), *emm77* (47%) ( $p < 0.001$  for each), *emm83* (34%), and *emm87* (26%) ( $p < 0.05$  for each), that were more common among these patients (table 2.). Furthermore, patients with arthritis were less prone to be infected by *emm28* isolates (5%,  $p < 0.05$ ), and cellulitis was more often caused by either *emm87* (32%,  $p < 0.0001$ ) or *emm83* (30%,  $p < 0.05$ ), as compared to infections caused by types other than the 10 most prevalent. Though puerperal sepsis was caused by 16 different types and only 8% of *emm28* were patients with puerperal sepsis a clear correlation with *emm28* was noted (31% of cases,  $p < 0.001$ ). Other *emm* types significantly involved in causing puerperal sepsis were *emm1*, *emm89* and *emm87* (4% each,  $p < 0.001$  and  $< 0.05$  respectively).

Data regarding risk factors, as well as *emm*-type, were available for 2796 patients (table 3). Patients with diabetes were statistically more prone to an infection caused by either *emm81* ( $p < 0.001$ ) or *emm12* ( $p < 0.05$ ), as compared to “other types” in the logistic regression analysis.

Information on *emm*-type distribution among patients who were injecting drug users (IDU) was available for 359 of 471 (76%) cases, a majority of these (93%) was identified in the UK. The ten most prevalent types among these patients were, in descending order: *emm83*, 87, 82,

89, 81, 43, 33, 101, 1 and 53, accounting for 70% of these infections. Conversely, as many as 70% of *emm33*, *emm82* and *emm83*, and 54% of the *emm43* infections were IDU related.

Among 242 health care associated infections (HAI), the same types as the over all ten most prevalent ones caused the majority of infections (71%). However, *emm1* and *emm3* infections were less commonly related to surgery before disease onset, as determined by the regression model ( $p>0.05$  for each).

Among patients with chicken pox, the probabilities for *emm1* and *emm12* were high ( $p<0.001$  each), which is in concordance with the high frequency of both types among children.

**Case fatality rates and *emm*-types.** Overall, the CFR over 7 days among cases with typed isolates was 19% and highest among infections caused by *emm3* (36%), followed by *emm5*, 1, 43, and 77 (table 2). Furthermore, the highest CFRs were, as expected, noted among cases with STSS (44%) and NF (31%), and as already mentioned correlated to *emm1* and 3 infections. For patients with cellulitis, the overall CFR was 18%, but considerably higher for infectious caused by *emm77*, *emm3* (33%  $p<0.001$  each), or *emm1* (25%,  $p<0.05$ ) isolates. Among infections without focus the overall CFR was 15%, and the deaths predominantly caused by *emm3* (32%), *emm83* (19%), *emm87* (17%), *emm1* (16%), and *emm 28* (15%) infections (table 2).

**SAg genes patterns and *emm*-types.** As expected, *speB*, *speF*, and *speG*, were detected in the vast majority of strains, though *speG* was lacking among *emm4* and *emm77* isolates from several countries.

Data regarding *speA* and *speC* was available for 2321 isolates. Overall, 30% and 54 % were positive for *speA* and *speC*, respectively. As shown in table 4, *speA* was primarily associated with *emm1* and *emm3* ( $p<0.001$  for both), whereas *speC* was common in several other types such as *emm4*, 5, 6, 28 and 77 ( $p<0.001$  for each), *emm18* ( $p< 0.01$ ). Both *emm1* and *emm3* harboured

1 *speC* to a lesser extent ( $p<0.001$  for both) and the same was true for *emm* 81 and 12 isolates  
2 ( $p<0.05$  for both). The *speA* gene was less prevalent among Finnish and Swedish strains (10% and  
3 13%, respectively), ascribable to the *emm*-type distribution in these countries where both *emm*1  
4 and 3 isolates were less common than in the other countries (figure 1). However, among *emm*1 and  
5 *emm*3 isolates from the Czech Republic, Denmark and Finland frequencies of *speA* were lower,  
6 about 70% and 50%, for each type respectively, as compared to more than 90% among these  
7 isolates from remaining countries (data not shown). Conversely, the high proportion of *emm*28 in  
8 Finland was reflected in an overall higher prevalence of *speC* positive isolates (80%).

9         The presence of *speI* was investigated in more than 800 isolates from five countries, and  
10 only one percent of these isolates harboured the gene. The gene *speH* was detected in 10% of 1667  
11 isolates tested, most notably in *emm*12 (65%;  $p<0.001$ ) and *emm*81 (19%;  $p<0.01$ ) (table 4). The  
12 highest prevalence of *speH* among *emm*12 isolates was noted for Swedish (97%) and UK (91%)  
13 isolates, but surprisingly, *speH* was not detected among *emm*12 isolates from either Denmark or  
14 Finland. The gene *ssa* was detected in 31% of tested isolates, primarily among *emm*3 and *emm*4  
15 ( $p<0.001$  for both) but also among *emm*87 isolates ( $p<0.05$ ). However, *ssa* was less frequently  
16 found among *emm*1, *emm*81, *emm*89 ( $p<0.001$  each) and *emm*6 ( $p<0.05$ ) isolates (table 4).

## DISCUSSION

In the present paper clinical and microbiological data obtained from patients with severe GAS infections from the eleven Strep-EURO participating countries are presented. The number of characterized isolates (4353) exceeds any previous European study. Strikingly, the overall distribution of the most prevalent *emm*-types agreed closely with recent data reported from the US where *emm*-types 1, 3, 28, 12, and 89 accounted for 55% of invasive isolates collected over a period of four years (2000-2004) (35). However, the country-specific *emm*-type distributions differed markedly, as exemplified by. *emm*87 though overall highly represented, essentially confined to the UK (figure 1). Differences in type proportions were also noted between neighbouring countries, like Denmark, Finland, and Sweden. In Sweden, high rates of *emm*81 and *emm*89 was seen, accounting for 30% of isolates, whereas *emm*28 was the most prevalent type in Denmark (26%), and *emm*89 only accounted for 7% of cases (30). In Finland, 45% of all isolates were *emm*28, being the only country with such a large proportion of a single type. Isolates of *emm*3, in addition to *emm*1, have previously been shown to be of major role in invasive GAS disease (19, 46, 48). However, in Finland, the number of *emm*3 isolates was negligible (3 cases), and a low prevalence of this type was also noted in Greece, the Czech Republic and Sweden (3-4%). As shown in the Swedish study (18) *emm*-types of invasive cases essentially agreed with those recorded among cases with non-invasive GAS disease. Though non-invasive isolates were not studied in other participating countries, the country-specific type distributions may to a large extent reflect ongoing epidemic waves, herd immunity (39) or population mobility (11), as previously seen for streptococcal disease (39).

There were significant differences between genders regarding some particular types. For example, *emm*28 and *emm*87 were overrepresented among female cases. The role of *emm*28

isolates in puerperal fever has already been recognized (2, 32), as this type are known to express R28, which is related to the Rib protein in group B streptococci, the major cause of neonatal infections (38, 39). Recently it was shown that the gene encoding R28 is located on a 37.4-kb region (region of difference – RD2) similar in content and organization to a region described in group B streptococci, apparently acquired by horizontal gene transfer and enabling *emm28* strains to often cause puerperal sepsis (24, 49). Since *emm87* was not among those types carrying RD2 (e.g. M2, 4, 48, 77, 124), it is of interest to investigate whether *emm87* isolates may harbor similar pathogenic factors. In contrast, *emm*-types 83, 81, and 43, were associated with intravenous drug use and found preferentially among male patients (68%, 62%, and 61% respectively). Interestingly, also a predominance of *emm81* isolates among male patients with skin involvement were found in Sweden (18).

It is known that no *emm*-type can be uniquely associated to a particular disease, though there is evidence correlating certain types, e.g. *emm1* and *emm3* with the most severe GAS diseases NF and STSS (12, 31, 43, 44), or *emm28* with puerperal sepsis (36). However, in our material 50% of all STSS cases and 55% of NF cases were caused by types other than *emm1* and *emm3* respectively, and in Sweden no *emm3* strain was involved in STSS, indicating that most types of GAS may have the potential to give rise to these severe manifestations. However, the mortality associated with either *emm1* or *emm3*, whether causing STSS, NF or puerperal sepsis, clearly exceed that of remaining types which, in agreement with previous studies, demonstrates these two types as particularly virulent.

Over the years, the number of GAS SAgS identified have increased, and also the knowledge on their role in disease pathogenesis (8, 10, 14). The disease severity is also determined by many other GAS virulence factors (41) and is clearly host dependent (28, 34). In the present

study, a high occurrence of *speA* was found for isolates of *emm*-types 1 and 3, types that were often involved in severe infections, and also for the less frequent type *emm43*; these *emm*-types were associated with high CFRs (29%, 36%, and 21%, respectively). However, *emm5* and *emm18* cases had high CFRs (30%, and 21%, respectively), though these types lacked *speA* but harboured *speC* at high proportions (both 91%). In addition, the presence of *speC* was common in several prevalent types such as *emm4*, 6, 28, 77, 18, 81 and 12.

The *emm*-types included in the 26-valent vaccine now in clinical trial (17) were generally well represented in the present study (figure 3A). Within Strep-EURO, 16 of the vaccine types accounted for 69% of isolates, though proportion of coverage varied among participating countries (figure 3B), and the prevalence of some *emm*-types changed temporally, which could be at least partly related to epidemic waves (6), type substitution due to herd immunity, or population mobility (11). Nevertheless, the total number of *emm*-types detected exceeded one hundred, and expansion of non-vaccine types (1) and higher risk of infection by non-vaccine types(37), as is the case for recent pneumococcal experience after introduction of vaccination posing an obvious challenge to attempts of type-specific vaccine development.

In conclusion, among 104 GAS *emm*-types identified during the present project, 45 were involved in causing STSS and/or NF. A major role of *emm1* and *emm3* isolates in these severe entities, also found in previous studies was confirmed; however, a number of other types also caused high mortality rates, suggesting similar pathogenic potential. In general, the SAg gene repertoire of isolates appeared to correlate with *emm*-type in a complex pattern, precluding definite conclusions on the role of individual SAg for severe disease. The data here presented, demonstrating high mortality and devastating consequences of the invasive manifestations in

- 1 particular, should be of value for preventive work, including ongoing attempts at creating vaccine
- 2 prophylaxis against GAS disease.

## **Acknowledgements**

We direct our sincere thanks to all the clinicians and microbiologists across the eleven participating countries who took the time to report cases to their respective country leads. We would also like to thank Helena Pettersson at the Department of Epidemiology (Swedish Institute for Infectious Disease Control) for statistical analysis and advice. The following additional members of country teams also contributed significantly to the project: Cyprus - Maria Alexandrou, Yiannis Ioannou, Eleni Konteaitou; Czech Republic - Radmila Dousova, Iveta Mouchova; Finland - Sari Rantala, Petri Ruutu; France – Gislène Collobert; Germany - Claudia Brandt; Greece - Angeliki Stathi, Anastasia Pangalis; Italy - Marco Pataracchia, Simona Recchia; Sweden - Hans Tapper, Ulrich von Pawel-Rammingen, Madeleine Kais, Christina Johansson, Gunnel Mölleberg, Ingrid Andersson; UK - Chenchal Dhami

*Financial support.* EU Fifth Framework Research Programme (QLK2.CT.2002.013)

## References

1. **Aguiar, S. I., I. Serrano, F. R. Pinto, J. Melo-Cristino, and M. Ramirez.** 2008. Changes in *Streptococcus pneumoniae* serotypes causing invasive disease with non-universal vaccination coverage of the seven-valent conjugate vaccine. *Clin Microbiol Infect* **14**:835-43.
2. **Areschoug, T., F. Carlsson, M. Stalhammar-Carlemalm, and G. Lindahl.** 2004. Host-pathogen interactions in *Streptococcus pyogenes* infections, with special reference to puerperal fever and a comment on vaccine development. *Vaccine* **22 Suppl 1**:S9-S14.
3. **Beachey, E. H., and J. M. Seyer.** 1986. Protective and nonprotective epitopes of chemically synthesized peptides of the NH<sub>2</sub>-terminal region of type 6 streptococcal M protein. *J Immunol* **136**:2287-92.
4. **Beachey, E. H., J. M. Seyer, J. B. Dale, W. A. Simpson, and A. H. Kang.** 1981. Type-specific protective immunity evoked by synthetic peptide of *Streptococcus pyogenes* M protein. *Nature* **292**:457-9.
5. **Beall, B., R. Facklam, and T. Thompson.** 1996. Sequencing *emm*-specific PCR products for routine and accurate typing of group A streptococci. *J Clin Microbiol* **34**:953-8.
6. **Beres, S. B., G. L. Sylva, D. E. Sturdevant, C. N. Granville, M. Liu, S. M. Ricklefs, A. R. Whitney, L. D. Parkins, N. P. Hoe, G. J. Adams, D. E. Low, F. R. DeLeo, A. McGeer, and J. M. Musser.** 2004. Genome-wide molecular dissection of serotype M3 group A *Streptococcus* strains causing two epidemics of invasive infections. *Proc Natl Acad Sci U S A* **101**:11833-8.
7. **Bessen, D. E., C. M. Sotir, T. L. Readdy, and S. K. Hollingshead.** 1996. Genetic correlates of throat and skin isolates of group A streptococci. *J Infect Dis* **173**:896-900.

- 1 8. **Bisno, A. L., M. O. Brito, and C. M. Collins.** 2003. Molecular basis of group A  
2 streptococcal virulence. *Lancet Infect Dis* **3**:191-200.
- 3 9. **Carapetis, J. R., A. C. Steer, E. K. Mulholland, and M. Weber.** 2005. The global  
4 burden of group A streptococcal diseases. *Lancet Infect Dis* **5**:685-94.
- 5 10. **Chatellier, S., N. Ihendyane, R. G. Kansal, F. Khambaty, H. Basma, A. Norrby-  
6 Teglund, D. E. Low, A. McGeer, and M. Kotb.** 2000. Genetic relatedness and  
7 superantigen expression in group A streptococcus serotype M1 isolates from patients with  
8 severe and nonsevere invasive diseases. *Infect Immun* **68**:3523-34.
- 9 11. **Cleary, P., D. Johnson, and L. Wannamaker.** 1979. Genetic variation in the M antigen  
10 of group A streptococci: reassortment of type-specific markers and possible antigenic drift.  
11 *J Infect Dis* **140**:747-57.
- 12 12. **Colman, G., A. Tanna, A. Efstratiou, and E. T. Gaworzewska.** 1993. The serotypes of  
13 *Streptococcus pyogenes* present in Britain during 1980-1990 and their association with  
14 disease. *J Med Microbiol* **39**:165-78.
- 15 13. **Creti, R., M. Imperi, L. Baldassarri, M. Pataracchia, S. Recchia, G. Alfarone, and G.  
16 Orefici.** 2007. *emm* Types, virulence factors, and antibiotic resistance of invasive  
17 *Streptococcus pyogenes* isolates from Italy: What has changed in 11 years? *J Clin*  
18 *Microbiol* **45**:2249-56.
- 19 14. **Cunningham, M. W.** 2000. Pathogenesis of group A streptococcal infections. *Clin*  
20 *Microbiol Rev* **13**:470-511.
- 21 15. **Dale, J. B.** 2008. Current status of group A streptococcal vaccine development. *Adv Exp*  
22 *Med Biol* **609**:53-63.

- 1 16. **Dale, J. B., and E. H. Beachey.** 1986. Localization of protective epitopes of the amino  
2 terminus of type 5 streptococcal M protein. *J Exp Med* **163**:1191-202.
- 3 17. **Dale, J. B., T. Penfound, E. Y. Chiang, V. Long, S. T. Shulman, and B. Beall.** 2005.  
4 Multivalent group A streptococcal vaccine elicits bactericidal antibodies against variant M  
5 subtypes. *Clin Diagn Lab Immunol* **12**:833-6.
- 6 18. **Darenberg, J., B. Luca-Harari, A. Jasir, A. Sandgren, H. Pettersson, C. Schalen, M.**  
7 **Norgren, V. Romanus, A. Norrby-Teglund, and B. H. Normark.** 2007. Molecular and  
8 clinical characteristics of invasive group A streptococcal infection in Sweden. *Clin Infect*  
9 *Dis* **45**:450-8.
- 10 19. **Davies, H. D., A. McGeer, B. Schwartz, K. Green, D. Cann, A. E. Simor, and D. E.**  
11 **Low.** 1996. Invasive group A streptococcal infections in Ontario, Canada. Ontario Group A  
12 Streptococcal Study Group. *N Engl J Med* **335**:547-54.
- 13 20. **Efstratiou, A.** 2000. Group A streptococci in the 1990s. *J Antimicrob Chemother* **45**  
14 **Suppl**:3-12.
- 15 21. **Facklam, R.** 2002. What happened to the streptococci: overview of taxonomic and  
16 nomenclature changes. *Clin Microbiol Rev* **15**:613-30.
- 17 22. **Facklam, R., B. Beall, A. Efstratiou, V. Fischetti, D. Johnson, E. Kaplan, P. Kriz, M.**  
18 **Lovgren, D. Martin, B. Schwartz, A. Totolian, D. Bessen, S. Hollingshead, F. Rubin, J.**  
19 **Scott, and G. Tyrrell.** 1999. emm typing and validation of provisional M types for group  
20 A streptococci. *Emerg Infect Dis* **5**:247-53.
- 21 23. **Fischetti, V. A., K. F. Jones, S. K. Hollingshead, and J. R. Scott.** 1988. Structure,  
22 function, and genetics of streptococcal M protein. *Rev Infect Dis* **10 Suppl 2**:S356-9.

24. **Green, N. M., S. Zhang, S. F. Porcella, M. J. Nagiec, K. D. Barbian, S. B. Beres, R. B. LeFebvre, and J. M. Musser.** 2005. Genome sequence of a serotype M28 strain of group A streptococcus: potential new insights into puerperal sepsis and bacterial disease specificity. *J Infect Dis* **192**:760-70.
25. **Johnson, D. R., and E. L. Kaplan.** 1993. A review of the correlation of T-agglutination patterns and M-protein typing and opacity factor production in the identification of group A streptococci. *J Med Microbiol* **38**:311-5.
26. **Johnson, D. R., E. L. Kaplan, A. VanGheem, R. R. Facklam, and B. Beall.** 2006. Characterization of group A streptococci (*Streptococcus pyogenes*): correlation of M-protein and *emm*-gene type with T-protein agglutination pattern and serum opacity factor. *J Med Microbiol* **55**:157-64.
27. **Jones, K. F., B. N. Manjula, K. H. Johnston, S. K. Hollingshead, J. R. Scott, and V. A. Fischetti.** 1985. Location of variable and conserved epitopes among the multiple serotypes of streptococcal M protein. *J Exp Med* **161**:623-8.
28. **Kotb, M., A. Norrby-Teglund, A. McGeer, H. El-Sherbini, M. T. Dorak, A. Khurshid, K. Green, J. Peeples, J. Wade, G. Thomson, B. Schwartz, and D. E. Low.** 2002. An immunogenetic and molecular basis for differences in outcomes of invasive group A streptococcal infections. *Nat Med* **8**:1398-404.
29. **Lamagni, T. L., J. Darenberg, B. Luca-Harari, T. Siljander, A. Efstratiou, B. Henriques-Normark, J. Vuopio-Varkila, A. Bouvet, R. Creti, K. Ekelund, M. Koliou, R. R. Reinert, A. Stathi, L. Strakova, V. Ungureanu, C. Schalen, and A. Jasir.** 2008. Epidemiology of severe *Streptococcus pyogenes* disease in Europe. *J Clin Microbiol* **46**:2359-67.

- 1 30. **Luca-Harari, B., K. Ekelund, M. van der Linden, M. Staum-Kaltoft, A. M.**  
2 **Hammerum, and A. Jasir.** 2008. Clinical and epidemiological aspects of invasive  
3 *Streptococcus pyogenes* infections in Denmark during 2003 and 2004. J Clin Microbiol  
4 **46**:79-86.
- 5 31. **Mencarelli, M., R. Corbisiero, M. G. Padula, I. Galgani, L. Stolzuoli, and C. Cellesi.**  
6 2005. Group A streptococcal infections: trend and strain emm typing in an area of central  
7 Italy, 1985-2002. Epidemiol Infect **133**:1107-11.
- 8 32. **Mihaila-Amrouche, L., A. Bouvet, and J. Loubinoux.** 2004. Clonal spread of *emm* type  
9 28 isolates of *Streptococcus pyogenes* that are multiresistant to antibiotics. J Clin Microbiol  
10 **42**:3844-6.
- 11 33. **Neal, S., B. Beall, K. Ekelund, B. Henriques-Normark, A. Jasir, D. Johnson, E.**  
12 **Kaplan, M. Lovgren, R. R. Reinert, and A. Efstratiou.** 2007. International quality  
13 assurance study for characterization of *Streptococcus pyogenes*. J Clin Microbiol **45**:1175-  
14 9.
- 15 34. **Norrby-Teglund, A., G. T. Nepom, and M. Kotb.** 2002. Differential presentation of  
16 group A streptococcal superantigens by HLA class II DQ and DR alleles. Eur J Immunol  
17 **32**:2570-7.
- 18 35. **O'Loughlin, R. E., A. Roberson, P. R. Cieslak, R. Lynfield, K. Gershman, A. Craig, B.**  
19 **A. Albanese, M. M. Farley, N. L. Barrett, N. L. Spina, B. Beall, L. H. Harrison, A.**  
20 **Reingold, and C. Van Beneden.** 2007. The epidemiology of invasive group A  
21 streptococcal infection and potential vaccine implications: United States, 2000-2004. Clin  
22 Infect Dis **45**:853-62.

- 1 36. **Raymond, J., L. Schlegel, F. Garnier, and A. Bouvet.** 2005. Molecular characterization  
2 of *Streptococcus pyogenes* isolates to investigate an outbreak of puerperal sepsis. Infect  
3 Control Hosp Epidemiol **26**:455-61.
- 4 37. **Spindler, C., J. Hedlund, A. Jasir, B. H. Normark, and A. Ortqvist.** 2008. Effects of a  
5 large-scale introduction of the pneumococcal polysaccharide vaccine among elderly  
6 persons in Stockholm, Sweden. Vaccine **26**:5541-6.
- 7 38. **Stalhammar-Carlemalm, M., T. Areschoug, C. Larsson, and G. Lindahl.** 2000. Cross-  
8 protection between group A and group B streptococci due to cross-reacting surface  
9 proteins. J Infect Dis **182**:142-9.
- 10 39. **Stalhammar-Carlemalm, M., T. Areschoug, C. Larsson, and G. Lindahl.** 1999. The  
11 R28 protein of *Streptococcus pyogenes* is related to several group B streptococcal surface  
12 proteins, confers protective immunity and promotes binding to human epithelial cells. Mol  
13 Microbiol **33**:208-19.
- 14 40. **Stevens, D. L.** 1999. The flesh-eating bacterium: what's next? J Infect Dis **179 Suppl**  
15 **2**:S366-74.
- 16 41. **Stevens, D. L.** 1992. Invasive group A streptococcus infections. Clin Infect Dis **14**:2-11.
- 17 42. **Stevens, D. L.** 1995. Streptococcal toxic-shock syndrome: spectrum of disease,  
18 pathogenesis, and new concepts in treatment. Emerg Infect Dis **1**:69-78.
- 19 43. **Strakova, L., J. Motlova, P. Urbaskova, and P. Krizova.** 2004. [Surveillance of serious  
20 diseases caused by group A streptococci in the Czech Republic in 2003--the Strep-EURO  
21 project]. Epidemiol Mikrobiol Imunol **53**:106-11.

- 1 44. **Svensson, N., S. Oberg, B. Henriques, S. Holm, G. Kallenius, V. Romanus, and J.**  
2 **Giesecke.** 2000. Invasive group A streptococcal infections in Sweden in 1994 and 1995:  
3 epidemiology and clinical spectrum. *Scand J Infect Dis* **32**:609-14.
- 4 45. **The Working, Group on Severe Streptococcal Infections.** 1993. Defining the group A  
5 streptococcal toxic shock syndrome. Rationale and consensus definition. . *JAMA* **269**:390-  
6 1.
- 7 46. **Tyrrell, G. J., M. Lovgren, B. Kress, and K. Grimsrud.** 2005. Invasive group A  
8 streptococcal disease in Alberta, Canada (2000 to 2002). *J Clin Microbiol* **43**:1678-83.
- 9 47. **Wahl, R. U., R. Lutticken, S. Stanzel, M. van der Linden, and R. R. Reinert.** 2007.  
10 Epidemiology of invasive *Streptococcus pyogenes* infections in Germany, 1996-2002:  
11 results from a voluntary laboratory surveillance system. *Clin Microbiol Infect.*
- 12 48. **Vlaminckx, B., W. van Pelt, L. Schouls, A. van Silfhout, C. Elzenaar, E. Mascini, J.**  
13 **Verhoef, and J. Schellekens.** 2004. Epidemiological features of invasive and noninvasive  
14 group A streptococcal disease in the Netherlands, 1992-1996. *Eur J Clin Microbiol Infect*  
15 *Dis* **23**:434-44.
- 16 49. **Zhang, S., N. M. Green, I. Sitkiewicz, R. B. Lefebvre, and J. M. Musser.** 2006.  
17 Identification and characterization of an antigen I/II family protein produced by group A  
18 *Streptococcus*. *Infect Immun* **74**:4200-13.
- 19

**Table 1.** Type distribution among invasive GAS isolates collected within eleven Strep-EURO participating countries during 2003–2004

| <i>emm</i> -type | No. of isolates | No. of T/ <i>emm</i> -type combinations | T-type (no. of isolates)*                                                                                 |
|------------------|-----------------|-----------------------------------------|-----------------------------------------------------------------------------------------------------------|
| 1                | 819             | 5                                       | 1 (802); 3 (1); 3/13/B3264 (1); 4 (2); NT (8); NA (5)                                                     |
| 2                | 35              | 5                                       | 2 (6); 2/28 (13); 12 (1); 28 (1); NT (4)                                                                  |
| 3                | 431             | 8                                       | 3 (175); 3/13 (1); 3/13/B3264, (180); 3/B3264, (5); B3264 (1), 13/B3264 (1),13 (1); NT (52); NA (15)      |
| 4                | 199             | 3                                       | 4 (181); 2/4 (1); NT (13); NA (4)                                                                         |
| 5                | 98              | 5                                       | 5 (67); 5/27 (1); 5/27/44 (4); 11 (1); NT (15); NA (10)                                                   |
| 6                | 91              | 3                                       | 6 (77); 12 (1); NT (9); NA (4)                                                                            |
| 8                | 2               | 2                                       | 8 (1); NT (1)                                                                                             |
| 9                | 21              | 3                                       | 9 (17); 5/12 (1); NT (3)                                                                                  |
| 11               | 48              | 3                                       | 11 (33); 8/11 (1); NT (11); NA (3)                                                                        |
| 12               | 227             | 3                                       | 12 (216); 11/12/27 (1); NT (9); NA (1)                                                                    |
| 13               | 1               | 1                                       | 13 (1)                                                                                                    |
| 15               | 1               | 1                                       | 15/17/19/23/47 (1)                                                                                        |
| 18               | 66              | 3                                       | 8/25/imp19 (2); 3/B3264 (1); NT (23); NA (40)                                                             |
| 22               | 72              | 6                                       | 12 (57); 12/3/13/B3264 (1); 3/13 (1); 4 (1); 5/12 (1); NT (7); NA (4)                                     |
| 24               | 1               | 1                                       | NT                                                                                                        |
| 25               | 8               | 3                                       | 25 (2); 14 (2); 1 (4)                                                                                     |
| 26               | 3               | 1                                       | NT (3)                                                                                                    |
| 27G              | 10              | 4                                       | 5/27/44 (4); 5/27 (1); 5 (3); NT (2)                                                                      |
| 28               | 505             | 9                                       | 28 (458); 2/28 (1); 28/11 (1); 28/11/12 (2); 3/13/B3264 (1), 3/B3264 (2); 12 (2); 14 (1); NT (31); NA (6) |
| 29               | 8               | 1                                       | 4 (1); NT (6); NA (1)                                                                                     |
| 30               | 1               | 1                                       | NT                                                                                                        |
| 32               | 5               | 1                                       | 23 (5)                                                                                                    |
| 33               | 35              | 4                                       | 3/13/B3264 (20), 3 (4), 13 (6); NT (5)                                                                    |
| 36               | 1               | 1                                       | NA                                                                                                        |
| 41               | 4               | 3                                       | 3/13/B3264 (2); 6 (1); NT (1)                                                                             |
| 43               | 51              | 5                                       | 3/13/B3264 (17); 3 (10); 28 (1); 15/17/19/23/47 (1); NT (9); NA (13)                                      |
| 44/61            | 30              | 7                                       | 5 (17); 5/27/44 (2), 27/44 (1); 12/27/44 (1); 11 (3); 8/11 (1); NT (2); NA (3)                            |
| 48               | 1               | 1                                       | 4/28                                                                                                      |
| 49               | 16              | 4                                       | 14 (12); 3/B3264 (1); 8/25/imp19 (1); NT (1); NA (1)                                                      |
| 50/62            | 4               | 2                                       | 12 (3); NT (1)                                                                                            |
| 52               | 3               | 2                                       | 3/13/B3264 (1); NT (2)                                                                                    |
| 53               | 25              | 7                                       | 3/13/B3264 (5), 3 (2); 3/13/B3264/28/8 (1); 8/25 (1), 8/25/imp19 (1); 28 (3); NT (12)                     |
| 55               | 1               | 1                                       | 8/25/imp19 (1)                                                                                            |
| 57               | 1               | 1                                       | NT                                                                                                        |
| 58               | 22              | 5                                       | 8/25/imp19 (11), 8/25 (1), 8 (2); 2 (1); NT (4); NA (3)                                                   |
| 59               | 5               | 4                                       | 11 (2); 8/25/imp19 (1); 14/25 (1); NT (1)                                                                 |
| 60               | 5               | 2                                       | 4 (4); B3264 (1)                                                                                          |
| 63               | 3               | 1                                       | 4 (3)                                                                                                     |
| 64               | 5               | 2                                       | 3 (1); NT (4)                                                                                             |
| 65               | 3               | 2                                       | 3/B3264 (2); 8/25/imp19 (1)                                                                               |
| 66               | 8               | 2                                       | 12 (5); NT (3)                                                                                            |
| 67               | 1               | 1                                       | 3 (1)                                                                                                     |
| 68               | 16              | 6                                       | 3/13/B3264 (11), 3/B3264 (1), B3264 (1); 8 (1); 12 (1); 8/25/imp19 (1)                                    |
| 69               | 2               | 2                                       | 3/13/B3264 (1), 3/B3264 (1)                                                                               |
| 70               | 1               | 1                                       | 5/11/27 (1)                                                                                               |
| 71               | 1               | 1                                       | 8/25/imp19 (1)                                                                                            |

|           |     |    |                                                                                                                                       |
|-----------|-----|----|---------------------------------------------------------------------------------------------------------------------------------------|
| 73        | 37  | 6  | 3/13/B3264 (18), 13/B3264 (1), B3264 (5), 3 (2), 13 (8); NT (2); NA (1)                                                               |
| 74        | 3   | 1  | NT (3)                                                                                                                                |
| 75        | 64  | 6  | 8/25/Imp19 (33), 8/25 (10), 8 (1), 25 (15); <u>12 (1)</u> ; NT (3); NA (1)                                                            |
| 76        | 15  | 4  | 12 (8); 22 (2); <u>23 (1)</u> ; 8 (1); NA (3)                                                                                         |
| 77        | 97  | 7  | 11 (4); 13 (29), 3/13/B3264 (4), B3264 (2); 13/28 (20); 28 (30);; NT (5); NA (3)                                                      |
| 78        | 39  | 5  | 11 (32); 11/12/27 (1); 3/B3264 (1); 5/27/44 (1); NT (2); NA (2)                                                                       |
| 79        | 21  | 8  | <u>1 (1)</u> ; 11 (3); 12 (1); <u>13 (1)</u> ; 8 (5); 8/25/Imp19 (1); <u>B3264 (1)</u> ; NT (6); NA (2)                               |
| 80        | 5   | 3  | 3/13/B32647 (2); 14/8/25 (1); NT (2)                                                                                                  |
| 81        | 143 | 11 | 12 (20); 3/13/B3264 (44), 13/B3264 (1), B3264 (24), 3/13 (1), 3 (4); 5/17/19/23/47 (2); 3/12/B3264 (2); 4 (1); 8 (4); NT (32); NA (8) |
| 82        | 58  | 5  | 12 (1); 5 (42), 5/27 (1), 5/27/44 (1)                                                                                                 |
| 83        | 150 | 7  | 13 (7), 3 (20), 3/13 (1), 3/13/B3264 (78), <u>8 (1)</u> ; 8/25 (2); NT (22); NA (19)                                                  |
| 84        | 3   | 2  | 8/25/Imp19 (1); NT (2)                                                                                                                |
| 85        | 10  | 5  | <u>12 (1)</u> ; 3 (5), 3/13 (1); <u>3/9 (2)</u> ; NT (1)                                                                              |
| 87        | 256 | 6  | 11 (1); 28 (242); 3/13/B3264 (1); 4 (1); 8/25/Imp19 (1); NT (4); NA (6)                                                               |
| 88        | 11  | 3  | 8/25/Imp19 (6); 8 (4); NT (1)                                                                                                         |
| 89        | 343 | 9  | 11 (2); 13 (3); 28 (1); 3 (15), 3/13/B3264 (262), 3/B3264 (12), B3264 (21); <u>8 (1)</u> ; NT (16); NA (10)                           |
| 90        | 4   | 2  | 3/13/B3264 (2); NT (1); NA (1)                                                                                                        |
| 91        | 12  | 5  | 3 (1), 13 (1); 25/Imp19 (1); 5 (2); NT (4); NA (3)                                                                                    |
| 92        | 4   | 3  | 8/25/Imp19 (2); <u>Imp19 (2)</u> ; NT (1)                                                                                             |
| 93        | 6   | 3  | <u>14 (2)</u> ; 3/13/B3264 (1); NT (3)                                                                                                |
| 94        | 11  | 4  | 3/13/B3264 (9), 3/B3264 (1); NT (1)                                                                                                   |
| 95        | 10  | 1  | 6 (3); <u>8/25 (1)</u> , 8/25/Imp19 (1); NT (1); NA (4)                                                                               |
| 96        | 2   | 1  | <u>13 (2)</u>                                                                                                                         |
| 97        | 1   | 3  | 14                                                                                                                                    |
| 100       | 3   | 5  | 3/13 (1); 8/25/Imp19 (1); NT (1)                                                                                                      |
| 101       | 17  | 6  | 13 (1); 28 (1); 3 (3); 3/13/B3264 (6); NT (3); NA (3)                                                                                 |
| 102       | 13  | 1  | 12 (1); 13/B3264 (1), 3 (2), 3/13/B3264 (7); <u>4 (1)</u> ; NT (1)                                                                    |
| 103       | 3   | 3  | NT (2); NA (1)                                                                                                                        |
| 104       | 5   | 2  | 11 (1); <u>8/11 (1)</u> ; NT (3)                                                                                                      |
| 106       | 2   | 3  | 5 (1); NT (1)                                                                                                                         |
| 108       | 9   | 1  | 5/12/27 (1); 6 (3); NT (5)                                                                                                            |
| 109       | 1   | 3  | 6 (1)                                                                                                                                 |
| 110       | 6   | 1  | 12 (2); <u>8 (1)</u> ; NT (2); NA (1)                                                                                                 |
| 111       | 2   | 2  | NT (1); NA (1)                                                                                                                        |
| 112       | 4   | 1  | 8 (1); <u>13 (1)</u> ; NA (2)                                                                                                         |
| 113       | 1   | 1  | 12                                                                                                                                    |
| 115       | 1   | 2  | NT                                                                                                                                    |
| 117       | 5   | 8  | <u>28 (1)</u> ; NT (4)                                                                                                                |
| 118       | 34  | 1  | 3/13/B3264 (23), 3/13 (3), B3264 (2), 3 (1), 13 (1); 6 (1); <u>11 (1)</u> ; NT (2)                                                    |
| 119       | 2   | 1  | NT (2)                                                                                                                                |
| 120       | 2   | 2  | 3 (2)                                                                                                                                 |
| 122       | 2   | 1  | 3/13/B3264 (1); NT (1)                                                                                                                |
| 124       | 2   | 1  | NT (2)                                                                                                                                |
| emmst369  | 1   | 1  | 14                                                                                                                                    |
| emmst1389 | 2   | 1  | <u>3/13/B3264 (2)</u>                                                                                                                 |
| emmst2037 | 1   | 1  | <u>8/25/Imp19</u>                                                                                                                     |
| emmst2147 | 1   | 1  | <u>5/27/44</u>                                                                                                                        |
| emmst2460 | 1   | 1  | 3/13/B3264                                                                                                                            |
| emmst2904 | 2   | 1  | NT                                                                                                                                    |
| emmst3757 | 2   | 1  | NT                                                                                                                                    |
| emmst6735 | 1   | 1  | <u>5/27/44</u>                                                                                                                        |

|                     |      |  |     |                                                                                                             |
|---------------------|------|--|-----|-------------------------------------------------------------------------------------------------------------|
| <i>emmst</i> 11014  | 2    |  | 1   | 4 (1); NA (1)                                                                                               |
| <i>emmst</i> D633   | 1    |  | 1   | 3/13/B3264                                                                                                  |
| <i>emmst</i> NS1033 | 20   |  | 3   | 28 (6); 8/25/ <i>Imp</i> 19 (1); NT (2); NA (11)                                                            |
| <i>emmst</i> G6     | 1    |  | 1   | 8/25/ <i>Imp</i> 19                                                                                         |
| <i>emmst</i> G1750  | 1    |  | 1   | NT                                                                                                          |
| <i>emmst</i> G62647 | 1    |  | 1   | 12                                                                                                          |
| <i>emmst</i> 221    | 1    |  | 1   | NT                                                                                                          |
| <i>emmst</i> 4986   | 1    |  | 1   | 4                                                                                                           |
| NA                  | 1168 |  |     | 3/8/B3264 (2), 3/B3264 (2), B3264 (5); 3/9/13 (1); 5 (1); 8 (2); 8/25/ <i>Imp</i> 19 (2); NT (6); NA (1147) |
| Total               | 5521 |  | 314 | 4171                                                                                                        |

**Note.** Underlined are uncommon T/*emm*-type combinations, according to the CDC homepage and previous publication (26).

T-types belonging to the same pool are separated by comma (,) and distinct patterns by semicolon (;)

NA, not available; NT, not typable; *emmst*, *emm* sequence type not yet assigned (as described at the CDC homepage).

\* The number of isolates of each specific T/*emm*-type combination is described within brackets.

**Table 2.** Disease manifestations and case fatality rates (CFR) for 15 most prevalent *emm*-types.

|                                   | emm-<br>typed | emm-type no./% (CFR %) |                |               |               |               |               |               |               |               |               |               |               |               |              |               |                |
|-----------------------------------|---------------|------------------------|----------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|--------------|---------------|----------------|
|                                   |               | 1                      | 28             | 3             | 89            | 87            | 12            | 4             | 83            | 81            | 5             | 77            | 6             | 18            | 75           | 43            | Other          |
| Total with clin. info.<br>CFR (%) | 3458<br>(19)  | 694<br>(29)            | 356<br>(14)    | 333<br>(36)   | 276<br>(13)   | 217<br>(19)   | 191<br>(17)   | 161<br>(10)   | 113<br>(8)    | 121<br>(10)   | 74<br>(30)    | 77<br>(20)    | 75<br>(18)    | 51<br>(21)    | 43<br>(9)    | 38<br>(21)    | 584<br>(11)    |
| No focus                          | 859<br>(15)   | 117/17<br>(16)         | 109/31<br>(15) | 70/21<br>(32) | 88/32<br>(14) | 57/26<br>(17) | 42/22<br>(13) | 40/25<br>(9)  | 38/34<br>(19) | 55/45<br>(2)  | 16/22<br>(40) | 36/47<br>(12) | 16/21<br>(13) | 8/16<br>(0)   | 13/30<br>(8) | 7/18<br>(33)  | 147/25<br>(10) |
| STSS                              | 476<br>(44)   | 174/25<br>(47)         | 32/9<br>(40)   | 83/25<br>(49) | 18/7<br>(47)  | 12/6<br>(42)  | 27/14<br>(50) | 20/12<br>(33) | 6/5<br>(33)   | 14/12<br>(30) | 4/5<br>(0)    | 7/9<br>(57)   | 9/12<br>(60)  | 8/16<br>(63)  | 3/7<br>(0)   | 1/≤1<br>(0)   | 58/10<br>(3)   |
| NF                                | 296<br>(31)   | 92/13<br>(36)          | 18/5<br>(13)   | 40/12<br>(51) | 16/6<br>(36)  | 5/2<br>(50)   | 13/7<br>(39)  | 19/12<br>(13) | 5/4<br>(0)    | 19/16<br>(21) | 7/9<br>(43)   | 8/10<br>(25)  | 4/5<br>(0)    | 5/10<br>(75)  | 2/5<br>(0)   | 1/≤1<br>(0)   | 42/7<br>(13)   |
| Cellulitis                        | 865<br>(18)   | 177/26<br>(25)         | 74/21<br>(18)  | 95/29<br>(33) | 72/26<br>(17) | 71/33<br>(14) | 49/26<br>(17) | 36/22<br>(9)  | 34/30<br>(0)  | 20/17<br>(6)  | 21/28<br>(14) | 14/18<br>(33) | 15/20<br>(0)  | 12/24<br>(20) | 8/19<br>(14) | 16/42<br>(18) | 151/26<br>(10) |
| Arthritis                         | 302<br>(10)   | 65/9<br>(17)           | 18/5<br>(18)   | 38/11<br>(17) | 27/10<br>(8)  | 12/6<br>(0)   | 22/12<br>(0)  | 16/10<br>(0)  | 10/9<br>(0)   | 11/9<br>(0)   | 4/5<br>(0)    | 9/12<br>(22)  | 4/5<br>(0)    | 3/9<br>(0)    | 6/14<br>(0)  | 2/5<br>(0)    | 55/9<br>(9)    |
| Puerperal sepsis                  | 96<br>(4)     | 15/2<br>(18)           | 30/8<br>(0)    | 7/2<br>(17)   | 11/4<br>(0)   | 9/4<br>(0)    | 4/2<br>(0)    | 3/2<br>(0)    | -/-<br>(-)    | -/-<br>(-)    | -/-<br>(-)    | 1/1<br>(-)    | 2/3<br>(-)    | 1/1<br>(-)    | -/-<br>(-)   | -/-<br>(-)    | 13/2<br>(-)    |
| Meningitis                        | 59<br>(24)    | 16/2<br>(36)           | 8/2<br>(0)     | 11/19<br>(56) | 2/1<br>(0)    | 2/1<br>(0)    | 3/2<br>(0)    | 3/1<br>(33)   | -/-<br>(-)    | -/-<br>(-)    | 1/1<br>(0)    | 1/1<br>(0)    | 2/2<br>(0)    | -/-<br>(-)    | -/-<br>(-)   | -/-<br>(-)    | 10/2<br>(22)   |
| Other                             | 835<br>(26)   | 189/27<br>(39)         | 86/24<br>(22)  | 71/21<br>(53) | 57/21<br>(4)  | 48/22<br>(28) | 51/27<br>(20) | 33/20<br>(8)  | 23/20<br>(5)  | 14/12<br>(31) | 24/32<br>(36) | 9/12<br>(60)  | 32/43<br>(35) | 20/39<br>(24) | 14/32<br>(9) | 11/29<br>(22) | 153/26<br>(12) |

**Note.** STSS, streptococcal toxic shock syndrome; NF, necrotizing fasciitis.

**Table 3.** Risk factor data of cases caused by 15 most prevalent *emm*-types.

|                                    |      | <i>emm</i> -type (no/%) |          |         |         |         |         |         |         |         |        |        |        |        |        |        |
|------------------------------------|------|-------------------------|----------|---------|---------|---------|---------|---------|---------|---------|--------|--------|--------|--------|--------|--------|
| Total                              |      | 1                       | 28       | 3       | 89      | 87      | 12      | 4       | 83      | 81      | 5      | 77     | 6      | 18     | 75     | 43     |
| Cases with risk factor information | 2796 | 525/18.8                | 287/10.3 | 278/9.9 | 238/8.5 | 190/6.8 | 154/5.5 | 137/4.9 | 109/3.9 | 106/3.8 | 61/2.2 | 65/2.3 | 64/2.3 | 40/1.4 | 27/1.0 | 37/1.3 |
| Diabetes                           | 232  | 38/16.4                 | 27/11.6  | 31/13.4 | 21/9.1  | 14/6.0  | 16/6.9  | 10/4.3  | 6/2.6   | 23/9.9  | 5/2.2  | 8/3.4  | 4/1.7  | 3/1.3  | 0/0    | 5/2.2  |
| IDU                                | 359  | 10/2.8                  | 4/1.1    | 7/1.9   | 24/6.7  | 37/10.3 | 8/2.2   | 5/1.4   | 74/20.6 | 20/5.6  | 2/0.6  | 2/0.6  | 1/0.3  | 1/0.3  | 3/0.8  | 20/5.6 |
| Chicken pox                        | 72   | 29/40.3                 | 3/4.2    | 6/8.3   | 1/1.4   | 4/5.6   | 13/18.1 | 4/5.6   | 0/0     | 0/0     | 0/0    | 1/1.4  | 3/4.2  | 0/0    | 1/1.4  | 0/0    |
| Immunocomprom.                     | 478  | 92/19.2                 | 45/9.4   | 50/10.5 | 44/9.2  | 31/6.5  | 31/6.5  | 20/4.2  | 12/2.5  | 20/4.2  | 18/3.8 | 16/3.3 | 10/2.1 | 10/2.1 | 7/1.5  | 6/1.3  |
| Skin lesions                       | 593  | 128/21.6                | 54/9.1   | 57/9.6  | 46/7.8  | 49/8.3  | 35/5.9  | 25/4.2  | 24/4.0  | 17/2.9  | 11/1.9 | 16/2.7 | 6/1.0  | 5/0.8  | 5/0.8  | 10/1.7 |
| Surgery                            | 151  | 24/15.9                 | 21/13.9  | 10/6.6  | 12/7.9  | 14/9.3  | 10/6.6  | 9/6.0   | 2/1.3   | 4/2.6   | 6/4.0  | 2/1.3  | 6/4.0  | 5/3.3  | 1/0.7  | 3/2.0  |
| HAI*                               | 242  | 37/15.3                 | 39/16.1  | 16/6.6  | 22/9.1  | 18/7.4  | 17/7.0  | 15/6.2  | 2/0.8   | 5/2.1   | 9/3.7  | 2/0.8  | 7/2.9  | 5/2.1  | 3/1.2  | 4/1.7  |
| Other                              | 782  | 148/18.9                | 112/14.3 | 70/9.0  | 75/9.6  | 24/3.1  | 39/5.0  | 35/4.5  | 11/1.4  | 53/6.8  | 11/1.4 | 35/4.5 | 23/2.9 | 8/1.0  | 11/1.4 | 1/0.1  |
| None reported                      | 613  | 137/22.3                | 73/11.9  | 85/13.9 | 54/8.8  | 44/7.2  | 31/5.1  | 49/8.0  | 5/0.8   | 9/1.5   | 17/2.8 | 8/1.3  | 16/2.6 | 14/2.3 | 3/0.5  | 2/0.3  |

**Note.** HAI, health care associated infections; IDU, injecting drug users.

**Table 4.** Presence of superantigens as related to clinical presentation and *emm*-type

|                   | speA    |    | speC    |    | speF    |    | speG    |     | speH   |    | speI  |   | speJ   |    | smeZ    |    | ssa     |    |
|-------------------|---------|----|---------|----|---------|----|---------|-----|--------|----|-------|---|--------|----|---------|----|---------|----|
|                   | a       | %  | a       | %  | a       | %  | a       | %   | a      | %  | a     | % | a      | %  | a       | %  | a       | %  |
| No focus          | 87/475  | 18 | 295/475 | 62 | 415/422 | 98 | 402/436 | 92  | 61/437 | 14 | 0/76  | 0 | 74/404 | 18 | 316/403 | 78 | 72/443  | 16 |
| STSS              | 187/363 | 52 | 138/363 | 38 | 240/265 | 91 | 218/238 | 92  | 18/238 | 8  | 0/74  | 0 | 15/123 | 12 | 59/133  | 52 | 66/294  | 22 |
| NF                | 93/224  | 42 | 83/224  | 37 | 134/145 | 92 | 130/146 | 89  | 9/145  | 6  | 0/47  | 0 | 15/99  | 15 | 56/105  | 53 | 37/172  | 22 |
| Cellulitis        | 128/371 | 35 | 168/371 | 45 | 179/205 | 87 | 194/198 | 98  | 13/198 | 7  | 3/151 | 2 | 9/153  | 6  | 17/162  | 11 | 38/225  | 17 |
| Arthritis         | 57/175  | 33 | 91/175  | 52 | 122/130 | 94 | 122/137 | 89  | 19/137 | 14 | 0/39  | 0 | 15/112 | 13 | 75/114  | 66 | 28/142  | 20 |
| Puerperal sepsis  | 15/61   | 25 | 41/61   | 67 | 48/54   | 89 | 50/52   | 96  | 3/52   | 8  | 1/18  | 6 | 3/31   | 10 | 15/33   | 46 | 8/58    | 14 |
| Meningitis        | 23/43   | 53 | 20/43   | 47 | 31/33   | 94 | 24/25   | 96  | 2/25   | 8  | 1/15  | 7 | 3/15   | 20 | 5/16    | 31 | 12/40   | 30 |
| Other clin. pres. | 135/347 | 39 | 174/347 | 50 | 244/258 | 95 | 184/191 | 96  | 15/191 | 8  | 2/146 | 1 | 10/149 | 7  | 25/163  | 15 | 28/281  | 10 |
| M/ <i>emm</i> 1   | 400/456 | 88 | 88/456  | 19 | 285/329 | 87 | 280/289 | 97  | 5/289  | 2  | 1/143 | 1 | 35/203 | 17 | 67/208  | 32 | 79/362  | 22 |
| M/ <i>emm</i> 28  | 31/387  | 8  | 343/387 | 89 | 343/355 | 97 | 311/322 | 97  | 15/322 | 5  | 1/199 | 1 | 72/289 | 25 | 106/291 | 36 | 114/370 | 31 |
| M/ <i>emm</i> 3   | 153/182 | 84 | 34/182  | 19 | 123/128 | 96 | 107/109 | 98  | 7/109  | 6  | 1/66  | 2 | 1/86   | 1  | 25/86   | 29 | 114/133 | 86 |
| M/ <i>emm</i> 89  | 9/169   | 5  | 88/169  | 52 | 140/146 | 96 | 136/138 | 99  | 5/138  | 4  | 0/47  | 0 | 2/131  | 2  | 86/131  | 66 | 24/148  | 16 |
| M/ <i>emm</i> 87  | 4/52    | 8  | 38/52   | 73 | 20/34   | 59 | 34/34   | 100 | 1/34   | 3  | 0/24  | 0 | 8/33   | 24 | 10/33   | 30 | 18/35   | 51 |
| M/ <i>emm</i> 12  | 11/129  | 9  | 83/129  | 64 | 89/90   | 99 | 82/83   | 99  | 54/83  | 65 | 2/35  | 6 | 4/72   | 6  | 47/75   | 63 | 25/95   | 26 |
| M/ <i>emm</i> 4   | 8/120   | 7  | 111/120 | 93 | 82/92   | 89 | 52/88   | 59  | 3/88   | 3  | 0/26  | 0 | 3/66   | 5  | 44/68   | 65 | 81/96   | 84 |
| M/ <i>emm</i> 83  | 7/34    | 21 | 16/34   | 47 | 25/27   | 93 | 26/26   | 100 | 0/26   | 0  | 0/17  | 0 | 0/25   | 0  | 9/26    | 35 | 4/27    | 15 |
| M/ <i>emm</i> 81  | 2/112   | 2  | 49/112  | 44 | 91/95   | 96 | 102/105 | 97  | 20/104 | 19 | 0/30  | 0 | 6/99   | 6  | 48/102  | 47 | 6/105   | 6  |
| M/ <i>emm</i> 5   | 4/23    | 17 | 21/23   | 91 | 6/9     | 67 | 8/8     | 100 | 0/8    | 0  | 0/6   | 0 | 0/8    | 0  | 2/8     | 25 | 0/10    | 0  |
| M/ <i>emm</i> 77  | 3/76    | 4  | 57/76   | 75 | 60/62   | 97 | 27/59   | 46  | 4/60   | 7  | 0/20  | 0 | 0/55   | 0  | 35/55   | 64 | 13/67   | 19 |
| M/ <i>emm</i> 6   | 8/44    | 18 | 39/44   | 89 | 28/34   | 82 | 27/28   | 96  | 2/28   | 7  | 1/11  | 9 | 4/20   | 20 | 10/20   | 50 | 8/34    | 24 |
| M/ <i>emm</i> 18  | 11/22   | 50 | 20/22   | 91 | 5/11    | 46 | 11/13   | 85  | 1/13   | 8  | 0/7   | 0 | 0/11   | 0  | 3/11    | 27 | 1/14    | 7  |
| M/ <i>emm</i> 75  | 3/43    | 7  | 23/43   | 53 | 30/37   | 81 | 28/34   | 82  | 0/34   | 0  | 0/19  | 0 | 0/24   | 0  | 8/26    | 31 | 7/38    | 18 |

**NOTE.** STSS, streptococcal toxic shock syndrome; NF, necrotizing fasciitis.

a – positive isolates/no. of tested isolates

Fig. 1A





Fig. 1B



**Fig. 2**




Fig. 3A





Fig. 3B



**Figure 1.** *emm*-type distribution among the ten participating Strep-EURO countries in years 2003-2004. Numbers of isolates presented for each country are indicated beneath each bar. A) The distribution of the ten most prevalent *emm*-types within each country and overall, with and without UK isolates. The 10 overall prevalent types are indicated in the figure, within the bar for “Total”, and other types indicated for each country. B) The distribution of the ten overall most prevalent *emm*- types. Types of more than 5% of all types within a country are indicated by numbers in the figure. Abbreviations for countries: CZE, the Czech Republic; DNK, Denmark; FIN, Finland; FRA, France; GER, Germany; GRC, Greece; ITA, Italy; ROU, Roumania; SWE, Sweden; UK, the United Kingdom.

**Figure 2.** Overall seasonal fluctuation of the ten most prevalent *emm*-types among the Strep-EURO participating countries. Percentages are calculated by number of each major type divided to total number of isolates per month.

**Figure 3.** Distribution of *emm*-types among Strep-EURO invasive GAS cases, with special regard to coverage by a 26-valent candidate vaccine. Types hypothetically covered or not by the vaccine candidate are indicated.

A) Prevalence of 30 most common *emm*-types. Among these, 16 (accounting for 69% of reported cases) are included in the 26-valent vaccine (since subtypes were not assessed, vaccine subtype *emm*1.2 is not considered in the present discussion). Vaccine types *emm*14, *emm*19, and *emm*114 were not encountered. Other 1= other types included in the vaccine (6 *emm*-types); Other 2= remaining types not covered by the vaccine (70 different *emm*-types)

B) Country-specific *emm*-type proportions based on potential coverage by the 26-valent vaccine. Numbers of *emm*-types potentially covered or not are indicated below the graph for each country.

- 1 Abbreviations for countries: CZE, the Check Republic; DNK, Denmark; FIN, Finland; FRA,
- 2 France; GER, Germay; GRC, Greece; ITA, Italy; ROU, Roumania; SWE; Sweden; UK, the
- 3 United Kingdom.