



LUND UNIVERSITY

Waterlogging after clear-cutting turns boreal forest soils into sources of methane

Vestin, Patrik; Mölder, Meelis; Hellström, Margareta; Sundqvist, Elin; Klemedtsson, Leif; Lindroth, Anders

Published in:

[Publication information missing]

2011

[Link to publication](#)

Citation for published version (APA):

Vestin, P., Mölder, M., Hellström, M., Sundqvist, E., Klemedtsson, L., & Lindroth, A. (2011). Waterlogging after clear-cutting turns boreal forest soils into sources of methane. *[Publication information missing]*. http://www.ileaps.org/sci_conf_book/pdf/20110415140706_Patrik_Vestin_Waterlogging_after_clear-felling_turns_boreal_forest_soils_into_sources_of_methane_iLEAPS_2011.pdf

Total number of authors:

6

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

WATERLOGGING AFTER CLEAR-CUTTING TURNS BOREAL FOREST SOILS INTO SOURCES OF METHANE

Patrik Vestin (1), Anders Lindroth (1), Meelis Mölder (1), Margareta Hellström (1), Elin Sundqvist (1), Leif Klemedtsson (2)

(1) Department of Earth and Ecosystem Sciences, Lund University, Sweden

(2) Department of Plant and Environmental Sciences, Gothenburg University, Sweden

Patrik.Vestin@nateko.lu.se

Clear-cutting and subsequent site preparation is common forest management practice in Sweden. According to the Swedish National Forest Inventory [1], final fellings were carried out on 170.000 ha during 2009. The net effects of final fellings on greenhouse gas fluxes are not well understood.

Increased substrate availability for decomposers following harvest may result in higher carbon dioxide (CO₂) emissions from soils and in increased nitrogen mineralization (e.g. [2]). This may be further enhanced by increased soil temperatures after site preparation. In addition, removal of trees causes reduced evapotranspiration and subsequently, a raised ground water table. Boreal forest soils are normally sinks of atmospheric methane (CH₄) (e.g. [3]; [4]), with soil water content, nitrogen availability [5] and soil temperature (see review [6] and references therein) as main factors controlling CH₄ exchange between the biosphere and the atmosphere.

The net effects of clear-cutting on CH₄, CO₂ and H₂O fluxes were studied at Norunda forest in central Sweden. Micrometeorological measurements (i.e. flux-gradient and eddy covariance) allowed for quantification of CO₂, CH₄ and H₂O fluxes at two plots at a new clear-cut during the growing season of 2010. Soil chamber measurements (CO₂, CH₄ and H₂O) were carried out in the adjacent forest stand during the main growing season and at the clear-cut during October-November 2010.

The clear-cut became waterlogged after harvest and preliminary results indicate a switch from a weak CH₄ sink to a significant CH₄ source at both plots. Daily average fluxes during the period 20 May through 30 November were in the order of -6.3 – 142.5 $\mu\text{mol m}^{-2}\text{hr}^{-1}$ (fig.1) with mean values of 39.0 $\mu\text{mol m}^{-2}\text{hr}^{-1}$ (plot 1) and 16.3 $\mu\text{mol m}^{-2}\text{hr}^{-1}$ (plot 2). The preliminary results also indicate increased soil temperatures at disturbed microsites and enhanced CO₂ emissions. Data will be further analyzed and presented at the conference.




Fig 1. Daily average CH₄ fluxes at plot 1 (circles) and plot 2 (triangles) measured with the flux-gradient technique at the Norunda clear-cut during the period 20 May through 30 November 2010. Both plots are, on average, significant CH₄ sources. The decreasing trend from mid-June through late July coincides with a period with low amounts of precipitation and decreasing soil water content.

References

- [1] Swedish Forest Agency, 2010. *Swedish Statistical Yearbook of Forestry 2010*. Swedish Forest Agency, Huskvarna, Sweden, 386 pp.
- [2] Chen, H., Harmon, M.E., Sexton, J. et al., 2002. Fine-root decomposition and N dynamics in coniferous forests of the Pacific Northwest, U.S.A. *Canadian Journal of Forest Research*, 32, 320-331.
- [3] Crill, P.M., 1991. Seasonal patterns of methane uptake and carbon dioxide release by a temperate woodland soil. *Global Biogeochemical Cycles*, 5, 319-334.
- [4] Kähkönen, M.A., Wittmann, C, Ilvesniemi, H, Westman, C.J., Salkinoja-Salonen, M.S., 2002. Mineralization of detritus and oxidation of methane in acid boreal coniferous forest soils: seasonal and vertical distribution and effects of clear-cut. *Soil Biology & Biochemistry*, 34, 1191-1200.
- [5] Steudler, P.A., Bowden, R.D., Melillo, J.M., Aber, J.D. 1989. Influence of nitrogen fertilization on methane uptake in temperate forest soils. *Nature*, 341, 314-316.
- [6] Le Mer, J, Roger, P. 2001. Production, oxidation, emission and consumption of methane by soils: A review. *European Journal of Soil Biology*, 37, 25–50.