



LUND UNIVERSITY

The Arabidopsis 14-3-3 family -target protein specificity and expression of isoforms

Arkell, Annika

2010

[Link to publication](#)

Citation for published version (APA):

Arkell, A. (2010). *The Arabidopsis 14-3-3 family -target protein specificity and expression of isoforms*. [Licentiate Thesis, Biochemistry and Structural Biology]. Lund University.

Total number of authors:

1

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

The Arabidopsis 14-3-3 family

-target protein specificity and expression of isoforms

Annika Arkell

Department of Biochemistry and Structural Biology
Lund University, Sweden
2010

Akademisk avhandling för avläggande av filosofie licentiatexamen vid Naturvetenskapliga fakulteten vid Lunds universitet. Avhandlingen kommer att försvaras vid ett officiellt seminarie på Kemicentrum, Getingevägen 60, Lund, Hörsal E, tisdagen den 19 oktober 2010, kl 10.15.



LUND
UNIVERSITY

The Arabidopsis 14-3-3 family – target protein specificity and expression of isoforms
© 2010 Annika Arkell and the respective publishers

DEPARTMENT OF BIOCHEMISTRY AND STRUCTURAL BIOLOGY
LUND UNIVERSITY
P.O. BOX 124
SE-221 00 LUND
SWEDEN

ISBN 978-91-7422-254-8

Printed by Media-Tryck, Lund, Sweden

Organization LUND UNIVERSITY Department of Biochemistry and structural biology P.O. Box 124 SE-221 00 Lund	Document name	
	Date of issue October 2010	
	CODEN:	
Author(s) Annika Arkell	Sponsoring organization	
Title and subtitle The Arabidopsis 14-3-3 family – target protein specificity and expression of isoforms		
Abstract 14-3-3 proteins comprise a family of highly conserved proteins. 14-3-3 proteins have been found in all organisms examined except for members of the prokaryotic kingdom. 14-3-3s are involved in numerous processes in the cell and they typically bind to phosphorylated motifs in other proteins and regulate their activities. In plants, 14-3-3 proteins are recognized as key regulators of primary metabolism and membrane transport. In Arabidopsis, there are 15 genes coding for 14-3-3s and hence several 14-3-3 isoforms may be present simultaneously in the plant. The aim of my work has been to understand why there are so many 14-3-3 isoforms. To investigate if there is specificity in 14-3-3/target protein interaction, the H ⁺ -ATPase/14-3-3 interaction was used as a model system. The study indicated some specificity but also a wide redundancy. To further analyse the question of specificity at different levels promoter:GUS fusions were utilized. The results clearly indicate a developmental, cell-, tissue- and organ-specific expression for all of the 14-3-3 isoforms in Arabidopsis. There is not a single case where the promoter of one isoform shows an expression that is identical to the expression of another isoform.		
Key words 14-3-3, H ⁺ -ATPase, Arabidopsis, GUS, protein-protein interaction, expression		
Classification system and/or index terms (if any)		
Supplementary bibliographical information		Language English
ISSN and key title		ISBN 978-91-7422-254-8
Recipient's notes	Number of pages 61	Price
	Security classification	

Distribution by (name and address) Department of Biochemistry and Structural Biology
P.O. Box 124, SE-221 00 Lund, Sweden

I, the undersigned, being the copyright owner of the abstract of the above-mentioned dissertation, hereby grant to all reference sources permission to publish and disseminate the abstract of the above-mentioned dissertation.

Signature _____ Date _____

ACKNOWLEDGEMENTS

I wish to thank:

My supervisors Christer Larsson and Marianne Sommarin for great support

Magnus Alsterfjord for supervision and team work in the lab

Adine Karlsson for all help in the lab

All past and present members of Plant Biochemistry and Biochemistry

Mum and dad for always believing in me

My sons Jonatan and Alexander, you make life worth living

My husband Niklas, for everything. I love you

CONTENTS

1. INTRODUCTION.....	1
2. STRUCTURE AND FUNCTION.....	3
3. 14-3-3 IN PLANTS.....	11
4. ISOFORM SPECIFICITY IN TARGET PROTEIN INTERACTIONS	15
4.1 THE H ⁺ -ATPASE/14-3-3-INTERACTION	16
5. SPECIFICITY IN EXPRESSION	19
5.1 PROMOTER ANALYSES OF 14-3-3 ISOFORMS	20
5.2 RESULTS FROM THE PROMOTER:GUS ANALYSIS.....	21
6. SUMMARY AND FUTURE WORK	31
7. POPULÄRVETENSKAPLIG SAMMANFATTNING PÅ SVENSKA.....	33
8. REFERENCES.....	35

THIS THESIS IS BASED ON THE FOLLOWING PUBLICATIONS:

Paper I

Alsterfjord M, Sehnke P C, Arkell A, Larsson H, Sventenlid F, Rosenquist M, Ferl R J, Sommarin M and Larsson C (2004) Plasma membrane H⁺-ATPase and 14-3-3 isoforms of Arabidopsis leaves: Evidence for isoform specificity in the 14-3-3/H⁺-ATPase interaction. *Plant Cell Physiol* 45: 1202-1210

Paper II

Alsterfjord M, Arkell A, Larsson C and Sommarin M (2010) Expression patterns of Arabidopsis 14-3-3 isoforms. Manuscript in preparation.

1

INTRODUCTION

The 14-3-3 proteins constitute a family of highly conserved proteins with a molecular mass of about 30 kDa. The first 14-3-3 proteins were discovered in 1967 as part of an examination of brain tissue proteins (Moore and Perez 1967). The 14-3-3s were given their name based on their fraction number on DEAE-cellulose chromatography and migration position on starch gel electrophoresis. For a long time 14-3-3s were thought to be brain-specific proteins but in 1992 14-3-3s were found in several plants (Brandt *et al* 1992, Hirsch *et al* 1992, Lu *et al* 1992) as well as in the yeast *Saccharomyces cerevisiae* (van Heusden *et al* 1992). 14-3-3s have now been found in all organisms examined except for members of the prokaryotic kingdom. In unicellular organisms as yeast there are only few 14-3-3 isoforms whereas in multicellular organisms, such as *Arabidopsis*, there may be as many as 15 isoforms (van Heusden *et al* 1995, Rosenquist *et al* 2000, Rosenquist *et al* 2001).

The first functional properties of 14-3-3s described were their ability to bind to and activate tyrosine and tryptophane hydroxylase in bovine brain in the presence of Ca^{2+} /calmodulin-dependent protein kinase type II (Ichimura *et al* 1988). Over the following years more functions were discovered, such as inhibition of protein kinase C in sheep brain, activation of Raf in *Xenopus* oocytes and binding to cruciform DNA in humans (Toker *et al* 1990, Aitken *et al* 1995, Todd *et al* 1998). The first 14-3-3 protein identified in plants was found to be involved in gene regulation (Lu *et al* 1992). Now, 14-3-3 proteins are

recognized as key regulators of primary metabolism and membrane transport in plants (Bachmann *et al* 1996, Moorhead *et al* 1996, Toroser *et al* 1998, Huber *et al* 2002). For eukaryotes in general, 14-3-3s have been found to be involved in numerous processes in the cell (Finnie *et al* 1999, Ferl *et al* 2002, Roberts 2003) and over 700 binding partners have been identified and the total number of potential plant 14-3-3 targets is more than 300 (MacKintosh 2004, Oecking *et al* 2009).

The aim of my work has been to understand why multicellular organisms need so many 14-3-3 isoforms. One possibility is that there is specificity in the 14-3-3/target protein interaction, which we have investigated in the model plant *Arabidopsis thaliana* using the plasma membrane H⁺-ATPase/14-3-3 interaction as a model system. Another possibility is that there is a specific expression of isoforms (Daugherty *et al* 1996), which we also have investigated in *Arabidopsis*. This was done using promoter:GUS fusions, which has yielded information on the cell-, tissue- and organ-specific distribution of most of the fifteen 14-3-3 isoforms, as well as on developmental regulation of 14-3-3 expression.

2

STRUCTURE AND FUNCTION

The 14-3-3 proteins are small acidic proteins, which are highly conserved even between kingdoms (Paul *et al* 2009) (Figure 1).

The 14-3-3 proteins assemble as stable homo- or heterodimers (Roberts 2000, Jones *et al* 1995) and, for example, 14-3-3 omega is shown to dimerize with at least 10 of the other 14-3-3 isoforms in Arabidopsis (Chang *et al* 2009). All 14-3-3 proteins appear to share a similar tertiary structure, first defined for the human isoforms tau (Xiao *et al* 1995) and zeta (Liu *et al* 1995) (Figure 2A). Each polypeptide is organized into nine anti-parallel α -helices, each separated by a short loop. The four N-terminal helices lie in a planar array and create an extensive dimerization surface. Parts of helices 1 and 2 from one monomer and parts of helices 3 and 4 from the other monomer form the dimerization domain (Liu *et al* 1995, Xiao *et al* 1995, Aitken 2002). The amino acids in the dimerization domains are not completely conserved which might indicate differences in dimer formation between isoforms.

The crystal structure has been solved for several mammalian and plant 14-3-3s (Liu *et al* 1995, Xiao *et al* 1995, Würtele *et al* 2003) and the extreme conservation of the central core region of the 14-3-3s make it very likely that this structure is a common feature of all 14-3-3s in all eukaryotes. However, all of the known crystal structures fail to resolve the N and C termini, which (along with several small regions within the molecule) are highly divergent among isoforms. Thus it is possible to consider the model as generally applicable to all plant 14-3-3s while recognizing that divergent areas might well contribute to specific structures and regulatory functions (Paul *et al* 2008).




Figure 2 **A** Structure of the human 14-3-3 zeta dimer. The numbers indicate helix numbers. Blue and yellow parts on helices indicate dimerization domains and the nuclear export signal (NES), respectively (compare Figure 1) (modified from Liu *et al* 1995). **B** A dimeric 14-3-3 protein binding two peptides (in yellow), mimicking the C-terminal end of the plasma membrane H^+ -ATPase from tobacco (Würtele *et al* 2003) .

14-3-3s typically bind to phosphorylated motifs in their target proteins. Thus, the α -helices forms the walls of an amphipathic groove, large enough for a phosphopeptide chain of a target protein to fit in (Figure 2B). The amino acid residues exposed in the binding groove are highly conserved and each subunit of the dimer is able to bind one target protein independently of the target in the other subunit (Liu *et al* 1995, Obsil *et al* 2001, Rittinger *et al* 1999). The C-terminal region of a 14-3-3 was shown to change in conformation when phosphorylated (Obsilova *et al* 2004). This relatively unconserved region of the 14-3-3 has been hypothesized to play an autoinhibitory role in ligand binding (Obsilova *et al* 2004, Truong *et al* 2002) through its high content of negatively charged amino acid residues, possibly in an isoform-specific manner (Figures 1 and 3).

The C-terminal end is the part of the 14-3-3s that differs most, both in sequence and length, and the length of Arabidopsis C termini vary from 8 to 31 amino acids, counting from the last conserved residue (Figure 3). Also, the amino acids facing the outside of the 14-3-3 molecule are relatively less conserved (Liu *et al* 1995, Xiao *et al* 1995).

Pi	S G D G N G N K T D C
Epsilon	S D L N E E G D E R T H C A D E P Q D E N
Omicron	S D L E E G C G K
Iota	S D L P E D G G E D N I K T E E S K Q E Q A K P A D A T E N
Mu	S D I S E E G G D A H K T N G S A K P G A G G D D A E
Kappa	S D M Q E Q N D E A
Lambda	S D M Q E Q N D E A
Omega	S D M Q D A A D E I K E A A A P K P T E E Q Q
Chi	S D M Q D V A D D I K E A A P A A K P A D E Q Q S
Phi	S D M Q D E S P E E I K E A A A P K P A E Q E I
Psi	S D M T D E A C D E I K E A S K P D C A E
Nu	S D I N D E A C G D E I K R A S K H E P E E G K P A E T C Q
Upsilon	S D L N D E A G D D I K E A P K E V Q K V D E Q A Q P P S Q

Figure 3 The C-terminal amino acid sequences from the last conserved residue of all 14-3-3s in Arabidopsis except the putative products of *grf14* (*xi*) and *grf15*. *grf* is short for G-box regulating factor (see section 3). *grf14* is truncated N-terminally of this domain. Acidic amino acid residues are indicated in bold (modified from Alsterfjord 2006).

The amino acid sequence motifs that 14-3-3s bind to are usually phosphorylated but also non-phosphorylated motifs have been identified. These non-phosphorylated motifs contain negatively charged amino acids which replace and mimic the negatively charged phosphogroup. The phosphorylated motifs have been divided into three modes (Ganguly *et al* 2005, Coblitz *et al* 2006), as shown in Figure 4.




Figure 4 The phosphorylated motifs that 14-3-3 bind to have been divided into three modes (mode 1, 2 and 3). pS may be replaced by pT.

The proline in mode 1 and 2 is needed to bend the peptide to exit the binding groove (Rittinger *et al* 1999) (Figure 5). The mode 3 motifs are also called C-terminal binding motifs, since they constitute the C termini of the target proteins (Ganguly *et al* 2005). The motifs in mode 3 do not need a proline to bend the peptide since the target protein ends in the binding groove (Figure 5).

The C-terminal motif was first found in the plasma membrane proton (H⁺)-ATPase of plants, Y/H-pT-V/L/I-COOH, which constitutes the last three amino acids in Arabidopsis plant plasma membrane H⁺-ATPase isoforms 1 to 11 (Olsson *et al* 1998, Fuglsang *et al* 1999, Svennelid *et al* 1999). In 2003, a crystallized structure of 14-3-3 from tobacco, together with the fungal toxin fusicoccin and a phosphopeptide mimicking the mode 3 binding motif of a tobacco H⁺-ATPase, showed that the same residues in the binding groove of the 14-3-3 protein are involved in binding the phosphothreonine in mode 3 as in mode 1 and 2 binding to phosphoserine (Würtele *et al* 2003, Ganguly *et al* 2005, Coblitz *et al* 2006). Fusicoccin occupies the hydrophobic end of the binding groove, which is not occupied in mode 3 (Figure 5) and enhances binding strength 100-fold (Würtele *et al* 2003).




Figure 5 The different modes of 14-3-3 binding (Modified from Coblitz *et al* 2006 and Alsterfjord 2006). FC=Fusicoccin.

All 14-3-3 proteins contain a nuclear export signal (NES) (Figure 2). This domain can bind to the chromosome maintenance region 1 (Crm1), which interacts with the nuclear pore and in this way the NES-containing proteins will be exported from the nucleus (Fornerod *et al* 1997, Fukuda *et al* 1997, Ossareh-Nazari *et al* 1997, Stade *et al* 1997). The NES domain in 14-3-3 proteins includes amino acids that are also involved in target binding and thereby the target will compete with the Crm1 (Rittinger *et al* 1999). Even if one NES domain is occupied by a target protein, the 14-3-3 proteins assemble as dimers and there will thus be one free NES domain. 14-3-3s can be found in the nucleus and this indicates that the NES signal can be hidden (Cutler *et al* 2000, van Zeijl *et al* 2000). When the NES signal is exposed, the 14-3-3 protein will be exported from the nucleus together with its target. This is a way of removing proteins, such as transcription factors, from the nucleus, as a response to other regulatory mechanisms (Fornerod *et al* 1997, Fukuda *et al* 1997, Ossareh-Nazari *et al* 1997, Stade *et al* 1997).

The number of identified physiological functions involving 14-3-3 proteins has increased rapidly since the discovery of 14-3-3s in 1967 (Moore and Perez 1967). Some of the physiological functions involving 14-3-3 proteins are regulation of signaling pathways, apoptosis, cell cycle entry, intracellular trafficking and metabolism (Darling *et al* 2005) (Figure 6).




Figure 6 Some of the numerous physiological functions that involves 14-3-3 proteins (modified from Darling *et al* 2005 and Johnson *et al* 2010).

Figure 7 shows the various proposed mechanisms of action for 14-3-3 proteins (Darling *et al* 2005). **(I)** 14-3-3 proteins have a rigid structure which leads to deformation of the target protein with little or no change in the structure of the 14-3-3 dimer (Yaffe 2002, Obsil *et al* 2001). The deformation of the target protein will lead to a change in activity, an increase or a decrease. Plant ATP synthases in chloroplasts and mitochondria are examples of proteins that are negatively regulated by this mechanism (Bunney *et al* 2001, Moorhead *et al* 1999) whereas the plant plasma membrane H^+ -ATPase is upregulated upon 14-3-3 binding (Jahn *et al* 1997, Oecking *et al* 1997, Olsson *et al* 1998, Fuglsang *et al* 1999, Sveneslid *et al* 1999). **(II)** 14-3-3s can mask a region of a protein and in that way e.g. hinder protein/protein interaction. **(III)** A 14-3-3 can hold two phosphoproteins close together, stabilizing their interaction. **(IV)** 14-3-3s can bind to phosphorylated targets and prevent either dephosphorylation or proteolysis. One example of prevention of proteolysis is 14-3-3 binding to phosphorylated plant nitrate reductase (Weiner and Kaiser 1999, Cotelle *et al* 2000). **(V)** Binding to 14-3-3 can increase the nuclear export or decrease nuclear import for the target protein (Muslin and Xing 2000).




Figure 7 The various proposed mechanisms of action that 14-3-3 can take on its targets (Darling *et al* 2005).

Thus, the functional diversity of 14-3-3 proteins is very high, and may even have facilitated evolutionary changes in target protein families (Johnson *et al* 2010).

3

14-3-3 IN PLANTS

14-3-3s were first discovered in plants (spinach, pea, barley and *Arabidopsis*) in 1992 (Brandt *et al* 1992, de Vetten *et al* 1992, Hirsch *et al* 1992, Lu *et al* 1992).

Numerous data for 14-3-3 functions and 14-3-3 interactants in plants have come from the study of *Arabidopsis thaliana*, a well-characterized model organism first described by Laibach in 1943. *Arabidopsis* has five chromosomes and approximately 26,000 genes and was fully sequenced in 2000 (*Arabidopsis* Genome Initiative). The life cycle of *Arabidopsis* from germination to mature seed is approximately 6 weeks, facilitating its study. It is also susceptible to genetic manipulation with mutagens or *Agrobacterium tumefaciens* infection to create transgenic *Arabidopsis* plants. *Arabidopsis* has 15 genes for 14-3-3s (Rosenquist *et al* 2001) which may be divided into two groups dependent on their exon patterns, the non-epsilon group and the epsilon group (Figure 8).




Figure 8 Gene maps of all Arabidopsis 14-3-3 genes. Exons are indicated as black boxes and introns as white boxes. Exon and intron sizes are indicated with the number of bases within each box. The genes can be divided into two groups based on the exon patterns: *grfs* 1 to 8 (the non-epsilon group) and *grfs* 9 to 14 (the epsilon group). *grf15* is aligned N-terminally with its closest neighbors, *grfs* 3 and 5 (modified from Rosenquist *et al* 2001). *Grfs* stands for G-box regulating factors.

The G-box is a common regulatory element found in many plant gene promoters, and historically 14-3-3s have been named GF14 proteins in Arabidopsis, because the protein was first identified to be a “G-box factor 14-3-3 homologue” (de Vetten *et al* 1992). The 14-3-3 genes in Arabidopsis are named *grfs* which stands for G-box regulating factors. Experiments showed that a plant 14-3-3 was able to function as a mammalian 14-3-3 (Lu *et al* 1994) providing evidence of the conserved nature of 14-3-3s. In 1994, 14-3-3 was identified as the binding protein for the fungal toxin fusaric acid (Korthout and de Boer 1994, Marra *et al* 1994, Oecking *et al* 1994). The binding protein was later shown to be a complex of the plasma membrane H⁺-ATPase and 14-3-3 (Jahn *et al* 1997, Oecking *et al* 1997), and binding of 14-3-3 was shown to activate the plasma membrane H⁺-ATPase.

Since 1992, 14-3-3s have been discovered to be involved in numerous processes in plants, such as metabolism, the cell cycle, apoptosis, signal transduction and intracellular trafficking.

Thus, 14-3-3 proteins were found to interact with the mitochondrial and chloroplastic ATP synthase, negatively regulating the ATP synthesis in both organelles (Bunney *et al* 2001).

In anti-sense experiments, down-regulation of specific 14-3-3 isoforms resulted in increase in leaf starch accumulation. The starch synthase III family was identified as a possible 14-3-3 target as all members of the family contain a 14-3-3 binding motif. The interaction between the starch synthase III member, DU1 and 14-3-3 was demonstrated, confirming a role for 14-3-3 proteins in regulation of starch synthesis (Sehnke *et al* 2001).

Regulation of key enzymes such as sucrose phosphate synthase, starch synthase (Toroser *et al* 1998), nitrate reductase (Bachmann *et al* 1996, Moorhead *et al* 1996), the plasma membrane H⁺-ATPase as well as the mitochondrial and chloroplast ATP synthases implies that 14-3-3 is essential for regulating carbon and nitrogen metabolism in plants (Kulma *et al* 2004, Harthill *et al* 2006, Huber *et al* 2002).

14-3-3s also play diverse roles during seed germination. Two major hormones, abscisic acid (ABA) and gibberellins have opposite functions during germination. Gibberellins generally promote germination, whereas ABA inhibits germination and has a role in inducing and maintaining dormancy. ABA seems to mediate its effect by promotion of the *Em* (embryo) gene, and in the absence of *Em* transcripts, maize produces embryos that germinate while still attached to the parent plant. By use of a yeast two-hybrid system, 14-3-3 dimers were demonstrated to provide a structural link between elements of the transcriptional protein complex and the *Em* promoter to inhibit germination (Schultz *et al* 1998). RSG is a tobacco plant bZIP transcription factor that regulates shoot growth by altering transcription of genes required for gibberellin synthesis. A yeast two-hybrid screen demonstrated that RSG interacts with several isoforms of tobacco 14-3-3 in a phosphorylation-dependent manner (Igarashi *et al* 2001, Ishida *et al* 2004). If a point mutation is done in the phosphorylated motif, the transcriptional activity of RSG is increased due to transcription factor accumulation within the nucleus. In wild-type cells, RSG is distributed throughout the cell. 14-3-3 therefore negatively regulates gibberellin signaling by localizing RSG outside the nucleus.

A recent study has demonstrated 14-3-3s to be an essential part of brassinosteroid (BR) signaling (Gampala *et al* 2007, Ryu *et al* 2007). Many components in this signaling pathway are already well known in Arabidopsis, including the cell surface receptor kinase BRASSINOSTEROID-INSENSITIVE 1 (BRI1) and its coreceptor BRI1-ASSOCIATED RECEPTOR KINASE 1 (BAK1) as well as the transcription factor BRASSINAZOLE-RESISTANT 1 (BZR1), which directly regulates BR responsive gene expression (Gendron *et al* 2007). The recent studies showed 14-3-3 to bind the BZR1 transcription factor upon phosphorylation by an intracellular kinase which is active in the absence of BR (Gampala *et al* 2007, Ryu *et al* 2007). Comparable to RSG, 14-3-3 association mediates the cytoplasmic retention and/or nuclear export of BZR1, thus efficiently inhibiting its function.

4

ISOFORM SPECIFICITY IN TARGET PROTEIN INTERACTIONS

In multicellular organisms there is a relatively large number of 14-3-3 isoforms and the question arises as to the reason for this. One possibility is that there is specificity in the 14-3-3/target protein interaction, which we have investigated in the model plant *Arabidopsis thaliana* using the plasma membrane H⁺-ATPase/14-3-3 interaction as a model system. Another possibility is that there is a developmental, cell-, tissue- or organ-specific expression of isoforms (Daugherty *et al* 1996), which we also have investigated in *Arabidopsis* (see section 5).

14-3-3 isoform specificity has been shown for some targets. Thus, both plasma membrane H⁺-ATPase (see section 4.1, Rosenquist *et al* 2000, Emi *et al* 2001) and nitrate reductase (Bachmann *et al* 1996) show binding specificity to 14-3-3 isoforms in *Arabidopsis*. In barley, 14-3-3B and C are efficient inhibitors of nitrate reductase whereas 14-3-3A is not (Sinnige *et al* 2005). Also the phototropin receptor kinase 1 (phot1) shows specificity in *Arabidopsis*. 14-3-3 binding to phot1 is limited to non-epsilon 14-3-3 isoforms (Sullivan *et al* 2009).

The C terminus of 14-3-3s may act as an autoinhibitory domain that interferes with ligand binding (Truong *et al* 2002, Shen *et al* 2003, Kubala *et al* 2004, Silhan *et al* 2004). All 14-3-3 C termini contain acidic amino acids (Figure 3) which may mimic phosphorylated target motifs and thus compete with proper target binding. The parts of 14-3-3 that bind to a target protein are very conserved but the C terminus is the part of the 14-3-3 that differs most. This feature may be one factor leading to isoform specificity.

Posttranslational modification (proteolytic cleavage or phosphorylation) could be another way to increase specificity of 14-3-3s (Fuller *et al* 2006). Proteolytic cleavage was shown in barley (van Zeijl *et al* 2000) and also the 43 kD band in Figure 10 is probably a proteolytic cleavage product of a 14-3-3 dimer (Bernfur and Alsterfjord, personal communication). 14-3-3s may also be posttranslationally modified by phosphorylation. Phosphorylation sites in the dimerization domain of human 14-3-3s have been identified (Aitken 2002).

4.1 The H⁺-ATPase/14-3-3-interaction

The plasma membrane H⁺-ATPase couples ATP hydrolysis to proton transport. This creates the pH and potential difference across the plasma membrane required by secondary transporters whose activity is directly dependent upon the proton motive force. In plants, the plasma membrane H⁺-ATPase also participates in other functions essential for normal plant growth such as salt tolerance, intracellular pH regulation and cellular expansion (Morsomme and Boutry 2000, Palmgren 2001). Given these multiple physiological roles and the high ATP consumption, the H⁺-ATPase has to be tightly regulated. The C terminus of the H⁺-ATPase acts as an autoinhibitory domain (Palmgren *et al* 1991) and when it is phosphorylated 14-3-3 can bind, the autoinhibitory domain is displaced and the activity of the H⁺ pump is increased (Jahn *et al* 1997, Oecking *et al* 1997, Olsson *et al* 1998, Fuglsang *et al* 1999, Sannelid *et al* 1999).

Fusicoccin is a wilt-inducing toxin produced by the fungus *Fusicoccum amygdale*. The natural hosts of *F. amygdale* are almond and peach trees. The toxin is commonly used in plasma membrane H⁺-ATPase experiments as it causes an “irreversible” 14-3-3 binding and thus gives a stable H⁺-ATPase/14-3-3 complex (Jahn *et al* 1997, Oecking *et al* 1997, Fullone *et al* 1998, Würtele *et al* 2003).

Similar to the Arabidopsis 14-3-3 proteins the Arabidopsis plasma membrane H⁺-ATPase belongs to a large gene family, with 12 predicted isoforms of which 11 are expressed (Arango *et al* 2003). The plasma membrane H⁺-ATPase is the main 14-3-3 target in the plasma membrane and during conditions requiring full activation of H⁺ pumping several percent of total cellular 14-3-3 may be involved in activation of the H⁺-ATPase (Paper I). Considering the large number of 14-3-3 and H⁺-ATPase isoforms in Arabidopsis, specificity in binding may exist between isoforms. This assumption is supported by the large differences in

amino acids in the C-terminal binding motif of the H⁺-ATPase isoforms, which may affect binding (Figure 9). There is also another motif in the C terminus of the H⁺-ATPase (DIE/DID/DLE)) that has been suggested as a nonphosphorylated binding motif for 14-3-3 (Visconti *et al* 2003) and this motif is not found in H⁺-ATPase isoform 10 (AHA10) which may indicate a difference in binding properties (Figure 9).

```

AHA1  LKGLDIDTAGH-HYTV
AHA2  LKGLDIETPS--HYTV
AHA3  LKGLDIETAG--HYTV
AHA4  LKGVDIETIQQ-AYTV
AHA5  LKGLDIDTIQQ-HYTV
AHA6  LKGLDIDNLNQ-HYTV
AHA8  LKGLDIDTIQQ-HYTV
AHA11 LKGLDIETIQQ-AYTV
AHA7  LKGYDLEDPNSNNYTI
AHA9  QKGLDIEAIQQ-HYTL
AHA10 LKQIDQRMIRA-AHTV

```

Figure 9 Amino acid sequences of the C termini of the 11 expressed Arabidopsis H⁺-ATPase (AHA) isoforms harbouring the 14-3-3 binding motifs. The most conserved part of the mode 3 motif is indicated in red, the more variable beginning of the motif is in green and a proposed nonphosphorylated binding motif (Visconti *et al* 2003) is in yellow (modified from Alsterfjord 2006).

Using 12 of the Arabidopsis 14-3-3 isoforms, we could show that they all bind to the H⁺-ATPase present in isolated Arabidopsis plasma membranes (Paper I). This was however not the case *in vivo*. Using 14-3-3 isoform-specific antibodies, all isoforms tested (all but omicron, iota, psi and pi for which we do not have access to isoform-specific antibodies) except kappa and phi were identified in the supernatant (Figure 10).




Figure 10 *14-3-3* isoforms in plasma membrane and supernatant fractions visualized by isoform-specific antibodies. Single arrows indicate the position of *14-3-3* monomers and double arrows the position of dimers (modified from Paper I and Alsterfjord 2006). The 43 kD band is probably a proteolytic cleavage product of the *14-3-3* dimer (Bernfur and Alsterfjord, personal communication).

However, the plasma membrane fraction lacked not only kappa and phi but also chi and upsilon in the absence of fusicoccin (Figure 10, control). Thus, we could detect differences in distribution of the *14-3-3*s between isolated plasma membranes from Arabidopsis leaves and a supernatant fraction, representing all soluble proteins. In the presence of fusicoccin which increases the H^+ -ATPase/*14-3-3* interaction 100-fold (Würtele *et al* 2004) also phi and upsilon were attached to the plasma membrane, but not chi. Thus, phi and upsilon are accessible to the plasma membrane but have either a low affinity for the plasma membrane H^+ -ATPase isoforms present or are, in the absence of fusicoccin, occupied by other targets for which they have higher affinity. Chi may be localized in another compartment than the cytosol and may therefore not be available to the plasma membrane. Altogether this suggests that there is some isoform specificity in the *14-3-3*/ H^+ -ATPase interaction *in vivo* (Paper I).

5

SPECIFICITY IN EXPRESSION

A developmental, cell-, tissue- or organ-specific expression is suggested by the large number of 14-3-3 isoforms in multicellular organisms compared to the very few in unicellular organisms (Rosenquist *et al* 2000, Alsterfjord *et al* 2004). An organ-specific expression was shown for Arabidopsis 14-3-3 iota, which was only expressed in the flower whereas Arabidopsis omicron was expressed in leaf, root and flower (Rosenquist *et al* 2001). Arabidopsis 14-3-3 mu was shown to be expressed in all tissues and developmental stages (Kuromori and Yamamoto 2000) and transcripts for all of the Arabidopsis 14-3-3 isoforms except pi, iota and psi are present in leaves (Paper I). Also in mammals some 14-3-3 isoforms are widely expressed whereas others show a more specific expression. Human 14-3-3 zeta is present in high levels in the brain grey matter, 14-3-3 gamma is specific for the central nervous system, 14-3-3 epsilon is found in the pineal gland and the retina, and 14-3-3 tau is only present in glial cells (Watanabe *et al* 1993, Takahashi 2003).

There are also differences in subcellular localization between isoforms. Arabidopsis 14-3-3 epsilon, mu, nu and upsilon are present in both the chloroplast and cytoplasm (Sehnke *et al* 2000) and three isoforms in barley were detected in mitochondria (Bunney *et al* 2001), and it has been suggested that some 14-3-3 subcellular localization is driven by both isoform specificity and target interactions (Paul *et al* 2005).

5.1 Promoter analyses of 14-3-3 isoforms

Agrobacterium tumefaciens infection was used to create transgenic Arabidopsis plants to see if there is specificity in developmental expression and localization of the different 14-3-3 isoforms. *A. tumefaciens* has the ability to insert its natural so called T-DNA into the nuclear genome of Arabidopsis. The specific T-DNA is chosen by two short sequences, left border and right border. The DNA fragment that is to be inserted into the genome of the plant is cloned between the left and the right border together with a marker and thus an artificial T-DNA is created and the *A. tumefaciens* can introduce it into the genome of the plant.

The methods to get *A. tumefaciens* to insert the T-DNA into the plant have rapidly changed and today it is relatively simple to get transgenic plants. The Arabidopsis flower is simply dipped into a solution of *A. tumefaciens* approximately five days before the flowers open, the so called floral-dip method (Desfeux *et al* 2000).

The *Escherichia coli uidA* gene encoding β -glucuronidase (GUS) is one of the most effective reporter gene systems used for evaluating transient and stable transformation in plants. Since its description by Jefferson *et al* (1987), the GUS gene fusion system has found extensive application in plant gene expression studies because of the enzyme stability and high sensitivity and suitability of the assay to detection by fluorometric, spectrophotometric or histochemical techniques. The GUS protein is a 68kD homo-tetramer that catalyzes the hydrolysis of β -glucuronides. In most eukaryotic organisms, these are formed to detoxify and excrete xenobiotic and endogenous waste products (Fior *et al* 2009).

To see if there is specificity in expression and localization 1,5kb upstream of the start codon of thirteen of the fifteen 14-3-3 isoforms have been fused to reporter genes coding for GUS and enhanced green fluorescent protein (EGFP) (the EGFP was not used in this work) (Figure 11). This 1,5 kb region represents the 14-3-3 promoter. This construct is transformed into Arabidopsis with *A. tumefaciens*. When the transgenic plants are incubated with the substrate 5-bromo-4-chloro-3-indolyl β -D-glucuronide cyclohexylammonium salt (X-gluc) a blue precipitation is created where the 14-3-3 isoform normally is expressed. This precipitation can easily be seen by the eye or in a microscope. To see if there is any difference in expression of the promoter when the plant matures, samples were taken at four different developmental stages, seedlings (one week old), adult leaves (three weeks old), flowers (four to five weeks old) and siliques (six weeks old).




Figure 11 Schematic view of the promoter-reporter genes fusion used in expression analyses of Arabidopsis 14-3-3 promoters (Alsterfjord 2006).

5.2 Results from the promoter:GUS analysis

The data presented below demonstrate a specific developmental cell-, tissue and organ distribution of the Arabidopsis 14-3-3 isoforms. Arabidopsis transformed with the 14-3-3 promoter-EGFP-GUS construct makes the study of both tissue distribution and developmental regulation possible. The results support the idea that the various 14-3-3 isoforms have separate and distinctive tissue distribution, suggesting that plant 14-3-3 isoforms may indeed have specific roles within individual tissues (Paper II).




Figure 12 A schematic Arabidopsis thaliana flower(A) (modified from geochembio.com) and plant (B) (modified from Winter et al 2007) with different tissues indicated.




Figure 13 *GUS-staining detecting promoter activity of the Arabidopsis 14-3-3 isoforms in the epsilon group. The promoter of pi shows activity in hydathodes, xi shows promoter activity in the hypocotyls of seedlings, the omicron promoter is active in the petioles of leaves and the promoter of epsilon shows expression especially in vascular tissue. The promoter of iota is expressed in early stamens exclusively and the mu promoter shows expression in green tissue, especially in vascular tissue. Compare figure 12.*

5.2.1 The epsilon group

The promoters of the epsilon group show a more specific expression than the promoters of the non-epsilon group (Figures 13 and 15).

The Arabidopsis 14-3-3 **pi** shows promoter activity in hydathodes, pollen and seeds only.

The isoform **xi** has never been shown to be expressed, however the promoter shows activity in the hypocotyls of seedlings, in roots, in pollen and in seeds.

The **omicron** promoter shows an expression pattern which is different from all the other Arabidopsis 14-3-3 promoters. Activity is found in internodes of vascular tissue, in leaves and in the petioles of leaves. The promoter of omicron also shows expression in pollen and seeds.

The promoter of **epsilon** shows activity in young tissue, especially in vascular tissue, in roots (it is the only isoform that shows expression in the root buds) and in stipules.

The promoter of **iota** is exclusively active in early stamens but not anywhere else in the plant.

The promoter of **mu** shows expression in the green tissues of the plant, especially in the vascular tissues, but also in roots and sepals.




Figure 14 *GUS-staining detecting promoter activity of the Arabidopsis 14-3-3 isoforms in the non-epsilon group. The kappa promoter shows much activity in the flower and in cauline leaf hydathodes, the lambda promoter is active primarily in vascular tissue. The phi promoter shows activity in young tissues, the psi promoter is mainly active in the vascular tissue and the nu promoter shows activity in young tissues and decreases with age. The promoter of epsilon is active throughout the whole plant especially in the vascular tissue and the promoter of omega is active in the flower and the activity in carpels is decreasing but is still present in the siliques. The promoter of omega is also active in the roots with the exception of the root tip. Compare figure 12.*

5.2.2 The non-epsilon group

The promoters of the non-epsilon group are widely active and shows more random expression in leaves, roots and flowers than the promoters of the epsilon group (Figures 14 and 15).

The **kappa** promoter is mainly active in young tissues and in meristematic tissues of seedlings. The roots also show activity, especially the ends of the roots. Kappa is also the only isoform whose promoter shows activity in the cauline leaf hydathodes and it also shows much activity in the flower.

The promoter of **lambda** shows activity primarily in vascular tissue, both in green tissue and in roots. Although the amino acid sequences of kappa and lambda are very much alike they do not show the same expression pattern.

The **phi** promoter is active in the young tissue of seedlings and decreases with age, the activity remains in the vascular tissue whereas activities in other parts disappear in older tissue. The roots as well as the flower show some phi promoter activity.

The promoter of **psi** is mainly active in vascular tissues and in roots.

The **nu** promoter shows activity in young tissue which decreases with ageing, and then appears again in pollen and seeds.

The promoter of **epsilon** is active throughout the whole plant especially in the vascular tissue.

In seedlings, the promoter of **omega** is active in young tissues and especially in meristem and vascular tissue. There is also high activity in the roots, with the exception of the root tip. The omega promoter is also highly active in flowers. In early flowers the activity is limited to parts of the carpels. As the flower matures the activity is changed to include the stamens, especially pollen. The promoter activity in carpels is decreasing but is still present in siliques.

5.2.3 Conclusions regarding specificity of expression

Studied at this level there is clearly cell-, tissue- and organ-specific expression for all of the Arabidopsis 14-3-3 isoforms as well as specific developmental expression. There is not a single case where the promoter of one isoform shows an expression that is identical to the expression of another isoform (figure 12). The overall finding is that the promoters of the non-epsilon group show more random expression in leaves, roots and flowers and the promoters of the epsilon group show more specific expression. For example the promoter of iota shows expression only in the early stamens and the promoter of pi shows expression in hydathodes, pollen and seeds only.

The promoter of Arabidopsis 14-3-3 chi was earlier shown to be active in several tissues of the plant detected by a promoter:GUS fusion (Daugherty et al 1996). Promoter activity was shown in roots of seedlings and mature plants, in root hairs, in the whole bud of immature flowers, in the anthers, stigma and pollen of more mature flowers, in differential style and abscission zone of immature siliques and in mature siliques throughout the tissue. Promoter activity could also be detected in imbibed seeds.

We used the promoter:GUS fusion to see where in Arabidopsis the different 14-3-3 isoforms are expressed. It should be noted that a quantitative measure of the expression was not obtained due to the fact that inhibitors of GUS activity are ubiquitous in organ tissues of Arabidopsis, tobacco and rice (Fior *et al* 2009). In order to achieve reliable quantitative results, inhibitor activity should be routinely tested during quantitative GUS assays which was not done here.

Epsilon group

Non-epsilon group

Isoform Tissue	Pi	Xi	Epsilon	Omicron	Iota	Mu	Kappa	Lambda	Omega	Chi*	Phi	Psi	Nu	Upsilon
Primordial		+	+				+	+	+		+	+	+	+
Hypocotyls		+		+			+		+		+			+
Vascular hypocotyls		+	+			+		+	+		+	+		+
Petioles				+		+	+		+		+	+	+	+
Vascular petioles						+	+	+	+		+	+	+	+
Cotyledons				+			+				+			+
Vascular cotyledons			+	+		+	+	+			+	+		+
Leaves						+	+	+	+		+	+	+	+
Emerging leaves						+	+	+	+		+	+	+	+
Vascular leaves			+	+		+	+	+	+		+	+	+	+
Trichomes							+		+		+	+	+	+
Hydathodes	+		+	+		+	+	+			+			+
Roots		+		+		+		+	+	+	+	+		+
Vascular roots		+	+				+	+						
Lateral roots		+					+		+			+	+	+
Root branches		+	+											
Root tips		+	+									+	+	
Root buds			+											
Cauline leaf							+							
Stem							+				+		+	+
Stipules			+						+					
Sepals						+	+	+	+		+			
Stamen							+	+	+		+			+
Pollen	+	+		+	+		+		+	+	+		+	+
Carpels							+		+		+		+	+
Pedicels									+		+			+
Pedicel tips							+				+			+
Siliques							+		+	+	+			+
Seeds	+	+		+			+		+				+	+

Figure 14 Promoter expression of *Arabidopsis 14-3-3s* in plant organs at different developmental stages (Paper II). Compare Figure 12 . *Data from Daugherty et al 1996.

Our results are largely supported by the microarray data reflecting mRNA abundance, available at Genevestigator. Notably, these microarray data give a resolution at organ level and developmental stages, where as GUS staining gives a resolution at tissue and sometimes even cellular level.




Figure 16 A phylogenetic tree with topology representative for the *Arabidopsis* 14-3-3 protein family (modified from Rosenquist *et al* 2001 and Alsterfjord 2006).

As seen from Rosenquist *et al* (2001) some of the 14-3-3 isoforms are more closely related than others. For example, lambda and kappa are situated alone on one branch in the phylogenetic tree and also phi, chi and omega are situated on a common branch of the tree (Figure 16). The promoters of phi, chi and omega show similar expression, except that the promoter of chi does not show any expression in green tissues (Daugherty *et al* 1996). Also epsilon, psi and nu are close and they also show similar expression except that the promoter of psi

is not at all expressed in the reproductive tissues. However, the situation is different for kappa and lambda. Although these isoforms are situated close to each other in the phylogenetic tree their promoters do not always show a similar expression.

From the present work it is clear that a phylogenetic tree does not always reveal similarities or differences in where different isoforms are expressed, and this is because a phylogenetic tree is based on the amino acid sequences of the isoforms and not on promoter resemblance.

We have tried to compare the promoter regions but it has not been successful, since it is not easy to determine which bases in the promoter region that are important. A typical promoter contains a TATA-box and a CAAT-box. The function of a TATA-box is mainly the precise initiation of transcription. The CAAT-box is frequently focused on controlling transcription initiation. A typical promoter also harbours some special DNA sequences; cis-acting elements inhibiting or activating gene transcription by combining with the transcription factor (Gou *et al* 2010). Grondal *et al* (1990) were unable to identify conserved sequence element(s) by direct comparison of the promoter region of RNA polymerase I in *Crithidia fasciculata* and similar promoter regions in other eukaryotes, including the promoter region of the most closely related kinetoplastid species. When deciding what to use as the promoter region, a 1,5kb fragment is often chosen (Engelmann *et al* 2008). The fragment contains all the necessary information for the specific expression and it is not too long to work with. Barrero *et al* (2009) showed that a 850bp fragment upstream of the start codon of ZmMRP-1 is sufficient to direct GUS reporter gene activity in maize, but both Guo *et al* (2010) and Engelmann *et al* (2008) reported that they needed at least 1,5kb to capture the promoter of the calcium sensor gene CBL1 in *Ammopiptanthus mongolicus* and the glycine decarboxylase in *Flaveria trinervia* respectively .

6

SUMMARY AND FUTURE WORK

The 14-3-3 proteins are involved in a large number of processes and over 700 target proteins have been identified. In plants, 14-3-3s are involved in regulation of metabolism, membrane transport, and signal transduction. 14-3-3s have an important role in the regulation of nitrogen and carbon metabolism and in regulating the plasma membrane H^+ -ATPase.

To investigate if there is specificity in 14-3-3/target protein interaction the model system H^+ -ATPase/14-3-3 was used and it indicated some specificity but also a wide redundancy (Paper I). To further analyse the question of specificity the promoter:GUS fusion was utilized. The results clearly indicated developmental, cell-, tissue- and organ-specific expression for all of the Arabidopsis 14-3-3 isoforms. There is not a single case where the promoter of one isoform shows an expression that is identical to the expression of another isoform (Paper II). The results support the idea that 14-3-3s may indeed have specific roles within individual tissues.

It would be of interest to continue a sequence analysis of the different promoters to see if there are any similarities in the DNA sequence correlating to the similarities in expression between the different 14-3-3 isoforms.

Since promoter:GUS fusions of all isoforms of Arabidopsis H⁺-ATPase are also available, it would be of interest to compare the expression of these to the expression of the 14-3-3s and see if there is any correlation between expression of 14-3-3s and expression of one of the 14-3-3 proteins major targets, the H⁺-ATPase . For example, promoter activity of the Arabidopsis H⁺-ATPase 3 (AHA3) has been shown in vascular tissues similar to many 14-3-3s (DeWitt and Sussman, 1995) and the promoter of Arabidopsis H⁺-ATPase 10 has shown activity in developing seeds (Harper *et al* 1994).

7

POPULÄRVETENSKAPLIG SAMMANFATTNING PÅ SVENSKA

Allt levande är uppbyggt av celler, från encelliga organismer som t ex jäst till flercelliga organismer som t ex djur och växter. Alla celler regleras noggrant så att den inre miljön är konstant oavsett vad som händer i den yttre miljön. Cellerna har sensorer så att de alltid vet vad som händer runt dem och kan svara på detta. Dessa reglerande system består till stor del av stora molekyler som kallas proteiner. En sådan grupp av reglerande proteiner kallas 14-3-3.

14-3-3 proteiner utgör en familj av proteiner som är mycket lika varandra. De upptäcktes redan 1967 i hjärna från ko. Det konstiga namnet har 14-3-3 proteinerna fått från sättet de renades fram på. Från början trodde man att dessa proteiner bara fanns i hjärna men sedan hittades de också i andra vävnader hos djur. 1992 hittades 14-3-3 även i växter och i jäst. I encelliga organismer som tex jäst finns få varianter (isoformer) av 14-3-3 medan i djur och växter kan det finnas många. T ex så finns det 15 stycken i modellväxten Arabidopsis. 14-3-3 är med stor sannolikhet det protein som är involverat i flest processer i cellen och har visat sig interagera med mer än 700 andra sorters proteiner.

Växten *Arabidopsis* heter backtrav på svenska och används mycket inom växtforskningen. I växten har 14-3-3 visat sig vara involverade i många viktiga processer. Tidigare trodde man att de många olika varianterna av 14-3-3 hade samma uppgifter i cellen men nya data visar att de faktiskt kan ha specifika uppgifter trots att de är så lika varandra. Därför är det av intresse att t ex kartlägga var i växten som de olika varianterna finns.

Med hjälp av ett modellsystem där 14-3-3 reglerar ett annat protein som sitter i cellmembranet och pumpar protoner från insidan av cellen till utsidan har vi kunnat visa att det finns skillnader i hur de olika 14-3-3 varianterna binder till protonpumpen.

Det har också visat sig att olika isoformer av ett protein inte behöver finnas (uttryckas) i alla celler i en hel organism och i de fall där det finns många isoformer så är det av intresse att se om man kan hitta var de olika isoformerna uttrycks. Vi har med hjälp av genmodifierade *Arabidopsis* lyckats ta reda på var 13 av de 15 isoformerna av 14-3-3 uttrycks och visat att de finns på olika ställen i växten och att de då har olika uppgifter.

8

REFERENCES

Aitken A. (2002) Functional specificity in 14-3-3 isoform interactions through dimer formation and phosphorylation. Chromosome location of mammalian isoforms and variants. *Plant Mol Biol* 50, 993-1010

Aitken A, Howell S, Jones D, Madrazo J and Patel Y. (1995) 14-3-3 alpha and delta are phosphorylated forms of raf-activating 14-3-3 beta and zeta. In vivo stoichiometric phosphorylation in brain at a Ser-Pro-Glu-Lys motif. *J Biol Chem* 270, 5706-5709

Alsterfjord M. (2006) The Arabidopsis 14-3-3 family. Evolution, expression, localization and target specificity. Doctoral thesis. ISBN 91-7422-132-9

Alsterfjord M, Sehnke P C, Arkell A, Larsson H, Svennelid F, Rosenquist M, Ferl R J, Sommarin M and Larsson C. (2004) Plasma membrane H⁺-ATPase and 14-3-3 isoforms of Arabidopsis leaves: Evidence for isoform specificity in the 14-3-3/H⁺-ATPase interaction. *Plant Cell Physiol* 45, 1202-1210

Arango M, Gevaudant F, Oufattole M and Boutry M. (2003) The plasma membrane proton pump ATPase: the significance of gene subfamilies. *Planta* 216, 355-365

Bachmann M, Huber J L, Athwal G S, Wu K, Ferl R J and Huber S C. (1996) 14-3-3 proteins associate with the regulatory phosphorylation site of spinach leaf nitrate reductase in an isoform-specific manner and reduce dephosphorylation of Ser-543 by endogenous protein phosphatases. *FEBS Lett* 398, 26-30

Brandt J, Thordal-Christensen H, Vad K, Gregersen P L and Collinge D B. (1992) A pathogen-induced gene of barley encodes a protein showing high similarity to a protein kinase regulator. *Plant J* 2, 815-820

Bridges D and Moorhead G. (2004) 14-3-3: A number of functions for a numbered protein. *Science* 304, 242-252

Bunney T D, van Walraven H S and De Boer A H. (2001) 14-3-3 protein is a regulator of the mitochondrial and chloroplast ATP synthase. *Proc Natl Acad Sci U S A* 98:4249-4254

Bunney T D, Wijngaard P W and de Boer A H. (2002) 14-3-3 protein regulation of proton pumps and ion channels. *Plant Mol Biol* 50, 1041-1051

Chang I F, Curran A, Woolsey R, Quilici D, Cushman J C, Mittler R, Harmon A and Harper J F. (2009) Proteomic profiling of tandem affinity purified 14-3-3 protein complexes in *Arabidopsis thaliana*. *Proteomics* 9, 2967-2985

Coblitz B, Wu M, Shikano S, Li M. (2006) C-terminal binding: an expanded repertoire and function of 14-3-3 proteins. *FEBS Lett* 580, 1531-1535

Cotelle V, Meek S E, Provan F, Milne F C, Morrice N and MacKintosh C. (2000) 14-3-3s regulate global cleavage of their diverse binding partners in sugar-starved *Arabidopsis* cells. *EMBO J* 19, 2869-2876

Cutler S R, Ehrhardt D W, Griffiths J S, Somerville C R. (2000) Random GFP::cDNA fusions enable visualization of subcellular structures in cells of *Arabidopsis* at a high frequency. *Proc Natl Acad Sci U S A* 97, 3718-3723

Darling D L, Yingling J and Wynshaw-Boris A. (2005) Role of 14-3-3 proteins in eukaryotic signaling and development. *Curr Top Dev Biol* 68, 281-315

Daugherty C J, Rooney M F, Miller P W and Ferl R J. (1996) Molecular organization and tissue-specific expression of an Arabidopsis 14-3-3 gene. *Plant Cell* 8, 1239-1248

Desfeux C, Clough S J and Bent A F. (2000) Female reproductive tissues are the primary target of *Agrobacterium*-mediated transformation by the Arabidopsis floral-dip method. *Plant Physiol* 123, 895-904

de Vetten N C, Lu G and Ferl R J. (1992) A maize protein associated with the G-box binding complex has homology to brain regulatory proteins. *Plant Cell* 4, 1295-1307

DeWitt ND and Sussman MR. (1995) Immunocytological localization of an epitope-tagged plasma membrane proton pump (H^+ -ATPase) in phloem companion cells. *Plant Cell* 7, 2053-2067

Emi T, Kinoshita T and Shimazaki K. (2001) Specific binding of vf14-3-3a isoform to the plasma membrane H^+ -ATPase in response to blue light and fusicoccin in guard cells of broad bean. *Plant Physiol* 125, 1115-1125

Engelmann S, Wiludda C, Burscheidt J, Gowik U, Schlue U, Koczor M, Streubel M, Cossu R, Bauwe H and Westhoff P. (2008) The gene for the P-subunit of glycine decarboxylase from the C4 species *Flaveria trinervia*: Analysis of transcriptional control in transgenic *Flaveria bidentis* (C4) and *Arabidopsis* (C3). *Plant Physiol* 146, 1773-1785

Ferl R J, Manak M S and Reyes M F. (2002) The 14-3-3s. *Genome Biol Rev* 3, 3010

Finnie C, Borch J and Collinge D B. (1999) 14-3-3 proteins: eukaryotic regulatory proteins with many functions. *Plant Mol Biol* 40, 545-554

Fior S and Gerola PD. (2009) Impact of ubiquitous inhibitors on the GUS gene reporter system: evidence from the model plants *Arabidopsis*, tobacco and rice and correction methods for quantitative assays of transgenic and endogenous GUS. *Plant Methodes* 5:19-31

Fornerod M, Ohno M, Yoshida M, Mattaj I W. (1997) CRM1 is an export receptor for leucine-rich nuclear export signals. *Cell* 90, 1051-1060

Fuglsang A T, Visconti S, Drumm K, Jahn T, Stensballe A, Mattei B, Jensen O N, Aducci P, Palmgren M G. (1999) Binding of a 14-3-3 protein to the plasma membrane H⁺-ATPase AHA2 involves the three C-terminal residues Tyr(946)-Thr-Val and requires phosphorylation of Thr(947). *J Biol Chem* 274, 36774-36780

Fukuda M, Asano S, Nakamura T, Adachi M, Yoshida M, Yanagida M, Nishida E. (1997) CRM1 is responsible for intracellular transport mediated by the nuclear export signal. *Nature* 390, 308-311

Fuller B, Stevens SM Jr, Sehnke PC and Ferl RJ. (2006) Proteomic analysis of the 14-3-3 family in Arabidopsis. *Proteomics* 6, 3050-3059

Fullone M R, Visconti S, Marra M, Fogliano V and Aducci P. (1998) Fusicoccin effect on the in vivo interaction between plant 14-3-3 proteins and plasma membrane H⁺-ATPase. *J Biol Chem* 273, 7698-7702

Gampala SS, Kim TW, He JX, Tang W, Deng Z, Bai MY, Guan S, Lalonde S, Sun Y, Gendron JM, Chen H, Shibagaki N, Ferl RJ, Ehrhardt D, Chong K, Burlingame AL and Wang Z. (2007) An essential role for 14-3-3 proteins in brassinosteroid signal transduction in Arabidopsis. *Dev Cell* 13, 177-189

Ganguly S, Weller J L, Ho A, Chemineau P, Malpaux B, Klein D C. (2005) Melatonin synthesis: 14-3-3-dependent activation and inhibition of arylalkylamine N-acetyltransferase by phosphoserine-205. *Proc Natl Acad Sci U S A* 102, 1222-1227

Gendron JM, Wang ZY. (2007) Multiple mechanisms modulate brassinosteroid signaling. *Curr Opin Plant Biol* 10, 436-441

Grondal E J, Evers R and Cornelissen A W. (1990) Identification and sequence analysis of the ribosomal DNA promoter region of *Crithidia fasciculata*. *Nucleic Acids Research* 18(6), 1333-1338

Guo L, Yu Y, Xia X and Yin W. (2010) Identification and functional characterisation of the promoter of the calcium sensor gene CBL1 from the xerophyte *Ammopiptanthus mongolicus*. *BMC Plant Biol* 10:18

Harper JF, Manney L and Sussman MR. (1994) The plasma membrane H⁺-ATPase gene family in *Arabidopsis*: genomic sequence of AHA10 which is expressed primarily in developing seeds. *Mol Gen Genet* 244, 572-587

Harthill JE, Meek SE, Morrice N, Peggie MW, Borch J, Wong BH and MacKintosh C. (2006) Phosphorylation and 14-3-3 binding of *Arabidopsis* trehalose-phosphate synthase 5 in response to 2-deoxyglucose. *Plant J* 47, 211-223

Hirsch S, Aitken A, Bertsch U and Soll J. (1992) A plant homologue to mammalian brain 14-3-3 protein and protein kinase C inhibitor. *FEBS Lett* 296, 222-224

Huber S C, MacKintosh C and Kaiser W M. (2002) Metabolic enzymes as targets for 14-3-3 proteins. *Plant Mol Biol* 50, 1053-1063

Ichimura T, Isobe T, Okuyama T, Takahashi N, Araki K, Kuwano R and Takahashi Y. (1988) Molecular cloning of cDNA coding for brain-specific 14-3-3 protein, a protein kinase-dependent activator of tyrosine and tryptophan hydroxylases. *Proc Natl Acad Sci USA* 85, 7084-7088

Igarashi D, Ishida S, Fukazawa J and Takahashi Y. (2001) 14-3-3 proteins regulate intracellular localization of the bZIP transcriptional activator RSG. *Plant Cell* 13, 2483-2497

Ishida S, Fukazawa J, Yuasa T and Takahashi Y. (2004) Involvement of 14-3-3 signaling protein binding in the functional regulation of the transcriptional activator REPRESSION OF SHOOT GROWTH by gibberellins. *Plant Cell* 16, 2641-2651

Jahn T, Fuglsang A T, Olsson A, Bruntrup I M, Collinge D B, Volkmann D, Sommarin M, Palmgren M G and Larsson C. (1997) The 14-3-3 protein interacts directly with the C-terminal region of the plant plasma membrane H⁺-ATPase. *Plant Cell* 9, 1805-1814

Jefferson RA, Kavanagh TA and Bevan MW. (1987) GUS fusions: β -glucuronidase as a sensitive and versatile gene fusion marker in higher plants. *EMBO* 6, 3901-3907

Johnson C, Crowter S, Stafford M, Campbel D, Toth R and MacKintosh C. (2010) Bioinformatic and experimental survey of 14-3-3-binding sites. *Biochem J* 427, 69-78

Jones D H, Ley S, Aitken A. (1995) Isoforms of 14-3-3 protein can form homo- and heterodimers in vivo and in vitro: implications for function as adapter proteins. *FEBS Lett* 368, 55-58

Korthout H A and de Boer A H. (1994) A fusicoccin binding protein belongs to the family of 14-3-3 brain protein homologs. *Plant Cell* 6, 1681-1692

Kubala M, Obsil T, Obsilova V, Lansky Z and Amler. (2004) Protein modelling combined with spectroscopic techniques: an attractive quick alternative to obtain structural information. *Physiol Res* 53 Suppl 1, 187-197

Kulma A, Villadsen D, Campbell DG, Meek SE, Harthill JE, Nielsen TH and MacKintosh C. (2004) Phosphorylation and 14-3-3 binding of Arabidopsis 6-phosphofructo-2-kinase/fructose-2,6-bisphosphatase. *Plant J* 37, 654-667

Kuromori T and Yamamoto M. (2000) Members of the Arabidopsis 14-3-3 gene family trans-complement two types of defects in fission yeast. *Plant Science* 158, 155-161

Liu D, Bienkowska J, Petosa C, Collier R J, Fu H, Liddington R. (1995) Crystal structure of the zeta isoform of the 14-3-3 protein. *Nature* 376, 191-194

Lu G, DeLisle A J, de Vetten N C and Ferl R J. (1992) Brain proteins in plants: an Arabidopsis homolog to neurotransmitter pathway activators is part of a DNA binding complex. *Proc Natl Acad Sci U S A* 89, 11490-11494

Lu G, de Vetten N C, Sehnke P C, Isobe T, Ichimura T, Fu H, van Heusden G P and Ferl R J. (1994) A single Arabidopsis GF14 isoform possesses biochemical characteristics of diverse 14-3-3 homologues. *Plant Mol Biol* 25, 659-667

MacKintosh C. (2004) Dynamic interactions between 14-3-3 proteins and phosphoproteins regulate diverse cellular processes. *Biochem J* 381, 329-342

Marra M, Fullone M R, Fogliano V, Pen J, Mattei M, Masi S and Aducci P. (1994) The 30-kilodalton protein present in purified fusicoccin receptor preparations is a 14-3-3-like protein. *Plant Physiol* 106, 1497-1501

Moore B W and Perez V J. (1967) Specific acid proteins in the nervous system. *Physiological and biochemical aspects of nervous integration*. Carlson, F. D. Ed 343-359

Moorhead G, Douglas P, Cotelle V, Harthill J, Morrice N, Meek S, Deiting U, Stitt M, Scarabel M, Aitken A and MacKintosh C. (1999) Phosphorylation-dependent interactions between enzymes of plant metabolism and 14-3-3 proteins. *Plant J* 18, 1-12

Moorhead G, Douglas P, Morrice N, Scarabel M, Aitken A and MacKintosh C. (1996) Phosphorylated nitrate reductase from spinach leaves is inhibited by 14-3-3 proteins and activated by fusicoccin. *Curr Biol* 6, 1104-1113

Morsomme P and Boutry M. (2000) The plant plasma membrane H⁺-ATPase: structure, function and regulation. *Biochim Biophys Acta* 1465, 1-16

Muslin A J and Xing H. (2000) 14-3-3 proteins: Regulation of subcellular localization by molecular interference. *Cell Signal* 12, 703-709

Muslin A J, Tanner J W, Allen P M, Shaw A S. (1996) Interaction of 14-3-3 with signaling proteins is mediated by the recognition of phosphoserine. *Cell* 84, 889-897

Obsil T, Ghirlando R, Klein D C, Ganguly S and Dyda F. (2001) Crystal structure of the 14-3-3 ζ :serotonin N-acetyltransferase complex: A role for scaffolding in enzyme regulation. *Cell* 105, 257-267

- Obsilova V, Herman P, Vecer J, Sulc M, Teisinger J, Obsil T. (2004) 14-3-3 ζ C-terminal stretch changes its conformation upon ligand binding and phosphorylation at Thr232. *J Biol Chem* 279, 4531-4540
- Oecking C, Eckerskorn C and Weiler E W. (1994) The fusicoccin receptor of plants is a member of the 14-3-3 superfamily of eukaryotic regulatory proteins. *FEBS Lett* 352, 163-166
- Oecking C and Jaspert N. (2009) Plant 14-3-3 proteins catch up with their mammalian orthologs. *Curr Opin Plant Biol* 12, 760-765
- Oecking C, Piotrowski M, Hagemeyer J and Hagemann K. (1997) Topology and target interaction of the fusicoccin-binding 14-3-3 homologs of *Commelina communis*. *Plant J* 12, 441-453
- Olsson A, Svennelid F, Ek B, Sommarin M, Larsson C. (1998) A phosphothreonine residue at the C-terminal end of the plasma membrane H⁺-ATPase is protected by fusicoccin-induced 14-3-3 binding. *Plant Physiol* 118, 551-555
- Ossareh-Nazari B, Bachelerie F, Dargemont C. (1997) Evidence for a role of CRM1 in signal-mediated nuclear transport export, *Science* 278, 141-144
- Palmgren MG, Sommarin M, Serrano R and Larsson C. (1991) Identification of an autoinhibitory domain in the C-terminal region of the plant plasma membrane H⁺-ATPase. *J Biol Chem* 266 (30), 20470-20475
- Palmgren M G. (2001) Plant plasma membrane H⁺-ATPase: Powerhouses for nutrient uptake. *Annu Rev Plant Mol Biol* 52, 817-845
- Paul AL, Folta KM and Ferl RJ. (2008) 14-3-3 proteins, red light and photoperiodic flowering. A point of connection? *Plant signaling & behavior* 3, 511-515

Paul AL, Liu L, McClung S, Laughner B, Chen S and Ferl RJ. (2009) Comparative interactomics: Analysis of Arabidopsis 14-3-3 complexes reveals highly Conserved 14-3-3 interactions between humans and plants. *J Prot Res* 8, 1913-1924

Paul AL, Sehnke P C and Ferl R J. (2005) Isoform-specific subcellular localization among 14-3-3 proteins in Arabidopsis seems to be driven by client interactions. *Mol Biol Cell* 16, 1735-1743

Rittinger K, Budman J, Xu J, Volinia S, Cantley L C, Smerdon S J, Gamblin S J, Yaffe M B. (1999) Structural analysis of 14-3-3 phosphopeptide complexes identifies a dual role for the nuclear export signal of 14-3-3 in ligand binding. *Mol Cell* 4, 153-166

Roberts M R. (2000) Regulatory 14-3-3 protein-protein interactions in plant cells. *Curr Opin Plant Biol* 3, 400-405

Roberts M R. (2003) 14-3-3 proteins find new partners in plant cell signaling. *Trends Plant Sci* 8, 218-223

Rosenquist M, Alsterfjord M, Larsson C, Sommarin M. (2001) Data mining the Arabidopsis genome reveals fifteen 14-3-3 genes: Expression is demonstrated for two out of five novel genes. *Plant Physiol* 127, 142-149

Rosenquist M, Sehnke P, Ferl R J, Sommarin M and Larsson C. (2000) Evolution of the 14-3-3 protein family: does the large number of isoforms in multicellular organisms reflect functional specificity? *J Mol Evol* 51, 446-458

Ryu H, Kim K, Cho H, Park J, Choe S and Hwang I. (2007) Nucleocytoplasmic shuttling of BZR1 mediated by phosphorylation is essential in Arabidopsis brassinosteroid signaling. *Plant Cell* 19, 2749-2762

Schultz T F, Medina J, Hill A and Quatrano R S. (1998) 14-3-3 proteins are part of an abscisic acid-VIVIPAROUS1 (VP1) response complex in the Em promoter and interact with VP1 and EmBP1. *Plant Cell* 10, 837-847

Sehnke P C, Chung H J, Wu K and Ferl R J. (2001) Regulation of starch accumulation by granule-associated plant 14-3-3 proteins. *Proc Natl Acad Sci U S A* 98, 765-770

Sehnke P C, Henry R, Cline K and Ferl R J. (2000) Interaction of a plant 14-3-3 protein with the signal peptide of a thylakoid-targeted chloroplast precursor protein and the presence of 14-3-3 isoforms in the chloroplast stroma. *Plant Physiol* 122, 235-242

Shen W, Clark A C and Huber S C. (2003) The C-terminal tail of Arabidopsis 14-3-3 ω functions as an autoinhibitor and may contain a tenth α -helix. *Plant J* 34, 473-484

Silhan J, Obsilova V, Vecer J, Herman P, Sulc M, Teisinger J and Obsil T. (2004) 14-3-3 protein C-terminal stretch occupies ligand binding groove and is displaced by phosphopeptide binding. *J Biol Chem* 279, 49113-49119

Sinnige M P, Roobeek I, Bunney T D, Visser A J W G, Mol J N M, and de Boer A H. (2005) Single amino acid variation in barley 14-3-3 proteins leads to functional isoform specificity in the regulation of nitrate reductase. *Plant J* 44, 1001-1009

Stade K, Ford C S, Guthrie C, Weis K. (1997) Exportin 1 (Crm1) is an essential nuclear export factor. *Cell* 90, 1041-1050

Sullivan S, Thomson CE, Kaiserli E and Chrisitie JM. (2009) Interaction specificity of Arabidopsis 14-3-3 proteins with phototropin receptor kinase. *FEBS Lett* 583, 2187-2193

Svennelid F, Olsson A, Piotrowski M, Rosenquist M, Ottman C, Larsson C, Oecking C, Sommarin M. (1999) Phosphorylation of Thr-948 at the C terminus of the plasma membrane H⁺-ATPase creates a binding site for the regulatory 14-3-3 protein. *Plant Cell* 11, 2379-2391

Takahashi Y. (2003) The 14-3-3 proteins: gene, gene expression and function. *Neurochem Res* 28, 1265-1273

Todd A, Cossons N, Aitjen A, Price G B and Zannis-Hadjopoulos M. (1998) Human cruciform binding protein belongs to the 14-3-3 family. *Biochemistry* 37, 14317-14325

Toker A, Ellis C A, Sellers L A and Aitken A. (1990) Protein kinase C inhibitor proteins. Purification from sheep brain and sequence similarity to lipocortins and 14-3-3 protein. *Eur J Biochem* 191, 421-429

Toroser D, Athwal G S and Huber S C. (1998) Site-specific regulatory interaction between spinach leaf sucrose-phosphate synthase and 14-3-3 proteins. *FEBS Lett* 435, 110-114

Truong A B, Masters S C and Yang H, Fu H. (2002) Role of the 14-3-3 C-terminal loop in ligand interaction. *Proteins* 49, 321-325

van Heusden G P, Griffiths D J, Ford J C, Chin A W T F, Schrader P A, Carr A M and Steensma H Y. (1995) The 14-3-3 proteins encoded by the BMH1 and BMH2 genes are essential in the yeast *Saccharomyces cerevisiae* and can be replaced by a plant homologue. *Eur J Biochem* 229, 45-53

van Heusden G P, Wenzel T J, Legendijk E L, Steensma H Y and van den Berg J A. (1992) Characterization of the yeast BMH1 gene encoding a putative protein homologous to mammalian protein kinase II activators and protein kinase C inhibitors. *FEBS Lett* 302, 145-150

van Zeijl M J, Testerink C, Kijne J W, Wang M. (2000) Subcellular differences in post-translational modification of barley 14-3-3 proteins. *FEBS Lett* 473, 292-296

Visconti S, Camoni L, Fullone M R, Lalle M, Marra M and Aducci P. (2003) Mutational analysis of the interaction between 14-3-3 proteins and plant plasma membrane H⁺-ATPase. *J Biol Chem* 278, 8172-8178

Watanabe M, Isibe T, Ishimura T, Kuwano R, Takahashi Y and Kondo H. (1993) Molecular cloning of rat cDNA for beta and gamma subtypes of 14-3-3 protein and developmental changes in expression of their mRNAs in the nervous system. *Mol Brain Res* 17, 135-146

Weiner H and Kaiser W M. (1999) 14-3-3 proteins control proteolysis of nitrate reductase in spinach leaves. *FEBS Lett* 455, 75-78

Winter D, Ben Vinegar B, Nahal H, Ammar R, Wilson GV, Provart NJ. (2007) An “Electronic Fluorescent Pictograph” Browser for Exploring and Analyzing Large-Scale Biological Data Sets. *PLoS One* 2, 718-726

Würtele M, Jelich-Ottmann C, Wittinghofer A, Oecking C. (2003) Structural view of a fungal toxin acting on a 14-3-3 regulatory complex. *EMBO J* 22, 987-994

Xiao B, Smerdon S J, Jones D H, Dodson G G, Soneji Y, Aitken A, Gamblin S J. (1995) Structure of a 14-3-3 protein and implications for coordination of multiple signalling pathways. *Nature* 376, 188-191

Yaffe M B. (2002) How do 14-3-3 proteins work? – Gatekeeper phosphorylation and the molecular anvil hypothesis. *FEBS Lett* 513, 53-57

Yaffe M B, Rittinger K, Volinia S, Caron P R, Aitken A, Leffers H, Gamblin S J, Smerdon S J, Cantley L C. (1997) The structural basis for 14-3-3:phosphopeptide binding specificity. *Cell* 91, 961-971

Yao Y, Du Y, Jiang L and Liu J. (2007) Molecular analysis and expression patterns of the 14-3-3 gene family in *Oryza sativa*. *J Biochem Mol Biol* 40, 349-357

