

The effects of VDT data entry work on operators

Gao, Chuansi; Lu, Demao; She, Qiyuan; Cai, Rongtai; Yang, Lei; Zhang, Guogao Published in: **Ergonomics**

1990

Link to publication

Citation for published version (APA): Gao, C., Lu, D., She, Q., Cai, R., Yang, L., & Zhang, G. (1990). The effects of VDT data entry work on operators. Ergonomics, 33(7), 917-924.

Total number of authors:

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.

 • You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: https://creativecommons.org/licenses/

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

The effects of VDT data entry work on operators

Abstract

The before/after study of physiological and biochemical parameters was used to delineate the effects of VDT data entry work on operators. Twenty-nine healthy Chinese students were chosen and divided at random into the simple and the complicated data entry group. The subjects were instructed to work as quickly and correctly as possible according to the 'Data Entry Work Programme' for 150min. Work performance (correct entry) was automatically recorded once every IOmin. The before/after parameters were tested respectively. The results showed that performance fluctuated over time. It decreased obviously after 50–60 min of work, followed by a rebound, and there was a terminal motivation phenomenon at the end of the test, which was associated with the auto-arousal and cerebral compensatory effort. Changes in physiological parameters revealed that operators were fatigued after data entry work. The adrenaline excretion in urine showed a tendency to increase after simple data entry work. The noradrenaline excretion showed a tendency to decrease after complicated data entry work. The differences in performance, diastolic blood pressure in a standing position and neurobehaviour between two groups indicated that much stress was experienced when performing complicated data entry work.