



LUND UNIVERSITY

On the occurrence of the Alpine Swift *Apus melba* in Nigeria.

Tobler, Michael; Naurin, Sara

Published in:
Malimbus

2008

[Link to publication](#)

Citation for published version (APA):

Tobler, M., & Naurin, S. (2008). On the occurrence of the Alpine Swift *Apus melba* in Nigeria. *Malimbus*, 30(2), 167-168.

Total number of authors:

2

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

- MOREAU, R.E. (1972) *The Palaearctic-African Bird Migration Systems*. Academic Press, London.
- MOREL, G.J. & MOREL, M.-Y. (1988) Liste des oiseaux de Guinée. *Malimbus* 10: 143–176.
- MOREL, G.J. & MOREL, M.-Y. (1990) *Les Oiseaux de Sénégal*. ORSTOM, Paris.
- RIEGERT, J., SEDLÁČEK, O. & HUTTERER, R. (in press) Diet of sympatric African Grass Owl (*Tyto capensis*) and Spotted Eagle Owl (*Bubo africanus*) in the Bamenda Highlands, NW Cameroon. *Afr. J. Ecol.*
- RODWELL, S.P., SAUVAGE, A., RUMSEY, S.J.R. & BRÄULICH, A. (1996) An annotated checklist of birds occurring at the Parc National des Oiseaux du Djoudj in Senegal, 1984–1994. *Malimbus* 18: 74–111.
- SAUVAGE, A. & RODWELL, S.P. (1998) Notable observations of birds in Senegal (excluding Parc National des Oiseaux du Djoudj), 1984–1994. *Malimbus* 20: 75–122.
- SMALLEY, M.E. (1983) The Marsh Owl *Tyto capensis*: a wet season migrant to the Gambia. *Malimbus* 5: 31–33.
- VELMALA, W. & GUSTAFSSON, R. (2003) Two new raptors for Nigeria and other raptor observations at Lake Chad. *Malimbus* 25: 52–55.

Received 23 February 2008

Revised 6 May 2008

Jan Riegert¹, Marcin Antczak² & Drahomíra Fainová¹

¹Univ. of South Bohemia, Faculty of Science, Dept of Zoology, Branišovská 31, 370 05 České Budějovice, Czech Republic

²Adam Mickiewicz Univ., Dept of Behavioural Ecology, Umultowska 89, PL-61 614 Poznań, Poland

On the occurrence of the Alpine Swift *Apus melba* in Nigeria

The Alpine Swift *Apus melba* is a not uncommon to rare Palaearctic winter visitor to W Africa (Keith *et al.* 1988), whose winter distribution is not adequately known in the region (Borrow & Demey 2001). For Nigeria, the known distribution extends only to about 7° N. However, there are two records from Cameroon, at 4–5° N, close to the Nigerian border (Borrow & Demey 2001). Here, we add four observations of large flocks of Alpine Swifts from the Cross River National Park, in SE Nigeria, suggesting that the species may regularly winter in this region. Observations were part of an ornithological survey in the Okwangwo Division of the Cross River NP, conducted in the surroundings of Bashu Okpambe village (c. 6°6'N, 9°8'E) from 1 to 26 Nov 2006.

Our observations are as follows (with minimum number of birds recorded): two flocks, of 40 and 30 birds, 16 Nov; 50 birds, 20 Nov; five birds, 24 Nov. The first and

the third observations were made on hilltops (c. 200–300 m above sea level), whereas the others were of birds hunting over the village and its adjacent farmland. Identification of the species was based on size (considerably larger than the other swifts they were with, see below), the characteristic call (typical trilling call, well known to the authors from experience in central Europe), plumage characteristics (overall brown with a white throat patch and a large white patch on the belly; the two white patches separated by a smaller brown band) and flight (relatively slow wing beats compared to the smaller swifts). The larger flocks on 16 and 20 Nov also contained 1–5 Little Swifts *Apus affinis* and 1–5 Common or African Black Swifts *A. apus* or *A. barbatus*. As it has recently been suggested that the Alpine Swift may not actually breed in W Africa (Dowsett & Dowsett-Lemaire 2005), we assume that the observed birds were Palearctic visitors from NW Africa or Europe. We do not know whether this species occurs in this region throughout the winter.

We are grateful to the Swedish Ornithological Society, Ångpanneföreningen and Mr A.P. Leventis for financing the survey, and Drs Phil Hall and Ulf Ottosson for helping with logistics. This is publication 20 from the A.P. Leventis Ornithological Institute.

References

- BORROW, N. & DEMEY, R. (2001) *Birds of Western Africa*. Christopher Helm, London.
- DOWSETT, R.J. & DOWSETT-LEMAIRE, F. (2005) Additions to the avifauna of Mali. *Bull. Afr. Bird Club* 12: 119–124.
- KEITH, S., URBAN, E.K. & FRY, C.H. (1988) *The Birds of Africa*, vol. 3. Academic Press, London.

Received 14 February 2007

Revised 26 November 2007

Michael Tobler & Sara Naurin

Dept of Animal Ecology, Lund University, Ecology Building, 223 62 Lund, Sweden
<michael.tobler@zooekol.lu.se>, <sara.naurin@zooekol.lu.se>

African Barred Owlet *Glaucidium capense* new to Togo

Between 21h00 and 22h00 on 25 July 1989 I heard a series of calls noted as “drrrr-drrrr repeated many times” at Djodji (7°40'N, 0°35'E), Togo. The calls, often repeated during the hour I listened, were coming from the interior of thick secondary forest beside a track leading to the River Gban-Houa, also known as the River Wawa, where it acts as the border between Togo and Ghana. The forested area was populated with trees such as *Antiaris africana* and *Chlorophora excelsa*, but was patchy with some areas being logged. At the time I was unable to identify the memorable calls, which I