



# LUND UNIVERSITY

## Continuous multi-plot measurements of CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O and H<sub>2</sub>O in a managed boreal forest : the importance of accounting for all greenhouse gases

Vestin, Patrik; Mölder, Meelis; Sundqvist, Elin; Båth, Anders; Lehner, Irene; Weslien, Per; Klemedtsson, Leif; Lindroth, Anders

2015

[Link to publication](#)

### *Citation for published version (APA):*

Vestin, P., Mölder, M., Sundqvist, E., Båth, A., Lehner, I., Weslien, P., Klemedtsson, L., & Lindroth, A. (2015). *Continuous multi-plot measurements of CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O and H<sub>2</sub>O in a managed boreal forest : the importance of accounting for all greenhouse gases*. Abstract from AGU (American Geophysical Union) Fall Meeting, 2015, San Francisco, United States.

### *Total number of authors:*

8

### **General rights**

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117  
221 00 Lund  
+46 46-222 00 00

## **Continuous multi-plot measurements of CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O and H<sub>2</sub>O in a managed boreal forest – The importance of accounting for all greenhouse gases**

Patrik Vestin<sup>1</sup>, Meelis Mölder<sup>1</sup>, Elin Sundqvist<sup>1</sup>, Anders Båth<sup>2</sup>, Irene Lehner<sup>2</sup>, Per Weslien<sup>3</sup>, Leif Klemetsson<sup>3</sup>, Anders Lindroth<sup>1</sup>

<sup>1</sup>Department of Physical Geography and Ecosystem Science, Lund University, Sweden, <sup>2</sup>Centre for Environmental and Climate Research, Lund University, Sweden, <sup>3</sup>Department of Earth Sciences, Gothenburg University, Sweden

In order to assess the effects of different management practices on the exchange of greenhouse gases (GHG), it is desirable to perform repeated and parallel measurements on both experimental and control plots. Here we demonstrate how a system combining eddy covariance and gradient techniques can be used to perform this assessment in a managed forest ecosystem.

The net effects of clear-cutting and stump harvesting on GHG fluxes were studied at the ICOS site Norunda, Sweden. Micrometeorological measurements (i.e., flux-gradient measurements in 3 m tall towers) allowed for quantification of CO<sub>2</sub>, CH<sub>4</sub> and H<sub>2</sub>O fluxes (from May 2010) as well as N<sub>2</sub>O and H<sub>2</sub>O fluxes (from June 2011) at two stump harvested plots and two control plots. There was one wetter and one drier plot of each treatment. Air was continuously sampled at two heights in the towers and gas concentrations were analyzed for CH<sub>4</sub>, CO<sub>2</sub>, H<sub>2</sub>O (LGR DLT-100, Los Gatos Research) and N<sub>2</sub>O, H<sub>2</sub>O (QCL Mini Monitor, Aerodyne Research). Friction velocities and sensible heat fluxes were measured by sonic anemometers (Gill Windmaster, Gill Instruments Ltd). Automatic chamber measurements (CO<sub>2</sub>, CH<sub>4</sub>, H<sub>2</sub>O) were carried out in the adjacent forest stand and at the clear-cut during 2010.

Average CO<sub>2</sub> emissions for the first year ranged between 14.4–20.2 ton CO<sub>2</sub> ha<sup>-1</sup> yr<sup>-1</sup>. The clear-cut became waterlogged after harvest and a comparison of flux-gradient data and chamber data (from the adjacent forest stand) indicated a switch from a weak CH<sub>4</sub> sink to a significant source at all plots. The CH<sub>4</sub> emissions ranged between 0.8–4.5 ton CO<sub>2</sub>-eq. ha<sup>-1</sup> yr<sup>-1</sup>. N<sub>2</sub>O emissions ranged between 0.4–2.6 ton CO<sub>2</sub>-eq. ha<sup>-1</sup> yr<sup>-1</sup>. Enhanced N<sub>2</sub>O emission on the drier stump harvested plot was the only clear treatment effect on GHG fluxes that was observed. Mean CH<sub>4</sub> and N<sub>2</sub>O emissions for the first year of measurements amounted up to 29% and 20% of the mean annual CO<sub>2</sub> emissions, respectively. This highlights the importance of including all GHG when assessing the climate impacts of different forest management options.

Our results show that continuous multi-plot measurements of the main GHGs are possible also at sites where GHG fluxes are low, at a reasonable cost and with reduced plot inter-comparison uncertainties.