



LUND UNIVERSITY

PIXE and Other Applications of Nuclear Physics

Akselsson, Roland

1978

[Link to publication](#)

Citation for published version (APA):

Akselsson, R. (1978). *PIXE and Other Applications of Nuclear Physics*. (Lund Institute of Technology, Department of Nuclear Physics). [Publisher information missing].

Total number of authors:

1

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

LUND INSTITUTE OF TECHNOLOGY

DEPARTMENT OF NUCLEAR PHYSICS

Sölvegatan 14

S-223 62 LUND, Sweden

PIXE AND OTHER APPLICATIONS OF NUCLEAR PHYSICS

R. AKSELSSON AND THE PIXE-GROUP

THE PIXE-PART OF THE REPORT FROM THE DEPARTMENT
OF NUCLEAR PHYSICS, UNIVERSITY OF LUND AND
LUND INSTITUTE OF TECHNOLOGY, SEPTEMBER 1978.

4 PIXE AND OTHER APPLICATIONS OF NUCLEAR PHYSICS

A project at the Department of Nuclear Physics, Lund, Institute of Technology, partly supported by NFR through the contacts

- | | | | |
|----|-----------|------------------|---------------|
| 1. | F7487-004 | Roland Akselsson | salary for RA |
| 2. | F7487-005 | Roland Akselsson | material |

and indirectly through

- | | | | |
|----|----------------|---------------------|--|
| 3. | 02-056, 02-057 | Sven A E Johansson, | salaries and material partly covering the expenses for research at the Pelletron laboratory. |
|----|----------------|---------------------|--|

Table of contents - see appendix 1.

I Introduction

PIXE is the acronym for Particle Induced X-ray Emission Analysis. In this method for trace element analysis, MeV particles bombard the sample to be analysed. Characteristic X-rays are induced, which are detected with an energy-dispersive high-resolution detector, e.g. a Si(Li)-detector. PIXE is a fast multielemental method with very low detection limits in small samples. For elements lighter than sulphur, absorption in the sample makes the quantification uncertain or impossible. However, there are nuclear methods, which may be used simultaneously to PIXE, giving the mass of low-Z elements.

The properties of PIXE mentioned above, including those of complementing nuclear methods, make the PIXE method unique among methods for elemental analysis. New types of important experiments e.g. in the domains of environmental health, biology and technique may be planned and performed, if PIXE is fully developed and used. However, then physicists have to 1) further develop the method, 2) be responsible for the analysis at least in the introductory period and

3) cooperate with scientists from other fields pointing out the possibilities of PIXE and actively taking part in planning, sampling and analysis. The last point is important because PIXE has special properties which make it strong - when those properties are used.

There are very good conditions for a successful PIXE-programme in Lund, if NFR supports an active group of physicists around PIXE.

The activities of the PIXE-group may be divided in the programme for applied nuclear and atomic physics and the programme for technical hygiene. In this report the emphasis is laid on the first programme, as it is the part under expert evaluation.

II History with the most important results

In 1970 the Lund-group was publishing the first brief paper about PIXE /Ref 1/. Another paper from the same group /Ref 3/ pointed out fields of applications. In exploration of the basic properties of PIXE, the variation of K-ionization cross section with protonenergy and atomic number were carefully investigated and a universal expression for these cross sections was derived /Ref 6/. Such a formula is very useful for computer evaluation of masses of elements in thin samples because in calibration of the analysis system the formula allows us to interpolate and extrapolate from a few elements analyzed at one energy. A key component in the analysis is the computer unravelling of the pulse height spectrum. This procedure should be fast, accurate and should normally not require interaction with an operator. Kaufmann and Akselsson designed a first version of such a programme, which used a physical model of the processes taking place in the detector /Ref 34/. The universal formula is still more important for thick sample analysis, a field also thoroughly evaluated at our laboratory /Ref 20/. Many important procedures for accurate analysis have been presented e g beam mapping /Ref 2/, elimination of charging /Ref 25/ and mass

calibration /Ref 70/. The unique possibilities offered by an accelerator laboratory in allowing simultaneous use of particle elastic scattering analysis, nuclear reaction and PIXE have been partly explored /Refs 5, 27, 87, 101/ and already the $^{19}\text{F}(\text{p},\alpha\gamma)^{16}\text{O}$ reactions is routinely in use simultaneous to PIXE in our laboratory. An extensive review of PIXE is given by Johansson and Johansson /Ref 50/.

As seen from the list of publications (Chapter VI), a considerable amount of work has been carried out by members of the PIXE-group in Lund concerning transport of airborne particulates and concerning welding aerosols. In these fields multielemental analysis of many samples is necessary and it is a great advantage, if small samples are sufficient for accurate results. Thus, PIXE has opened new possibilities for important research in fields of environmental hygiene /Refs 8, 9, 14, 15, 19, 28, 42, 43, 49, 51, 52, 56, 59, 62, 78, 81-84, 86-88, 102, 105, 106, 110, 112, 113/.

A single orifice cascade impactor has been found which gives size-fractionated aerosol samples, suitable for PIXE-analyses. The impactor is of Battelle design, however it has been modified to include one impactor stage more, to include an after-filter for the smallest particles ($<0.25\mu\text{m}$ aerodynamic diameter), and to have two stages rotating eccentrically in order to spread the sample on a larger area.

Another sampler, which has been used and will be used, is the "streaker". It is a long filter with a moving, small, sucking orifice arranged to give time resolution (2h resolution and about one week per sample). This sampler is designed at Florida State University especially for PIXE analysis.

In aerosols generated in welding on stainless steel insoluble chromium VI is a key factor for health effects. In the project "Characterization of Welding Aerosols", we have developed a procedure for quantification of soluble and insoluble chromium VI as well as soluble and insoluble chromium III /Ref 108/. ESCA, TEM, the DPC-method and PIXE are used in this procedure.

Although the accelerator change in Lund 1974-1976 hampered our activities considerably for almost three years, we now have a laboratory well suited for applications. The status of current arrangement is reported in appendix 2.

Many useful international contacts have been created. The First International Conference on PIXE and its Applications was held in Lund, August 23-26, 1976. Many researchers from different parts of the world have visited our laboratory for longer or shorter periods. Likewise, members of the PIXE-group have visited many PIXE laboratoires and participated in many conferences.

III Economy 77/78

The activities of the PIXE-group were supported 77/78 by

1. NFR (The Swedish National Science Research Council) by contract

F 7487-004 - salary for Roland Akselsson	100 kkr
F 7487-005 - material, travels	6 kkr

and indirectly by

- | | |
|--|---------|
| 02-056 - research at the Pelletron and VdG laboratories in Lund | |
| 2. ASF (The Swedish Work Environmental Fund)
74/109:5 - airborne particulates from welding operations | 305 kkr |
| 3. SNV (The National Swedish Environment Protection Board) | |
| 7-147/77 - aerosol composition | 75 kkr |
| 7-303/77 - multivariate statistical analysis of aerosol data | 25 kkr |
| 4. Miscellaneous sources (for special applications) approximatively | 60 kkr |

In addition, basic facilities and some important service functions from the University of Lund have been used. The Department of Environmental Health has contributed with salaries corresponding to about 1 man/year. The lack of support for work

IV Status, September 1978

a) Persons engaged in the PIXE-group

Member of the PIXE-group	Titles	Engagement in PIXE-work	Main source of salary *
Mats Ahlberg	Tekn dr	1/2	Dept of Env Health
Roland Akxelsson	Ph D	1/1	NFR
Erling Andersson	-	parttime	-
Mats Bohgard	Civ ing	1/1	PIXE-group
Lars Eric Carlsson	Civ ing	1/1	PIXE-group
Hans Christer Hansson	Civ ing	1/1	SNV
Eva Marta Johansson	-	1/2	PIXE-group
Gerd Johansson	Civ ing	1/1	ASF
Katarina Johansson	B S	1/2	ASF
Sven Johansson	Ph D	parttime	LTH
Erik Karlsson	Ing	parttime	LTH
Hans Lannefors	Civ ing	1/1	SNV
Klas Malmqvist	Civ ing	1/1	ASF
Bengt Martinsson	-	parttime	-
Jan Pallon	Civ ing	1/2	PIXE-group
Bertil Rudell	Civ ing	1/2	Dept of Env Health
Knut Sjöberg	Staff (mechanics)	3/4	LTH
Leif Spanier	Civ ing	-	-

* NFR Swedish Natural Science Research Council

SNV National Swedish Environment Protection Board

ASF Swedish Work Environment Fund

LTH Lund Institute of Technology

Only Professor Sven Johansson, Mr Erik Karlsson and Mr Knut Sjöberg have permanent positions.

b) Experimental arrangements - see appendix 2.

c) Applications

PIXE is a method especially suitable for detecting small amounts of elements in small samples. Other features of PIXE are its multielemental character and its speed. These properties make it very favourable to use PIXE in investigations of airborne particles, because in field measurements it is desirable to have small sampling devices and because often a large number of samples has to be analysed for several elements. In projects supported by the Swedish Work Environmental Fund, the National Swedish Environment Protection Board and others, 1 l/min single-orifice cascade impactors are used for size fractionation.

Analysis of trace elements in water is another field of great promise. Some preliminary investigations indicate that detection limit in the 0.01 - 1 ppb range may be obtained for several elements. Procedures have to be optimized and consolidated.

Biological samples constitute a third type of samples which may be analysed by PIXE. In an investigation concerning the lead content in teeth of children, the possibility to guarantee a narrow beam has proved to be very advantageous.

Summaries of several applied projects in progress are given in appendix 3.

V Plans (Letter of intent)

a) Objectives of the PIXE-group in Lund

The objectives of the PIXE-group is to develop and apply the unique possibilities, which an accelerator laboratory

offers for trace elemental analyses of high priority in fields as technology, environmental hygiene, medicine and biology. Another important goal is to train physicists. The volume of this training-programme has to be in some harmony with the labour-market for doctors.

The activities of the PIXE-group may be divided in two programmes - on one side "PIXE research and development" and on the other side "Technical Hygiene". The same persons, with a few exceptions, are engaged in the two programmes today, but as the tasks get more complex, it may be feasible and necessary with further specialization.

b) The programme for PIXE research and development

In the following fourteen paragraphs fourteen important projects are briefly presented.

1. One of the major advantages with PIXE compared to other X-ray methods, is the possibility of simultaneous backscattering analysis (PESA - Particle Elastic Scattering Analysis). The mass resolution of PESA is not good for 2.5 MeV protons, but it improves considerably if α -particles are used. Such a change should not significantly impair the properties of PIXE. Backscattering would increase the number of analysed elements per run and also give information about the total mass of the sample. This information (mass/cm^2) is also of great value in evaluating the PIXE-spectrum. During the autumn 1978, α -particles may be obtained from the Pelletron. Then we plan to optimize the α -particle energy and the analyzing arrangement for combined PIXE/PESA analysis. The usefulness of other nuclear reaction will also be investigated. There is also a challenging task in developing a computer-programme for fast and accurate unravelling of backscattering pulse-height spectra.

2. Another unique advantage of PIXE is the possibility of focusing the beam to obtain low detection limits in $10\mu\text{m}$ samples. By cooperation with the Department of Physical Biology in Uppsala, a microbeam arrangement is planned to be set up during 1978/79 in Lund. The halo-effect will be studied and minimized. Feasibility tests for several applications are also planned.

3. For some applications the resolution of semiconductor detectors is not sufficiently good. Then at low X-ray energies crystal spectrometers may be useful. Design and construction of a curved crystal spectrometer has started and will be finished during 1979. The first version will be used to study L- and M-line relative intensities in the 3-7 keV range. These values are crucial for the resolution of pulse-height spectra. The variation of these intensities as a function of particle energy is also interesting from a theoretical point of view. The crystal spectrometer may also be used to resolve interferences in routine PIXE/PESA analysis. The combination of microbeam and curved crystal may offer further advantages, since crystal spectrometry is favourable for point sources.

4. With another crystal still lower X-ray energies may be detected. Then it will be interesting to investigate energy shifts and identify them with the chemical compounds of the sample. It is often very important to know the chemical compounds in samples especially in biological samples and it may be possible to obtain some information from PIXE-runs about that.

5. Bremsstrahlung and characteristic X-rays from the backing is often a limiting factor. By using crystalline material as backing and by mounting the sample in certain angles these bothering contributions may be significantly reduced. Some preliminary investigations will be done during 78/79.

6. X-ray absorption and particle slowing-down effects may be significant and good corrections are crucial for the accuracy of analysis. Helpful information may be obtained from the continuous part of the X-ray spectra and from the PESA-spectrum. Such information may also be obtained in separate runs, where the transmission of the sample for certain X-ray energies is measured. We plan to do systematic investigation of absorption and slowing-down effects.

7. Today PIXE is used mostly at proton energies above 1.5 MeV. However, although the cross sections are lower at lower energies, it may be feasible to use lower proton energies in certain cases, e g for elements having X-rays in the 2-3 keV range. Also smaller accelerators may be used, which implies lower cost and a wider use of PIXE. Since there is a lack of basic information about cross sections for X-ray production and bremsstrahlung production for 0.6-1.2 MeV protons, we plan to investigate these areas and survey the properties of PIXE at low proton energies.

8. A larger detector is ordered and a closer geometry will be arranged in order to achieve lower detection limits with less intense beams.

9. An on-line computer and an off-line computer will be ordered during the fiscal year 78/79. Then there will be a strong need for designing procedures for efficient analysis giving maximum information per time unit. Almost all the projects mentioned above (IV b 1-8) involve computer routines. The count rate from the different detectors (X-ray detector, particle detector and the crystal spectrometer) should be optimized automatically and the parameters registered by the computer. All the information obtained should be synthesized in the final calculations of the sample composition. For some applications it is crucial to implement further effects in the physical model, which forms the base for the computer programme currently used for pulseheight spectrum resolution. Such effects are e.g. particle slowing-down, pile-up tails on the peaks in the spectra.

10. By mixing complexbinding compounds into water, adsorb the complexes on active carbon, filter off the carbon and make pellets of the carbon, detection limits in the range of 0.01 to 1 ppb are obtained for several elements. We plan to optimize the technique for different kind of water and measure the yield for different element. Also, it would be desirable to modify the technique, as only small volumes

of water is needed, for other liquids in limited supply.

11. Detection limits on small samples of proteins should be investigated since 1) many biochemical fractionating techniques give small amounts of proteins in certain fractions, for which elemental information is of great interest 2) in the study of metabolism as well as in biological sampling (an interesting manner to obtain information about health hazards due to exposure), elemental analysis of small protein samples is often required.

12. Elemental concentrations in saliva from exposed welders will be analysed to investigate the feasibility of saliva as an indicator of retention.

13. Microbeam applications are planned in cooperation with the Department of Physical Biology, Uppsala (Ulf Lind - biological samples) and FOA (Lars Henrik Andersson - 10 μ m particles).

14. An investigation of the applicability of nuclear and X-ray techniques to analysis of mineralogic samples is planned in cooperation with Svenska Gruvföreningen and Boliden Metall AB.

c) The programme for technical hygiene

As this part is outside the area under evaluation, only a brief listing of planned projects will be given.

1. We plan to test and further modify our cascade impactor in order to get the loading on the last stages spread out on a larger area. This would decrease bounce-off effects and also allow longer sampling times without introducing problems with too thick samples. As mentioned above, a prototype is already built and is partly in use.

2. Using our experiences of the Tallahassee-streaker, we plan to design a sampling device giving size fractionation (two fractions) and time resolution. By using PIXE for the analysis it should be possible to design a small, maybe

portable, sampler with 15 minutes resolution for work environment and an outdoor sampler which could be steered by weather conditions or a clock.

3. In order to test the efficiency of sampling devices, a good facility for aerosol generation should be provided.
4. We plan to follow up the project "welding aerosol characterization" with cooperative projects in which welding procedures are optimized. In such optimizations weld quality, economy and work environment should be taken into consideration.
5. Long range transport of airborne particulates.
6. Variation in urban aerosol composition. Sampling strategy will be considered in the first phase.
7. Special applications of trace elemental water analysis.
8. Continued cooperation with the Department of Environmental Health, e g in studies of lung deposition and toxicology.

d) Proposals for NFR

PIXE has been applied to analysis of sampled airborne particulates since several years and the feasibility of PIXE for this kind of samples is well established. As seen above there are several research and developmental projects planned which could further strengthen the power of accelerator-based trace elemental analysis. These methods for trace elemental analysis offer unique characteristics which make new types of important investigations possible.

The potential for further meaningful development is one strong reason for NFR to support the programme for PIXE research and development. Another reason is that such a support would be beneficial and probably necessary for a successful programme for technical hygiene as PIXE still is new as a method for trace element analysis.

To assure a progressing programme for PIXE research and development, we suggest NFR to support a person (e g 1:e forskn ing from July 1980) who would be responsible for the analysing system and the development of computer routines as mentioned in section IV b 9.

For electronics, chemicals, standards, material, travel, temporary hands, we ask for 100 000 Sw cr/year.

There is also a need for at least some basic aerosol research in Sweden, because there are many projects going on in which aerosol sampling and/or measurements take place. The programme of technical hygiene may be well suited as an embryo. Because of symbiotic reactions between the programmes, a development of the technical hygiene programme to include basic aerosol physics would guarantee high quality, ambient and work environment aerosol investigations and an efficient use of the analytical capability of the PIXE-method. We suggest that our visiting committee should urge NFR to investigate these ideas.

Appendix 1Table of contents

PIXE and other applications to Nuclear Physics to trace elemental analysis.

I Introduction

II History with the most important results

III Economy 77/78

IV Status, September 1978

a) Persons engaged in the PIXE-group

b) Experimental arrangements

c) Applications

V Plans (Letter of intent)

a) Objections of the PIXE-group in Lund

b) The programme for PIXE research and development

c) The programme for technical hygiene

d) Proposals for NFR

VI References (this section is transferred to end of the report from the Pelletron laboratory)

Appendix 1 Table of contents

Appendix 2 Experimental arrangements

Appendix 3 Summaries of current applied projects

References 1, 3, 6, 33, 34, 38, 50, 70, 82, 102 and 108 are enclosed.

Appendix 2

Experimental arrangements

In the currently used routines for PIXE analyses at the Pelletron laboratory in Lund, Sweden, 2.5 MeV protons are used. The beam is made uniform by an aluminium diffuser foil, with thickness of 6 μm , placed about 50 cm before the sample. Carbon Collimators of different diameters may be rotated in the beam position, such that a homogeneous beam of a diameter of about 3, 4, 5, 6, 7 and 9 mm may be obtained. X-rays are detected in 135° and the sample is oriented so that the angle between the proton beam and the sample normal as well as the angle between the sample normal and the detected X-rays are 22.5° . The collimator nearest the sample is held at a positive potential (+30V) to keep secondary electrons from the collimator from entering the Faraday cup. An electron suppressor at -50V is inserted between the +30V-collimator and the sample to stop secondary electrons from the sample from escaping backwards through the collimator. The linearity of the current integration has been tested from 0.5 nA to 200 nA.

The X-rays from the sample pass an optional absorber, a 6 μm Mylar window of the sample chamber and a 13 μm Be-window of the detector-house before reaching the detector. The Si(Li)-detector used has an area of 10 mm^2 and a depth of 3 mm. The resolution is about 175 eV at 6.4 keV. The pulseheight spectra recorded on a ND 4420 are written on magnetic tape and unravelled by a computer program, HEX, at the main computer center of Lund (Univac 1108).

Samples are mounted in slideframes (2" x 2") and loaded in a box with space for 40 slides. By a pushbutton at the operator's desk, samples may be changed.

In routine analysis the beamtime per sample is 2 - 4 minutes with a countrate of 1000 pulses/s. To be able to increase the countrate and decrease the beamtime/sample a fast deflection

system has been constructed and it is now being tested. The rise-time of the 1000V deflection pulse is 30 ns.

For biological samples we have extracted the beam to an air or nitrogen filled chamber. A Kaptonfoil with a thickness of 7.5 μ m is used as an exit window.

Fluorine is measured simultaneously with the PIXE-analysis by detection of γ -rays from the reaction $^{19}\text{F}(p,\alpha)^{16}\text{O}$. A 5" x 4" NaI crystal is used and the detection limit is about 50 ng in our set up.

To get information about the low Z elements in the sample a silicon detector for backscattered protons has been mounted in the chamber. It is situated in 155 $^{\circ}$ and the area used is 3 mm 2 . The full width at half maximum of the system for 3.72 MeV protons is 27 keV. The preliminary choice of particle and particle energy for excitation (2.5 MeV protons) may be changed to increase the information from the backscattering.

In analysing thick nonconducting samples their charging introduces a severe increase of bremsstrahlung. A carbon filament with a grid for suppressing positive ions is included in the system. This solves the charging problem without introducing any noticeable contamination.

Appendix 3

Summaries of Current Applied Projects

September 1978

Characterization of welding aerosols

Introduction

The electric arc welding is a common method in industry. Welding aerosols, some of which may be hazardous, are formed during the welding. We are characterizing such aerosols by a reproducible sampling procedure and subsequent elemental analysis. By varying the welding current and voltage, information is gained about the particle formation processes in the welding processes.

Method

The welding aerosol produced is collected in a hood. Samples are taken for elemental and gravimetric analysis as well as particle size analysis. The elemental analysis is performed with the PIXE-method and simultaneous nuclear methods.

Results

The welding parameters play an important role in the aerosol formation and will affect the particle size distribution as well as the elemental composition. The most common elements are F, K, Ca, Ti, Cr, Mn, Fe, Ni and Zn, but the composition

is very dependent on the method used.

As most aerosol particles have an aerodynamic diameter in the interval 0.05-2 μm most of the aerosol may be deposited in the lung with its slow clearance rate. Finally the work so far has demonstrated that the PIXE-method is highly suitable for this kind of work.

Characterization of airborne dust from metal spraying processes

In metal spraying a material is heated to a molten condition and deposited to a workpiece by a high velocity gas stream. It can be used to produce coatings with high corrosion and wear resistance.

The health hazards, due to inhalation of the fume produced by the process, depends on the particle size distribution of the aerosol, the chemical composition of the particles and the dust concentration in the breathing zone of the workers. Metals, which can form cancerogenic compounds, as chromium and nickel, often exceed the hygienic threshold limit values.

PIXE is a superior method to determine the elemental composition of the small size - fractionated samples from a Battelle impactor and its multielemental nature make it possible to reveal unexpected elements in the fume.

When the fume contains chromium, the oxidation state is of importance for toxicity evaluations. Electron Spectroscopy for Chemical Analysis (ESCA) and spectrophotometry are used as complements to PIXE. The multielemental nature of PIXE is essential to eliminate the interferences from other elements in the samples.

The analytical techniques developed in this project, with PIXE as the basic method, can be used in the health hazard assessments of existing and coming processes in metallurgy

where toxic fumes is expected.

Airborne particulates in ambient air

Studies of the composition of the atmospheric aerosol are of basic importance for judging the air quality. Adverse effects on both man and the environment from airpollution are well established facts for several elements in heavily polluted areas. To prevent future damage, a great need for measurements of different components in the atmospheric aerosol has come up. The atmospheric aerosol consists of several components, both with a natural and an atmospheric origin. Our interest is focused on the elemental composition of the airborne particles. Conclusions on dominating sources, transportation properties and possible medical effects of the aerosol could be drawn when the elemental concentrations on different partice sizes are known.

Using a single-orifice cascade impactor of Battelle-design, the aerosol particles are collected according to aerodynamic size. The elemental composition on each size-fraction is determined by particle-induced X-ray emission (PIXE) analysis.

Studies on the atmospheric aerosol have so far included measurements on both urban and non-urban sites in the three projects described below. The urban aerosol was studied in a small industrial town (Landskrona) in the southern part of Sweden, using four measuring sites. By correlating the measured elemental concentrations with the winddirections it was concluded that emissions from the industrial area heavily increased the concentrations of the elements S, K, Ca, Ti, V, Mn, Fe, Zn and Pb in the town. Possible sources for all these elements were identified.

At the remote site of Velen in the South of Sweden (lat 59^o), samples were taken every third day during a one-year cycle. The results indicate a strong dependence of the elemental concentrations on the history of the air-mass. Concentrations

are a factor of 5-10 higher when the air has passed over the European continent or Great Britain and Sweden than when coming from north over Norway and Sweden.

The transport of pollutants from Great Britain and the European continent to Sweden has been studied using two sampling sites. One of them was situated at the westcoast and the other at the southcoast of Sweden. The results are now being evaluated.

Multivariate statistical (analytic) methods as cluster analysis and factor analysis, are now used to extract as much information as possible from the results. Conclusions on possible sources in different source areas and the transformation of the particles during their transport are two examples of the potential of these powerful statistical methods.

Determination of lead levels in the dentine of teeth using the PIXE method (Environmental Health)

The lead level in the dentine of a tooth is a measure of the lead exposure during the mineralisation of the tooth. Premolar teeth from children raised in cities (Stockholm and Malmö) and a rural area (Jönköping) in Sweden will be analyzed for lead with the PIXE method.

The teeth are split in two halves. One of the uncontaminated surfaces is irradiated with 2.5 MeV protons which are brought out to a nitrogen surrounding through a thin plastic foil (Kapton, 1 mg/cm^2) and collimated to a diameter of 1 mm. The number of protons hitting the sample is monitored with a surface barrier detector measuring protons backscattered from the exit foil. The nitrogen surrounding eliminates charging and reduces heating of the tooth. The small dimension of the beam and a moveable target holder assure that the same volume in the middle of the dentine is analysed in every tooth. With a set-up featuring a short detector-sample distance and a thick absorber to eliminate negative effects of the abundant calcium K-radiation a high sensitivity for lead is achieved with a moderate proton current.

Even in this application with only single element determination the PIXE method with simple sample treatment and rapid analysis is favourable. By using peak to background measurements the uncertainties in the concentration determinations are kept very low.

Preliminary results show that lead levels of city teeth are higher than those of rural area teeth. Anyhow, the levels seem to be low compared to those obtained by other investigations in other parts of the world. However, teeth from New York children will be analysed with our procedures to get further material for comparisons.

Lungdeposition (Environmental Health)

Man is exposed to airborne particulates in outdoor air as well as in work environment. Particles may be deposited in the human lung and induce toxic effects. At present, there are intense research activities taking place in many laboratories to study how the deposition depends on different parameters as e.g. particle size. Artificial aerosols, often radioactive, are used in those investigations. Frequently animals are used as models for homo. The low limits of detections given by the PIXE-method make it possible to perform deposition studies with aerosols normally encountered in work environment. This means lower exposure to involved persons and easier interpretation of results.

In this project size-fractionated particles are sampled from inhaled as well as exhaled air, while a welder is breathing welding smoke. The results seem to be in agreement with theoretical models. However, there are discrepancies for particles generated in welding operations using coated electrodes. Those particles are hygroscopic and grow in the high relative humidity of the lungs. At present, the growth of welding particles is studied in atmospheres simulating humidity and temperature of the human lung.

7:III PIXE AND OTHER APPLICATIONS OF NUCLEAR PHYSICS

Abstracts are marked with "A"

References from the last few years

- 1 T B Johansson, R Akselsson, S A E Johansson
X-ray analysis: elemental trace analysis at the 10^{-12} g level
Nucl Instr and Meth 84, (1970) 141-143
- 2 R Akselsson, T B Johansson
A beam mapping method
Nucl Instr and Meth 91, (1971) 663-664
- 3 T B Johansson, R Akselsson, S A E Johansson
Proton-induced X-ray emission spectroscopy in elemental trace analysis
Advances in X-ray Analysis, Vol 15 (1972) 373-387
- 4 R Akselsson
Protoninducerad karakteristisk röntgenstrålning
Dissertation, Lund 1973
- 5 J W Nelson, I Williams, T B Johansson, R E Van Grieken, K R Chapman, J W Winchester
Elemental analysis of aerosols using proton scattering
IEEE, Nuclear Section, Nov 1973
- 6 R Akselsson, T B Johansson
X-ray production by 1.5-11 MeV protons
Z Physik 266, (1974) 245
- A 7 R Akselsson, G Johansson, T B Johansson, K Malmqvist
Metallpartiklar i svetsrök, koncentration och storleksvariation
Läkarsällskapets Riksstämma, Arbetsmedicin nr 4, Nov 27-30, 1974

- 8 R Akselsson, G Johansson, K Malmqvist, J Frismark,
T B Johansson
Elemental abundance variation with particle size in
aerosols from welding operations
Proceedings of the Second International Conference on
Nuclear Methods in Environmental Research, July 29-31,
1974, Columbia, Missouri, USA, pp 395-401
- 9 R Akselsson, J W Nelson, J W Winchester
Aerosol composition studies using accelerator proton
bombardment
Proceedings of the Third Conference on Applications of
Small Accelerators, Oct 20-23, 1974, Denton, Texas,
USA, I, pp 139-147
- 10 G Johansson, K Malmqvist
Grundämnen i olika partikelstorleksfraktioner i svetsrök
Examensarbete för civilingenjörsexamen, LUNP 7402
- 11 T B Johansson
Multielement trace analysis by proton-induced X-ray
emission
Dissertation, Lund 1974
- 12 T B Johansson, M Ahlberg
Metallpartiklar i luftföroreningar. Storleksvariation
och koncentration
Läkarsällskapets Riksstämma, Omgivningshygien nr 8,
Nov 27-30 1974
- 13 T B Johansson, R E Van Grieken, J W Winchester
Trace elements in atmospheric aerosol in the Heidelberg
area, measured by instrumental neutron activation
analysis
Atm Env 7, (1974) 297-298
- 14 T B Johansson, R E Van Grieken, J W Winchester
Marine influences on aerosol composition in the coastal
zone
Journal de Recherches Atmosphériques 8 (1974) 761-776

- 15 T B Johansson, R E Van Grieken, J W Winchester
Interpretation of aerosol trace metal particle size
distribution
Proceedings of the Second International Conference on
Nuclear Methods in Environmental Research, July 29-31,
1974, Columbia, Missouri, USA, pp 356-365
- A 16 B Möller, R Akselsson, M Berlin, A Fischbein,
L Hammarström, T B Johansson
Tanddentinets halt av bly. Dosimeter för ackumulerad
blyexposition
Läkarsällskapets Riksstämma, Omgivningshygien nr 7,
Nov 27-30 1974
- A 17 R E Van Grieken, R Akselsson, T B Johansson, J W Nelson,
J W Winchester
Aerosol composition measurements using accelerator
proton beams
Presented at 32ème Congrès de GAMS, Groupement pour
l'Avancement des Methodes Spectroscopiques et Physico-
Chimiques d'Alalyse, Paris, 9-13 Dec 1974
- 18 R E Van Grieken, T B Johansson, J W Winchester
Trace metal fractionation effects between sea water and
aerosols from bubble bursting
Journal de Recherches Atmosphériques 8, (1974) 611-621
- 19 J W Winchester, D L Meinert, J W Nelson, T B Johansson,
R E Van Grieken, C Orsini, H C Kaufmann, R Akselsson
Trace metals in the St Louis aerosol
Proceedings of the Second International Conference on
Nuclear Methods in Environmental Research, July 29-31,
1974, Columbia, Missouri, USA, pp 385-394
- 20 M Ahlberg
Trace analysis of thick samples by proton-induced
X-ray emission
Dissertation, Lund 1975

- 21 M Ahlberg
Simple depth profile determination by proton-induced
X-ray emission
Nucl Instr and Methods 131, (1975) 381-384
- 22 M Ahlberg, R Akselsson, D Brune, J Lorenzen
Proton-induced X-ray analysis of steel surface for
microprobe purposes
Nucl Instr and Meth 123, (1975) 385-393
- A 23 M Ahlberg, R Akselsson, G Johansson, K Malmqvist,
T B Johansson
Proton-induced X-ray emission applied to thick samples
Swedish National Physics Conference, June 13-15, 1978,
Gothenburg, Sweden
- 24 M Ahlberg, J W Bawden, L E Hammarström, R Hellborg
Proton activation analysis studies on fluorine and
nitrogen content of developing dental enamel
Odont Revy 26, (1975) 267-279
- 25 M Ahlberg, G Johansson, K Malmqvist
Elimination of charging in the proton-induced X-ray
emission analysis of thick insulating samples
Nucl Instr and Meth 131, (1975) 377-379
- 26 R Akselsson, G G Desaedeleer, J W Winchester, K A Hardy,
J W Nelson
Bromine and lead relationships with particle size and
time along an urban freeway,
Summary of a paper presented at the International Nuclear
and Atomic Activation Conference and 19th Annual
Meeting on Analytical Chemistry in Nuclear Technology,
Gatlinburg, Tenn, Oct 14-16, 1975
Transactions Am Nucl Sci 21, (1975) 36-37

- 27 R Akselsson, J W Nelson, J W Winchester
Proton scattering for analysis of atmospheric particulate matter
Proceedings of the Conference on Nuclear Cross Sections and Technology, Washington DC, March 3-7, 1975 and
Bulletin American Physical Society, Ser II, 20, (1975) 155
- 28 R Akselsson, C Orsini, D L Meinert, T B Johansson,
R E Van Grieken, H C Kaufmann, K R Chapman, J W Nelson,
J W Winchester
Application of proton-induced X-ray emission analysis to the St Louis regional air pollution study
Advances in X-ray Analysis 18, (1975) 588-597
- A 29 K A Hardy, R Akselsson, J W Nelson, J W Winchester
Preliminary study of air particulates in Miami, Florida, USA, using proton-induced X-ray emission
Abstract to the Anaheim Meeting of the American Physical Society, Jan 29-Febr 1, 1975
- A 30 G Johansson, M Ahlberg, R Akselsson, T B Johansson,
K Malmqvist
Heavy metals in welding aerosols and young human teeth dentine determined by proton-induced X-ray emission,
Swedish National Physics Conference, June 10-12, 1975,
Gothenburg, Sweden
- A 31 G Johansson, R Akselsson, T B Johansson, K Malmqvist
Karaktärisering av svetsrök och expositionsundersökningar på svetsarbetsplatser
Läkarsällskapetets Riksstämman, Omgivningshygien nr 5,
Nov 26-29, 1975, Stockholm
- 32 T B Johansson, M Ahlberg, R Akselsson, G Johansson,
K Malmqvist
Analytical use of proton-induced X-ray emission,
Abstract of a paper presented at the International Nuclear and Atomic Activation Analysis Conference and the 19th Annual Meeting on Analytical Chemistry in Nuclear Technology, October 14-16, 1975, Gatlinburg, Tenn, USA, Transactions Am Nucl Sci 21, (1975) 31

- 33 T B Johansson, R E Van Grieken, J W Nelson, J W Winchester
Elemental trace analysis of small samples by proton-induced
X-ray emission
Analytical Chemistry 47, (1975) 855-860
- 34 H C Kaufmann, R Akselsson
Non-linear least squares analysis of proton-induced
X-ray emission data
Advances in X-ray Analysis 18, (1975) 353-361
- 35 K Malmqvist, G Johansson, R Akselsson, T B Johansson
Metallanalyser av svetsrök
Report LUNP 7508
- 36 R E Van Grieken, T B Johansson, J W Winchester, L A Odom
Micro-determination of zirconium-hafnium ratios in
zircons by proton-induced X-ray emission
Z Anal Chem 275, (1975) 343-348
- 37 M Ahlberg
Matrix effects in trace analysis by proton-induced
X-ray emission
Proceedings of the Fourth International Conference on the
Scientific and Industrial Applications of small
Accelerators, Denton, Texas, 1976, 106-109
- 38 M Ahlberg, R Akselsson
Proton-induced X-ray emission in the trace analyses of
human tooth enamel and dentine
International Journal of Applied Radiation and
Isotopes 27, (1976) 279-290
- 39 M Ahlberg, R Akselsson, B Forkman, G Rausing
Gold traces on wedge-shaped artefacts from late neolithic
of South Scandinavia analysed by proton-induced X-ray
emission spectroscopy
Archaeometry 18, (1976) 39-49

- 40 M Ahlberg, A P Woodward Jr, A C D Leslie, J W Winchester
Sampling apparatus for lung deposition studies of
aerosols
FSU Technical Report, Tallahassee, Fla, 1976
- 41 R Akselsson, M Bohgard, L E Carlsson, G Johansson
PIXE-metoden - en analysmetod lämpad för yrkeshygieniska
undersökningar
Contribution to 25th Scandinavian Industrial Hygiene
Conference, Beito, Norway, Oct 17-20, 1976
- 42 K R Akselsson, G G Desaedeleer, T B Johansson,
J W Winchester
Particle size distribution and human respiratory
deposition of trace metals in indoor work environments
Ann Occup Hyg 19, (1976) 225-238
- 43 R Akselsson, K A Hardy, G G Desaedeleer, J W Winchester,
W W Berg, T B Vanderwood, J W Nelson
X-ray technique for aerosol sulphur baseline assessment
along an urban freeway
Advances in X-ray analysis 19, (1976) 415-425
- 44 R Akselsson, G Johansson, K Johansson, K Malmqvist,
K Sjöberg, M Ahlberg, JE Andersson, M Bohgard, LE Carlsson,
J Frismark, H Lannefors, T B Johansson, B Rudell, B Zaman
S A E Johansson
Lägesrapport för anslag Dnr 74/109, Arbetarskyddsfonden
per den 1 maj 1976, LUNP 7606
- A 45 R Akselsson, G Johansson, K Malmqvist, M Bohgard,
L E Carlsson, B Zahman
Proton-induced X-ray emission and work environment
Contribution to the Conference on the Effect of Toxic
Metals on Man and Environment, June 15-18, 1976,
Luleå, Sweden

- 46 M Bohgard, L-E Carlsson
En uppställning för röntgenfluorescensanalys med ett
röntgenterapiaggregat
LUNP 7605, 1976
- A 47 M Bohgard, L Jangida, J Frismark, R Akselsson
ESCAanalyser av krom i svetsrök
Läkaresällskapets Riksstämma, Dec 1-4, 1976,
Stockholm, Sweden
- A 48 J Frismark, R Akselsson
Karaktärisering av svetsrök med elektronmikroskop
Läkaresällskapets Riksstämma, Dec 1-4, 1976,
Stockholm, Sweden
- 49 K A Hardy, K R Akselsson, J W Nelson, J W Winchester
Elemental constituents of the Miami aerosol as a
function of particle size
Env Sci Techn 10, (1976) 176-182
- 50 S A E Johansson, T B Johansson
Analytical application of particle induced X-ray
emission
Nucl Instr and Meth 137, (1976) 473-516
- 51 T B Johansson, M Ahlberg, R Akselsson, G Johansson,
K Malmqvist
Analytical use of proton-induced X-ray emission
Journal of Radioanalytical Chemistry 32, (1976) 207-217
- 52 T B Johansson, R E Van Grieken, J W Winchester
Elemental abundance variation with particle size in
North Florida Aerosols
Journal of Geophysical Research 81, (1976) 1039
- 53 H C Kaufmann, K R Akselsson, W J Courtney
REX: A computer program for PIXE spectrum resolution of
aerosols
Advances in X-ray Analysis 19, (1976) 355-366

- 54 H Lannefors
Förslag till förbättring av energiupplösningen vid
PIXE-analys med hjälp av kristallspektrometer
LUNP 7603, 1976
- 55 H Lannefors
Grundämnen i olika partikelstorleksfraktioner i
Landskronaluft
Internal report, 1976
- 56 H Lannefors, T B Johansson
Analysis of heavy metals by proton-induced X-ray
emission
Contribution to the World Meteorological Organization
Technical Conference on Atmospheric Pollution
Measurement Techniques, Oct 11-15, 1976, Gothenburg, Sweden
- 57 A C D Leslie, J W Winchester, F W Leysieffer, M S Ahlberg
Prediction of health effects of pollution aerosol
Proceedings of the Tenth National Conference on Trace
Substances in Environmental Health, University of
Missouri, Columbia, 1976
- A 58 K Malmqvist, G Johansson, L E Carlsson, K R Akselsson
X-ray absorption in PIXE analysis of nonuniform samples
Contribution to the International Conference on Particle
Induced X-ray Emission and its Analytical Applications,
Aug 23-26, 1976, Lund, Sweden
- 59 J W Nelson, B Jensen, G G Desaedeleer, R Akselsson,
J W Winchester
Automatic time sequence filter sampling of aerosols for
rapid multi-elemental analysis by proton-induced X-ray
emission
Advances in X-ray Analysis 19, (1976) 403-413
- A 60 B Rudell, R Akselsson, M Berlin
Deposition av svetsrökspartiklar i lunga
Läkarsällskapets Riksstämma, Dec 1-4, 1976, Stockholm,
Sweden

- 61 J Sheline, R Akselsson, J W Winchester
Trace element similarity groups in North Florida
Spanish moss
Journal Geophysical Research 81, (1976) 1047-1050
- 62 R E Van Grieken, T B Johansson, K R Akselsson,
J W Winchester, J W Nelson, K R Chapman
Geophysical applicability of aerosol size distribution
measurements using cascade impactors and proton-induced
X-ray emission
Atm Env 10, (1976) 571-576
- 63 M Ahlberg
Enhancement in PIXE analysis
Nucl Instr and Meth 142, (1977) 61-65
- 64 M Ahlberg
Comparison of yield versus depth for particle induced
and photon induced X-ray emission analysis
Nucl Instr and Meth 146, (1977) 465-467
- 65 M Ahlberg
PIXE-metoden, dess karaktärsdrag och tillämpningar
VI Nordiska Spårelementkonferensen, Aug 23-26, 1977,
Södertälje, Sweden, invited
- 66 M S Ahlberg, A C D Leslie, J W Winchester
Aerosol particle response during inhalation by humans
National Meeting, 20-25 March, 1977, New Orleans, USA,
invited
- 67 M S Ahlberg, W Winchester, C D Leslie, B O Dahlbäck
Sampling exhaled aerosol particle size fractions under
controlled humidity and temperature conditions
The Tenth Aerosol Technology Meeting, Sept 26-28, 1977,
Albuquerque and Los Alamos, New Mexico, USA

- 68 R Akselsson
PIXE och tillämpningar inom arbetsmiljöområdet
Contribution to the STU-seminar "Provtagningsmetodik
och analysteknik inom arbetsmiljöområdet nu och i
framtiden, Feb 15-16, 1977, Mariehamn, Finland", invited
- 69 R Akselsson, G Johansson, K Johansson, K Malmqvist,
K Sjöberg, M Bohgard, L-E Carlsson, H Lannefors,
J Frismark, M Ahlberg, B Rudell, B Jangida, S A E Johansson
Luftföroreningar vid svetsning
Lägesrapport för anslag Dnr 74/109, Arbetarskyddsfonden
den 1 mars 1977, LUTFD2/TFKF-3005/1-57/1977
- 70 K R Akselsson, S A E Johansson, T B Johansson
Mass calibration in PIXE analyses,
In "X-ray fluorescence analysis of environmental
samples" 175-180, ed T G Dzubay (Ann Arbor Science
Publishers, Inc Ann Arbor 1977), invited
- A 71 L-E Carlsson, M Bohgard, G Johansson, K Malmqvist,
R Akselsson
Karakterisering av svetsrök
Läkarsällskapets Riksstämma, Nov 30 - Dec 3, 1977,
Stockholm, Sweden
- A 72 G Desaedeleer, R Akselsson, W Winchester, R Hardy,
J W Nelson
Elemental analysis of roadway aerosol in the Los Angeles
bay area by proton-induced X-ray emission analysis
Summary of presentation at the American Chemical Society
173rd National Meeting, New Orleans, March 20-25, 1977
- 73 G G Desaedeleer, J W Winchester, K R Akselsson
Monitoring aerosol elemental composition in particle
size fractions for predicting human respiratory uptake
Nucl Instr and Meth 142, (1977) 97-99
- A 74 G Desaedeleer, J W Winchester, R Akselsson
Toxicological implications of aerosol size distribution
measurements using cascade impactors and elemental analysis
by particle induced X-ray emission
American Chemical Society 173rd National Meeting, New
Orleans. March 20-25. 1977

- 75 B Erlandsson, S Mattsson
Medically used radionuclides in sewage sludge
Water, Air and Soil Pollution, vol 9, in press
- 76 B Erlandsson, S Mattsson
Medicinskt använda radionuklider i rötslam och vatten
från Malmö kommuns reningsverk i Sjölanda
Report to Malmö kommuns Vatten- och Avloppsverk, Oct, 1977
- A 77 B Erlandsson, S Mattsson, T Bramstång
Medicinskt använda radionuklider i rötslam från Malmö
kommuns reningsverk i Sjölanda
Svensk förening för medicinsk radiologi, contribution 17
- 78 G Johansson, K Malmqvist, R Akselsson
Analysis of welding aerosol by simultaneous use of
particle induced X-ray emission and γ -ray detection
for fluorine analysis
International Symposium on the Control of Air Pollution
in the Working Environment, Sept 6-8, 1977, Stockholm,
Sweden
- 79 H C Kaufmann, K R Akselsson, W J Courtney
REX - a computer-programme for PIXE analysis
Nucl Instr and Meth 142, (1977) 251-257
- A 80 H Lannefors
Study of elemental composition of different size
fractions in an Arctic aerosol
Workshop on sources and significance of natural and
man-made aerosols in the Arctic, April 1977, Lilleström,
Norge
- 81 H Lannefors, R Akselsson
Ambient air contamination by micron and submicron
particles from welding operations
Bulletin of environmental contamination and toxicology,
17, (1977) 521-527

- 82 H Lannefors, T B Johansson, L Granat, B Rudell
Elemental concentrations and particle size distributions
in an atmospheric background aerosol
Nucl Instr and Meth 142, (1977) 105
- 83 A C D Leslie, M S Ahlberg, J W Winchester, J W Nelson
Aerosol characterization for sulfate health effects
assessment in Florida
Proceedings of the XIth International Conference on
Trace Substances in Environmental Health, University of
Missouri, Columbia, 1977
- 84 C Q Orsini, H C Kaufmann, K R Akselsson, J W Winchester,
J W Nelson
Variation of elemental composition with particle size
in the St Louis aerosol
Nucl Instr and Meth 142, (1977) 91-96
- 85 M S Ahlberg, F C Adams
Experimental comparison of photon and particle induced
X-ray emission analysis of air particulate matter
X-ray Spectrometry 7, (1978) 73-80
- 86 M S Ahlberg, J W Winchester
Dependence of aerosol sulphur particle size on relative
humidity
Atm Env 12, (1978) 1631-1632
- 87 M S Ahlberg, A C D Leslie, J W Winchester
Characteristics of sulphur aerosol in Florida as
determined by PIXE analysis
Atm Env 12, (1978) 773-777
- 88 M S Ahlberg, A C D Leslie, J W Winchester
The chemical state of particulate sulphur in ambient
aerosols determined by PIXE analysis
Nucl Instr and Meth 149, (1978) 451-455

- 89 R Akselsson
Provtagning och analys av tungmetaller
Contribution to "Symposium om tillförsel av mikrokomponenter i Norden, Oct 3-5, 1978, Ekerö", invited
- 90 R Akselsson
Applications of particle induced X-ray emission (PIXE)
Proceedings of the Nordic Spring Symposium on Atomic Inner Shell Phenomena, April 17-21, 1978, Geilo, Norway,
In press, invited
- A 91 K R Akselsson, S A E Johansson
Elemental analysis at the PIXE laboratory in Lund
Swedish National Physics Conference, June 13-15, 1978, Linköping, Sweden, invited
- A 92 K R Akselsson, S A E Johansson
PIXE analysis research in Lund
Abstract for an invited talk at the Conference on Applications of Small Accelerators, Denton, Texas, Nov 6-8, 1978, invited
- A 93 M Bohgard, G Johansson, K Johansson, K Malmqvist, K R Akselsson
Characterization of welding aerosols at the PIXE laboratory in Lund
Swedish National Physics Conference, June 13-15, 1978, Linköping, Sweden
- A 94 L-E Carlsson, M Ahlberg, H-C Hansson, E M Johansson, H Lannefors, J Pallon, K R Akselsson
Applications of particle-induced X-ray emission analysis
Swedish National Physics Conference, June 13-15, 1978, Linköping, Sweden
- 95 H-C Hansson
Tungmetallkoncentrationer i ett renluftsområde under ett år
Contribution to "Symposium om tillförsel av mikrokomponenter i Norden, Oct 3-5, 1978, Ekerö, Sweden"

- 96 G Johansson, R Akselsson, M Bohgard, L-E Carlsson,
H-C Hansson, H Lannefors, K Malmqvist
Particle-induced X-ray emission - a useful analytical
method
Proceedings of the Fourth SAC Conference on Analytical
Chemistry, July 17-22, 1977, Birmingham, Great Britain,
Proc Analyt Div Chem Soc 15, (1978) 24-27
- A 97 G Johansson, K Malmqvist, M Bogard, R Akselsson
Karakterisering av svetsrök - en lägesrapport över
svetsprojektet i Lund
Contribution to "27 Nordiske Arbetshygieniske Möde,
Nov 19-22, 1978"
- 98 G Johansson, K Malmqvist, M Bohgard, L-E Carlsson,
K Johansson, K Sjöberg, R Akselsson, S Johansson
Luftföroreningar vid svetsning
LUTFD2/(TFKF-3010)/1-30/1978
- 99 H Lannefors, H-C Hansson
Luftburna partiklars grundämnessammansättning i olika
storleksklasser i Landskronaluften
LUTFD2/(TFKF-3013)/1-32/(1978)
- 100 A C D Leslie, M S Ahlberg, J W Winchester, J W Nelson
Aerosol characterization for sulphur oxide health
effects assessment
Atm Env 12, (1978) 729-734
- 101 K Malmqvist, G Johansson, R Akselsson
Determination of heavy metals and fluorine in airborne
particulates in an indoor environment by simultaneous
use of PIXE and γ -ray detection
Proceedings of the Fourth SAC Conference on Analytical
Chemistry July 17-22, 1977, Birmingham, Great Britain
Proc Analyt Div Chem Soc 15, (1978) 13-16

- 102 K Malmqvist, G Johansson, R Akselsson
Proton-induced X-rays and gamma-rays for the analysis
of welding fumes
Contribution to the American Industrial Hygiene
Conference, May 7-12, 1978, Los Angeles, USA
- 103 J Pallon
Kammare för PIXE analys av biologiska prov vid
atmosfärstryck
Internal Report in press (1978)
- 104 The PIXE-group in Lund
Trace elemental analysis by means of heavy particle
induced X-rays
Progress Report for IAEA, Research Agreement No 1459/
R2/CF, Jan 1975-Dec 1977
- 105 M S Ahlberg, H J Annegarn, B O Dahlbäck, A-C D Leslie,
J W Winchester
Particle size dependent aerosol deposition in the
human respiratory tract
Trace Substances in Environmental Health - XII, in press
- 106 M S Ahlberg, A C D Leslie, J W Winchester
Environmental and occupational health analyses using
proton-induced X-ray emission
Proceedings of the Symposium on Electron Microscopy
and X-ray Applications to Environmental and Occupational
Health Analyses, April 25-28, 1977, Denver, Colorado,
Ann Arbor Science Publishers, in press
- A 107 K R Akselsson, M Bohgard, L-E Carlsson, H C Hansson,
E M Johansson, G Johansson, H Lannefors, K Malmqvist
PIXE analysis of environmental samples
Abstract for "XXI Colloquium Spectroscopicum Internationale,
July 1-6, 1979, Cambridge"
- 108 M Bohgard, B L Jangida, K R Akselsson
An analytical procedure for determining chromium in
samples of airborne dust
Submitted to Annals of Occup Hygiene

- A 109 M Bohgard, G Johansson, K Malmqvist, K R Akselsson
Characterization of work environment aerosols by PIXE
Abstract for "XXI Colloquium Spectroscopicum
Internationale, July 1-6, Cambridge, 1979"
- 110 G G Desaeleer, J W Winchester, K R Akselsson
Aerosol particle size relationships along a Los Angeles
freeway
To be published
- 111 J Frismark, M Bohgard, K R Akselsson
Characterization of welding aerosols
To be published
- 112 J W Winchester, G G Desaeleer, K R Akselsson
Time variations in elemental composition of atmospheric
particulates along a Los Angeles freeway
To be published
- 113 J W Winchester, A C D Leslie, M S Ahlberg
A north-south concentration gradient of aerosol sulphur
in Florida
Proceedings of the 4th Joint Conference on Sensing
of Environmental Pollutants, Nov 6-11, 1977, New Orleans,
USA, in press