



# LUND UNIVERSITY

## Aerosol deposition of AuNP into biological fluids produce specific biomolecule corona

Svensson, Christian; Messing, Maria; Schollin, Alexander; Deppert, Knut; Mueller, Bengt; Pagels, Joakim; Rissler, Jenny; Bohgard, Mats; Linse, Sara; Cedervall, Tommy

2011

[Link to publication](#)

### *Citation for published version (APA):*

Svensson, C., Messing, M., Schollin, A., Deppert, K., Mueller, B., Pagels, J., Rissler, J., Bohgard, M., Linse, S., & Cedervall, T. (2011). *Aerosol deposition of AuNP into biological fluids produce specific biomolecule corona*. 1-2. Abstract from 5th international symposium on Nanotechnology occupational and environmental health, Boston, MA, United States.

### *Total number of authors:*

10

### **General rights**

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117  
221 00 Lund  
+46 46-222 00 00

## **Aerosol deposition of AuNP into biological fluids produce specific biomolecule corona**

Christian R. Svensson<sup>1\*</sup>, Maria E. Messing<sup>2\*</sup>, Alexander Schollin<sup>3\*</sup>, Knut Deppert<sup>2\*</sup>, Bengt O. Mueller<sup>2\*</sup>, Joakim Pagels<sup>1\*</sup>, Jenny Rissler<sup>1\*</sup>, Mats Bohgard<sup>1\*</sup>, Sara Linse<sup>3\*</sup>, Tommy Cedervall<sup>3\*</sup>

\*<sup>1</sup>Lund University / Ergonomics and Aerosol technology, Box 118, Sweden, Lund 221 00, Tel.: + 46-46 222 00 00; Fax: +46-46 222 47 20; E-mail: [Christian.Svensson@design.lth.se](mailto:Christian.Svensson@design.lth.se)

\*<sup>2</sup>Lunds University / Department of Solid State Physics

\*<sup>3</sup>Lunds University / Biochemistry and Structural Biology

**Keywords:** *nanoparticles, characterization, protein corona, biomolecules, toxicity*

Due to the special properties of nanoparticles, concern has been raised with regards to their potential effects in biological systems. When biomolecules bind to the particle surface a dynamic protein/biomolecule corona form (Cedervall, 2007). This particle/biomolecule complex is believed to be what the cells actually perceive and in addition to particle properties may be a major determinant of particle toxicity (Lynch, 2009). We present a method to investigate the composition of the biomolecule/protein corona on model nanoparticles in different physiological fluids, porcine serum and lung fluid. Gold nanoparticle (AuNP) agglomerates were generated as aerosols by high temperature evaporation-condensation furnace. The agglomerates where sintered to spherical shape and characterized with regards to mobility size, mass and with transmission electron microscopy. The sintered AuNPs (median 60 nm) were deposited into solutions of bovine serum albumin (BSA) and homocysteine using an electrostatic precipitator. The AuNPs was in addition to mass-mobility characterized, in gas phase, analyzed with dynamic light scattering in suspension. AuNPs in solution were mixed with porcine serum or porcine lung fluid. Using gel electrophoresis (SDS-PAGE) the protein corona was determined for AuNPs mixed with the physiological fluids. SDS-PAGE results indicate that the corona is similar regardless of surfactant, BSA or homocysteine. Also the corona is different between particles mixed with lung fluid or blood-serum.

In conclusion: We have shown that the protein / biomolecule corona can be studied using model particles generated in the aerosol phase and stabilized with BSA and homocysteine in suspension, and that deposition directly into serum and lung fluid is also possible. However, the observed corona differ between particles administered to porcine blood serum and lung fluid. This work was supported by the Nanometer

Structure Consortium at Lund University (nmC@LU) and the Swedish research council FAS through project 2009-1291 and the FAS-centre METALUND.

Cedervall, T et al. (2007). PNAS, 104, 7, 2050-2055.

Lynch, I et al. (2009). Nat Nanotechnol, 4, 577-80.