

# LUND UNIVERSITY

# Tests of CALIFA Barrel modules at CCB in Krakow

Park, Joochun; Heiss, B.; Hartig, A.-L.; Alvarez-Pol, H.; Bruni, G.; Casarejos, E.; Cederkäll, Joakim; Cortina-Gil, D.; Díaz Fernández, P.; Galaviz, D.; Gernhäuser, R.; Golubev, Pavel; Heinz, A.; Johansson, H. T.; Klenze, P.; Knyazev, Alexander; Kröll, T.; Nilsson, T.; Perea, A.; Rhee, H. B.; Tengblad, O.; Teubig, P.; Zieblinski, M.; Califa Collaboration Published in: **GSI-FAIR Scientific Report** 

DOI: 10.15120/GR-2018-1

2018

# Link to publication

## Citation for published version (APA):

Park, J., Heiss, B., Hartig, A.-L., Alvarez-Pol, H., Bruni, G., Casarejos, E., Cederkäll, J., Cortina-Gil, D., Díaz Fernández, P., Galaviz, D., Gernhäuser, R., Golubev, P., Heinz, A., Johansson, H. T., Klenze, P., Knyazev, A., Kröll, T., Nilsson, T., Perea, A., ... Califa Collaboration (2018). Tests of CALIFA Barrel modules at CCB in Krakow. In *GSI-FAIR Scientific Report* (GSI-FAIR Scientific Report; Vol. 2018, No. 1). GSI Helmholtzzentrum für Schwerionenforschung. https://doi.org/10.15120/GR-2018-1

Total number of authors: 24

#### **General rights**

Unless other specific re-use rights are stated the following general rights apply: Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the

 Isgal requirements associated with these rights.
 Users may download and print one copy of any publication from the public portal for the purpose of private study or research.

- You may not further distribute the material or use it for any profit-making activity or commercial gain
  You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: https://creativecommons.org/licenses/

#### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

**PO Box 117** 221 00 Lund +46 46-222 00 00

# Tests of CALIFA Barrel modules at CCB in Kraków

J. Park<sup>1</sup>, B. Heiss<sup>2</sup>, A.-L. Hartig<sup>3</sup>, H. Alvarez-Pol<sup>4</sup>, G. Bruni<sup>5</sup>, E. Casarejos<sup>6</sup>, J. Cederkäll<sup>1</sup>, D. Cortina-Gil<sup>4</sup>, P. Díaz Fernández<sup>5</sup>, D. Galaviz<sup>7</sup>, R. Gernhäuser<sup>2</sup>, P. Golubev<sup>1</sup>, A. Heinz<sup>5</sup>, H. T. Johansson<sup>5</sup>, P. Klenze<sup>2</sup>, A. Knyazev<sup>1</sup>, T. Kröll<sup>3</sup>, T. Nilsson<sup>5</sup>, A. Perea<sup>8</sup>, H.-B. Rhee<sup>3</sup>, O. Tengblad<sup>8</sup>, P. Teubig<sup>7</sup>, M. Zieblinski<sup>9</sup>, for the CALIFA working group of the R<sup>3</sup>B collaboration

<sup>1</sup>Lund University, <sup>2</sup>TU München, <sup>3</sup>TU Darmstadt, <sup>4</sup>University of Santiago de Compostela, <sup>5</sup>Chalmers University of Technology, <sup>6</sup>Universidade de Vigo, <sup>7</sup>LIP Lisbon, <sup>8</sup>CSIC Madrid, <sup>9</sup>IFJ PAN Kraków

CALIFA (CALorimeter for In-Flight gamma-ray and pArticle detection) [1] is dedicated to the detection, tracking and energy determination of light charged particles and  $\gamma$  rays emerging from R<sup>3</sup>B experiments. Significant progress has been made on the Barrel section [2] of CALIFA, which consists of ~2000 CsI(Tl) scintillator crystals that provide a high angular resolution needed for effective Doppler corrections.

For the Phase-0 experiments at GSI in 2018, a substantial part of the CALIFA Barrel detectors will be installed and operated in the Demonstrator [3] configuration. An extensive preparatory work in detector performance, stability and data acquisition is required to provide a sound and effective experimental campaign at GSI. To that end, in-beam tests at the Bronowice Cyclotron Center (CCB) at IFJ PAN in Kraków were carried out. The cyclotron at CCB provided proton beams with energies up to 225 MeV, which enabled experiments for (p,2p) quasifree scattering (QFS) reactions on stable targets.



Figure 1: Detectors seen during mounting at CCB for the (p,2p) experiment. Three CALIFA Barrel petals (one placed below the target later), two DSSSD tracking detectors, and one CEPA4 detector were tested.

The experimental setup at CCB is shown in Fig. 1. Two petals were positioned on a horizontal plane with a polar angular coverage of  $25^{\circ} < \theta < 58^{\circ}$  to detect the scattered particles. One petal was deployed at its nominal CALIFA position (43° <  $\theta$  < 82°) to detect  $\gamma$  rays. Two doublesided silicon strip detectors (DSSSDs) were also employed as particle trackers. A CEPA4 [4] prototype detector for the CALIFA Forward Endcap was placed at  $\theta$  ~ 11° for its performance test. Data from the petals and the DSSSD trackers were read out with the MBS system featuring FEBEX3B cards [5]. The calibration data from a  $^{60}$ Co y-ray source analysed was with the UCESB/R<sup>3</sup>BRoot software [6].

A laminar water jet, with a small diameter of 0.46 mm was used to provide well-defined tracking correlations for the reactions at the target, and give an energy straggling below the expected resolution limit for the total energy reconstruction of the scattered protons with CALIFA. Other targets such as graphite, polypropylene, <sup>112,124</sup>Sn and <sup>208</sup>Pb were also irradiated. The proton beam energy was 200 MeV for over 95% of the total beam time. Other beam energies ranged from 70 to 225 MeV.

A preliminary analysis of (p,2p) data from the water target revealed a  $\gamma$ -ray peak at 6.3 MeV (see Fig. 2), corresponding to the known transition from the first excited  $3/2^-$  state to the  $1/2^-$  ground state in <sup>15</sup>N. More sophisticated energy calibration, event selection cuts and simulations are being developed to refine and extend the analysis to the data from other targets. We acknowledge the local support at CCB with logistics and beam delivery.



Figure 2:  $\gamma$ -ray energy spectrum following (p,2p) reactions on the water target. The 6.3-MeV transition in <sup>15</sup>N is clearly visible. The inset shows the proton tracking data from a DSSSD-petal pair used in the event selection cut.

## References

- [1] D. Cortina-Gil et al., NDS 120 (2014) 99.
- [2] H. Alvarez-Pol et al., NIM A 767 (2014) 453.
- [3] B. Pietras et al., NIM A 814 (2016) 56.
- [4] E. Nacher et al., NIM A 769 (2015) 105.
- [5] T. Le Bleis et al., GSI Scientific Report (2013) 201.
- [6] D. Bertini, J. Phys. Conf. Ser. 331 (2011) 032036.

Experiment beamline: none

**Experiment collaboration:** NUSTAR-R3B

## Experiment proposal: none

Accelerator infrastructure: CCB at IFJ PAN in Kraków PSP codes: none

Grants: EU H2020 contract No. 654002, BMBF 05P15WOFNA

Strategic university co-operation with: none