



LUND UNIVERSITY

Change of voice in puberty in choir girls

Pedersen, M.F.; Moller, S.; Krabbe, S.; Munk, E.; Bennet, P.; Kitzing, Peter

Published in:
Acta Oto-Laryngologica

1984

[Link to publication](#)

Citation for published version (APA):

Pedersen, M. F., Moller, S., Krabbe, S., Munk, E., Bennet, P., & Kitzing, P. (1984). Change of voice in puberty in choir girls. *Acta Oto-Laryngologica*, 412, 46-49.

Total number of authors:

6

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

Change of Voice in Puberty in Choir Girls

A Cross-sectional Study of Electroglottographical Measured Fundamental Frequency in Reading Compared with Phonetograms, Pubertal Stages and Endocrine Analysis

M.F. PEDERSEN, S. MØLLER, S. KRABBE, E. MUNK, P. BENNET and P. KITZING

From Copenhagen Singing School, Skt. Annae Gymnasium, Department of Paediatrics, Rigshospitalet, University of Copenhagen, Hormone and Biostatistical Departments, Statens Seruminstitut, Copenhagen, Denmark, and Department of Phoniatics, University ENT Clinic General Hospital, Malmö, Sweden

Pedersen M F, Møller S, Krabbe S, Munk E, Bennet P and Kitzing P. Change of voice in puberty in choir girls. *Acta Otolaryngol (Stockh) 1984; 412: 46-49.*

Forty-seven girls in Copenhagen Singing School were chosen by stratification on class level from their 3rd to 12th school year. Examination was carried out of fundamental frequency (Fo) in a reading situation with registration of about 2000 electroglottographical signals with means expressed in Hz and with 95% deviation in semitones, phonetograms according to UEP standard proposal with analysis of conditions of intensity together with lower, middle and upper frequency. By comparison with evaluation of state of puberty and androgen and oestrogen status it was found that Fo was significantly correlated to age ($r=0.40$) and to Fmin ($r=0.51$) and Fmiddle ($r=0.45$). Furthermore, the phonetogram area was significantly correlated to age ($r=0.65$). This corresponds to $P < 0.01$ for all values. There was no significant correlation between change of voice and hormonal conditions. Supplementary examinations using multivariate analysis will be carried out as it was previously done for boys.

M.F. Pedersen, Copenhagen Singing School, Sjøllør Blvd. 135, DK-2500 Valby, Denmark.

Puberty conditions have always been of interest in connection with singing. It has been documented earlier that fundamental frequency (Fo) decreases with age^{1, 2, 3, 4}, while the tone range increases^{5, 6}. The difference between boys and girls is not sufficiently elucidated⁷. Since better methods have appeared for analysis of both voice conditions and hormonal changes we wanted to obtain an impression of the possibilities of predicting the occurrence of voice puberty in girls. This has been possible to a certain degree in boys, in whom the Fo was always correlated to a total serum testosterone < 10 nmol/l.

MATERIALS AND METHODS

Forty-seven girls aged 8-19 years from the Copenhagen Singing School participated in our cross-sectional examination. The girls were chosen by stratification with similar numbers from each school class. The girls are admitted to the school at the 3rd class level with criteria of good musicality according to standardized examination and a good voice. They can remain there until the 12th class level. All participants were healthy and had vocal cords without organic changes and none took medicine. An informal written consent from the girls' homes, including information on menarche, was obtained. Blood sampling for hormone status was carried out on day 3-6 in the menstruation circle. Development of puberty was evaluated with height, weight, pubes hair and mamma development according to Tanner¹⁰. Stroboscopy was performed on all the girls in order to exclude presence of abnormalities. Fo was recorded by means of electroglottography together with the vocal range by reading of a phonetically well balanced text. A computer program based on consecutive analyses of a mean of 2000 electroglottographical cycles states the mean Fo in Hz and the 95% deviation in semitones. The equipment is commercially available. Phonetograms were made by a phoniatician and an experienced singing master according to proposed international standard¹¹. Calculation of the area was performed in cm^2 on the

standard paper. Further examinations were carried out on a phonetograph developed by a Danish engineering firm. The hormone investigation was performed at Statens Seruminstitut. The observations were recorded with geometric means in three age groups.

All observations were logarithmically transformed in order to obtain a normal distribution. A one-way analysis of variance was performed and correlation co-efficients were calculated for all parametres.

Equipment

Electroglottograph FJ Electronics type 830
 Microdatamate Digisys type 2205, Cybernex model LGD-1
 Spectral analyzer, Nicolet 440 A mini ubequitous
 Sound level meter with microphone, Brüel and Kjaer type 2205
 Oscilloscope Tectronix type 510 3n, storage with polaroid camera
 Text: IPA book "The North Wind and the Sun"
 Stroboscope: Brüel and Kjaer type 5066
 Phonetograph Voice Profile instruments type 830I

RESULTS

Height and weight corresponded to other investigations of Danish and Swedish school-children^{12, 13}

Fig. 1.




Fig. 1 illustrates phonetograms at different age levels:

- 8.9 years: The time when the untrained girl gets into the choir
- 11.7 years: Typical child's voice with register change at fis^1 (between 330-392 Hz)
- 13.8 years: Absolutely well functioning girl's voice without break of register at weak intensities
- 14.8 years: Change of register at gis^1 (between 392-440) and d^2 (between 524-660) evaluated as puberty voice by singing master.

Table 1

Geometric means of androgen oestrogen pubertal stage and voice

| Age | | 8.6-12.9 | 13-15.9 | 16-19.8 |
|---|---------------|--------------|---------|---------|
| Number | | 22 | 14 | 11 |
| Oestrone | (nmol/l) | 78 (10 < 40) | 113 | 113 |
| Oestradiol | (nmol/l) | 103(9 < 40) | 185 | 167 |
| Testosterone | (nmol/l) | 0.576 | 0.74 | 0.94 |
| Oestrone sulphate | 887 (3 < 200) | 2261 | 2521 | |
| Dihydro epi androsterone | (nmol/l) | 2990 | 3500 | 7200 |
| Androsten dione | (nmol/l) | 1.39 | 3.19 | 3.43 |
| Menarche (No.) | | +4 | -3 | +all |
| Pubic hair (stage) | | 1-4 | 2-5 | 4-6 |
| Mamma development (stage) | | 1-4 | 2-5 | 5 |
| Fundamental frequency (Fo) in speech (Hz) | | 257 | 248 | 197 |
| Voice range (semitones) in speech | | 3.2 | 3.2 | 5.2 |
| Voice range (semitones) in singing | | 23 | 30 | 38 |
| Phonetogram area (cm ²) | | 17.5 | 22.2 | 28.3 |
| Lower biological tone (Hz) | | 166 | 153 | 145 |
| Middle biological tone (Hz) | | 434 | 406 | 413 |
| Higher biological tone (Hz) | | 1134 | 1136 | 1263 |

Table 1 shows results in three age a groups of the parametres in question. Fo was correlated to age ($r=0.40$, $p < 0.001$) and to Fmin ($r= 0.51$, $P < 0.001$) and Fmiddle ($r=0.45$, $P < 0.01$). Correlation between phonetogram area and age was $r=0.65$, $P < 0.001$. Analysis of intensity conditions in the phonetograms and the other results showed to significant circumstances. No correlation between hormone status and Fo during reading was found in this study. However, supplementary multivariate statistical analyses of hormone values in relation to voice parametres will be carried out. This may possibly show predictive values.

DISCUSSION

Fo was significantly related to age showing a mean decrease from 257 to 197 Hz and was related to the lower biological tone in the phonetogram with $P < 0.001$. The phonetogram was similarly significantly related to age with the same significance in these girls who were all maximally trained in singing. 2-3 years after menarche a change of the areas was seen at the same time as the girls were moved from child to adult choir.

Fo and phonetography should be included as sex characteristics in routine analyses of puberty due to the accuracy of the method and the reproducibility of the phonetograms¹⁴. They also provide a better possibility of evaluating the pathological deviation¹⁵.

REFERENCES

1. Firch, J.L., Holbrook, A. Modal vocal fundamental frequency of young adults. Arch. Otolaryng. 1970; 92: 379-382.
2. Hollien, H., Paul, P. A second evaluation of the speaking fundamental frequency characteristics of post-adolescent girls. Lang. Speech 1969; 12: 119-124.
3. Bennett, S. A 3-year longitudinal study of school-aged children's fundamental frequencies. J. Speech Hear. Res. 1983; 26: 137-142.

4. McCglone, R.E., McCglone, J. Speaking fundamental frequency of eight-year old girls. *Folia Phoniatr* (Basel) 1972; 24: 313-317.
5. Hartlieb, K. Der Umfang der Jugendstimme. *Folia Phoniatr*. (Basel) 1957; 9: 225-239.
6. van Oordt, H.W.A., Drost, H.A. Development of the frequency range of the voice in children. *Folia Phoniatr*. (Basel) 1963; 15: 289-298.
7. Ingrisano, D., Weismer, G., Schuckers, G. Sex identification of preschool children's voices. *Folia Phoniatr*. (Basel) 1980; 32: 61-69.
8. Pedersen, M.F., Kitzing, P., Krabbe, S., Heramb, S. The change of voice during puberty in 11 to 18 years old choir singers measured with electropuberty. *Acta Orolaryngol* (Stockh) 1982; 386: 189-192.
9. Pedersen, M.F., Munk, E., Bennet, P., Møller, S. The change of voice during puberty in choir singers measured with phonetograms and compared to androgen status together with other phenomena of puberty. *Proc. Tenth Int. Congr. Phon. Sci.* 1983; p. 604-608.
10. Tanner, J.M. *Growth and adolescence*, 2nd ed. Blackwell Scient. Publ. Oxford 1962.
11. Seidner, W., Schutte, H.K. Standardisierungsvorschlag Stimmfeldmessung/Phonetographie. *Congr. Proc. IXth Congr. UEP, Amsterdam* 1981, p. 88-94.
12. Karlberg, P., Taranger, J. The somatic development of children in a Swedish urban community. *Acta Paediatr. Scand.* 1976; Suppl. 258.
13. Andersen, E. Skeletal maturation of Danish school children in relation to height, sexual development, and social conditions. *Acta Paediatr. Scand.* 1968; Suppl. 185.
14. Hägg, U., Taranger, J. Menarche and voice change as indicators of the pubertal growth spurt. *Octa Odeontol. Scand.* 1980; 38: 179-186.
15. Cohen, S.R., Thompson, J.W., Geller, K.A., Birns, J.W. Voice change in the pediatric patient. A differential diagnosis. *Ann. Otol. Rhinol. Laryngol.* 1983; 92: 437-443.