



LUND UNIVERSITY

Methodology for studies on respiratory and cardiovascular effects of humans at occupational exposure to airborne nanoparticles

Bohgard, Mats; Nielsen, Jörn; Tinnerberg, Håkan; Hagerman, Inger; Berglund, Margareta; Swietlicki, Erik; Gudmundsson, Anders; Pagels, Joakim; Löndahl, Jakob; Nilsson, Erik; Deppert, Knut

Published in:

Book of extended abstracts: 7th Intl Aerosol Conf, St. Paul, Mn, USA (2006)

2006

[Link to publication](#)

Citation for published version (APA):

Bohgard, M., Nielsen, J., Tinnerberg, H., Hagerman, I., Berglund, M., Swietlicki, E., Gudmundsson, A., Pagels, J., Löndahl, J., Nilsson, E., & Deppert, K. (2006). Methodology for studies on respiratory and cardiovascular effects of humans at occupational exposure to airborne nanoparticles. In *Book of extended abstracts: 7th Intl Aerosol Conf, St. Paul, Mn, USA (2006)*

Total number of authors:

11

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

Methodology for Studies on Respiratory and Cardiovascular Effects of Humans at Occupational Exposure to Airborne Nanoparticles

Mats Bohgard¹, Jörn Nielsen², Håkan Tinnerberg², Inger Hagerman³, Margareta Berglund³, Erik Swietlicki⁴, Anders Gudmundsson¹, Joakim Pagels¹, Jakob Löndahl², Erik Nilsson¹, Knut Deppert⁵

¹Ergonomics and Aerosol Technology, Lund University, P.O.Box 118, SE- 221 00 Lund, Sweden

²Occupational and Environmental Health, Lund University, University Hospital, SE 221 85 Lund, Sweden

³Cardiology, Karolinska Institute, Karolinska University Hospital Huddinge, SE 141 86, Stockholm, Sweden

⁴Nuclear Physics, Lund University, P.O.Box 118, SE- 221 00 Lund, Sweden

⁵Solid State Physics, Lund University, P.O.Box 118, SE- 221 00 Lund, Sweden

INTRODUCTION

In production environments, airborne particles have been a threat to workers' health for several hundred years. In 1713 the Italian physician Ramazzini published *De Morbis Artificum* with descriptions of health hazards in a number of different occupations. He observed, for example, that many workers who were exposed to airborne mineral particles (e.g. stone-cutters and quarrymen) got the disease we today know as *silicosis*.

Today the effects on health and environment of the increasing industrial applications of nanotechnology are addressed (Oberdorster et al., 2005). Exploitation of nanotechnology can increase emissions of nano-sized particles into the air in production, manufacturing treatment, handling, usage, waste disposal and recycling of various materials and as the result of accidents such as fires.

Awareness of the health impact of airborne nanoparticles particles is also increasing. Studies on health effects of air pollution show that airborne particles can cause a variety of diseases and are responsible for increased morbidity and mortality (e.g. respiratory and cardiovascular diseases) in polluted areas (Ezzati et al., 2002; Brook et al., 2004; WHO, 2002).

Most of the knowledge about occupational health hazards has been achieved by observations of effects on humans. The aim of this work was to develop and recommend methodology based on direct observations and measurements of effects on humans for dose-response and dose-effect determination at occupational exposures to nanoparticles. The methodology combines competence and techniques from Aerosol physics and technology, Occupational Hygiene, Occupational Medicine, Nanotechnology and Cardiology.

METHODS

The methodology includes measurements and examinations of humans in two model environments 1) Work places where significant exposures are expected to occur and 2) A controlled chamber environment for

exposure provocations. The exposure levels in the chamber are not exceeding the levels found in the subjects' places of work.

Workplace studies

Characterisation of the workplace concerning exposure to nanoparticles: The aerosol characterisation techniques include: dry particle size distributions (3-10 000 nm) measured with differential mobility analyser – condensation particle counter (DMA/CPC) and time-of-flight technique (Aerodynamic Particle Sizer, APS), hygroscopic properties with a Tandem DMA-system (20-500 nm), and filter sampling with subsequent elemental analysis and electron microscopy. An instrument RESPI (Löndahl et al. 2006) will be used to measure, on-line and in-situ, the particle dose for particle diameters 10-1000 nm, to the airways

Examination:

A) Basal examination: Volunteers of the exposed workers are asked to complete a questionnaire about symptoms from airways, the cardiovascular system, smoking habits, atopy, medical and work history. A general physical examination including pulmonary auscultation is performed. A skin prick-test with 13 common allergens is carried out. Each subject performs spirometry and those who complain of asthmatic symptoms will also be tested with methacholine.

B) Diary study: For 3 weeks after having been off work for holiday, the workers record in a diary symptoms from their eyes, airways and cardiovascular systems and results from peak-flow measurements. Before and after the diary study, the workers' are sampled blood for fibrinogen and C-reactive protein analysis. In a subgroup nasal lavage is performed and analysed for mediators and inflammatory markers (Nielsen et al., 1992).

The exposure is assessed for each individual worker based on the aerosol measurements performed and on information about work tasks during the period. The individual exposure assessment will be performed for

several variables regarding particle characteristics as well as peak and number of peak exposures.

Chamber studies

A specially designed stainless steel aerosol chamber (Pagels, 2005) with a floor area of 3x3 m² and height 2.4 m is used for controlled human provocation studies (each individual is exposed for 8 hours). Particles to be used (from work place) are generated into the chamber. Particle concentrations can be controlled as well as ventilation rate, temperature and relative humidity. Aerosol measurement systems (DMA/CPC and tandem-DMA) are connected to the chamber and are used to determine size distribution, number concentration hygroscopicity determination. Filter sampling devices are used for obtaining samples for electron microscopy and elemental analysis. The RESPI instrument is used to determine particle deposition to the airways.

Medical examinations are carried out before, after four hours, and after the provocation. Symptoms from eyes and airways are registered according to a symptom score model. Furthermore, spirometry, acoustic rhinometry, nasal lavage (Nielsen et al., 1994), induced sputum and breath condensate will be obtained. A venous blood sample is obtained before, after, and 24 hours after the provocation for analysis of fibrinogen, C-reactive protein, TNF- α and ICAM-1. From the nasal lavage and induced sputum, cells are separated for differential count and the supernatant is later analysed for markers of inflammation and mediators such as albumin, ECP, ICAM-1, Substance P, TNF- α , IL 1, and IL8. The breath condensate is also analysed for markers of inflammation.

ECG is recorded at rest. Influence of nanoparticles to affect the autonomic regulation of the heart will be studied by analysis of the heart rate variability (HRV) (Heart Rate Variability. Standards of Measurement, Physiological Interpretation and Clinical Use, 1996) Time series of ECG are collected and changes in the sympathetic and the parasympathetic nervous system are studied both in frequency and time domain. Frequency analysis of changes of HRV due to sympathetic tonus are shown in the low frequencies (0.04-0.15 Hz) and to parasympathetic tonus in the high frequencies (0.15- 0.4 Hz), where effects of respiratory frequency also can be observed.

RESULTS

Various parts of the methodology have been successfully applied for nanoparticles from combustion and thermal working processes. Preliminary results from a pilot study of HRV at exposure to an aerosol from burning candle (3-4 10⁴ particles/cm³, median diameter: 12 nm) showed changes in both low and high frequencies compared to low particle concentration.

Keywords: nanoparticles, occupational health

REFERENCES

- Brook R.D., Franklin B., Cascio W., Hong Y.L., Howard G., Lipsett M., Luepker R., Mittleman M., Samet J., Smith S.C. & Tager I. (2004) Air pollution and cardiovascular disease - A statement for healthcare professionals from the expert panel on population and prevention science of the American Heart Association. *Circulation* 109, 2655-2671.
- Ezzati M., Lopez A.D., Rodgers A., Vander Hoorn S. & Murray C.J.L. (2002) Selected major risk factors and global and regional burden of disease. *Lancet* 360, 1347-1360.
- Heart Rate Variability. (1996) Standards of Measurement, Physiological Interpretation and Clinical Use. *Task Force of the European Society of Cardiology and the North American Society of Pacing and Electrophysiology. Circulation* 93:1043-1065.
- Löndahl J., Pagels J., Swietlicki E., Zhou J., Ketzel M., Massling A. & Bohgard M. (2006) A Set-up for Field Studies of Respiratory Tract Deposition of Fine and Ultra-fine Particles in Humans, *in press J Aerosol Science*
- Nielsen J, Welinder H, Horstmann, Skerfving S. (1992) Allergy versus methyltetrahydrophthalic anhydride in epoxy resin workers. *British Journal of Industrial medicine* 49: 769-775.
- Nielsen J, Welinder H, Ottosson H, Bensryd I, Venge P, Skerfving S. (1994) Nasal challenge shows pathogenetic relevance of specific IgE serum antibodies for nasal symptoms caused by hexahydrophthalic anhydride. *Clinical and Experimental Allergy*; 24: 440-449.
- Oberdorster G., Oberdorster E. & Oberdorster J. (2005) Nanotoxicology: An emerging discipline evolving from studies of ultrafine particles. *Environmental Health Perspectives* 113, 823-839
- Pagels J. (2005) *Fine and Ultrafine Particles from Combustion Sources*. Doctoral thesis, Lund University
- Ramazzini B. (1713) *De Morbis Artificum* (2nd edition) Translated by W. C. Wright (1940) *Diseases of Workers*, University of Chicago Press, Chicago
- WHO. (2002) *World Health Report 2002*. Geneva, World Health Organization

The work was supported by the Swedish Council for Working Life and Social Research and the Swedish Research Council for Environment, Agricultural Sciences and Spatial Planning.