Evaluation of some integrals relevant to multiple scattering by randomly distributed obstacles

Kristensson, Gerhard

2014

CODEN:LUTEDX/(TEAT-7228)/1-16/(2014)

Revision No. 2: November 2014

Evaluation of some integrals relevant to multiple scattering by randomly distributed obstacles

Gerhard Kristensson

Electromagnetic Theory
Department of Electrical and Information Technology
Lund University
Sweden

Gerhard Kristensson
Gerhard.Kristensson@eit.lth.se

Department of Electrical and Information Technology
Electromagnetic Theory
Lund University
P.O. Box 118
SE-221 00 Lund
Sweden

Editor: Gerhard Kristensson

Abstract

This paper analyzes and solves an integral and its indefinite Fourier transform of importance in multiple scattering problems of randomly distributed scatterers. The integrand contains a radiating spherical wave, and the two-dimensional domain of integration excludes a circular region of varying size. A solution of the integral in terms of radiating spherical waves is demonstrated. The method employs the Erdélyi operators, which leads to a recursion relation. This recursion relation is solved in terms of a finite sum of radiating spherical waves. The solution of the indefinite Fourier transform of the integral contains the indefinite Fourier transforms of the Legendre polynomials, which are solved by a closed formula.

1 Introduction

In recent years, the electromagnetic scattering problem by randomly distributed objects has been successfully formulated and solved. Some important contributions in the field are found in *e.g.*, [3–8, 10, 11, 13, 16–19, 21–25]. These references refer to various aspects of the topic, and more references can be found in these papers. The topic is also treated in several textbooks, see *e.g.*, [12, 14, 20], which can be consulted for a comprehensive treatment of the various multiple scattering theories.

Of critical importance for the solution of a specific scattering problem with hole-corrections (HC) is an integral of the form [9, 18, 20]

$$I_l(z) = \frac{k^2}{2\pi} \iint\limits_{\mathbb{R}^2} H(r-a) h_l^{(1)}(kr) P_l(\cos\theta)\, \mathrm{d}x\, \mathrm{d}y, \quad z \in \mathbb{R} \tag{1.1}$$

where $H(x)$ denotes the Heaviside function, $h_l^{(1)}(kr)$ the spherical Hankel function, and $P_l(x)$ the Legendre polynomial of order l, respectively. We have also adopted the spherical coordinates, $r = \sqrt{x^2+y^2+z^2}$ and θ ($\cos\theta = z/r$), and the wave number k. The domain of integration is the plane $z =$ constant, excluding the sphere of radius $a > 0$ at the center, see Figure 1. For a given value of $|z| \leq a$, the radius of the excluded circle is $\sqrt{a^2 - z^2}$. For $|z| \geq a$ the integration is the entire x-y plane. This integral, for a given $a > 0$, is a non-trivial function of $z \in \mathbb{R}$. To ensure convergence of the integral at infinity, we assume the wave number k has an arbitrarily small imaginary part. The explicit solution of this integral, as a function of z and the index $l = 0, 1, 2, \ldots$, is the aim of this paper, and the goal is to express the solutions in a form that is attractive from a numerical computation point of view.

The solution of the integral $I_l(z)$ is developed in Sections 2 and 3. The indefinite Fourier transform of $I_l(z)$ is also essential for a successful solution of the multiple scattering problem with hole-corrections, and this analysis is found in Sections 4 and 5. The paper is concluded with a short summary in Section 6.



Figure 1: The geometry of the integration domain — the plane z = constant (dotted line), and the exclusion volume — the sphere of radius a located at the origin (in gray).

2 The integral $I_l(z)$

Rewrite the integral $I_l(z)$ in (1.1) in cylindrical coordinates and perform the integration in the azimuthal angle. We get from (1.1)

$$I_l(z) = k^2 \int_{h(z)}^{\infty} h_l^{(1)}\left(k\sqrt{\rho^2+z^2}\right) P_l\left(z/\sqrt{\rho^2+z^2}\right) \rho \, \mathrm{d}\rho, \quad z \in \mathbb{R} \tag{2.1}$$

where

$$h(z) = \begin{cases} \sqrt{a^2-z^2}, & -a \le z \le a \\ 0, & |z| > a \end{cases}$$

From the parity of the Legendre polynomials, $P_l(-x) = (-1)^l P_l(x)$, we see that also $I_l(-z) = (-1)^l I_l(z)$. Thus, it suffices to evaluate the integral for $z > 0$. In particular, $I_l(0) = 0$, if l is an odd integer. From (2.1) we also easily compute the integral for $l = 0$, *viz.*,

$$I_0(z) = \begin{cases} \mathrm{e}^{-\mathrm{i}kz}, & z \le -a \\ \mathrm{i}kah_0^{(1)}(ka) = \mathrm{e}^{\mathrm{i}ka}, & -a \le z \le a \\ \mathrm{e}^{\mathrm{i}kz}, & z \ge a \end{cases}$$

2.1 Solution outside the interval $[-a, a]$

In the interval $z > a$, the integral is evaluated with the use of the transformation of the outgoing scalar spherical wave in terms of planar waves [2, p. 180], *i.e.*, for a

general value of $z \neq 0$

$$\begin{aligned} & h_l^{(1)}\left(k\sqrt{\rho^2+z^2}\right) P_l\left(\pm|z|/\sqrt{\rho^2+z^2}\right) \\ & = \frac{\mathrm{i}^{-l}}{2\pi} \iint\limits_{\mathbb{R}^2} P_l\left(\pm k_z/k\right) \mathrm{e}^{\mathrm{i}\boldsymbol{k}_\mathrm{t}\cdot\boldsymbol{\rho}+\mathrm{i}k_z|z|} \frac{k}{k_z} \frac{\mathrm{d}k_x\,\mathrm{d}k_y}{k^2}, \quad z \gtrless 0 \end{aligned}$$

where $\boldsymbol{\rho} = x\hat{\boldsymbol{x}} + y\hat{\boldsymbol{y}}$, $\boldsymbol{k}_\mathrm{t} = k_x\hat{\boldsymbol{x}} + k_y\hat{\boldsymbol{y}}$, $k_\mathrm{t} = |\boldsymbol{k}_\mathrm{t}|$, and k_z is defined by

$$k_z = \left(k^2 - k_\mathrm{t}^2\right)^{1/2} = \begin{cases} \sqrt{k^2 - k_\mathrm{t}^2} \text{ for } k_\mathrm{t} < k \\ \mathrm{i}\sqrt{k_\mathrm{t}^2 - k^2} \text{ for } k_\mathrm{t} > k \end{cases}$$

For $z > a$, we get from (1.1)

$$\begin{aligned} I_l(z) &= \frac{k^2}{2\pi} \iint\limits_{\mathbb{R}^2} \frac{\mathrm{i}^{-l}}{2\pi} \left(\iint\limits_{\mathbb{R}^2} P_l\left(k_z/k\right) \mathrm{e}^{\mathrm{i}\boldsymbol{k}_\mathrm{t}\cdot\boldsymbol{\rho}+\mathrm{i}k_z|z|} \frac{k}{k_z} \frac{\mathrm{d}k_x\,\mathrm{d}k_y}{k^2} \right) \mathrm{d}x\,\mathrm{d}y \\ &= \mathrm{i}^{-l} \iint\limits_{\mathbb{R}^2} P_l\left(k_z/k\right) \mathrm{e}^{\mathrm{i}k_z|z|} \delta(\boldsymbol{k}_\mathrm{t}) \frac{k}{k_z}\, \mathrm{d}k_x\,\mathrm{d}k_y = \mathrm{i}^{-l}\mathrm{e}^{\mathrm{i}kz} \end{aligned}$$

by orthogonality or completeness of the planar waves.[1] This implies that the integral for $z > a$ is

$$I_l(z) = \mathrm{i}^{-l}\mathrm{e}^{\mathrm{i}kz}, \quad z > a$$

and consequently, by parity, or analogous calculations

$$I_l(z) = \mathrm{i}^{l}\mathrm{e}^{-\mathrm{i}kz}, \quad z < -a$$

We observe that the integral outside the interval $[-a, a]$ is not singular as $a \to 0$. In fact, the module is constant 1.

3 Solution of the integral $I_l(\eta)$, $-a \leq z \leq a$

We have already obtained a solution of the integral in the interval $|z| > a$, and we now concentrate on finding a solution of the integral in the interval $-a \leq z \leq a$.

The Erdélyi operators $\mathcal{Y}_n^m$ in Ref. 12 are instrumental in finding a closed formula for the integral $I_l(z)$. From [12, Th. 3.13], we have the following very useful result:

$$D\left(h_l^{(1)}(kr)P_l(\cos\theta)\right) = \frac{l+1}{2l+1} h_{l+1}^{(1)}(kr)P_{l+1}(\cos\theta) - \frac{l}{2l+1} h_{l-1}^{(1)}(kr)P_{l-1}(\cos\theta)$$

where $D = -k^{-1}(\partial/\partial z)$. The D operator and the Erdélyi operators are related by $\mathcal{Y}_1^0 = \sqrt{\frac{3}{4\pi}} D_1^0 = \sqrt{\frac{3}{4\pi}} D$.

[1] To ensure convergence of the integral at infinity, assume the wave number k has an arbitrary small, positive imaginary part.

Apply the differential operator D to the integral $I_l(z)$ in (2.1), and use the relation above. We obtain, since $h'(z)h(z) = -z$, the following recursion relation:[2]

$$DI_l(z) = -kzh_l^{(1)}(ka)P_l(z/a) + \frac{l+1}{2l+1}I_{l+1}(z) - \frac{l}{2l+1}I_{l-1}(z), \quad -a \le z \le a$$

with initial condition $I_0(z) = \mathrm{i}kah_0^{(1)}(ka)$.

In the dimensionless variables $\eta = kz$ and $\xi = ka > 0$, this leads to the recursion relation, $l = 0, 1, 2, \ldots$ (note the mild change in notation)

$$I_{l+1}(\eta) = \frac{2l+1}{l+1}\xi h_l^{(1)}(\xi)\frac{\eta}{\xi}P_l(\eta/\xi) - \frac{2l+1}{l+1}\frac{\mathrm{d}}{\mathrm{d}\eta}I_l(\eta) + \frac{l}{l+1}I_{l-1}(\eta), \quad -\xi \le \eta \le \xi$$

The recursion relation is conveniently put in a more generic form by introducing the variable $x = \eta/\xi \in [-1, 1]$. The dependent variable is now x, and ξ is a parameter. Retaining the same notation for the integral, but with a change of the independent variable, we get

$$I_{l+1}(x) = \frac{2l+1}{l+1}\xi h_l^{(1)}(\xi)xP_l(x) - \frac{2l+1}{\xi(l+1)}I_l'(x) + \frac{l}{l+1}I_{l-1}(x), \quad -1 \le x \le 1$$

The following proposition states the surprisingly simple and elegant solution of this recursion relation.

Proposition 3.1. *The recursion relation*

$$I_{l+1}(x) = \frac{2l+1}{l+1}\xi h_l^{(1)}(\xi)xP_l(x) - \frac{2l+1}{\xi(l+1)}I_l'(x) + \frac{l}{l+1}I_{l-1}(x), \quad l = 0, 1, 2, \ldots \tag{3.1}$$

with initial condition

$$I_0(z) = \mathrm{i}\xi h_0^{(1)}(\xi)$$

has the solution

$$\begin{aligned} I_l(x) &= -\xi h_{l+1}^{(1)}(\xi)P_l(x) \\ &\quad + \sum_{k=0}^{[l/2]}(-1)^k\left(\xi h_{l+1-2k}^{(1)}(\xi) + \xi h_{l-1-2k}^{(1)}(\xi)\right)P_{l-2k}(x), \quad l = 0, 1, 2, \ldots \end{aligned} \tag{3.2}$$

[2] Outside the interval $z \in [-a, a]$ the recursion relation reads

$$I_{l+1}(z) = \frac{2l+1}{l+1}DI_l(z) + \frac{l}{l+1}I_{l-1}(z), \quad I_0(z) = \mathrm{e}^{\mathrm{i}kz} \quad z \ge a$$

which is easily solved by induction over the integer l. The result is

$$I_l(z) = \mathrm{i}^{-l}\mathrm{e}^{\mathrm{i}kz}, \quad z \ge a$$

in agreement with the result above.

Proof. We prove the proposition by induction over the integer l. The recursion relation (3.2) is true for $l = 0$, due to the properties of the spherical Hankel functions [15, 10.16.1]. We have from (3.2)

$$I_0(x) = \xi h_{-1}^{(1)}(\xi) = \xi \left(\frac{\pi}{2\xi}\right)^{1/2} H_{-1/2}^{(1)}(\xi) = \mathrm{i}\xi \left(\frac{\pi}{2\xi}\right)^{1/2} H_{1/2}^{(1)}(\xi) = \mathrm{i}\xi h_0^{(1)}(\xi)$$

Now assume the solution (3.2) holds for all integers less than or equal to l, and we want to prove that it holds for $l+1$. We have from (3.1) and the induction assumption

$$\begin{aligned}
I_{l+1}(x) &= \frac{2l+1}{l+1}\xi h_l^{(1)}(\xi) x P_l(x) - \frac{2l+1}{\xi(l+1)} I_l'(x) + \frac{l}{l+1} I_{l-1}(x) \\
&= \xi h_l^{(1)}(\xi) P_{l+1}(x) + \frac{2l+1}{\xi(l+1)} \xi h_{l+1}^{(1)}(\xi) P_l'(x) \\
&- \frac{2l+1}{\xi(l+1)} \sum_{k=0}^{[l/2]} (-1)^k \left(\xi h_{l+1-2k}^{(1)}(\xi) + \xi h_{l-1-2k}^{(1)}(\xi)\right) P_{l-2k}'(x) \\
&+ \frac{l}{l+1} \sum_{k=0}^{[(l-1)/2]} (-1)^k \left(\xi h_{l-2k}^{(1)}(\xi) + \xi h_{l-2-2k}^{(1)}(\xi)\right) P_{l-1-2k}(x)
\end{aligned}$$

where we used the following recursion relation for the Legendre polynomials:

$$(2l+1) x P_l(x) = (l+1) P_{l+1}(x) + l P_{l-1}(x)$$

We conclude that $I_{l+1}(x)$ is a polynomial in x of the order $l+1$, and therefore can be expanded in a series of Legendre polynomials. The form is

$$I_{l+1}(x) = \sum_{n=0}^{[(l+1)/2]} a_n P_{l+1-2n}(x)$$

where a_n depends on l and ξ. The coefficients a_n are determined by orthogonality of the Legendre polynomials.

$$a_n = \frac{2l+3-4n}{2} \int_{-1}^{1} I_{l+1}(x) P_{l+1-2n}(x) \, \mathrm{d}x$$

The first coefficient is special.

$$a_0 = \xi h_l^{(1)}(\xi) = -\xi h_{l+2}^{(1)}(\xi) + \left(\xi h_{l+2}^{(1)}(\xi) + \xi h_l^{(1)}(\xi)\right)$$

Proceed in the same way with the remaining coefficients, $n = 1, 2, \ldots, [(l+1)/2]$.

$$\begin{aligned}
a_n &= \frac{2l+1}{l+1} \frac{2l+3-4n}{2} h_{l+1}^{(1)}(\xi) I_{l,l+1-2n} \\
&- \frac{2l+1}{l+1} \frac{2l+3-4n}{2} \sum_{k=0}^{[l/2]} (-1)^k \left(h_{l+1-2k}^{(1)}(\xi) + h_{l-1-2k}^{(1)}(\xi)\right) I_{l-2k,l+1-2n} \\
&+ \frac{l}{l+1} \sum_{k=0}^{[(l-1)/2]} (-1)^k \left(\xi h_{l-2k}^{(1)}(\xi) + \xi h_{l-2-2k}^{(1)}(\xi)\right) \delta_{k,n-1}
\end{aligned}$$

where we used the notion

$$I_{k,n} = \int_{-1}^{1} P_k'(x)P_n(x)\ \mathrm{d}x = \begin{cases} 0, & 0 \le k \le n \\ 1-(-1)^{k+n}, & 0 \le n < k \end{cases}$$

Use this result, and the following recursion relation for the spherical Hankel functions:

$$(2l+1)h_l^{(1)}(\xi) = \xi h_{l+1}^{(1)}(\xi) + \xi h_{l-1}^{(1)}(\xi) \tag{3.3}$$

We get

$$\begin{aligned} a_n &= \frac{2l+1}{l+1}(2l+3-4n)\Bigg(h_{l+1}^{(1)}(\xi) - \underbrace{\sum_{k=0}^{n-1}(-1)^k\left(h_{l+1-2k}^{(1)}(\xi)+h_{l-1-2k}^{(1)}(\xi)\right)}_{=h_{l+1}^{(1)}(\xi)+(-1)^{n-1}h_{l+1-2n}^{(1)}(\xi)}\Bigg) \\ &\quad + \frac{l}{l+1}(-1)^{n-1}\left(\xi h_{l+2-2n}^{(1)}(\xi)+\xi h_{l-2n}^{(1)}(\xi)\right) \\ &= \frac{2l+1}{l+1}(-1)^n(2l+3-4n)h_{l+1-2n}^{(1)}(\xi) - \frac{l}{l+1}(-1)^n\left(\xi h_{l+2-2n}^{(1)}(\xi)+\xi h_{l-2n}^{(1)}(\xi)\right) \\ &= (-1)^n\left(\xi h_{l+2-2n}^{(1)}(\xi)+\xi h_{l-2n}^{(1)}(\xi)\right) \end{aligned}$$

Collecting the results gives

$$\begin{aligned} I_{l+1}(x) &= \sum_{n=0}^{[(l+1)/2]} a_n P_{l+1-2n}(x) \\ &= -\xi h_{l+2}^{(1)}(\xi)P_{l+1}(x) + \sum_{n=0}^{[(l+1)/2]}(-1)^n\left(\xi h_{l+2-2n}^{(1)}(\xi)+\xi h_{l-2n}^{(1)}(\xi)\right)P_{l+1-2n}(x) \end{aligned}$$

which is the statement (3.2) for $l+1$, and the proposition is proved. □

Alternative expressions of the integral $I(z)$ in the interval $z \in [-a,a]$ can be found. The following corollary shows some.

Corollary 3.1. *The integral $I(z)$ in Proposition 3.1 has the following alternative expressions:*

$$I_l(x) = -\xi h_{l+1}^{(1)}(\xi)P_l(x) + \sum_{k=0}^{[l/2]}(-1)^k(2l-4k+1)h_{l-2k}^{(1)}(\xi)P_{l-2k}(x), \quad l=0,1,2,\ldots \tag{3.4}$$

and

$$\begin{aligned} I_l(x) &= \mathrm{i}^{1-l}\xi h_0^{(1)}(\xi)P_{l-2[l/2]}(x) \\ &\quad + \sum_{k=0}^{[l/2]-1}(-1)^k\xi h_{l-2k-1}^{(1)}(\xi)\left(P_{l-2k}(x)-P_{l-2k-2}(x)\right), \quad l=0,1,2,\ldots \end{aligned} \tag{3.5}$$

and

$$I_l(x) = \mathrm{i}^{1-l}\xi h_0^{(1)}(\xi)P_{l-2[l/2]}(x) - \sum_{k=0}^{[l/2]-1}(-1)^k\xi h_{l-2k-1}^{(1)}(\xi)\frac{2l-4k-1}{(l-2k-1)(l-2k)}P'_{l-2k-1}(x), \quad l=0,1,2,\ldots \tag{3.6}$$

where the two last sums are zero for $l = 0, 1$.

Proof. The solution in (3.4) is equivalent to (3.2), which is easily seen since the spherical Hankel functions $h_l^{(1)}(\xi)$ satisfy the recursion relation (3.3). The representation in (3.5) is simply a rearrangement of the sum in (3.2). We obtain from (3.2) ($l = 0, 1, 2, \ldots$)

$$\begin{aligned} I_l(x) &= \sum_{k=0}^{[l/2]-1}(-1)^k\xi h_{l-1-2k}^{(1)}(\xi)\left(P_{l-2k}(x) - P_{l-2-2k}(x)\right) \\ &\qquad + (-1)^{[l/2]}\xi h_{l-1-2[l/2]}^{(1)}(\xi)P_{l-2[l/2]}(x) \\ &= \sum_{k=0}^{[l/2]-1}(-1)^k\xi h_{l-1-2k}^{(1)}(\xi)\left(P_{l-2k}(x) - P_{l-2-2k}(x)\right) \\ &\qquad + \mathrm{i}^{1-l}\xi h_0^{(1)}(\xi)P_{l-2[l/2]}(x) \end{aligned}$$

where we used [15, 10.16.1]

$$h_{-1}^{(1)}(\xi) = \mathrm{i}h_0^{(1)}(\xi)$$

Finally, the relation (3.6) from (3.5) with the use of the recursion relation

$$l(l+1)\left(P_{l+1}(x) - P_{l-1}(x)\right) = -(2l+1)(1-x^2)P'_l(x)$$

□

In the original variables z and a, we have

$$I_l(z) = \mathrm{i}^{1-l}kah_0^{(1)}(ka)P_{l-2[l/2]}(z/a) + \sum_{n=0}^{[l/2]-1}(-1)^n kah_{l-2n-1}^{(1)}(ka)\left(P_{l-2n}(z/a) - P_{l-2n-2}(z/a)\right), \quad l=0,1,2,\ldots$$

or

$$I_l(z) = -kah_{l+1}^{(1)}(ka)P_l(z/a) + \sum_{k=0}^{[l/2]}(-1)^k(2l-4k+1)h_{l-2k}^{(1)}(ka)P_{l-2k}(z/a), \quad l=0,1,2,\ldots$$

and we see that the integral $I_l(z)$ can be written as a finite sum of spherical waves (except the first term). The most singular term in powers of ka is of the order $(ka)^{1-l}$ (order $O(1)$ if $l = 0$), which is most easily seen from the representation in (3.5).

4 Fourier transform of $I_l(z)$

The indefinite Fourier transform of the function $I_l(z)$ has also importance in the analysis of [9]. More specifically, our goal in this section is to compute

$$\widehat{I}_l^{\pm}(z) = k\int_{z_0}^{z} I_l(t)\mathrm{e}^{\pm \mathrm{i}kt}\,\mathrm{d}t, \quad z \geq z_0, \quad l = 0, 1, 2, \ldots \tag{4.1}$$

where z_0 is a fixed number such that $z_0 < -a$.

The function $I_l(t)$ has explicit forms in the three intervals $[z_0, -a]$, $(-a, a)$, and $[a, \infty)$. The explicit forms are:

$$I_l(t) = \mathrm{i}^l \mathrm{e}^{-\mathrm{i}kt}, \quad t \leq -a$$

and in the interval $t \in (-a, a)$ as a finite sum of spherical waves

$$\begin{aligned} I_l(t) &= \mathrm{i}^{1-l} kah_0^{(1)}(ka) P_{l-2[l/2]}(t/a) \\ &\quad + \sum_{n=0}^{[l/2]-1} (-1)^n kah_{l-2n-1}^{(1)}(ka)\left(P_{l-2n}(t/a) - P_{l-2n-2}(t/a)\right) \end{aligned}$$

In the interval $t \geq a$

$$I_l(t) = \mathrm{i}^{-l}\mathrm{e}^{\mathrm{i}kt}$$

To compute the indefinite Fourier transform we need to calculate the function

$$h_l^{\pm}(z) = k\int_{-a}^{z} P_l(t/a)\mathrm{e}^{\pm \mathrm{i}kt}\,\mathrm{d}t = ka\int_{-1}^{z/a} P_l(t)\mathrm{e}^{\pm \mathrm{i}kat}\,\mathrm{d}t, \quad |z| \leq a \tag{4.2}$$

For $z = a$ the integral is a spherical Bessel function, *viz.*,

$$h_l^{\pm}(a) = k\int_{-a}^{a} P_l(t/a)\mathrm{e}^{\pm \mathrm{i}kt}\,\mathrm{d}t = ka\int_{-1}^{1} P_l(t)\mathrm{e}^{\pm \mathrm{i}kat}\,\mathrm{d}t = 2ka(\pm\mathrm{i})^l j_l(ka)$$

We divide the interval $[z_0, z]$ in three parts. In the interval $z_0 \leq z < -a$, we have

$$\widehat{I}_l^{\pm}(z) = \mathrm{i}^l k\int_{z_0}^{z} \mathrm{e}^{\mathrm{i}(\pm 1-1)kt}\,\mathrm{d}t = \mathrm{i}^l \begin{cases} k(z - z_0) \\ \frac{1}{2\mathrm{i}}\left(\mathrm{e}^{-2\mathrm{i}kz_0} - \mathrm{e}^{-2\mathrm{i}kz}\right) \end{cases}$$

and in the interval $-a < z < a$, we have

$$\begin{aligned} \widehat{I}_l^{\pm}(z) &= \mathrm{i}^l \begin{cases} k(-a - z_0) \\ \frac{1}{2\mathrm{i}}\left(\mathrm{e}^{-2\mathrm{i}kz_0} - \mathrm{e}^{2\mathrm{i}ka}\right) \end{cases} + \mathrm{i}^{1-l} kah_0^{(1)}(ka) h_{l-2[l/2]}^{\pm}(z) \\ &\quad + \sum_{n=0}^{[l/2]-1} (-1)^n kah_{l-2n-1}^{(1)}(ka)\left(h_{l-2n}^{\pm}(z) - h_{l-2n-2}^{\pm}(z)\right) \end{aligned}$$

and in the interval $a < z$, we have

$$\widehat{I}_l^{\pm}(z) = \mathrm{i}^l \begin{cases} k(-a-z_0) \\ \frac{1}{2\mathrm{i}}\left(\mathrm{e}^{-2\mathrm{i}kz_0} - \mathrm{e}^{2\mathrm{i}ka}\right) \end{cases} + \mathrm{i}^{1-l} 2(ka)^2 h_0^{(1)}(ka)(\pm\mathrm{i})^{l-2[l/2]} j_{l-2[l/2]}(ka)$$
$$+ 2(ka)^2(\pm\mathrm{i})^l \sum_{n=0}^{[l/2]-1} h_{l-2n-1}^{(1)}(ka)\left(j_{l-2n}(ka) + j_{l-2n-2}(ka)\right)$$
$$+ \mathrm{i}^{-l} \begin{cases} \frac{1}{2\mathrm{i}}\left(\mathrm{e}^{2\mathrm{i}kz} - \mathrm{e}^{2\mathrm{i}ka}\right) \\ k(z-a) \end{cases}$$

5 Indefinite integral of Legendre polynomials

It remains to find an effective method to compute the functions $h_l^{\pm}(z)$ in (4.2). To this end, define

$$h_l(\eta,\zeta) = \int_{-1}^{\eta} P_l(t)\mathrm{e}^{\mathrm{i}\zeta t}\,\mathrm{d}t, \quad |\eta| \le 1 \tag{5.1}$$

We see that $h_l(1,\zeta) = 2\mathrm{i}^l j_l(\zeta)$. In terms of the functions $h_l(\eta,\zeta)$, the functions $h_l^{\pm}(z)$ are

$$h_l^{\pm}(z) = kah_l(z/a, \pm ka)$$

Our ambition in this section is to find an efficient method to compute the integrals in (5.1). We express the function $h_l(\eta,\zeta)$ as a recursion relation.

5.1 Solution by recursion

The following recursion relation of Legendre polynomials is useful:

$$P_l(t) = \frac{1}{2l+1}\left(P'_{l+1}(t) - P'_{l-1}(t)\right)$$

Integration by parts then implies ($P_l(-1) = (-1)^l$)

$$h_l(\eta,\zeta) = \int_{-1}^{\eta} P_l(t)\mathrm{e}^{\mathrm{i}\zeta t}\,\mathrm{d}t = \frac{1}{2l+1}\int_{-1}^{\eta}\left(P'_{l+1}(t) - P'_{l-1}(t)\right)\mathrm{e}^{\mathrm{i}\zeta t}\,\mathrm{d}t$$
$$= \frac{1}{2l+1}\left(P_{l+1}(\eta) - P_{l-1}(\eta)\right)\mathrm{e}^{\mathrm{i}\zeta\eta} - \frac{\mathrm{i}\zeta}{2l+1}\left(h_{l+1}(\eta,\zeta) - h_{l-1}(\eta,\zeta)\right)$$

or solving for $h_{l+1}(\eta,\zeta)$

$$h_{l+1}(\eta,\zeta) = \frac{1}{\mathrm{i}\zeta}\left(P_{l+1}(\eta) - P_{l-1}(\eta)\right)\mathrm{e}^{\mathrm{i}\zeta\eta} - \frac{2l+1}{\mathrm{i}\zeta}h_l(\eta,\zeta) + h_{l-1}(\eta,\zeta), \quad l = 1,2,3,\ldots$$

The functions $h_l(\eta,\zeta)$ can therefore be found by iteration with starting values

$$h_0(\eta,\zeta) = \frac{1}{\mathrm{i}\zeta}\left(\mathrm{e}^{\mathrm{i}\zeta\eta} - \mathrm{e}^{-\mathrm{i}\zeta}\right) = \frac{1}{\mathrm{i}\zeta}P_0(\eta)\mathrm{e}^{\mathrm{i}\zeta\eta} + h_0^{(2)}(\zeta) = \eta h_0^{(1)}(\zeta\eta) + h_0^{(2)}(\zeta)$$

and

$$h_1(\eta,\zeta) = \frac{1}{\mathrm{i}\zeta}\left(\eta \mathrm{e}^{\mathrm{i}\zeta\eta} + \mathrm{e}^{-\mathrm{i}\zeta}\right) + \frac{1}{\zeta^2}\left(\mathrm{e}^{\mathrm{i}\zeta\eta} - \mathrm{e}^{-\mathrm{i}\zeta}\right)$$
$$= \frac{1}{\mathrm{i}\zeta}\left(P_1(\eta) - \frac{1}{\mathrm{i}\zeta}P_0(\eta)\right)\mathrm{e}^{\mathrm{i}\zeta\eta} + \mathrm{i}h_1^{(2)}(\zeta) = \mathrm{i}\eta^2 h_1^{(1)}(\zeta\eta) + \mathrm{i}h_1^{(2)}(\zeta)$$

To find the general solution to this recursion scheme, we start by solving the homogeneous difference equation.

Lemma 5.1. *The solution to the homogeneous difference equation*

$$a_{l+1} + \frac{2l+1}{\mathrm{i}\zeta}a_l - a_{l-1} = 0, \quad l = 1,2,3,\ldots$$

given the initial values a_0 and a_1 is

$$a_l = -\frac{\zeta^2}{2\mathrm{i}}\left(a_0 h_0^{(2)'}(\zeta) - \mathrm{i}a_1 h_0^{(2)}(\zeta)\right)\mathrm{i}^l h_l^{(1)}(\zeta)$$
$$+ \frac{\zeta^2}{2\mathrm{i}}\left(a_0 h_0^{(1)'}(\zeta) - \mathrm{i}a_1 h_0^{(1)}(\zeta)\right)\mathrm{i}^l h_l^{(2)}(\zeta), \quad l = 2,3,4,\ldots$$

Proof. Two linearly independent solutions to the homogeneous difference equation in the lemma are $\mathrm{i}^l h_l^{(1)}(\zeta)$ and $\mathrm{i}^l h_l^{(2)}(\zeta)$, which is easily proved by the recursion relation $f_{l+1}(z) - (2l+1)f_l(z)/z + f_{l-1}(z) = 0$, where $f_l(z)$ is any spherical Bessel or Hankel function. The general solution therefore is

$$a_l = c_1 \mathrm{i}^l h_l^{(1)}(\zeta) + c_2 \mathrm{i}^l h_l^{(2)}(\zeta), \quad l = 2,3,4,\ldots$$

where c_1 and c_2 are constants determined by the starting values a_0 and a_1. Explicitly, we get

$$\begin{cases} c_1 h_0^{(1)}(\zeta) + c_2 h_0^{(2)}(\zeta) = a_0 \\ c_1 \mathrm{i} h_1^{(1)}(\zeta) + c_2 \mathrm{i} h_1^{(2)}(\zeta) = a_1 \end{cases}$$

with solution

$$\begin{cases} c_1 = -\dfrac{\zeta^2}{2\mathrm{i}}\left(a_0 h_0^{(2)'}(\zeta) - \mathrm{i}a_1 h_0^{(2)}(\zeta)\right) \\ c_2 = \dfrac{\zeta^2}{2\mathrm{i}}\left(a_0 h_0^{(1)'}(\zeta) - \mathrm{i}a_1 h_0^{(1)}(\zeta)\right) \end{cases}$$

where we used the Wronskian of the spherical Hankel functions.

$$h_n^{(2)}(z)h_n^{(1)'}(z) - h_n^{(2)'}(z)h_n^{(1)}(z) = \frac{2\mathrm{i}}{z^2}$$

and $h_0^{(1,2)'}(z) = -h_1^{(1,2)}(z)$. This completes the proof of the lemma. □

We are now ready for the solution to the inhomogeneous difference equation in $h_l(\eta,\zeta)$ above. We formulate this as a lemma.

Lemma 5.2. *Define an iteration scheme by*

$$h_{l+1}(\eta,\zeta) = \frac{1}{\mathrm{i}\zeta}\left(P_{l+1}(\eta) - P_{l-1}(\eta)\right)\mathrm{e}^{\mathrm{i}\zeta\eta} - \frac{2l+1}{\mathrm{i}\zeta}h_l(\eta,\zeta) + h_{l-1}(\eta,\zeta), \quad l = 1,2,3,\ldots$$

with starting values

$$h_0(\eta,\zeta) = \eta h_0^{(1)}(\zeta\eta) + h_0^{(2)}(\zeta)$$

and

$$h_1(\eta,\zeta) = \mathrm{i}\left(\eta^2 h_1^{(1)}(\zeta\eta) + h_1^{(2)}(\zeta)\right)$$

The solution is

$$h_l(\eta,\zeta) = f_l(\eta,\zeta)\mathrm{e}^{\mathrm{i}\zeta\eta} + \mathrm{i}^l h_l^{(2)}(\zeta), \quad l = 0,1,2,3,\ldots$$

where

$$f_l(\eta,\zeta) = \mathrm{i}^l h_l^{(1)}(\zeta)\left\{\sum_{k=1}^{l}\frac{1}{\zeta h_{k-1}^{(1)}(\zeta)h_k^{(1)}(\zeta)}\left(-\sum_{n=0}^{k}\mathrm{i}^{-n+1}(2n+1)\frac{h_n^{(1)}(\zeta)}{\zeta}P_n(\eta)\right.\right.$$
$$\left.\left.+\,\mathrm{i}^{-k+2}h_k^{(1)}(\zeta)P_{k-1}(\eta) + \mathrm{i}^{-k+1}h_{k+1}^{(1)}(\zeta)P_k(\eta)\right) - \mathrm{i}\frac{P_0(\eta)}{\zeta h_0^{(1)}(\zeta)}\right\}, \quad l = 0,1,2,\ldots$$

Proof. We first subtract the part of the solution that contains the spherical Hankel function of the second kind $h_l^{(2)}(\zeta)$ and the exponential function $\mathrm{e}^{\mathrm{i}\zeta\eta}$. To this end, let $h_l(\eta,\zeta) = f_l(\eta,\zeta)\mathrm{e}^{\mathrm{i}\zeta\eta} + \mathrm{i}^l h_l^{(2)}(\zeta)$. The recursion relation for $f_l(\eta,\zeta)$ is easily found by the use of the recursion relation $h_{l+1}^{(2)}(z) = (2l+1)h_l^{(2)}(z)/z - h_{l-1}^{(2)}(z)$. We get the new difference equation

$$f_{l+1}(\eta,\zeta) = \frac{1}{\mathrm{i}\zeta}\left(P_{l+1}(\eta) - P_{l-1}(\eta)\right) - \frac{2l+1}{\mathrm{i}\zeta}f_l(\eta,\zeta) + f_{l-1}(\eta,\zeta), \quad l = 1,2,3,\ldots$$

with starting values

$$f_0(\eta,\zeta) = \frac{1}{\mathrm{i}\zeta}P_0(\eta)$$

and

$$f_1(\eta,\zeta) = \frac{1}{\mathrm{i}\zeta}\left(P_1(\eta) - \frac{1}{\mathrm{i}\zeta}P_0(\eta)\right)$$

To simplify the notation, we put the difference equation in a standard form [1].

$$a_{n+2} + p_1(n)a_{n+1} + p_0(n)a_n = q(n), \quad n = 1,2,\ldots$$

where

$$\begin{cases} a_n = f_{n-1}(\eta,\zeta) \\ p_1(n) = \dfrac{2n+1}{\mathrm{i}\zeta} \\ p_0(n) = -1 \\ q(n) = \dfrac{1}{\mathrm{i}\zeta}\left(P_{n+1}(\eta) - P_{n-1}(\eta)\right) \end{cases}$$

with initial values

$$\begin{cases} a_1 = \dfrac{1}{\mathrm{i}\zeta} P_0(\eta) \\ a_2 = \dfrac{1}{\mathrm{i}\zeta}\left(P_1(\eta) - \dfrac{1}{\mathrm{i}\zeta} P_0(\eta)\right) \end{cases}$$

A solution to the homogeneous difference equation is (see Lemma 5.1)

$$y_l = \mathrm{i}^{l-1} h_{l-1}^{(1)}(\zeta)$$

The final solution then is [1], $(n = 3, 4, \ldots)$

$$a_n = \left(\sum_{k=1}^{n-1}\prod_{j=1}^{k-1}\frac{p_0(j)y_j}{y_{j+2}}\left(\sum_{l=1}^{k-1}\frac{q(l)}{y_{l+2}}\left[\prod_{m=1}^{l}\frac{p_0(m)y_m}{y_{m+2}}\right]^{-1} + \frac{a_2}{y_2} - \frac{a_1}{y_1}\right) + \frac{a_1}{y_1}\right) y_n$$

Insert the explicit values, and we obtain

$$\begin{aligned} f_l(\eta,\zeta) = \Bigg\{ \sum_{k=1}^{l} \frac{1}{\zeta h_{k-1}^{(1)}(\zeta) h_k^{(1)}(\zeta)} \Bigg(-\sum_{n=1}^{k-1} h_n^{(1)}(\zeta) \frac{P_{n+1}(\eta) - P_{n-1}(\eta)}{\mathrm{i}^n} + \mathrm{i} h_1^{(1)}(\zeta) P_0(\eta) \\ - h_0^{(1)}(\zeta)\left(P_1(\eta) + \mathrm{i}\frac{1}{\zeta}P_0(\eta)\right)\Bigg) - \mathrm{i}\frac{P_0(\eta)}{\zeta h_0^{(1)}(\zeta)}\Bigg\} \mathrm{i}^l h_l^{(1)}(\zeta), \quad l = 2, 3, 4, \ldots \end{aligned}$$

This relation holds also for $l = 0, 1$, provided the sums with upper limit smaller than the lower limit are interpreted as zero.

We now simplify the sum in this expression.

$$\begin{aligned} S &= -\sum_{n=1}^{k-1} \mathrm{i}^{-n} h_n^{(1)}(\zeta)\left(P_{n+1}(\eta) - P_{n-1}(\eta)\right) + \mathrm{i} h_1^{(1)}(\zeta) P_0(\eta) - h_0^{(1)}(\zeta) P_1(\eta) \\ &= \mathrm{i} h_1^{(1)}(\zeta)\left(P_2(\eta) - P_0(\eta)\right) + h_2^{(1)}(\zeta)\left(P_3(\eta) - P_1(\eta)\right) - \mathrm{i} h_3^{(1)}(\zeta)\left(P_4(\eta) - P_2(\eta)\right) \\ &\quad + \ldots - \mathrm{i}^{-k+2} h_{k-2}^{(1)}(\zeta)\left(P_{k-1}(\eta) - P_{k-3}(\eta)\right) - \mathrm{i}^{-k+1} h_{k-1}^{(1)}(\zeta)\left(P_k(\eta) - P_{k-2}(\eta)\right) \\ &\quad + \mathrm{i} h_1^{(1)}(\zeta) P_0(\eta) - h_0^{(1)}(\zeta) P_1(\eta) \\ &= -\left(h_0^{(1)}(\zeta) + h_2^{(1)}(\zeta)\right) P_1(\eta) + \mathrm{i}\left(h_1^{(1)}(\zeta) + h_3^{(1)}(\zeta)\right) P_2(\eta) \\ &\quad + \left(h_2^{(1)}(\zeta) + h_4^{(1)}(\zeta)\right) P_3(\eta) - \mathrm{i}\left(h_3^{(1)}(\zeta) + h_5^{(1)}(\zeta)\right) P_4(\eta) + \ldots \\ &\quad - \mathrm{i}^{-k+2}\left(h_{k-2}^{(1)}(\zeta) + h_k^{(1)}(\zeta)\right) P_{k-1}(\eta) - \mathrm{i}^{-k+1}\left(h_{k-1}^{(1)}(\zeta) + h_{k+1}^{(1)}(\zeta)\right) P_k(\eta) \\ &\quad + \mathrm{i}^{-k+2} h_k^{(1)}(\zeta) P_{k-1}(\eta) + \mathrm{i}^{-k+1} h_{k+1}^{(1)}(\zeta) P_k(\eta) \end{aligned}$$

The recursion relation $h_{l+1}^{(1)}(z) + h_{l-1}^{(1)}(z) = (2l+1) h_l^{(1)}(z)/z$ implies

$$\begin{aligned} S &= -3\frac{h_1^{(1)}(\zeta)}{\zeta} P_1(\eta) + 5\mathrm{i}\frac{h_2^{(1)}(\zeta)}{\zeta} P_2(\eta) + 7\frac{h_3^{(1)}(\zeta)}{\zeta} P_3(\eta) - 9\mathrm{i}\frac{h_4^{(1)}(\zeta)}{\zeta} P_4(\eta) + \ldots \\ &\quad - \mathrm{i}^{-k+1}(2k+1)\frac{h_k^{(1)}(\zeta)}{\zeta} P_k(\eta) + \mathrm{i}^{-k+2} h_k^{(1)}(\zeta) P_{k-1}(\eta) + \mathrm{i}^{-k+1} h_{k+1}^{(1)}(\zeta) P_k(\eta) \\ &= -\sum_{n=1}^{k} \mathrm{i}^{-n+1}(2n+1)\frac{h_n^{(1)}(\zeta)}{\zeta} P_n(\eta) + \mathrm{i}^{-k+2} h_k^{(1)}(\zeta) P_{k-1}(\eta) + \mathrm{i}^{-k+1} h_{k+1}^{(1)}(\zeta) P_k(\eta) \end{aligned}$$

which gives

$$f_l(\eta,\zeta) = \left\{ \sum_{k=1}^{l} \frac{1}{\zeta h_{k-1}^{(1)}(\zeta) h_k^{(1)}(\zeta)} \left(-\sum_{n=1}^{k} \mathrm{i}^{-n+1}(2n+1)\frac{h_n^{(1)}(\zeta)}{\zeta} P_n(\eta) \right.\right.$$
$$\left. + \mathrm{i}^{-k+2} h_k^{(1)}(\zeta) P_{k-1}(\eta) + \mathrm{i}^{-k+1} h_{k+1}^{(1)}(\zeta) P_k(\eta) - \mathrm{i}\frac{h_0^{(1)}(\zeta)}{\zeta} P_0(\eta) \right)$$
$$\left. - \mathrm{i}\frac{P_0(\eta)}{\zeta h_0^{(1)}(\zeta)} \right\} \mathrm{i}^l h_l^{(1)}(\zeta)$$

or

$$f_l(\eta,\zeta) = \left\{ \sum_{k=1}^{l} \frac{1}{\zeta h_{k-1}^{(1)}(\zeta) h_k^{(1)}(\zeta)} \left(-\sum_{n=0}^{k} \mathrm{i}^{-n+1}(2n+1)\frac{h_n^{(1)}(\zeta)}{\zeta} P_n(\eta) \right.\right.$$
$$\left.\left. + \mathrm{i}^{-k+2} h_k^{(1)}(\zeta) P_{k-1}(\eta) + \mathrm{i}^{-k+1} h_{k+1}^{(1)}(\zeta) P_k(\eta) \right) - \mathrm{i}\frac{P_0(\eta)}{\zeta h_0^{(1)}(\zeta)} \right\} \mathrm{i}^l h_l^{(1)}(\zeta)$$

This completes the lemma. □

In conclusion, the functions $h_l^{\pm}(z)$ defined in (4.2) can be expressed in the function $h(\eta,\zeta)$ in (5.1). Specifically, we have

$$h_l^{\pm}(z) = kah_l(z/a, \pm ka)$$

6 Summary and explicit terms

This paper contains an evaluation of a non-trivial integral that occurs in the formulation of scattering by randomly distributed obstacles.

To summarize, the integral $I_l(z)$ in (1.1) has been solved and the solution outside the interval $[-a,a]$ is a simple exponential function in kz, while inside the interval $[-a,a]$, the solution can be found in a finite series of spherical waves. The finite sum of spherical waves depends on the two parameters kz and ka, or, more precisely, the parameter ka and a polynomial of the order l in the parameter z/a. Several equivalent solutions are presented in the paper, one of them is ($l = 0, 1, 2, \ldots$)

$$I_l(z) = \begin{cases} \mathrm{i}^l \mathrm{e}^{-\mathrm{i}kz}, \quad z \le -a \\ \mathrm{i}^{1-l} kah_0^{(1)}(ka) P_{l-2[l/2]}(z/a) \\ \quad + \displaystyle\sum_{n=0}^{[l/2]-1} (-1)^n kah_{l-2n-1}^{(1)}(ka) \left(P_{l-2n}(z/a) - P_{l-2n-2}(z/a) \right), \quad z \in [-a,a] \\ \mathrm{i}^{-l} \mathrm{e}^{\mathrm{i}kz}, \quad z \ge a \end{cases}$$

The first integrals, $l = 0, 1, 2$, are of interest for low-frequency expansions. For $l = 0$ the integral is

$$I_0(z) = \begin{cases} \mathrm{e}^{-\mathrm{i}kz}, & z \le -a \\ \mathrm{e}^{\mathrm{i}ka}, & z \in [-a,a] \\ \mathrm{e}^{\mathrm{i}kz}, & z \ge a \end{cases}$$

and for $l = 1$ the result is

$$I_1(z) = \begin{cases} \mathrm{i}\mathrm{e}^{-\mathrm{i}kz}, & z \le -a \\ -\mathrm{i}\mathrm{e}^{\mathrm{i}ka}\dfrac{z}{a}, & z \in [-a, a] \\ -\mathrm{i}\mathrm{e}^{\mathrm{i}kz}, & z \ge a \end{cases}$$

For $l = 2$ the result is

$$I_2(z) = \begin{cases} -\mathrm{e}^{-\mathrm{i}kz}, & z \le -a \\ \mathrm{e}^{\mathrm{i}ka}\dfrac{(ka)^2(3\mathrm{i} + ka) - 3(\mathrm{i} + ka)(kz)^2}{2(ka)^3}, & z \in [-a, a] \\ -\mathrm{e}^{\mathrm{i}kz}, & z \ge a \end{cases}$$

and we notice that the integral contains a polynomial in z/a of order l.

Moreover, the indefinite Fourier transform of $I_l(z)$ has also been investigated. More precisely, the integral, see (4.1)

$$\widehat{I}_l^{\pm}(z) = k \int_{z_0}^{z} I_l(t)\mathrm{e}^{\pm \mathrm{i}kt}\, \mathrm{d}t, \quad z \ge z_0, \quad l = 0, 1, 2, \ldots$$

is shown to have a solution expressed in spherical waves.

Acknowledgement

The author is grateful to an anonymous reviewer for pointing out the possibility to use the Erdélyi operators $\mathcal{Y}_n^m$ in Ref. 12 in the solution of the integral $I_l(z)$.These operators systemized the solution considerably.

References

[1] C. M. Bender and S. A. Orszag. *Advanced Mathematical Methods for Scientists and Engineers.* McGraw-Hill, New York, 1978.

[2] A. Boström, G. Kristensson, and S. Ström. Transformation properties of plane, spherical and cylindrical scalar and vector wave functions. In V. V. Varadan, A. Lakhtakia, and V. K. Varadan, editors, *Field Representations and Introduction to Scattering*, Acoustic, Electromagnetic and Elastic Wave Scattering, chapter 4, pages 165–210. Elsevier Science Publishers, Amsterdam, 1991.

[3] V. Bringi, T. Seliga, V. Varadan, and V. Varadan. Bulk propagation characteristics of discrete random media. In *Multiple scattering and waves in random media; Proceedings of the Workshop, Blacksburg, VA, March 24-26, 1980.(A82-27720 12-70) Amsterdam, North-Holland Publishing Co.*, volume 1, pages 43–75, 1981.

[4] V. Bringi, V. Varadan, and V. Varadan. Coherent wave attenuation by a random distribution of particles. *Radio Science*, **17**(5), 946–952, 1982.

[5] W. C. Chew, J. A. Friedrich, and R. Geiger. A multiple scattering solution for the effective permittivity of a sphere mixture. *Geoscience and Remote Sensing, IEEE Transactions on*, **28**(2), 207–214, 1990.

[6] K. Ding and L. Tsang. Effective propagation constants in media with densely distributed dielectric particles of multiple sizes and permittivities. *Progress in Electromagnetics Research*, **1**(3), 241–295, 1989.

[7] J. Fikioris and P. Waterman. Multiple scattering of waves. III. The electromagnetic case. *J. Quant. Spectrosc. Radiat. Transfer*, **123**, 8–16, 2013.

[8] A. Ishimaru and Y. Kuga. Attenuation constant of a coherent field in a dense distribution of particles. *JOSA*, **72**(10), 1317–1320, 1982.

[9] G. Kristensson. Coherent scattering by a collection of randomly located obstacles — An alternative integral equation formulation. *J. Quant. Spectrosc. Radiat. Transfer*, **164**(0), 97–108, 2015.

[10] P. Lloyd and M. Berry. Wave propagation through an assembly of spheres: IV. Relations between different multiple scattering theories. *Proceedings of the Physical Society*, **91**(3), 678, 1967.

[11] C. Mandt, Y. Kuga, L. Tsang, and A. Ishimaru. Microwave propagation and scattering in a dense distribution of non-tenuous spheres: experiment and theory. *Waves in Random Media*, **2**(3), 225–234, 1992.

[12] P. A. Martin. *Multiple Scattering: Interaction of Time-Harmonic Waves with N Obstacles*, volume 107 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, U.K., 2006.

[13] M. I. Mishchenko, L. Liu, D. W. Mackowski, B. Cairns, and G. Videen. Multiple scattering by random particulate media: exact 3D results. *Opt. Express*, **15**(6), 2822–2836, 2007.

[14] M. I. Mishchenko, L. D. Travis, and A. A. Lacis. *Multiple scattering of light by particles: radiative transfer and coherent backscattering*. Cambridge University Press, Cambridge, U.K., 2006.

[15] F. W. J. Olver, D. W. Lozier, R. F. Boisvert, and C. W. Clark. *NIST Handbook of mathematical functions*. Cambridge University Press, New York, 2010.

[16] B. Peterson and S. Ström. T-matrix for electromagnetic scattering from an arbitrary number of scatterers and representations of E(3). *Phys. Rev. D*, **8**, 3661–3678, 1973.

[17] W. Ren. Multiple-scattering formalism for general discrete random composites. *Phys. Rev. E*, **49**, 2362–2367, Mar 1994.

[18] V. Tishkovets, E. Petrova, and M. Mishchenko. Scattering of electromagnetic waves by ensembles of particles and discrete random media. *Journal of Quantitative Spectroscopy and Radiative Transfer*, **112**, 2095–2127, 2011.

[19] L. Tsang and J. Kong. Effective propagation constants for coherent electromagnetic wave propagation in media embedded with dielectric scatters. *Journal of Applied Physics*, **53**(11), 7162–7173, 1982.

[20] L. Tsang and J. A. Kong. *Scattering of Electromagnetic Waves: Advanced Topics*. John Wiley & Sons, New York, 2001.

[21] V. Varadan, V. Bringi, and V. Varadan. Frequency dependent dielectric constants of discrete random media. In R. Burridge, S. Childress, and G. Papanicolaou, editors, *Macroscopic Properties of Disordered Media*, volume 154 of *Lecture Notes in Physics*, pages 272–284. Springer Berlin / Heidelberg, 1982.

[22] V. K. Varadan, V. N. Bringi, and V. V. Varadan. Coherent electromagnetic wave propagation through randomly distributed dielectric scatterers. *Phys. Rev. D*, **19**(8), 2480–2489, April 1979.

[23] V. Varadan, V. Bringi, V. Varadan, and A. Ishimaru. Multiple scattering theory for waves in discrete random media and comparison with experiments. *Radio science*, **18**(3), 321–327, 1983.

[24] V. Varadan, Y. Ma, and V. Varadan. Propagator model including multipole fields for discrete random media. *JOSA A*, **2**(12), 2195–2201, 1985.

[25] R. West, D. Gibbs, L. Tsang, and A. Fung. Comparison of optical scattering experiments and the quasi-crystalline approximation for dense media. *JOSA A*, **11**(6), 1854–1858, 1994.