



LUND UNIVERSITY

Project SSM2019-5225: "Marine 14C levels around the Swedish coast" – Additional gamma spectrometric measurements and ICP-MS analysis of brown algae (*Fucus* spp.)

Eriksson Stenström, Kristina; Mattsson, Sören

2021

Document Version:

Publisher's PDF, also known as Version of record

[Link to publication](#)

Citation for published version (APA):

Eriksson Stenström, K., & Mattsson, S. (2021). *Project SSM2019-5225: "Marine 14C levels around the Swedish coast" – Additional gamma spectrometric measurements and ICP-MS analysis of brown algae (Fucus spp.)*. (Lund University; No. Report BAR-2021/02), (Lund University; No. Report MA RADFYS 2021:02). Lund University.

Total number of authors:

2

Creative Commons License:

CC BY

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00



Faculty of Science

Department of Physics
Division of Nuclear Physics
*Biospheric and Anthropogenic
Radioactivity (BAR) Group*

Faculty of Medicine

Department of Translational Medicine
Medical Radiation Physics
*Environmental Radiology and Emergency
Preparedness Group*

Project SSM2019-5225:

“Marine ^{14}C levels around the Swedish coast” –
Additional gamma spectrometric measurements
and ICP-MS analysis of brown algae (*Fucus* spp.)

List of authors

| | |
|------------------------------------------|----------------------------------|
| Kristina Eriksson Stenström ¹ | kristina.stenstrom@nuclear.lu.se |
| Sören Mattsson ² | soren.mattsson@med.lu.se |

| | |
|------------------------------------|--------------------|
| ¹ Department of Physics | Report BAR-2021/02 |
| Division of Nuclear Physics | Lund 2021 |
| Professorsgatan 1 | |
| SE-223 63 Lund | |

| | |
|------------------------------------------------------|--------------------------|
| ² Department of Translational Medicine | Report MA RADFYS 2021:02 |
| Medical Radiation Physics | Malmö 2021 |
| Carl-Bertil Laurells gata 9 | |
| SE-205 02 Malmö | |

Summary

The results of the project SSM2019-5225, “Marine ^{14}C levels around the Swedish coast”, financed by the Swedish Radiation Safety Authority (SSM), are presented in the paper “Spatial and temporal variations of ^{14}C in *Fucus* spp. in Swedish coastal waters” published in the Journal of Environmental Radioactivity. The project SSM2019-5225 focussed on spatial variations in ^{14}C in *Fucus* spp. samples collected in 2020 along the entire Swedish west coast, in coastal waters in the very south of Sweden, and for the east coast up into the Gulf of Bothnia. In this report we present results from additional measurements of the *Fucus* spp. samples, using gamma-ray spectrometry (of ^7Be , ^{40}K , ^{60}Co , ^{137}Cs and ^{131}I) and inductively coupled plasma mass spectrometry (ICP-MS) (of Co, Ni, Cd, Gd, Hg and Pb). The gamma spectrometry showed in addition to the expected content of natural ^{40}K also ^{137}Cs and ^7Be , and at some places also ^{60}Co and ^{131}I . The ICP-MS analysis revealed a correlation between F^{14}C and Gd, and the highest values of F^{14}C and Gd were found close to Ringhals nuclear power plant. This strengthens the hypothesis that that a significant part of the observed peak in F^{14}C on the west coast originates from Ringhals nuclear power plant.

Abbreviations and notations

| | |
|-----------|----------------------------------------------|
| AMS | Accelerator Mass Spectrometry |
| d.w. | Dry weight |
| ICP-MS | Inductively Coupled Plasma Mass Spectrometry |
| NPP | Nuclear Power Plant |
| SD | Standard Deviation |
| SSM | Swedish Radiation Safety Authority |
| SUM | Standard Uncertainty of the Mean |
| $T_{1/2}$ | Physical half-life |
| w.w. | Wet weight |
| %RSD | Percent Relative Standard Deviation |

Contents

| | |
|-----------------------------------------|-----------|
| 1. Introduction | 5 |
| 2. Materials and Methods | 6 |
| Carbon-14 analysis | 6 |
| Gamma-ray spectrometry | 6 |
| ICP-MS | 6 |
| 3. Results and discussion..... | 8 |
| Carbon-14..... | 8 |
| Gamma spectrometry | 11 |
| ICP-MS | 17 |
| 4. Summary and conclusions | 21 |
| 5. Outlook | 21 |
| 6. References | 22 |
| Appendix.... | 25 |

1. Introduction

The results of the project SSM2019-5225, “Marine ^{14}C levels around the Swedish coast”, financed by the Swedish Radiation Safety Authority (SSM), are presented in the paper “Spatial and temporal variations of ^{14}C in *Fucus* spp. in Swedish coastal waters”, published in the Journal of Environmental Radioactivity [1]. That paper describes all objectives of the project, which were to answer the questions:

- Are there differences in ^{14}C levels between Swedish coastal waters?
- Are the ^{14}C levels in Swedish coastal waters influenced by discharged or dumped radioactive wastes? Can releases of ^{14}C from Sellafield (UK) or La Hague (France) be detected on the Swedish west coast?
- What are representative marine background levels of ^{14}C to be used by the different Swedish nuclear power plants? There are three nuclear power plants in the country, with a total of six reactors in operation. These NPPs are the Forsmark (3 reactors), Oskarshamn (1) and Ringhals (2) plants. The two reactors at the Barsebäck NPP were shut down in 1999 and 2005, respectively. These reactors are now undergoing decommissioning, together with Oskarshamn’s units 1 and 2. Apart from the reactors in operation or undergoing decommissioning, there are a number of other nuclear installations in Sweden, used to manufacture nuclear fuel and store spent nuclear fuel and radioactive waste.
- Do different parts of a specific brown algae show varying ^{14}C activity concentration (Bq/kg d.w.) and specific activity (Bq/kg C), e.g. depending on the age of the specific part of the algae?

The project SSM2019-5225 focussed on spatial variations in ^{14}C in *Fucus* spp. samples collected in 2020 along the entire Swedish west coast, in coastal waters in the very south of Sweden, and for the east coast up into Gulf of Bothnia. The paper [1] also includes ^{14}C data from another SSM project, SSM2020-797 (P.I. Sören Mattsson), providing temporal variations in ^{14}C at the sampling station Särö (56.76N, 12.63E) on the Swedish west coast during the period 1967-2020 (27 *Fucus* spp. samples from various years have been used in this report).

In this report we present additional measurements of the *Fucus* spp. samples collected in 2020, using gamma-ray spectrometry (of ^7Be , ^{40}K , ^{60}Co , ^{137}Cs , and ^{131}I) and inductively coupled plasma mass spectrometry (ICP-MS) (of Co, Ni, Cd, Gd, Hg, and Pb).

2. Materials and Methods

Fucus spp. samples were collected at 45 sampling sites (Table 1) between 3 March and 13 May 2020. Details about the sampling sites (including maps), sampling strategy and specific sampling dates can be found in [1].

Carbon-14 analysis

Methods or sample pre-treatment, sample preparation, measurement and analysis for ^{14}C can be found in [1].

Gamma-ray spectrometry

Prior to gamma analysis, the *Fucus* spp. samples (typically 1-2 kg w.w.) were dried in a heating cabinet at 70 °C, then grounded and put in either a 250 ml or a 60 ml sample beaker. The gamma spectrometric measurements were performed at Medical Radiation Physics in Malmö (Lund University) using different HPGe-gamma spectrometers in lead shielded cavities [2]. Measurement time was typically 24-48 hours. Pulse height distributions obtained for each sample were evaluated using an in-house evaluation sheet, which was cross-checked with the detector software evaluation program, Gamma Vision 7.1™. For further information, see Ref [2].

ICP-MS

0.5 g of each dried sample (consisting of leftovers from the samples prepared for ^{14}C analysis, fraction A: vesicles and above, i.e. the new vegetative fronds, mainly representing growth since last spring) was submitted to the Department of Biology (Instrumental Chemistry), Lund University, for sample digestion and to the Department of Geology, Lund University, for Inductively Coupled Mass Spectrometry (ICP-MS) analysis using a Bruker Aurora Elite ICP-MS instrument (see [3]). The stable isotopes analysed were ^{59}Co , ^{60}Ni , ^{114}Cd , ^{111}Cd , ^{157}Gd , ^{202}Hg and ^{208}Pb . The results reported are elemental concentrations, assuming that the isotopes have their natural abundance in the seaweed samples.

Analysis of a certified reference material (ERM-CD200 Bladder wrack) with specified element concentration of Pb, Cd, Hg, Se, As, Cu and Zn (from Unit for Reference materials, EC-Joint Research Center, Institute for Reference Materials and Methods, Geel, Belgium) showed agreement within estimated uncertainty intervals for Pb, Cd, Hg, Se and As (Cu and Zn were not analysed). For further details on quality assurance in the ICP-MS measurements, see [3].

Table 1 Sampling sites for the collection of *Fucus* spp. samples.

| Sea | Site number | Site name | Latitude | Longitude |
|-----------------|--------------------------|-----------------------------------|----------|-----------|
| Skagerrak | 1 | Strömstad, Båteviken | 58.95N | 11.13E |
| | 2 | Fjällbacka, Sälvik | 58.59N | 11.28E |
| | 3 | Smögen, Kleven | 58.35N | 11.22E |
| | 4 | Tjörn, Stockevik | 57.96N | 11.55E |
| Kattegat | 5 | Hällsvik | 57.70N | 11.74E |
| | 6 | Smarholmens badplats | 57.43N | 11.92E |
| | 7 | Frillesås | 57.31N | 12.15E |
| | 8 | Sallebacka | 57.28N | 12.14E |
| | 9 | Ringhals norr Gloppen | 57.27N | 12.11E |
| | 10 | Videbergshamn (Ringhals) | 57.25N | 12.11E |
| | 11 | Bua (Ringhals) | 57.24N | 12.10E |
| | 12 | Skeatången, Årnäshalvön | 57.20N | 12.17E |
| | 13 | Getterön | 57.12N | 12.21E |
| | 14 | Glommen | 56.94N | 12.36E |
| | 15 | Särdal | 56.76N | 12.63E |
| | 16 | Kattvik | 56.46N | 12.75E |
| | 17 | Mölle | 56.28N | 12.50E |
| Öresund Strait | 18 | Ålabodarna/Örenäs | 55.94N | 12.78E |
| | 19 | Barsebäcks camping | 55.77N | 12.92E |
| | 20 | Barsebäcks hamn Skansen | 55.75N | 12.90E |
| | 21 | Barsebäck between NPP and Skansen | 55.75N | 12.91E |
| | 22 | Barsebäck north of NPP | 55.75N | 12.92E |
| | 23 | Vikhög north fort | 55.73N | 12.95E |
| | 24 | Vikhög harbour | 55.73N | 12.96E |
| | 25 | Gamla Bjärred, Pilevägen | 55.71N | 13.03E |
| | 26 | Lomma | 55.68N | 13.06E |
| | 27 | Limhamn | 55.58N | 12.91E |
| 28 | Lilla Hammar (Höllviken) | 55.45N | 12.95E | |
| Baltic Sea | 29 | Smygehuks fyr | 55.34N | 13.35E |
| | 30 | Skillinge | 55.47N | 14.28E |
| | 31 | Hasslö | 56.09N | 15.46E |
| | 32 | Sandvik | 56.55N | 16.22E |
| | 33 | Dragskär | 57.31N | 16.57E |
| | 34 | Simpevarp, Åkvik | 57.41N | 16.64E |
| | 35 | Kråkelund | 57.45N | 16.75E |
| | 36 | Händelöp | 57.67N | 16.75E |
| | 37 | Jogersö | 58.66N | 17.05E |
| | 38 | Senneby, Sandviken, Vaddö | 59.98N | 18.88E |
| Gulf of Bothnia | 39 | Öregrund | 60.34N | 18.45E |
| | 40 | Valudden | 60.35N | 18.37E |
| | 41 | Stenskär | 60.36N | 18.33E |
| | 42 | Norrboda, Gräsö | 60.47N | 18.39E |
| | 43 | Bönan | 60.74N | 17.30E |
| | 44 | Hölick | 61.63N | 17.43E |
| | 45 | Svenskärs fiskeläge | 62.51N | 17.88E |

3. Results and discussion

Carbon-14

The results of the ^{14}C measurements are presented and discussed in [1] and briefly summarised here. The results are expressed as Fraction Modern, $F^{14}\text{C}$ (see [4, 5] for definition). The relation between specific activity of carbon, $\frac{A}{m_C}$, at the year of measurement, y , and $F^{14}\text{C}$ is given by [5]:

$$\frac{A}{m_C} = F^{14}\text{C} \cdot \left(\frac{1 + \frac{\delta^{13}\text{C}}{1000}}{0.975} \right)^2 \cdot e^{\frac{(1950-y)}{8267}} \cdot 226 \frac{\text{Bq}}{\text{kg C}}$$

where $\delta^{13}\text{C}$ is the isotope fractionation of the sample (typical values of various environmental sample types are presented in Table B4 in [2]). For the *Fucus* spp. samples in [1], $\delta^{13}\text{C}$ varied between -20.6‰ and -12.9‰ (measured in 13 samples). The advantage of using $F^{14}\text{C}$ instead of specific activity ($\frac{A}{m_C}$) in high precision measurements of ^{14}C is that:

- 1) $F^{14}\text{C}$ is independent of the year of measurement, whereas $\frac{A}{m_C}$ is not;
- 2) $F^{14}\text{C}$ eliminates the effect of isotope fractionation, making various environmental sample types directly comparable.

As an example, for $F^{14}\text{C} = 1$, $y = 2020$ and $\delta^{13}\text{C} = -16\text{‰}$:

$$\frac{A}{m_C} = 228.3 \text{ Bq (kg C)}^{-1}.$$

The main results were:

- 1) The ^{14}C levels in *Fucus* spp. were significantly higher on the west coast than on the east coast (see Figure 1, from [1]). $F^{14}\text{C}$ values in *Fucus* spp. were up to about 1.10 in Skagerrak, and about 1.01, close to atmospheric levels, in the Baltic Sea and the Gulf of Bothnia.
- 2) The highest value of $F^{14}\text{C}$ observed was close to the Ringhals nuclear power plants (NPP) on the west coast, $F^{14}\text{C} \approx 1.3$ (see Figure 1), from [1]). This is higher than expected in the terrestrial environment of Ringhals NPP.
- 3) $F^{14}\text{C}$ showed a strong positive correlation with salinity on the west coast, and both increased towards the north (see Figure 2). This indicates that ^{14}C may be carried from other anthropogenic sources (e.g. from the nuclear fuel reprocessing plants at La Hague and Sellafield).

- 4) The values of $F^{14}C$ in the SärDAL marine samples collected after the 1990s are clearly higher than $F^{14}C$ in clean air CO_2 , indicating contributions of ^{14}C of anthropogenic origin (see Figure 3).




Figure 1 $F^{14}C$ values obtained from samples of *Fucus* spp. and grass (the latter at Skillinge, Site 30) collected along the Swedish coast, starting at the northernmost site on the west coast (Strömstad, Site 1) and continuing to the northernmost site on the east coast (Svenskärs Fiskeläge, Site 45). Only data from samples composed of the top parts of the *Fucus* spp. individuals (corresponding to growth during 2020) are included. Samples collected within 10 km of the Swedish NPPs are highlighted in the grey areas. The dashed line shows the average $F^{14}C$ value of 26 terrestrial samples collected in southern Sweden for 2020, representing the atmospheric CO_2 reference value for 2020 (from Bernhardsson *et al.* [6]). Uncertainties represent 1 SD, obtained from repeated measurements of the same sample. *Picture from [1] with modifications.*




Figure 2 Left: Salinity measured at the time of sampling of *Fucus* spp. at the different sites in 2020. **Right:** Linear correlation between $F^{14}C$ and salinity for samples in the current study, divided into two sets separated by the border between Kattegat and the Öresund Strait. Sites 8-11 near Ringhals NPP have been excluded, as well as sites located close to all other NPPs (Sites 19-25, 34, 35 and 41; grey shading in Figure 1). The salinity at SärDAL (Site 15) was measured using a portable seawater refractometer RHS-10ATC. *Picture from [1] with modifications.*




Figure 3 ^{14}C in *Fucus* spp. from Särðal in the period 1967-2020. Uncertainty represents 1 SD. Values of F^{14}C in atmospheric CO_2 (atmospheric bomb pulse representative of the Northern Hemisphere) and oceans (modelled marine bomb pulse) are shown for comparison. Atmospheric data are Central European clean air CO_2 from Conen *et al.* [7], Hammer *et al.* [8], Levin and Kromer [9], and Levin *et al.* [10]. Modelled marine values up to 1996 are from Reimer *et al.* [11], representing average global oceanic water down to a depth of 75 m. Values from 1996 to 2020 are estimates based on linear regression of the marine data from 1987 to 1996. The periods of operation of the four reactors at Ringhals NPP, located ~70 km NNW of Särðal, are also given. *Picture from [1].*

Gamma spectrometry

Brown algae are known to display seasonal variations in activity concentrations of certain radionuclides, such as ^{137}Cs and ^{99}Tc [12-14]. The sampling in the current study was performed over a period of 2 months and 11 days. In an attempt to account for seasonal differences, sampling started in the south, and samples were collected from the northern sites during the latter part of the sampling period. The site Skillinge (in the very south-east of Sweden) was sampled at the very beginning (11 March 2020) as well as on the final day of sampling (13 May 2020) to assess seasonal variations. At Särö samples were taken 3 March and 2 May.

The results of the gamma spectrometry are presented in Table A1 in Appendix 1 and summarized in Table 2 below. Of particular interest in this project is to link increased levels of ^{14}C found at the west coast with increased levels of other radionuclides, which may contribute to source apportionment of ^{14}C (mainly to releases from Ringhals NPP and to from long-range transport e.g. from Sellafield and LaHague).

Table 2 Summary of the results from the gamma-ray spectrometry. A: Activity concentration. MDA: Minimum detectable activity concentration. N: Number of measurements. SD: standard deviation. SUM: Standard uncertainty of the mean.

| | ^7Be (Bq/kg d.w.) | | ^{40}K (Bq/kg d.w.) | | ^{60}Co (Bq/kg d.w.) | | ^{134}Cs (Bq/kg d.w.) | | ^{137}Cs (Bq/kg d.w.) | | ^{131}I (Bq/kg d.w.) | |
|--------|-------------------------------|-----|---------------------------------|-----|----------------------------------|-----|-----------------------------------|------|-----------------------------------|-----|----------------------------------|-----|
| | A | MDA | A | MDA | A | MDA | A | MDA | A | MDA | A | MDA |
| N | 46 | 46 | 47 | 47 | 3 | 47 | 42 | 46 | 47 | 47 | 22 | 47 |
| Mean | 30.6 | 6.3 | 934 | 20 | 3.1 | 1.8 | 1.5 | 1.0 | 7.9 | 1.4 | 5.7 | 0.8 |
| Median | 29.4 | 5.8 | 947 | 12 | 3.2 | 1.6 | 1.4 | 0.85 | 7.5 | 1.4 | 1.55 | 0.6 |
| SD | 12.3 | 2.5 | 165 | 16 | 0.4 | 1.0 | 0.5 | 0.6 | 5.8 | 0.8 | 11.3 | 0.5 |
| SUM | 1.8 | 0.4 | 24 | 2 | 0.2 | 0.1 | 0.1 | 0.1 | 0.8 | 0.1 | 2.4 | 0.1 |

^7Be

The results of the gamma spectrometry of ^7Be are shown in Figure 4. ^7Be is produced by interaction of cosmic rays with molecules of N, O and C in the atmosphere [15]. Once produced, ^7Be rapidly adsorbs onto aerosol particles in the stratosphere and troposphere and is further transported to the Earth's surface by dry and/or wet fallout. An almost linear relation between precipitation and ^7Be deposition has earlier been demonstrated [16, 17]. The mean yearly deposition in Sweden has been estimated to around 1500 Bq/m².

The difference between *Fucus serratus* and *F. vesiculosus* and between Särö, Skillinge and Hasselö (and there also for seaweed with and without ongrowth) is within the range of the variation. Long-term measurements in Särö show typical activity concentrations of 20-40 Bq/kg d.w. in *F. serratus* and 10-40 Bq/kg d.w. in *F. vesiculosus* [18]. Higher values, up to 100 Bq/kg were reached in 2018 and 2019. Still higher values were found in the environmental control programs for Ringhals with average values between 150 and 300 Bq/kg d.w. in 1995-2019 [19]. At Barsebäck the levels have varied between 19 and 150 Bq/kg in *F. vesiculosus* with an average of about 60 Bq/kg in the period 2011 to

2019 [19]. It would be interesting to further investigate the fraction of possible Ringhals produced ^7Be in *Fucus* taken in the immediate vicinity of the plant.




Figure 4 ^7Be in *Fucus* spp. collected in 2020. Samples collected within 10 km of the Swedish NPPs are highlighted in the grey areas (see Figure 1). Open circles: MDA.

^{40}K

Obtained activity concentrations for ^{40}K are shown in Figure 5. The mean value for the ^{40}K activity concentration is similar to that for the long-term measurements at Särdal. The mean value for *Fucus serratus* from Särdal during the period 1967-2020 was 885 ± 150 Bq/kg d.w. which corresponds to 29 g potassium per kg of dry seaweed [20]. There is a tendency of decreasing concentrations towards the Baltic Sea. *Fucus* spp. from the Baltic Sea showed a mean ^{40}K activity concentration of 817 ± 33 Bq/kg d.w. in the period 1978-1983 [21].




Figure 5 ^{40}K ($T_{1/2} = 1.28 \cdot 10^9$ years) in *Fucus* spp. at the various sites. Samples collected within 10 km of the Swedish NPPs are highlighted in the grey areas (see Figure 1). Open circles: MDA.

^{60}Co

Fucus vesiculosus and *F. serratus* have since long been used to map the spatial and temporal distribution of the activation products ^{60}Co , ^{58}Co , ^{54}Mn and ^{65}Zn around nuclear power plants [22, 23]. In the current measurement series, only ^{60}Co was detected and when detected, only in a few samples close to Barsebäck NPP (Figure 6), now under decommissioning. The levels were low and are probably explained by remobilization of ^{60}Co from sediments or fresh releases in connection with the ongoing decommissioning. The measuring points around the other nuclear power plants have been too far from the reactors to give detectable results.




Figure 6 ^{60}Co ($T_{1/2} = 5.3$ y) in *Fucus* spp. at the various sites. Samples collected within 10 km of the Swedish NPPs are highlighted in the grey areas (see Figure 1). Open circles: MDA.

^{137}Cs

^{137}Cs was higher on the east coast than on the west coast (Figure 7). Lower salinity is known to increase the uptake of ^{137}Cs in *Fucus* spp. [24, 25].




Figure 7 ^{137}Cs ($T_{1/2} = 30.2$ y) in *Fucus* spp. at the various sites. Samples collected within 10 km of the Swedish NPPs are highlighted in the grey areas (see Figure 1). Open circles: MDA.

Figure 8 shows the ^{137}Cs activity concentrations in *Fucus* spp. as a function of collection date for all sites. In general, the ^{137}Cs activity concentrations are higher towards the end of the sampling period (May) compared to the beginning of the sampling period (March). The site Skillinge (sampled at the very beginning as well as on the final day of sampling) shows almost a doubling of the ^{137}Cs activity concentration from March to May. At Särðal, the activity concentration (mean for *F. serratus* and *F. vesiculosus*) also almost doubled from 2.1 to 4.0 Bq/kg d.w. from March 3 to May 2, 2020. This is in accordance with the seasonal variations seen earlier [18].




Figure 8 ^{137}Cs ($T_{1/2} = 30.2$ y) in *Fucus* spp. as a function of collection date. Note the logarithmic scale.

¹³¹I

The activity concentrations of ¹³¹I in *Fucus vesiculosus* are increased around Barsebäck NPP compared to the other NPPs (see Figure 9). This is most likely a result of releases from the sewage treatment plants at Sjölanda (Malmö) and Källby (Lund) due to the presence of ¹³¹I in excreta from patients treated with ¹³¹I mainly for thyrotoxicosis and thyroid cancer). In the same way, the still higher level at station 5 (Hällsvik) is likely to be due to releases from patients at hospitals and patients who have returned home from the hospitals in Gothenburg and along the Göta älv. The sewage treatment plant Ryaverken serving Gothenburg and its surrounding municipalities was at least earlier known to have a very short delay time for the waste water [26].




Figure 9 ¹³¹I ($T_{1/2} = 8.0$ d) in *Fucus* spp. at the various sites. Samples collected within 10 km of the Swedish NPPs are highlighted in the grey areas (see Figure 1). Open circles: MDA.

Correlation between ¹⁴C and gamma emitters

A correlation matrix for F¹⁴C and the measured gamma emitters (except ⁶⁰Co due to the low number of data above the MDA) for all samples is presented in Table 3. A negative correlation is found between F¹⁴C and ¹³⁷Cs, which most likely is due to completely different sources resulting in higher F¹⁴C on the west coast and higher ¹³⁷Cs on the east coast. The high F¹⁴C on the west coast most likely originates from Ringhals NPP and Sellafield/La Hague, while the high ¹³⁷Cs on the east coast is Chernobyl-derived. In Table 4, only samples collected in Skagerrak and Kattegat are included in the correlation analysis: F¹⁴C shows no correlation to any of the gamma emitters. The same is valid when separately investigating the Öresund Strait, the Baltic Sea and the Gulf of Bothnia (Table 5).

Table 3 Correlation matrix for $F^{14}C$ and 7Be , ${}^{40}K$, ${}^{134}Cs$, ${}^{137}Cs$ and ${}^{131}I$ for *Fucus* spp. samples for all sites. p-values <0.05 are shaded in grey.

| | | 7Be | ${}^{40}K$ | ${}^{137}Cs$ | ${}^{131}I$ |
|--------------|---------------|----------|------------|--------------|-------------|
| $F^{14}C$ | Pearson Corr. | 0.21 | 0.04 | -0.54 | 2.0E-4 |
| | p-value | 0.16 | 0.809 | 1.5E-4 | 0.99 |
| 7Be | Pearson Corr. | | -0.119 | -0.15 | 0.55 |
| | p-value | | 0.469 | 0.31 | 0.01 |
| ${}^{40}K$ | Pearson Corr. | | | -0.10 | -0.03 |
| | p-value | | | 0.52 | 0.91 |
| ${}^{137}Cs$ | Pearson Corr. | | | | -0.02 |
| | p-value | | | | 0.91 |

Table 4 Correlation matrix for $F^{14}C$ and 7Be , ${}^{40}K$, ${}^{134}Cs$, ${}^{137}Cs$ and ${}^{131}I$ for *Fucus* spp. samples collected in Skagerrak and Kattegat. p-values <0.05 are shaded in grey.

| | | 7Be | ${}^{40}K$ | ${}^{137}Cs$ | ${}^{131}I$ |
|--------------|---------------|----------|------------|--------------|-------------|
| $F^{14}C$ | Pearson Corr. | 0.31 | 0.15 | 0.16 | -0.06 |
| | p-value | 0.20 | 0.56 | 0.53 | 0.87 |
| 7Be | Pearson Corr. | | -0.42 | -0.02 | 0.74 |
| | p-value | | 0.08 | 0.93 | 0.02 |
| ${}^{40}K$ | Pearson Corr. | | | 0.29 | -0.16 |
| | p-value | | | 0.25 | 0.67 |
| ${}^{137}Cs$ | Pearson Corr. | | | | 0.19 |
| | p-value | | | | 0.62 |

Table 5 Correlation matrix for $F^{14}C$ and 7Be , ${}^{40}K$, ${}^{134}Cs$, ${}^{137}Cs$ and ${}^{131}I$ for *Fucus* spp. samples collected in the Öresund Strait, the Baltic Sea and the Gulf of Bothnia. p-values <0.05 are shaded in grey (none).

| | | 7Be | ${}^{40}K$ | ${}^{137}Cs$ | ${}^{131}I$ |
|--------------|---------------|----------|------------|--------------|-------------|
| $F^{14}C$ | Pearson Corr. | 0.01 | -0.19 | 0.25 | -0.45 |
| | p-value | 0.97 | 0.34 | 0.21 | 0.14 |
| 7Be | Pearson Corr. | | 0.11 | -0.14 | 0.04 |
| | p-value | -- | 0.57 | 0.48 | 0.91 |
| ${}^{40}K$ | Pearson Corr. | | | -0.23 | 0.46 |
| | p-value | | -- | 0.23 | 0.12 |
| ${}^{137}Cs$ | Pearson Corr. | | | | 0.14 |
| | p-value | | | | 0.65 |

ICP-MS

The results of the ICP-MS analysis are presented in Table A2 in Appendix 1. The results are summarized in Table 6 and in Figure 10. Typical uncertainties in terms of relative standard deviation in percent (%RSD) were on average 5%-8% for most samples and isotopes analysed (⁵⁹Co, ⁶⁰Ni, ¹¹⁴Cd, ¹¹¹Cd, ¹⁵⁵Gd, ¹⁵⁷Gd, ²⁰⁸Pb). For ²⁰²Hg, the %RSD was generally higher than for the other metals (average %RSD: 59% with a standard deviation 40%). Forsberg et al. [27] have reported metal concentrations in *Fucus vesiculosus* from the archipelago of Stockholm. The ranges of concentrations for Ni, Cd, Pb and Co are of the same order of magnitude as results in Table 6. This is also the situation for the same metals in *Fucus vesiculosus* and *F. serratus* collected at Tjärnö (between Båteviken, Strömstad och Sälvik, Fjällbacka) in 2014 [28].

Table 6 Summary of the results from the ICP-MS measurements (elemental concentrations). %RSD: relative standard deviation. SD: standard deviation. SUM: Standard uncertainty of the mean. * From ¹¹⁴Cd. ** From ¹¹¹Cd.

| | Co (ppb) | % RSD | Ni (ppb) | % RSD | Cd* (ppb) | % RSD | Cd** (ppb) | % RSD | Gd (ppb) | % RSD | Hg (ppb) | % RSD | Pb (ppb) | % RSD |
|---------------|-------------|----------|-------------|----------|--------------|----------|---------------|----------|-------------|----------|-------------|----------|-------------|----------|
| Mean | 1725 | 5 | 7709 | 6 | 1271 | 7 | 1289 | 7 | 246 | 8 | 13 | 59 | 746 | 7 |
| Median | 1588 | | 6790 | | 1075 | | 1086 | | 162 | | 12 | | 444 | |
| SD | 893 | 1 | 4347 | 2 | 581 | 2 | 594 | 2 | 227 | 3 | 5 | 40 | 833 | 3 |
| SUM | 126 | | 615 | | 82 | | 84 | | 32 | | 1 | | 118 | |




Figure 10 Concentrations of various elements in *Fucus* spp. obtained with ICP-MS analysis. Relative standard deviations are typically 5-8 % for all elements except Hg (average %RSD 59%). Upper graph: all results. Lower graph: all results except Ni.

The average concentration was highest for Ni, followed by Co, Cd, Pb, Gd and Hg. The concentrations of Ni and Cd were highest in the northern Baltic Sea and in the Gulf of Bothnia. In the current report we focus on possible correlations between these elements and $F^{14}C$. The correlation between $F^{14}C$ and Co, Ni, Cd, Gd and Pb is presented in Table 7 (Hg has been excluded due to its high uncertainties). The only significant correlation (p -value < 0.05) for $F^{14}C$ can be seen for Gd. Other significant correlations can be seen between Co and Gd, Ni and Cd, and Gd and Pb.

Table 7 Correlation matrix for F¹⁴C and Co, Ni, Cd, Gd and Pb. p-values <0.05 are shaded in grey. *From ¹¹⁴Cd. **From ¹¹¹Cd.

| | | Co | Ni | Cd* | Cd** | Gd | Pb |
|------------------------|---------------|------|-------|----------|----------|----------|----------|
| F¹⁴C | Pearson Corr. | 0.25 | -0.15 | -0.14 | -0.15 | 0.29 | 0.08 |
| | p-value | 0.08 | 0.30 | 0.34 | 0.30 | 0.04 | 0.59 |
| Co | Pearson Corr. | | -0.15 | -0.13 | -0.13 | 0.58 | 0.19 |
| | p-value | | 0.28 | 0.38 | 0.37 | 8.63E-06 | 0.20 |
| Ni | Pearson Corr. | | | 0.71 | 0.71 | -0.21 | -0.22 |
| | p-value | | | 7.04E-09 | 7.04E-09 | 0.15 | 0.12 |
| Cd* | Pearson Corr. | | | | 1.00 | 0.04 | 0.09 |
| | p-value | | | | 7.04E-09 | 0.76 | 0.54 |
| Cd** | Pearson Corr. | | | | | 0.07 | 0.10 |
| | p-value | | | | | 0.62 | 0.50 |
| Gd | Pearson Corr. | | | | | | 0.45 |
| | p-value | | | | | | 7.04E-09 |

The correlation between F¹⁴C and Gd is further visualized in Figure 11. The highest Gd concentrations and F¹⁴C values are both found in Kattegat, close to Ringhals NPP, hence the correlation between the two. Gd is present in Ringhals NPP in fuel rods (as Gd₂O₃) [29]. The Gd concentration varies significantly at nearby sites in Skagerrak, Kattegat, the Baltic Sea and the Gulf of Bothnia. The variation in Gd concentration between nearby sites is less pronounced in southern Kattegat and in the Öresund Strait. Site 5 is located at the outflow of Göta älv and 24 is located north of the outflow from Sjölanda wastewater treatment plant and Höje å carrying the releases from the Källby wastewater treatment plant. Both these sites are characterized by lower salinity than the nearby sites (see Figure 2) and with higher Gd concentrations than in nearby sites. These increases in Gd concentration thus most likely stem from river runoff of Gd originating from the use of Gd-containing contrast agents used for magnetic resonance imaging in hospitals.




Figure 11 F¹⁴C and Gd concentration in *Fucus* spp. collected along the Swedish coast in 2020. S: Skagerrak.

4. Summary and conclusions

The gamma spectrometric measurements of *Fucus* spp. showed the well-known presence of naturally occurring ^{40}K (algae contain high concentrations of potassium), likewise naturally produced ^7Be among which there may also be a component caused by releases from the local nuclear power plants. The activation product ^{60}Co has been traced in the vicinity of the nuclear power plants in operation and in the vicinity of the decommissioned nuclear power plant in Barsebäck. The activity concentration of ^{137}Cs in *Fucus* spp. is affected by local emissions and above all by inflows from the North Sea and transport of Chernobyl ^{137}Cs from the Baltic Sea. The uptake of ^{137}Cs in *Fucus* spp. is strongly dependent on salinity and also on the season.

The correlation between F^{14}C and Gd, and in particular the finding that the highest values of F^{14}C and Gd are found close to Ringhals NPP, strengthens the hypothesis that that a significant fraction of the peak in F^{14}C on the west coast originates from Ringhals NPP.

5. Outlook

More information can most likely be extracted from the gamma and ICP-MS analyses. It would be interesting to do this in the context of a new project.

Acknowledgements and funding

We gratefully acknowledge the assistance of Mattias Jönsson, Guillaume Pédehontaa-Hiaa and Christian Bernhardsson, Medical Radiation Physics, Malmö, Lund University, in the sample preparation and measurements using gamma spectrometry. The project was funded by the Swedish Radiation Safety Authority (SSM), project SSM2019-5225.

6. References

1. Eriksson Stenström, K., Mattsson, S. *Spatial and temporal variations of ^{14}C in *Fucus* spp. in Swedish coastal waters*. Journal of Environmental Radioactivity Volume 242, February 2022, 106794, 2022.
2. Bernhardsson, C., Eriksson Stenström, K., Jönsson, M., Mattsson, S., Pedehontaa-Hiaa, G., Rääf, C., Sundin, K., et al. *Assessment of "Zero Point" radiation around the ESS facility*. Report MA RADFYS 2018:01, Report BAR-2018/04.
[https://portal.research.lu.se/portal/sv/publications/assessment-of-zero-point-radiation-around-the-ess-facility\(2153e07c-b465-4191-abc3-dbfcaf28b85b\).html](https://portal.research.lu.se/portal/sv/publications/assessment-of-zero-point-radiation-around-the-ess-facility(2153e07c-b465-4191-abc3-dbfcaf28b85b).html). 2018.
3. Pedehontaa-Hiaa, G., Bernhardsson, C., Barkauskas, V., Eriksson Stenström, K., Rääf, C., Mattsson, S. *Region-specific radioecological evaluation of releases of radionuclides from ESS - during operation and accidents*. SSM report of project SSM 2019-1010.
<https://www.stralsakerhetsmyndigheten.se/publikationer/rapporter/stralskydd/2021/202121/>. 2021.
4. Reimer, P.J., Brown, T.A., Reimer, R.W. *Discussion: Reporting and calibration of post-bomb ^{14}C data*. Radiocarbon, 46(3): 1299-1304, 2004.
5. Eriksson Stenström, K., Skog, G., Georgiadou, E., Genberg, J., Johansson, A. *A guide to radiocarbon units and calculations*. LUNFD6(NFFR-3111)/1-17/(2011). Lund University. Dep of Physics. Div of Nuclear Physics.
<http://lup.lub.lu.se/search/ws/files/5555659/2173661.pdf>. 2011.
6. Bernhardsson, C., Eriksson Stenström, K., Jönsson, M., Pedehontaa-Hiaa, G., Mattsson, S. *Radiological environmental monitoring at the ESS facility – Annual report 2020*. Report MA RADFYS 2021:01, Report BAR 2021/01. 2021.
7. Conen, F., Emmenegger, L., Leuenberger, M., Steger, D., Steinbacher, M. *"ICOS RI, 2020. ICOS ATC 14C Release, Jungfraujoch (10.0 m), 2016-01-04 2019-08-12"*. <https://hdl.handle.net/11676/X-IXPKZIO4DWX7wnesLQ7akY>; <http://calib.org/CALIBomb/>. 2019.
8. Hammer, S., Levin, I. *Monthly mean atmospheric D^{14}CO_2 at Jungfraujoch and Schauinsland from 1986 to 2016*. heiDATA: Heidelberg Research Data Repository [Distributor] V2 [Version]. <http://dx.doi.org/10.11588/data/10100>; www.calibomb.org. 2017.
9. Levin, I., Kromer, B. *The tropospheric $^{14}\text{CO}_2$ level in mid latitudes of the Northern Hemisphere (1959-2003)*. Radiocarbon, 46: 1261-1271, 2004.
10. Levin, I., Kromer, B., Hammer, S. *Atmospheric $\Delta^{14}\text{CO}_2$ trend in Western European background air from 2000 to 2012*. Tellus B: Chemical and Physical Meteorology, 65(1): 20092, 2013.
11. Reimer, P.J., Baillie, M.G., Bard, E., Bayliss, A., Beck, J.W., Blackwell, P.G., Ramsey, C.B., et al. *IntCal09 and Marine09 radiocarbon age calibration curves, 0–50,000 years cal BP*. Radiocarbon, 51(4): 1111-1150, 2009.
12. Carlson, L., Erlandsson, B. *Seasonal variation of radionuclides in *Fucus vesiculosus* L. from the Öresund, Southern Sweden*. Environmental pollution, 73(1): 53-70, 1991.
13. Lindahl, P., Ellmark, C., Gäfvert, T., Mattsson, S., Roos, P., Holm, E., Erlandsson, B. *Long-term study of ^{99}Tc in the marine environment on*

- the Swedish west coast*. Journal of Environmental Radioactivity, 67(2): 145-156, 2003.
14. Mattsson, S. *¹³⁷Cs in algae from the Swedish west coast, 1967-1983*, in *Radiation-risk-protection. Vol. 2. In: Radiation-risk-protection. (Ed. by A Kaul et al.). Fachverband für Strahlenschutz, Jülich, Germany, 1984, pp 901-904*. 1984.
 15. Lal, D., Peters, B. *Cosmic ray produced radioactivity on the Earth*, in *Kosmische Strahlung II/Cosmic Rays II*. 1967, Springer. p. 551-612.
 16. Ishikawa, Y., Murakami, H., Sekine, T., Yoshihara, K. *Precipitation scavenging studies of radionuclides in air using cosmogenic ⁷Be*. Journal of Environmental Radioactivity, 26(1): 19-36, 1995.
 17. Caillet, S., Arpagaus, P., Monna, F., Dominik, J. *Factors controlling ⁷Be and ²¹⁰Pb atmospheric deposition as revealed by sampling individual rain events in the region of Geneva, Switzerland*. Journal of Environmental Radioactivity, 53(2): 241-256, 2001.
 18. Ikäheimonen, T., Outola, I., Vartti, V.-P., Kotilainen, P. *Radioactivity in the Baltic Sea: inventories and temporal trends of ¹³⁷Cs and ⁹⁰Sr in water and sediments*. Journal of Radioanalytical and Nuclear Chemistry, 282(2): 419-425, 2009.
 19. Aquilonius, K. Personal communication. Swedish Radiation Safety Authority (SSM). 2021.
 20. Mattsson, S., Eriksson Stenström, K., Pédehontaa-Hiaa, G. *Long-term variations of radionuclides and metals in the marine environment of the Swedish west-coast studied using brown algae (Fucus serratus and F. vesiculosus)*. Draft report to SSM2020-797 and 2018-905. 2021.
 21. Holm, E., Rioseco, J., Mattsson, S. *Technetium-99 in the Baltic Sea, in Technetium in the Environment*. 1986, Springer. p. 61-68.
 22. Mattsson, S., Finck, R., Nilsson, M. *Distribution of activation products from Barsebäck nuclear power plant (Sweden) in the marine environment. Temporal and spatial variations as established by seaweed*. Environmental Pollution Series B, Chemical and Physical, 1(2): 105-115, 1980.
 23. Nilsson, M., Dahlgaard, H., Edgren, M., Holm, E., Mattsson, S., Notter, M. *Radionuclides in Fucus from inter-Scandinavian waters, in Impacts of radionuclide releases into the marine environment. In: International symposium on the impacts of radionuclide releases into the marine environment. Vienna, Austria, 6 - 10 Oct 1980. IAEA-SM--248/107 p. 501-513, IAEA. Vienna. International Atomic Energy Agency*. 1981.
 24. Aarkrog, A., *Bioindicator Studies in Nordic waters: Summary report of the NKA report REK-5B*. Nordic Council of Ministers. 8755011349, 1985.
 25. Carlson, L., Erlandsson, B. *Effects of salinity on the uptake of radionuclides by Fucus vesiculosus L.* Journal of Environmental Radioactivity, 13(4): 309-322, 1991.
 26. Erlandsson, B., Isaksson, M., Mattsson, S., Hemdal, B. *The urban contamination after the Chernobyl accident studied in sewage water and sludge from Lund and Gothenburg*. In: "On the validity of environmental transfer models", Proceedings BIOMOVs, Swedish Radiation Protection Institute, Stockholm, Sweden, pp 169-179, 1991.
 27. Forsberg, Å., Söderlund, S., Frank, A., Petersson, L., Pedersen, M. *Studies on metal content in the brown seaweed, Fucus vesiculosus, from the Archipelago of Stockholm*. Environmental pollution, 49(4): 245-263, 1988.

28. Olsson, J., Toth, G.B., Albers, E. *Biochemical composition of red, green and brown seaweeds on the Swedish west coast.* Journal of Applied Phycology, 32: 3305-3317, 2020.
29. Insulander Björk, K., Lau, C.W., Nylén, H., Sandberg, U. *Study of thorium-plutonium fuel for possible operating cycle extension in PWRs.* Science and Technology of Nuclear Installations, 2013, 2013.

Appendix

Table A1 Results of gamma spectrometry measurements. FV – *Fucus vesiculosus*. FS – *Fucus serratus*. FR – *Fucus radicans*.

| Site | Site name | Species | Sampling date YYY-MM-DD | Lab code | Detector | Cs-137 (Bq kg ⁻¹ d.w.) | | | Co-60 (Bq kg ⁻¹ d.w.) | | | Be-7 (Bq kg ⁻¹ d.w.) | | | I-131 (Bq kg ⁻¹ d.w.) | | | Ka-40 (Bq kg ⁻¹ d.w.) | | |
|------|---------------------------------------------------------------|---------|----------------------------|----------|----------|--------------------------------------|-----|-----|----------------------------------|----|-----|---------------------------------|------|-----|----------------------------------|-----|-----|----------------------------------|----|-----|
| | | | | | | A | SD | MDA | A | SD | MDA | A | SD | MDA | A | SD | MDA | A | SD | MDA |
| 1 | Båtevikens badplats. Strömstad | FV | 2020-04-08 | M645 | 5 | 1.6 | 0.3 | 1.1 | <MDA | | 1.3 | 30.4 | 1.6 | 3.2 | <MDA | | 0.5 | 1096 | 52 | 9 |
| 2 | Fjällbacka. Sälvik. norr om hamn. med påväxt | FV | 2020-04-08 | M644 | 5 | 1.6 | 0.2 | 0.7 | <MDA | | 0.8 | 31.0 | 1.8 | 4.9 | 0.3 | 0.6 | 0.3 | 1076 | 51 | 6 |
| 2 | Fjällbacka. Sälvik. hamn och söder om hamn. utan påväxt | FV | 2020-04-08 | M641 | 5 | 1.1 | 0.2 | 1.0 | <MDA | | 1.2 | 37.8 | 2.0 | 4.5 | <MDA | | 0.4 | 716 | 39 | 8 |
| 3 | Smögen. Kleven | FV | 2020-04-08 | M642 | 5 | 1.1 | 0.3 | 1.4 | <MDA | | 1.7 | 38.9 | 2.4 | 6.4 | <MDA | | 0.6 | 723 | 41 | 12 |
| 4 | Stockevik. Tjörn | FV | 2020-04-08 | M643 | 5 | 1.2 | 0.4 | 1.6 | <MDA | | 1.9 | 30.8 | 14.7 | 7.3 | <MDA | | 0.7 | 1127 | 54 | 13 |
| 5 | Hällsvik | FV | 2020-04-08 | M647 | 7 | 2.7 | 0.5 | 1.8 | <MDA | | 2.2 | 57.3 | 3.7 | 8.4 | 52.5 | 2.5 | 1.1 | 840 | 89 | 31 |
| 6 | Smarholmens badplats | FV | 2020-03-31 | M639 | 5 | 1.7 | 0.3 | 1.2 | <MDA | | 1.4 | 29.0 | 2.0 | 5.4 | <MDA | | 0.5 | 888 | 45 | 10 |
| 7 | Frillesås | FV | 2020-03-31 | M638 | 7 | 2.4 | 0.4 | 1.5 | <MDA | | 1.8 | 18.8 | 2.5 | 6.9 | 0.9 | 1.0 | 0.9 | 1136 | 88 | 25 |
| 8 | Sallebacka | FV | 2020-03-31 | M634 | 7 | 1.6 | 0.5 | 1.7 | <MDA | | 2.0 | 12.6 | 18.1 | 7.7 | <MDA | | 1.0 | 994 | 86 | 28 |
| 9 | Ringhals norr Gloppen | FV | 2020-03-31 | M635 | 7 | 2.5 | 0.5 | 1.8 | <MDA | | 2.2 | 53.3 | 3.5 | 8.3 | <MDA | | 1.1 | 890 | 78 | 30 |
| 10 | Videbergshamn (Ringhals) | FV | 2020-03-31 | M636 | 5 | 2.6 | 0.3 | 1.3 | <MDA | | 1.5 | 36.3 | 2.3 | 5.8 | <MDA | | 0.6 | 1084 | 53 | 11 |
| 11 | Bua (Ringhals) | FV | 2020-03-31 | M640 | 7 | 2.7 | 0.5 | 1.9 | <MDA | | 2.3 | 44.3 | 3.5 | 8.7 | <MDA | | 1.1 | 963 | 82 | 32 |
| 12 | Skeatången. Ärnäshalvön | FV | 2020-03-31 | M637 | 4 | 1.5 | 0.2 | 0.6 | <MDA | | 0.7 | 48.0 | 2.1 | 3.5 | 0.9 | 0.6 | 0.4 | 512 | 22 | 11 |
| 13 | Getterön | FV | 2020-03-24 | M630 | 4 | 2.0 | 0.2 | 0.6 | <MDA | | 0.8 | 18.3 | 1.6 | 3.8 | 1.0 | 0.9 | 0.5 | 706 | 28 | 12 |
| 14 | Glommen | FV | 2020-03-24 | M629 | 5 | 2.4 | 0.2 | 0.8 | <MDA | | 0.9 | 23.6 | 1.5 | 3.5 | 1.1 | 0.8 | 0.3 | 1020 | 49 | 7 |
| 15 | Särdal V-A | FV | 2020-03-03 | M612 | 5 | 2.4 | 0.2 | 0.8 | <MDA | | 0.9 | 29.8 | 1.6 | 3.5 | 1.7 | 0.4 | 0.3 | 1051 | 50 | 6 |
| 15 | Särdal S | FS | 2020-03-03 | M611 | 4 | 1.8 | 0.2 | 0.2 | <MDA | | 0.7 | 21.3 | 1.3 | 3.0 | 1.4 | 0.5 | 0.4 | 1016 | 37 | 9 |
| 17 | Mölle | FV | 2020-04-20 | M654 | 5 | 4.1 | 0.3 | 1.0 | <MDA | | 1.2 | 19.4 | 1.5 | 4.5 | 1.3 | 0.6 | 0.4 | 1012 | 50 | 8 |
| 18 | Ålabodarna/Örenäs | FV | 2020-03-25 | M626 | 5 | 6.4 | 0.4 | 1.2 | <MDA | | 1.3 | 14.4 | 1.5 | 5.1 | 1.1 | 0.4 | 0.5 | 1026 | 52 | 9 |

| | | | | | | | | | | | | | | | | | | | | |
|----|-----------------------------------|-------|------------|-------------------------|---|------|-----|-----|------|-----|-----|------|------|------|------|-----|-----|------|-----|----|
| 19 | Barsebäcks camping | FV | 2020-03-24 | M625 | 5 | 8.3 | 0.4 | 0.8 | <MDA | | 1.0 | 24.1 | 1.5 | 3.7 | 14.3 | 0.8 | 0.4 | 1093 | 55 | 7 |
| 20 | Barsebäcks hamn Skansen | FV | 2020-03-24 | M621 | 4 | 5.7 | 0.4 | 0.8 | 2.6 | 0.6 | 1.0 | 32.2 | 1.9 | 4.6 | 3.9 | 0.4 | 0.5 | 819 | 32 | 13 |
| 21 | Barsebäck between NPP and Skansen | FV | 2020-03-25 | N624 | 5 | 10.3 | 0.5 | 1.3 | 3.2 | 0.4 | 1.5 | 26.8 | 1.7 | 4.3 | 7.8 | 0.4 | 0.6 | 1121 | 54 | 10 |
| 22 | Barsebäck north of NPP | FV | 2020-03-25 | M627 | 7 | 10.2 | 0.8 | 2.5 | 3.4 | 0.8 | 3.1 | 44.3 | 2.4 | 5.7 | 5.2 | 0.8 | 1.5 | 1027 | 100 | 43 |
| 23 | Vikhög fort | FV | 2020-03-25 | M628 | 7 | 7.2 | 0.5 | 1.4 | <MDA | | 1.7 | 47.1 | 3.1 | 6.6 | 3.0 | 0.9 | 0.9 | 1129 | 103 | 24 |
| 24 | Vikhög hamn | FV | 2020-03-25 | M620 | 5 | 9.8 | 0.5 | 1.0 | <MDA | | 1.2 | 32.2 | 1.9 | 4.6 | 17.1 | 0.8 | 0.5 | 1083 | 53 | 9 |
| 25 | Gamla Bjärred. Pilevägen | FV | 2020-04-07 | M646 | 7 | 6.4 | 1.4 | 3.5 | <MDA | | 4.7 | 17.4 | 12.4 | 16.3 | <MDA | | 2.1 | 874 | 65 | 67 |
| 27 | Limhamn | FV | 2020-03-24 | M622 | 4 | 5.9 | 0.3 | 0.5 | <MDA | | 0.6 | 31.0 | 1.4 | 2.7 | 1.3 | 0.5 | 0.3 | 947 | 36 | 8 |
| 28 | Lilla Hammar (Höllviken) | FV | 2020-03-24 | M623 | 7 | 8.1 | 0.6 | 1.9 | <MDA | | 2.3 | 20.5 | 2.9 | 8.9 | <MDA | | 1.2 | 1207 | 104 | 33 |
| 29 | Smygehukuks fyr | FV | 2020-03-11 | M615 | 5 | 7.5 | 0.4 | 0.7 | <MDA | | 0.9 | 23.5 | 1.6 | 3.8 | 0.8 | 0.6 | 0.5 | 966 | 37 | 12 |
| 30 | Skillinge. 1A | FV | 2020-03-11 | M614 | 7 | 10.4 | 0.6 | 1.5 | <MDA | | 1.8 | 55.9 | 3.2 | 7.0 | <MDA | | 0.9 | 1010 | 89 | 25 |
| 30 | Skillinge | FV | 2020-05-13 | M674 | 7 | 19.9 | 0.9 | 1.4 | <MDA | | 1.7 | 42.5 | 3.1 | 6.6 | <MDA | | 0.9 | 1300 | 97 | 24 |
| 31 | Hasslö | FS | 2020-03-19 | M617 (FS) | 5 | 8.3 | 0.6 | 1.8 | <MDA | | 0.9 | 47.8 | 2.9 | 8.3 | <MDA | | 0.8 | 732 | 46 | 15 |
| 31 | Hasslö | FV | 2020-03-19 | M618 (FV) | 7 | 10.4 | 0.6 | 1.5 | <MDA | | 1.8 | 55.9 | 3.2 | 7.0 | <MDA | | 0.9 | 1010 | 89 | 25 |
| 31 | Hasslö | FV+FS | 2020-03-19 | M619 (FV with ongrowth) | 4 | 7.9 | 0.4 | 0.7 | <MDA | | 2.1 | 35.0 | 1.9 | 3.9 | 1.1 | 0.6 | 0.5 | 763 | 30 | 12 |
| 32 | Sandvik | FV | 2020-04-22 | M658 | 7 | 19.2 | 0.9 | 1.7 | <MDA | | 2.1 | 17.5 | 2.8 | 8.0 | | | 1.0 | 899 | 85 | 29 |
| 33 | Dragskär | FV | 2020-04-22 | M655 | 7 | 12.2 | 0.8 | 2.1 | <MDA | | 2.5 | 26.0 | 3.4 | 9.6 | 2.0 | 1.9 | 1.3 | 942 | 88 | 35 |
| 34 | Simpevarp Åkvik | FV | 2020-04-22 | M657 | 5 | 16.6 | 0.8 | 1.9 | <MDA | | 2.2 | 19.2 | 12.9 | 8.5 | <MDA | | 0.8 | 928 | 48 | 16 |
| 35 | Kräkelund | FV | 2020-04-22 | M656 | 7 | 12.2 | 0.8 | 2.1 | <MDA | | 2.5 | 26.7 | 3.5 | 9.6 | 2.4 | 2.3 | 1.3 | 942 | 88 | 35 |
| 36 | Händelöp | FV | 2020-04-22 | M659 | 5 | 11.1 | 0.5 | 1.2 | <MDA | | 1.4 | 11.0 | 1.6 | 5.5 | <MDA | | 0.5 | 1068 | 53 | 10 |
| 37 | Jogersö | FV | 2020-05-08 | M663 | 5 | 16.1 | 0.8 | 2.0 | <MDA | | 2.4 | 11.8 | 2.5 | 9.1 | <MDA | | 0.9 | 854 | 46 | 17 |
| 38 | Senneby. Sandviken. Väddö | FV | 2020-05-08 | M665 | 7 | 12.0 | 0.8 | 2.2 | <MDA | | 2.7 | 30.2 | 3.6 | 10.2 | <MDA | | 1.3 | 876 | 85 | 37 |
| 39 | Öregrund | FR | 2020-05-07 | M664 | 7 | 14.8 | 0.8 | 1.7 | <MDA | | 2.1 | 22.8 | 2.9 | 7.8 | <MDA | | 1.0 | 856 | 83 | 28 |
| 40 | Valudden | FR | 2020-05-08 | M661 | 7 | 14.9 | 1.5 | 4.5 | <MDA | | 5.8 | | | | <MDA | | 2.6 | 711 | 60 | 84 |
| 41 | Stenskär | FR | 2020-05-08 | M662 | 5 | 20.5 | 0.9 | 1.4 | <MDA | | 1.6 | 31.4 | 2.3 | 6.2 | <MDA | | 0.6 | 642 | 37 | 11 |
| 42 | Norrboda. Gräsö. A | FR | 2020-05-08 | M668 | 5 | 13.8 | 0.6 | 1.1 | <MDA | | 1.3 | 27.6 | 1.9 | 4.9 | <MDA | | 0.5 | 722 | 40 | 9 |
| 44 | Hölick | FR | 2020-05-07 | M667 | 5 | 14.9 | 0.7 | 1.2 | <MDA | | 1.4 | 25.7 | 2.1 | 5.3 | <MDA | | 0.5 | 788 | 42 | 10 |
| 45 | Svenskärs fiskeläge | FR | 2020-05-07 | M666 | 5 | 14.4 | 0.7 | 1.2 | <MDA | | 1.4 | 24.5 | 2.1 | 5.5 | 4.0 | 1.8 | 0.5 | 696 | 43 | 10 |

Table A2 ICP-MS data in *Fucus* spp. and grass, including details of sampling and sampling preparation conditions. N.D. – No data. *From ¹¹⁴Cd. **From ¹¹¹Cd.

| Site no | Site name | Sampling date YYY-MM-DD | Lab code | Co (ppb) | %RSD | Ni (ppb) | %RSD | Cd* (ppb) | %RSD | Cd** (ppb) | %RSD | Gd (ppb) | %RSD | Hg (ppb) | %RSD | Pb (ppb) | %RSD |
|---------|---------------------------------------------------------------|----------------------------|----------|----------|------|----------|------|-----------|------|------------|------|----------|------|----------|------|----------|------|
| 1 | Båteviken, Strömstad | 2020-04-08 | KS-1 | 2522 | 5.5 | 5972 | 7.4 | 1121 | 9.4 | 1088 | 11 | 223 | 4.1 | 21 | 98 | 346 | 5.3 |
| 2 | Fjällbacka, Sälvik, north of harbour, with ongrowth | 2020-04-08 | KS-3 | 2134 | 5.5 | 6090 | 7.6 | 1021 | 6.5 | 981 | 9.4 | 237 | 5.4 | 16 | 19 | 308 | 5.9 |
| 2 | Fjällbacka, Sälvik, harbour and south of harbour, no ongrowth | 2020-04-08 | KS-2 | 2777 | 5.4 | 5662 | 8.1 | 853 | 4.8 | 880 | 4.8 | 393 | 6.7 | 13 | 19 | 647 | 8.0 |
| 3 | Smögen, Kleven | 2020-04-08 | KS-4 | 1176 | 4.2 | 5076 | 11 | 1087 | 5.4 | 1081 | 4.0 | 155 | 3.7 | 16 | 44 | 3941 | 5.0 |
| 4 | Stockevik, Tjörn | 2020-04-08 | KS-5 | 1838 | 5.8 | 5063 | 7.1 | 994 | 7.1 | 1046 | 11 | 526 | 5.4 | 14 | 20 | 1775 | 7.3 |
| 5 | Hällsvik | 2020-04-08 | KS-7 | 2354 | 5.5 | 5133 | 2.7 | 1130 | 7.1 | 1156 | 6.9 | 730 | 6.4 | 27 | 14 | 3610 | 8.5 |
| 6 | Smarholmens badplats | 2020-03-31 | KS-8 | 1022 | 4.7 | 7545 | 6.2 | 1170 | 7.5 | 1196 | 6.7 | 61 | 12 | 12 | 65 | 161 | 9.2 |
| 7 | Frillesås | 2020-03-31 | KS-13 | 5570 | 2.9 | 6131 | 8.6 | 1558 | 3.0 | 1497 | 7.4 | 657 | 3.4 | 19 | 13 | 1079 | 4.1 |
| 8 | Sallebacka | 2020-03-31 | KS-6 | 1761 | 4.0 | 7374 | 5.2 | 1090 | 5.8 | 1110 | 7.4 | 164 | 8.9 | 18 | 19 | 426 | 5.5 |
| 9 | Ringhals, Gloppen | 2020-03-31 | KS-9 | 1656 | 7.2 | 8592 | 6.9 | 1285 | 7.7 | 1257 | 8.0 | 179 | 7.4 | 15 | 13 | 388 | 7.2 |
| 10 | Videbergshamn (Ringhals) | 2020-03-31 | KS-10 | 2962 | 8.0 | 4470 | 5.4 | 893 | 7.3 | 919 | 6.3 | 810 | 5.6 | 24 | 9.4 | 1066 | 5.6 |
| 11 | Bua (Ringhals) | 2020-03-31 | KS-11 | 2153 | 4.4 | 6684 | 4.4 | 1281 | 7.0 | 1272 | 5.3 | 361 | 4.6 | 18 | 20 | 546 | 8.1 |
| 12 | Skeatången, Årnäshalvön | 2020-03-31 | KS-12 | 3223 | 3.2 | 3919 | 3.4 | 1063 | 6.8 | 1058 | 5.6 | 906 | 6.0 | 15 | 16 | 811 | 5.2 |
| 13 | Getterön (M630) | 2020-03-24 | KS-14 | 1567 | 5.4 | 6273 | 7.0 | 918 | 7.4 | 904 | 5.1 | 71 | 6.7 | 18 | 27 | 183 | 5.0 |
| 14 | Glommen (M629) | 2020-03-24 | KS-15 | 2198 | 5.5 | 7596 | 8.1 | 1017 | 5.7 | 1005 | 6.5 | 116 | 8.0 | 18 | 19 | 335 | 8.4 |
| 15 | Särdal A | 2020-03-03 | KS-16 | 1436 | 2.6 | 7100 | 7.3 | 505 | 6.5 | 513 | 7.8 | 115 | 9.3 | 19 | 17 | 370 | 6.2 |
| 15 | Särdal b | 2020-03-03 | KS-17 | 861 | 6.7 | 2858 | 8.7 | 653 | 5.8 | 648 | 5.6 | 66 | 10 | 17 | 28 | 210 | 5.6 |
| 16 | Kattvik | 2020-04-20 | KS-18 | 1021 | 3.7 | 6226 | 6.5 | 886 | 8.1 | 855 | 5.2 | 47 | 8.6 | 14 | 12 | 163 | 8.4 |

| | | | | | | | | | | | | | | | | | |
|----|-----------------------------------|------------|-------|------|-----|-------|-----|------|-----|------|-----|-----|-----|----|-----|------|-----|
| 17 | Mölle | 2020-04-20 | KS-19 | 670 | 5.1 | 7110 | 8.6 | 958 | 14 | 951 | 8.8 | 21 | 12 | 12 | 26 | 159 | 6.8 |
| 18 | Ålabodarna | 2020-03-25 | KS-20 | 1383 | 5.9 | 7513 | 5.6 | 1329 | 8.8 | 1327 | 9.9 | 29 | 9.0 | 10 | 18 | 308 | 6.5 |
| 19 | Barsebäcks camping | 2020-03-24 | KS-21 | 1322 | 4.9 | 7969 | 8.6 | 938 | 7.6 | 879 | 8.3 | 62 | 10 | 11 | 22 | 503 | 5.7 |
| 20 | Barsebäck Skansen | 2020-03-24 | KS-23 | 914 | 6.1 | 7392 | 6.3 | 1018 | 7.9 | 1031 | 9.4 | 38 | 8.1 | 9 | 46 | 190 | 8.5 |
| 21 | Barsebäck between NPP and Skansen | 2020-03-25 | KS-22 | 1678 | 4.5 | 6052 | 2.8 | 662 | 8.6 | 670 | 7.9 | 95 | 6.0 | 9 | 23 | 624 | 6.3 |
| 22 | Barsebäck north of NPP | 2020-03-25 | KS-24 | 1796 | 6.7 | 6470 | 5.4 | 876 | 6.5 | 822 | 6.5 | 68 | 8.1 | 9 | 33 | 350 | 9.1 |
| 23 | Vikhög fort | 2020-03-25 | KS-25 | 1307 | 3.9 | 5665 | 5.1 | 909 | 10 | 950 | 10 | 93 | 9.8 | 16 | 66 | 405 | 6.8 |
| 24 | Vikhög hamn | 2020-03-25 | KS-26 | 1691 | 2.7 | 4982 | 2.9 | 791 | 6.4 | 812 | 5.6 | 273 | 9.6 | 15 | 64 | 853 | 6.7 |
| 25 | Gamla Bjärred, Pilevägen | 2020-04-07 | KS-27 | 1584 | 4.6 | 4934 | 6.8 | 1320 | 9.0 | 1318 | 7.7 | 154 | 7.4 | 16 | 50 | 693 | 8.0 |
| 26 | Lomma | 2020-03-24 | KS-28 | 1645 | 5.2 | 7523 | 3.2 | 1039 | 9.0 | 1083 | 6.9 | 74 | 13 | 12 | 65 | 347 | 9.2 |
| 27 | Limhamn | 2020-03-24 | KS-29 | 996 | 7.0 | 6998 | 3.2 | 1533 | 6.3 | 1526 | 7.6 | 101 | 14 | 12 | 68 | 1399 | 6.4 |
| 28 | Lilla Hammar | 2020-03-24 | KS-30 | 603 | 5.5 | 1348 | 3.1 | 1236 | 9.5 | 1214 | 6.4 | 16 | 15 | 5 | 114 | 105 | 20 |
| 29 | Smygehuks fyr | 2020-03-11 | KS-31 | 1437 | 4.2 | 4779 | 6.6 | 985 | 7.0 | 996 | 8.9 | 44 | 14 | 17 | 40 | 293 | 8.1 |
| 30 | Skillinge, 1A | 2020-03-11 | KS-32 | 3447 | 3.7 | 8437 | 4.5 | 940 | 6.2 | 895 | 8.3 | 161 | 5.5 | 8 | 83 | 921 | 7.6 |
| 30 | Skillinge, 2A | 2020-03-11 | KS-33 | 2881 | 4.4 | 10421 | 6.3 | 1009 | 8.6 | 1074 | 8.0 | 94 | 6.7 | 7 | 97 | 557 | 4.7 |
| 30 | Skillinge | 2020-05-13 | KS-34 | 2181 | 5.7 | 6923 | 2.5 | 806 | 7.9 | 866 | 6.4 | 142 | 7.4 | 9 | 79 | 793 | 7.1 |
| 31 | Hasslö | 2020-03-19 | KS-35 | 1972 | 3.7 | 8477 | 3.3 | 1138 | 7.6 | 1149 | 9.7 | 434 | 6.8 | 8 | 95 | 793 | 6.8 |
| 32 | Sandvik | 2020-04-22 | KS-36 | 2781 | 8.1 | 4446 | 5.5 | 941 | 9.5 | 1093 | 9.2 | 714 | 6.8 | 16 | 89 | 1673 | 6.5 |
| 33 | Dragskär | 2020-04-22 | KS-37 | 385 | 4.5 | 11237 | 5.1 | 1482 | 7.9 | 1499 | 6.6 | 44 | 17 | 7 | 94 | 155 | 7.4 |
| 34 | Simpevarp Åkvik | 2020-04-22 | KS-38 | 784 | 4.3 | 5957 | 4.9 | 798 | 9.1 | 828 | 6.1 | 419 | 6.4 | 9 | 90 | 265 | 11 |
| 35 | Kräkelund | 2020-04-22 | KS-39 | 1592 | 4.5 | 5937 | 3.6 | 1151 | 8.0 | 1209 | 7.1 | 179 | 8.9 | 5 | 107 | 348 | 6.8 |
| 36 | Händelöp | 2020-04-22 | KS-40 | 886 | 5.0 | 10705 | 6.6 | 1124 | 5.1 | 1093 | 5.8 | 87 | 8.6 | 5 | 115 | 205 | 12 |
| 37 | Jogersö | 2020-05-08 | KS-41 | 1817 | 3.6 | 4126 | 8.6 | 906 | 7.1 | 1021 | 5.2 | 504 | 4.5 | 10 | 60 | 881 | 8.7 |
| 38 | Senneby, Sandviken, Vaddö | 2020-05-08 | KS-42 | 1017 | 5.2 | 19110 | 4.3 | 2632 | 7.5 | 2676 | 5.2 | 175 | 9.4 | 8 | 129 | 384 | 5.5 |
| 39 | Öregrund | 2020-05-07 | KS-43 | 1983 | 4.7 | 20736 | 5.4 | 3219 | 5.8 | 3258 | 6.7 | 229 | 6.7 | 8 | 87 | 463 | 8.0 |
| 40 | Valudden | 2020-05-08 | KS-44 | 1559 | 4.8 | 11537 | 6.4 | 2158 | 8.6 | 2230 | 5.5 | 253 | 6.5 | 5 | 160 | 461 | 5.6 |

| | | | | | | | | | | | | | | | | | |
|-----------------------------------------|---------------------|------------|-------|-------------|----------|-------------|----------|-------------|----------|-------------|----------|------------|----------|-----------|-----------|------------|----------|
| 41 | Stenskär | 2020-05-08 | KS-45 | 1999 | 3.9 | 9179 | 3.9 | 1628 | 7.5 | 1785 | 6.6 | 578 | 6.1 | 8 | 81 | 967 | 3.1 |
| 42 | Norrboda, Gräsö, A | 2020-05-08 | KS-46 | 1197 | 3.5 | 14751 | 5.0 | 2378 | 8.5 | 2340 | 6.8 | 169 | 5.4 | 10 | 85 | 297 | 9.1 |
| 42 | Norrboda, Gräsö, B | 2020-05-08 | KS-47 | 1042 | 7.1 | 26045 | 5.4 | 2389 | 4.3 | 2505 | 4.9 | 121 | 6.4 | 4 | 147 | 141 | 10 |
| 43 | Bönan | 2020-05-07 | KS-48 | 1179 | 1.7 | 6619 | 6.1 | 1805 | 6.2 | 1839 | 5.2 | 439 | 5.7 | 9 | 109 | 1708 | 7.3 |
| 44 | Hölick | 2020-05-07 | KS-49 | 1018 | 5.0 | 7413 | 4.6 | 2353 | 4.8 | 2451 | 5.5 | 395 | 7.2 | 9 | 88 | 540 | 5.8 |
| 45 | Svenskärs fiskeläge | 2020-05-07 | KS-50 | 1288 | 6.1 | 6896 | 3.0 | 2571 | 4.7 | 2582 | 7.9 | 295 | 6.0 | 14 | 69 | 3157 | 6.5 |
| Mean | | | | 1725 | 5 | 7709 | 6 | 1271 | 7 | 1289 | 7 | 246 | 8 | 13 | 59 | 746 | 7 |
| Median | | | | 1588 | | 6790 | | 1075 | | 1086 | | 162 | | 12 | | 444 | |
| Standard deviation | | | | 893 | 1 | 4347 | 2 | 581 | 2 | 594 | 2 | 227 | 3 | 5 | 40 | 833 | 3 |
| Standard uncertainty of the mean | | | | 126 | 0 | 615 | 0 | 82 | 0 | 84 | 0 | 32 | 0 | 1 | 6 | 118 | 0 |