



LUND UNIVERSITY

Ionotropic receptors in the turnip moth *Agrotis segetum* respond to repellent medium-chain fatty acids

Hou, Xiaoqing; Zhang, Dan-Dan; Powell, Daniel; Andersson, Martin N; Löfstedt, Christer

Published in:
BMC Biology

DOI:
[10.1186/s12915-022-01235-0](https://doi.org/10.1186/s12915-022-01235-0)

2022

Document Version:
Publisher's PDF, also known as Version of record

[Link to publication](#)

Citation for published version (APA):

Hou, X., Zhang, D.-D., Powell, D., Andersson, M. N., & Löfstedt, C. (2022). Ionotropic receptors in the turnip moth *Agrotis segetum* respond to repellent medium-chain fatty acids. *BMC Biology*, 20, Article 34 (2022). <https://doi.org/10.1186/s12915-022-01235-0>

Total number of authors:
5

Creative Commons License:
CC BY-ND

General rights

Unless other specific re-use rights are stated the following general rights apply:
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

RESEARCH ARTICLE

Open Access



Ionotropic receptors in the turnip moth *Agrotis segetum* respond to repellent medium-chain fatty acids

Xiao-Qing Hou^{1,2†}, Dan-Dan Zhang^{1†}, Daniel Powell^{1,3}, Hong-Lei Wang¹, Martin N. Andersson¹ and Christer Löfstedt^{1*} 

Abstract

Background: In insects, airborne chemical signals are mainly detected by two receptor families, odorant receptors (ORs) and ionotropic receptors (IRs). Functions of ORs have been intensively investigated in Diptera and Lepidoptera, while the functions and evolution of the more ancient IR family remain largely unexplored beyond Diptera.

Results: Here, we identified a repertoire of 26 IRs from transcriptomes of female and male antennae, and ovipositors in the moth *Agrotis segetum*. We observed that a large clade formed by IR75p and IR75q expansions is closely related to the acid-sensing IRs identified in Diptera. We functionally assayed each of the five AseglIRs from this clade using *Xenopus* oocytes and found that two receptors responded to the tested ligands. AseglIR75p.1 responded to several compounds but hexanoic acid was revealed to be the primary ligand, and AseglIR75q.1 responded primarily to octanoic acid, and less so to nonanoic acid. It has been reported that the C₆-C₁₀ medium-chain fatty acids repel various insects including many drosophilids and mosquitos. We show that the C₆-C₁₀ medium-chain fatty acids elicited antennal responses of both sexes of *A. segetum*, while only octanoic acid had repellent effect to the moths in a behavioral assay. In addition, using fluorescence in situ hybridization, we demonstrated that the five IRs and their co-receptor AseglIR8a are not located in coeloconic sensilla as found in *Drosophila*, but in basiconic or trichoid sensilla.

Conclusions: Our results significantly expand the current knowledge of the insect IR family. Based on the functional data in combination with phylogenetic analysis, we propose that subfunctionalization after gene duplication plays an important role in the evolution of ligand specificities of the acid-sensing IRs in Lepidoptera.

Keywords: *Xenopus* oocytes, Fluorescence in situ hybridization, Acid-sensing, Repellent, Octanoic acid, Lepidoptera, Gene expression

Background

Insects have evolved the ability to accurately sense environmental chemical signals, which play critical roles for their survival and reproduction [1, 2]. The peripheral process of chemoreception relies on receptors that interact

with external molecular cues, triggering the transduction of chemical signals into electrical signals, which may ultimately result in a behavior [3, 4]. In insects, airborne chemicals are mainly detected by ligand-activated receptors from two families, namely the odorant receptors (ORs) [5] and ionotropic receptors (IRs) [6]. In *Drosophila*, these two receptor families are present in the dendritic membrane of the olfactory sensory neurons (OSNs) housed in different types of sensilla with distinct chemical preferences: ORs expressed predominantly in trichoid and basiconic sensilla detect, for example, pheromones

*Correspondence: christer.lofstedt@biol.lu.se

[†]Xiao-Qing Hou and Dan-Dan Zhang have shared first authorship

¹ Department of Biology, Lund University, Sölvegatan 37, 223 62 Lund, Sweden

Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

and plant volatiles, including alcohols, aldehydes, esters and aromatics; IRs, mainly expressed in coeloconic sensilla, primarily respond to acids and amines [6–9].

IRs originated from the ionotropic glutamate receptor (iGluR) gene family and share similar structure and mechanism of action as iGluRs. However, the expression pattern and function of the IR family differ from the three well-studied iGluR subfamilies; kainate, NMDA (N-methyl-D-aspartate) and AMPA (α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid) receptors that are expressed on the surface of synapses in animal nervous systems and play a role in signal transmission between neurons [6, 8]. Insect IRs are present in different tissues and are more diverse in function. IRs were first identified in the model insect *Drosophila melanogaster* as odorant sensing receptors [6], and subsequent studies extended our knowledge on their functions to the involvement in taste [10–13], temperature [14, 15], humidity [16–18] and even auditory sensing [19]. Similar to the ORs that are co-expressed with the conserved odorant receptor co-receptor (Orco) to form a functional ligand-gated ion channel [20–22], co-expression in neurons is also observed among insect IRs. These receptors display more complex combinations since at least three IRs (IR8a, IR25a, and IR76b) can function as co-receptors of specific tuning IRs. IR8a is normally co-expressed with IRs tuned to acids [23–26], while IR25a and IR76b are co-expressed with IRs responding to amines [8, 27]. In addition to the role in olfaction, IR25a was also reported as a co-receptor for IRs detecting temperature, humidity, and taste molecules [15, 16, 28].

Based on their phylogeny and expression profiles, *Drosophila* IRs are classified into two groups: antennal IRs (A-IRs) and divergent IRs (D-IRs). A-IRs are highly conserved across insects and mainly expressed in the antennae, whereas the D-IRs have radiated to different extents in different species and are present in various tissues [6, 8, 10, 25, 29–31]. A number of the so-called Lepidoptera-specific IRs (LS-IRs) are absent in *Drosophila* and were originally identified in Lepidoptera [32–37]. Despite the name, some of them are also present in Trichoptera and mosquitoes with overall high support values, although in some cases the orthology remains somewhat ambiguous [38, 39]. The LS-IRs do not represent a single monophyletic IR subfamily, because they are scattered across the phylogenetic tree, with some (IR1, IR75p, IR75q, and IR87a) located within the A-IR while others (IR7d, IR100b-j, and IR143) nested within the D-IR clade, and also include a putative pseudogene (IR2) [36, 37]. The large IR75p and IR75q expansions are of particular interest due to their close relationship with the characterized acid-sensing IRs [36, 37].

Over the past decade, much progress has been made in elucidating the functions of IRs in *Drosophila* and a few mosquito species. Although this major gene family spans across protostomes [29, 40], the functional information of IRs beyond dipterans is still limited, restricting our knowledge concerning the evolution and specialization of this ancient gene family. To our knowledge, only two IRs from the parasitoid wasp *Microplitis mediator* have been functionally investigated [41]. The second largest insect order, Lepidoptera, rely on their remarkable olfactory abilities to sense the external environment. However, the function of IRs in Lepidoptera remains largely unexplored, although recent studies in *Manduca sexta* and *Mythimna separata* suggested a role for the co-receptor IR8a in acid-sensing [9, 42]. In the present study, we report the first functional characterization of IRs in Lepidoptera. We show that two IRs from the turnip moth *Agrotis segetum* respond to medium-chain fatty acids and that these fatty acids elicit antennal responses, with octanoic acid being significantly repellent to moths in an olfactory bioassay. Further, we find that in this moth the acid-detecting IRs are expressed in basiconic or trichoid sensilla, but not in coeloconic sensilla as was found in *Drosophila*. Based on our phylogenetic analysis and receptor characterization, we suggest that subfunctionalization occurred in Lepidoptera after a gene duplication event in the IR75p/q clade.

Results

Characterization of the *A. segetum* IR repertoire from antennal and ovipositor transcriptomes

Studies on *M. sexta* have shown that in addition to the antennae, the ovipositor plays an important chemosensory role [43]. Hence, in the present study we profiled replicated transcriptomes of the antennae ($N = 4$ for both sexes) and ovipositors ($N = 2$) from *A. segetum* (“Aseg”) by mRNA sequencing (RNA-Seq) for an exhaustive search of IRs involved in chemosensation. This produced a total of over 50 Gb of sequence data with the yield of each sample reaching over 10 million paired-end (100 bp) reads. Clean reads from all 10 samples were assembled together to produce a transcriptome comprised of 83,645 non-redundant transcripts having a mean length of 976 bp and an N_{50} of 1788 bp.

In total, 26 AsegIR genes were identified, including the three co-receptors, IR8a, IR25a, and IR76b. Of the 26 assembled IR transcripts, 23 encoded full-length genes. We followed the nomenclature previously used to name lepidopteran IRs [36]. The raw sequence reads have been deposited in the SRA database at NCBI under the Bioproject accession number PRJNA707654 [44].

The IR75p and IR75q lineages grouped with acid-sensing IRs
We performed phylogenetic analyses of AsegIRs along with the IR sequences from *Bombyx mori* (Bombycidae), *Plutella xylostella* (Plutellidae), *Helicoverpa armigera* (Noctuidae), *D. melanogaster*, and *Anopheles gambiae* (both Diptera) (Fig. 1). The two IR classes, A-IRs and

D-IRs, identified in previous studies [29, 36, 37] can be recognized in the tree. A single AsegIR ortholog from all the 18 monophyletic groups of conserved A-IRs (IR1.1, IR1.2, IR21a, IR31a, IR40a, IR41a, IR60a, IR64a, IR68a, IR75d, IR75p, IR75p.1, IR75p.2, IR75q.1, IR75q.2, IR76b, IR77a, IR93a) was identified. Fewer AsegIRs




Fig. 1 Phylogeny of Lepidoptera IRs. The maximum-likelihood phylogenetic tree is based on a protein sequence alignment of ionotropic receptors (IRs) from *A. segetum*, *B. mori* [29], *P. xylostella* [36], *H. armigera* [36], *D. melanogaster* [6, 29], and *A. gambiae* [29, 45]. The tree was rooted with the lineage of IR8a and IR25a receptors. The IRs from different species are color coded as indicated in the figure. The monophyletic IR groups are noted on the tree, with the light blue color indicating "Lepidoptera-specific IR groups." The large IR7d, IR75p, and IR75q expansions are indicated by red branches, and the entire putative "expanded acid-sensing cluster" (IR75d however responding to an amine in *Drosophila*) is highlighted in yellow. The five IR75p and IR75q receptors functionally assayed in this study are marked with asterisks. Bootstrap support (100 replicates) values over 0.7 are shown at corresponding nodes

were identified in the D-IR class, which was expected because D-IRs are typically expressed in tissues other than antennae and ovipositors, and presumably mediate non-olfactory behaviors. The large expansions of IR7d, IR75p, and IR75q have been evident in almost all the reported antennal transcriptomes of Lepidoptera, and these IRs typically have relatively high levels of gene expression. The expression of IR75p and IR75q genes is predominantly restricted to the adult antennae, whereas IR7d genes seem to be generally distributed in different body parts of adults and larvae [35–37]. Notably, the lineage expansions IR75p and IR75q, along with IR1.1 and IR1.2, grouped within a larger cluster together with IR75a, IR64a, and IR31a, which have been reported as acid-sensing IRs in *Drosophila* and mosquitos [8, 24, 26], although IR75d in the neighboring expansion responded to an amine (pyrrolidine) in *Drosophila* [24].

AseglRs in 75p/q expansions were predominantly expressed in adult antennae

The expression levels of the 26 AseglRs were normalized across sequencing libraries using transcripts per million (TPM) scaling factor [46]. Surprisingly, one of the LS-IRs, AseglR7d.3, was found to be the most abundantly expressed IR, with much higher levels of expression than that of the three co-receptors, and with higher expression in ovipositors than in both male and female antennae. Among the three co-receptors, AseglR25a showed the highest expression, with higher TPM values compared to AseglR8a and AseglR76b. AseglR25a and AseglR76b had non-biased expression in the antennae of both sexes, while AseglR8a showed female-biased expression. AseglR75q.1 and AseglR75q.2 were identified from a large polycistronic transcript with comparatively high expression in both male and female antennae. The other three IRs in the IR75 clade, i.e., IR75p, IR75p.2 and especially IR75p.1, were expressed at much lower levels (Fig. 2A).

Quantitative PCR was conducted to further investigate the expression pattern of the AseglRs from the IR75p/q expansions and their putative co-receptor IR8a in various tissues of both sexes, including chemosensory organs (antennae and proboscises) of both sexes, female ovipositor, as well as male abdomen, head and thorax (without antennae), legs, and wings. GAPDH (glyceraldehyde 3-phosphate dehydrogenase) and RPS3

(ribosomal protein S3) were used as reference genes. The results showed that AseglRs from the IR75p/q expansions and AseglR8a were expressed predominantly in adult antennae of both sexes (one-way ANOVA followed by an LSD test, $p < 0.05$ for all genes), although IR75p.1 and IR75p.2 showed a broader tissue distribution, being weakly expressed in almost all assayed non-chemosensory tissues (Fig. 2B). Consistent with the results of the transcriptome analysis, AseglR8a had numerically higher expression level in female antennae than in male antennae, but the difference was not statistically significant (independent samples t -test, $t = -0.813$, d.f. = 4, two-sided $p = 0.462 > 0.05$); AseglR75q.1 and AseglR75q.2 were much more abundantly expressed in the antennae than the other IR genes in the expansion (one-way ANOVA followed by an LSD test, $p < 0.001$), while AseglR75p.1 had a lower expression level and was female-biased (independent samples t -test, $t = -13.441$, d.f. = 4, two-sided $p < 0.001$) (Fig. 2B).

AseglRs are expressed in basiconic or trichoid sensilla

In *Drosophila*, IRs are mainly located in the antennal coeloconic sensilla [6, 25], whereas it was previously unknown which sensillum type the IRs in moths are associated with. To address this question, we first studied the morphology and distribution of the sensilla on *A. segetum* antennae by scanning electron microscopy (SEM). The antennae of *A. segetum* are sexually dimorphic, with setaceous antennae in females and plumose antennae in males (Fig. 3A–D). There is one pair of branches on most of the segments of male antennae, whereas the segments towards the tip of the antennae are unbranched (Fig. 3B, D). The SEM images indicated that the trichoid sensillum was the most abundant type on both female and male antennae and it was the only sensillum type present on the branches of male antennae (Fig. 3E, F, H). Only two or three coeloconic sensilla were observed on each segment of the male antennae, including the unbranched segments and the stem parts of the branched segments, but not on the branches; they were more abundant (six observed) on each segment of the female antennae (Fig. 3E, F). Basiconic sensilla were present throughout the entire female antennae and male antennae (except for the branches) but in lower abundance compared to trichoid sensilla in both sexes (Fig. 3E, F).

(See figure on next page.)

Fig. 2 The expression profiles of AseglRs. **A** Heatmap with TPM value (transcripts per million) showing the expression levels of the AseglRs in different female and male antennal samples, as well as ovipositor samples. **B** The expression of AseglRs from IR75p and IR75q lineages and the co-receptor IR8a in different tissues and sexes, determined by qPCR. Three biological replicates, each including three technical replicates, were performed for each gene. M, male; F female; ant, antennae; prob, proboscis; ovi, ovipositors; ab, abdomens; h+t, head and thorax (without antennae); w, wings. Individual data values are shown as green filled circles. As AseglR75q.1 and AseglR75q.2 are located on the same transcript, it is not possible to distinguish the TPM value for each of them from the transcriptome data, whereas the qPCR results demonstrated that both were more abundant than the other three AseglRs in the IR75p/q expansion




Fig. 2 (See legend on previous page.)

To further explore the distribution of AsegIR75p/q and their putative co-receptor AsegIR8a on female and male antennae and to confirm their co-expression pattern, we performed two-color fluorescence in situ hybridization (FISH) with digoxigenin (DIG)-labelled antisense RNA probes for the five AsegIRs and biotin-labelled antisense probes for AsegIR8a. Corresponding sense probes were used as control. On female antennae, AsegIR8a was abundantly expressed across the whole segment, while signals of the five AsegIRs were much fewer (Fig. 4 and Fig. 5). On the male antennae, AsegIR8a and the five AsegIRs were predominantly expressed at the branched segments but not the unbranched segments towards the tip of the antennae (Fig. 4 and Fig. 5 and Additional file 1: Fig. S1). They were not expressed on the stretches of the branches (Additional file 1: Fig. S1) but distributed at the bases of the branches and at the stem parts (Fig. 4 and Fig. 5). Fewer signals of the five AsegIRs were detected compared to AsegIR8a signals, consistent with their expression levels. The two-color FISH confirmed that the five AsegIRs were indeed co-expressed with AsegIR8a in both sexes (Fig. 4 and Fig. 5). Intriguingly, all the five AsegIRs as well as AsegIR8a were in both sexes not expressed in coeloconic sensilla on the antennae as in *Drosophila* (Fig. 4 and Fig. 5 and Additional file 1: Fig. S1). On the female antennae, they seemed to be expressed in the OSNs located in basiconic sensilla. On the male antennae, however, because the bases of the branches have high density of hair-like sensilla, we were not able to distinguish from the FISH images whether the IRs were expressed under basiconic sensilla or trichoid sensilla.

IR75p.1 and IR75q.1 responded to medium-chain fatty acids in *Xenopus* oocytes

The five AsegIRs from the IR75p/q expansions were each co-expressed with AsegIR8a in *Xenopus* oocytes, and screened against a panel of 46 compounds, most of which are common plant volatiles, including acids, aldehydes, and alcohols. Oocytes co-expressing AsegIR75p.1/AsegIR8a showed responses to acids and aldehydes with exclusively C₅-C₇ straight chains at a concentration of 100 μ M. For the alcohol ligands, only C₆ unsaturated compounds elicited responses. The strongest response was elicited by hexanoic acid, followed by (Z)-3-hexenol, (E)-2-hexenal, and hexanal (Fig. 6A, B). Dose-response trials

indicated a higher sensitivity of AsegIR75p.1 to hexanoic acid compared to the other active ligands (Fig. 6B). Oocytes co-expressing AsegIR75q.1/AsegIR8a showed a strong response to octanoic acid and a weaker response to nonanoic acid at a concentration of 100 μ M. This receptor also responded slightly to heptanoic acid and octanal, with response magnitudes much lower than that to the primary ligand (Fig. 6A, C). The responses to octanoic acid and nonanoic acid were dose-dependent, with a threshold concentration at 1 μ M (Fig. 6C). The oocytes injected with the cRNAs of AsegIR75q.1, AsegIR75p.1, and AsegIR8a alone did not show any response to the tested compounds. In addition, the other three receptors in the same expansions, AsegIR75p, AsegIR75p.2, and AsegIR75q.2, did not respond to any of the tested compounds when co-expressed with AsegIR8a in oocytes.

Medium-chain fatty acids elicit antennal responses

Due to the responses of AsegIR75p.1 and AsegIR75q.1 to medium-chain fatty acids (C₆-C₁₀) in the oocytes, we further tested whether these acids could elicit antennal responses in the moths by gas chromatography coupled to electroantennographic detection (GC-EAD). Both female and male antennae showed clear responses to the tested acids (Fig. 7A). The responses to decanoic acid were smaller in both sexes, while the response magnitudes to the other tested acids were similar (one-way ANOVA followed by an LSD test, $p < 0.05$). Male moths showed larger responses to hexanoic acid and octanoic acid than female moths (independent samples *t*-test, $t = -3.721$, d.f. = 8, two-sided $p = 0.006$ for hexanoic acid; $t = -3.412$, d.f. = 8, two-sided $p = 0.009$ for octanoic acid), but the responses to the other tested acids were not significantly different between the two sexes (two-sided $p > 0.05$ in *t*-test) (Fig. 7B).

Octanoic acid is behaviorally repellent to *A. segetum*

In order to reveal the biological significance of the observed IR responses to fatty acids, we performed two-choice Y-tube bioassays. In an initial experiment, we tested whether the odor of honey solution (10%) was attractive to the adult moths. Among 51 individuals (29 females and 22 males), most of *A. segetum* from both sexes (female: $\chi^2 = 12.45$, d.f. = 1, $p = 0.0004$; male: $\chi^2 = 4.545$, d.f. = 1, $p = 0.033$) were attracted to the honey

(See figure on next page.)

Fig. 3 Scanning electron micrographs of *A. segetum* antennae. **A,B** Schematic diagrams and **C,D** overall SEM images showing the sexually dimorphic structures of female and male antennae of *A. segetum* moths. The distribution of coeloconic sensilla (yellow diamonds), trichoid sensilla (magenta triangles), and basiconic sensilla (green circles) are illustrated in the enlarged view of segments in the dashed frame. The stem part of male antennae, the branches, and unbranched part are labelled in the figure. **E,F** Antennal segments showing representative coeloconic, trichoid, and basiconic sensilla. **G** Close-up view of a coeloconic sensillum. **H** Coeloconic sensilla are absent on the branches of male antennae. **I** Close-up view of trichoid sensilla and basiconic sensilla. The labelled number in the scales represents the length between the first and last marks, and the distance between two marks represents a tenth of the labelled number




Fig. 3 (See legend on previous page.)

solution and only two males chose water. In another control experiment, cotton balls with 10% honey solution were placed in each of the two arms of the Y-tube and were equally attractive to both female and male *A. segetum* (Fig. 7C), demonstrating no side preference. In the treatment groups, individual female and male moths were given the choice between 10% honey solution and 10% honey solution loaded with one of the C₆-C₁₀ fatty acids at the dose of 100 µg. Our results indicated that among the five tested acids, only octanoic acid showed a repellent effect on both sexes, with significantly fewer individuals choosing the honey solution with octanoic acid compared to honey solution alone (female: $\chi^2 = 11.7647$, d.f. = 1, $p = 0.0006$; male: $\chi^2 = 9.3226$, d.f. = 1, $p = 0.0022$). The moths showed no significant preference/avoidance in treatments with other acids (Fig. 7C). The subsequent dose-response experiment showed that adults of both sexes were significantly less attracted to the 10% honey solution with octanoic acid above 1 µg. No significant difference was observed between the control and the dose of 100 ng octanoic acid in either sex (female: $\chi^2 = 0.1538$, d.f. = 1, $p = 0.6949$, male: $\chi^2 = 0.36$, d.f. = 1, $p = 0.5485$) (Fig. 7D).

Discussion

IRs have been well studied in *Drosophila* and in some mosquito species, whereas the function of this gene family remains poorly understood in other insects including the Lepidoptera. Through phylogenetic analysis of lepidopteran IRs, several large receptor lineage expansions are observed [36], among which the IR75p and IR75q lineages are the most salient due to their close relationship with acid-sensing IRs and enriched expression in adult antennae. The tandem duplications in these expansions were considered to be associated with the discrimination of structurally similar chemicals in high resolution [25, 35]. Here we present the first comprehensive investigation of the ligand specificities of IRs from the IR75p/q expansions.

We identified a set of 26 AsegIRs from the antennal and ovipositor transcriptomes of *A. segetum*. Based on the number of IRs identified in the antennal transcriptomes of other moth species and *Drosophila* (e.g., 17 in *Spodoptera littoralis*, 12 in *H. armigera* and 19 in *D. melanogaster*) [34, 47, 48], our dataset appears to include most of the IRs expected to be expressed in the antennae and ovipositor of *A. segetum*. This number is, however, lower compared to those identified from the genomes of

other species (e.g., 45 in *S. litura*, 39 in *H. armigera* and 66 in *D. melanogaster*) [29, 36, 37], which include the IRs expressed in the other body parts and pseudogenes. From the phylogenetic analysis, the IR75p/q expansions together with IR1.1 and 1.2 clades were grouped into a large cluster in which some receptors including IR31a, IR64a, and IR75a,b,c have been reported as acid sensors in *D. melanogaster*; IR64a.1 and IR64a.2 in the parasitoid wasp *M. mediator*; and IR75k in the mosquito *A. gambiae* [8, 24–26, 41, 49]. This suggested that the receptors within the IR75p/q expansions may also be involved in acid detection.

Based on the hypothesis of an “acid-sensing” cluster of receptors, all five AsegIRs within the IR75p/q expansions were functionally investigated in *Xenopus* oocytes. We characterized two functional IRs, with AsegIR75q.1 being specific for octanoic acid, and AsegIR75p.1 responding relatively broadly to C₅-C₇ straight chain acids and aldehydes, as well as unsaturated C₆ alcohols. On top of this in vitro test at the receptor protein level, our in vivo GC-EAD experiments showed that both sexes of *A. segetum* were able to detect C₆ to C₁₀ fatty acids. We further explored the behavioral significance of the responses to these fatty acids and found that only octanoic acid had a significant repellent effect on female and male *A. segetum* moths, indicating that AsegIR75q.1 might mediate the behavioral repellency of octanoic acid.

Fatty acids are common in nature and are normal constituents of vertebrate skin [50, 51]. It has been shown in a number of studies that the medium-chain fatty acids, in particular C₈-C₁₀, are highly repellent or toxic to many arthropods, including flies, mosquitos, ticks, ants, midges, and bees to name a few [50–55]. For instance, octanoic acid and hexanoic acid are highly abundant in the fruit of *Morinda citrifolia* and function as secondary defence compounds which are lethal or toxic to most drosophilids, with the exception of *D. sechellia* which has evolved physiological adaptations to specialize on the “toxic” *Morinda* fruit [56, 57]. Also, the C₆-C₁₀ saturated fatty acids act as negative signals for ovipositional responses of female mosquitos [58, 59]. In this study, we show that octanoic acid is also a repellent for a lepidopteran species, using a “reverse chemical ecology” approach. IR75q.1 is well conserved across the Lepidoptera, and further studies are needed to address the question of whether the repellency of octanoic acid is conserved in lepidopteran insects, or if it has a specific ecological importance to *A. segetum*. The biological

(See figure on next page.)

Fig. 4 Whole-mount fluorescence in situ hybridization (WM-FISH) showing the co-expression of AsegIR8a with AsegIR75p (A), AsegIR75p.1 (B), and AsegIR75p.2 (C) on adult female and male antennae. Neurons expressing AsegIR8a are indicated in green, AsegIR75p/p.1/p.2 signals are indicated in magenta, and the merged signals are shown in white. Red arrowheads point to the location of coeloconic sensilla. The moths used in WM-FISH were 3–5 days old after eclosion and unmated





Fig. 4 (See legend on previous page.)



significance of the response of AsegIR75p.1 to hexanoic acid for *Agrotis* moths is still elusive as this compound was behaviorally inactive in our assay. However, a recent study suggested that hexanoic acid deterred female hawk-moths (*M. sexta*) from ovipositing on plants [9].

We also explored the antennal distribution patterns of AsegIR75p/q and the co-receptor AsegIR8a, which showed signals from hair-like (basiconic or trichoid) sensilla, but not from coeloconic sensilla, on female and male antennae of *A. segetum*. In *D. melanogaster*, ten antennal IRs (IR31a, IR41a, IR75a, IR75b, IR75c, IR75d,

IR76a, IR76b, IR84a, and IR92a) are expressed in subsets of coeloconic sensilla on the antennae. The other four antennal IRs are not expressed in coeloconic sensilla but found in neurons of the arista (IR21a) and sacculus (IR40a, IR93a and IR64a) together with their co-receptors [6, 25, 26]. The two co-receptors, IR8a and IR25a, are broadly expressed in coeloconic sensilla in *D. melanogaster*; IR8a is also expressed in sacculus neurons while the IR25a is additionally detected, but more weakly so, in basiconic and trichoid sensilla as well as aristal and sacculus neurons [6, 25]. Electrophysiological recordings from *D. melanogaster* showed that the neurons housed in coeloconic sensilla detect acids, ammonia, and humidity, which indirectly suggests that the IRs define the response profiles of the coeloconic neurons [6, 7]. In the desert locust *Schistocerca gregaria*, IR25a and IR8a were shown to be expressed in coeloconic sensilla [60]. Beyond these two species, no additional evidence suggest that IRs are generally expressed in coeloconic sensilla. However, accumulating evidence indicates that in some insects, IRs might be expressed in non-coeloconic sensilla. In mosquitos, trichoid sensilla and grooved-peg sensilla both responded to amines and acids [61] and it was speculated that the responses were underlined by IRs expressed in these sensilla [25, 45]. In a basal insect species, *Lepismachilis y-signata* (Archaeognatha), in which IRs are the main olfactory receptors detecting different chemical classes of odors, IRs are expressed in OSNs underneath basiconic sensilla and no coeloconic sensilla were observed on the antennae [62]. It was suggested that the evolutionary origin of IRs might have been associated with basiconic sensilla and then their expression switched to coeloconic sensilla after the neopteran insects originated [62].


Based on the limited information regarding the sensory physiology of coeloconic sensilla in Lepidoptera, it seems that the response profiles of coeloconic, basiconic, and trichoid sensilla partly overlap despite their striking morphological differences. In *B. mori*, although some coeloconic sensilla were specialized towards C_4 – C_9 carboxylic acids, coeloconic sensilla responding to aldehydes and alcohols were also abundant; both coeloconic and


basiconic sensilla responded to hexanoic acid in *B. mori* and *M. sexta* [9, 63]. In addition, no thermo- or hygro-receptors were found in coeloconic sensilla of *B. mori* [63]. It is therefore likely that the expression of IRs is not restricted to coeloconic sensilla. Our study provides the first evidence from Lepidoptera that AsegIR75p/q and the co-receptor (AsegIR8a) are expressed in OSNs located in basiconic or trichoid sensilla, suggesting that the expression of IRs in specific sensillum types might be taxon-specific. This also suggests that the coeloconic structure is not a prerequisite for IR function in neopteran insects, at least not for acid-sensing IRs. However, further studies are needed to investigate at which taxonomic level or where in the insect phylogeny this change in expression pattern had occurred.

Gene duplications are considered an important mechanism through which organisms can acquire new genetic material for specific adaption and evolution of new functions [64, 65]. After gene duplication, the duplicated genes are less constrained by purifying selection and may start to accumulate mutations resulting in novel functions, or in most cases to accumulate loss-of-function mutations generating pseudogenes [65–67]. Gene subfunctionalization can also happen which means that both pre-existing and duplicated genes distinctively but complementarily retain some of their original function [65, 68]. Our phylogenetic tree shows that two mosquito (*A. gambiae*) IRs, AgamIR75k and AgamIR75l, clustered within the IR75p/q expansions and are closer to the IR75q lineage. AgamIR75k is also an acid-sensing receptor, showing primary responses to octanoic and nonanoic acids, and secondary responses to hexanoic, heptanoic, and decanoic acids [49]. The response spectrum of AgamIR75k seems to be a combination of the response profiles of AsegIR75p.1 and AsegIR75q.1, which indicates functional conservation of the IR75p/q expansions across different insect orders and also suggests that subfunctionalization has occurred after gene duplication in the IR75p/q expansions in Lepidoptera. The expression level of AsegIR75q.1 was much higher than that of AsegIR75p.1 and other IRs within the same clade, implying that the key ligand of AsegIR75q.1, octanoic acid, plays

(See figure on next page.)

Fig. 6 AsegIR75p.1 and AsegIR75q.1 respond to medium-chain fatty acids in *Xenopus* oocytes. **A** Response profiles of AsegIR75p.1 (left) and AsegIR75q.1 (right) to the full odor panel. Response magnitudes were normalized to the average response of primary ligand ($N \geq 4$). Error bars indicate the SE. **B,C** Upper panel: Representative current traces of oocytes upon successive exposures to 100 μ M stimuli. Each compound was applied at the time indicated by the arrowheads for 20 s. Lower panel: Dose-dependent responses, showing values normalized to the average response of the most active compound at 100 μ M ($N \geq 4$ for each ligand). Individual data values are listed in Additional file 3. The responses of AsegIR75p.1 to the four primary ligands at each dose were compared by one-way ANOVA followed by an LSD test. * in **B** indicates the doses at which the responses to hexanoic acid are significantly higher than to the other three ligands ($p < 0.05$). The responses of AsegIR75q.1 to the two primary ligands at each dose were compared by independent samples *t*-test. ** in **C** indicates the doses at which the responses to octanoic acid were significantly different from the response to nonanoic acid (at 1 mM concentration, $t = 9.055$, d.f. = 8, two-sided $p < 0.001$; at 100 μ M concentration, $t = 4.486$, d.f. = 8, two-sided $p = 0.002$)





a biologically important role for *A. segetum* and needs to be detected with high resolution. The other three receptors, AsegIR75p, AsegIR75p.2, and AsegIR75q.2, showed no response to any of the 46 tested compounds although they have higher expression levels than AsegIR75p.1, probably because the ligands of these receptors were not included in the test panel. An alternative explanation could be that nonfunctionalization occurred after the gene duplication.

Conclusions

We present the first comprehensive investigation of ligand specificities of IR75p and IR75q expansions and found that these IRs responded to medium-chain fatty acids. All the C₆–C₁₀ saturated fatty acids elicited antennal responses in the *A. segetum* moths, whereas only the primary ligand of AsegIR75q.1, octanoic acid, acted as a behavioral repellent. We also provide direct evidence that the acid-sensing receptor AsegIR75q.1 and

its co-receptor are expressed in basiconic or trichoid sensilla, indicating that the coeloconic structure is not a prerequisite for IR function in neopteran insects. Our study expands the acid-sensing IR cluster in insects and we propose that gene subfunctionalization may have played an important role after gene duplication events during olfactory specialization in the IR75p/q expansions in Lepidoptera.

Methods

Chemicals

A set of 46 odorants from different chemical classes were tested in this study, including acids, aldehydes, and alcohols. Most of them are common plant volatiles and can elicit responses of neurons housed in coeloconic sensilla in *Drosophila* and *B. mori* (Additional file 1: Table S1) [7, 63]. For oocyte recordings, stock solutions were prepared by dissolving each compound to 100 mM in dimethyl sulfoxide (DMSO), which were stored at -20°C . Before each experiment, the stock solution was diluted to indicated concentration in Ringer's buffer (96 mM NaCl, 2 mM KCl, 5 mM MgCl_2 , 0.8 mM CaCl_2 , 5 mM HEPES, pH 7.6) with the final stimuli containing 0.1% DMSO. Ringer's buffer containing 0.1% DMSO was used as negative control.

Insects

The *A. segetum* used in the experiments were from the continuous culture at the Pheromone Group, Department of Biology, Lund University. Larvae were reared on an artificial bean-based diet, under a 16L:8D photoperiod, 50% relative humidity, and a temperature of 25°C . Pupae were sexed and females and males kept separately, and emerged adults were fed with 10% honey solution. Individuals used in all experiments were unmated.

RNA extraction, transcriptome sequencing, assembly, and annotation

Male and female antennae, and ovipositors were collected from 3-day-old virgin *A. segetum* under a stereomicroscope. In total, four antennal samples for each sex and two ovipositor samples were prepared. Each antennal sample contained 5 pairs of antennae and each ovipositor sample contained 40 ovipositors. The collected tissues were immediately frozen on dry ice and then stored at -80°C until use. The samples were homogenized, and the total RNA was extracted and purified using RNeasy Micro Kit (Qiagen, GmbH, Hilden, Germany). The quality and quantity of the DNase-treated RNA samples were verified using a BioAnalyzer (Agilent).

Library preparation using the Illumina TruSeq RNA poly-A selection kit and sequencing over two lanes on an Illumina HiSeq2500 with Rapid SBS Kit v2 chemistry was performed at SciLifeLab (Stockholm, Sweden). Quality

of the raw read data was assessed using FastQC (<http://www.bioinformatics.babraham.ac.uk/projects/fastqc/>). Trimming of adaptors, filtering of low-quality reads, and removing contaminants were performed using Trimmomatic (v0.36) [69] with a custom screening database. Clean reads were assembled using Trinity (v2.8.2) with default settings. Contigs from the Trinity output were clustered to remove redundancy using CD-HIT-EST (v 4.6.8) [70] with a sequence identity threshold of 0.95. The non-redundant transcript assembly was screened for putative protein-coding regions using the program TransDecoder (release 5.0.1) (<http://transdecoder.github.io/>) with the option of retaining hits to the PfamA domain database [71]. Completeness was assessed using the BUSCO (v3.0.2b) [72] tool searching against the insecta_odb9 database of 1658 genes. Clean read data from each pooled sample was then mapped to the non-redundant transcripts using the align_and_estimate_abundance.pl script from the Trinity v2.8.2 software package using default parameters except for `--est_method RSEM --aln_method bowtie2 --trinity_mode`. Reads not mapping in pairs were discarded from DE analysis. The output was analyzed using the Bioconductor [73] package DESeq2 (v1.22.2) [74] in R.

Initial functional annotation of assembled sequences was performed by blasting against the pooled database of nr (NCBI non-redundant protein sequences), as well as KOG/COG (Clusters of Orthologous Groups of proteins) and Swiss-Prot with threshold E-value $< 1\text{e}-5$. Additional blast searches were performed using the identified *A. segetum* IR genes as queries and the transcriptome as a custom database to ensure that all IR genes were discovered. The IRs were labelled following the nomenclature convention of lepidopteran IRs [36].

Phylogenetic analysis

A phylogenetic tree was built based on the amino acid sequences of IRs from *A. segetum*, *B. mori*, *P. xylostella*, *H. armigera*, *D. melanogaster*, and *A. gambiae*. The sequence alignment was performed using ClustalW, a built-in plugin with default settings in Geneious R9 (Biomatters Ltd. Auckland, New Zealand). To ensure the quality of the alignment and subsequent construction of the tree, short sequences (< 300 amino acids) were excluded. A maximum-likelihood tree was constructed using MEGA 7 with the WAG+G+F substitution model that was chosen using "Find best protein model" in MEGA 7. The tree was rooted with the IR8a and IR25a receptor lineage and further viewed and edited using FigTree 1.4.4.

Quantitative PCR (qPCR)

For analysis of tissue expression patterns of IR8a and the five AseglRs from the IR75p and IR75q lineages, tissues

were dissected from 3–5-day-old virgin moths, including chemosensory organs (antennae and proboscises) of both sexes, female ovipositors, as well as male abdomens, heads+thoraxes (without antennae), legs, and wings. Total RNA of these tissues was extracted using the RNeasy Mini kit or RNeasy Plus Micro kit (Qiagen, GmbH, Hilden, Germany) depending on the weight of starting material and treated with DNase. First-strand cDNAs were synthesized from 250 ng total RNA with the SuperScript™ IV First-Strand Synthesis System (Invitrogen, Carlsbad, CA, USA) following the manufacturer's instruction. Gene expression patterns were examined by qPCR on a Stratagene Mx3005P Real-Time PCR System (Agilent technologies), with GADPH and RPS3 as reference genes. Primers used for qPCR are shown in Additional file 1: Table S2 and the primer efficiencies were validated by standard curves with 10X serial dilutions of the templates. Three biological replicates were performed in separate plates, with three technical replicates on each plate. The reaction was carried out in a 20 µL system containing 1 µL cDNA template, 10 µL Power SYBR Green PCR 2X Master Mix (Life Technologies), and 0.3 µM of each primer. The thermal cycling parameters consisted of an initial denaturation step at 95 °C for 10 min, followed by 45 cycles of 95 °C for 15 s and 60 °C for 1 min. No-template controls were run in parallel for each primer pair. A subsequent melting curve analysis was performed to ensure the primer specificity. The relative expression level of each gene to the average of reference genes was calculated by the comparative CT method [75]. The relative expression levels were log₁₀ transformed prior to statistical analysis.

Scanning electron microscopy (SEM)

The antennae from 2–3-day-old adults were dissected and airdried. The dried samples were carefully mounted on SEM stubs and then sputter-coated with gold (Cesington 108 auto, 45 s, 20 mA). The preparations were examined by a scanning electron microscope (SEM; Hitachi SU3500) at 5 kV in the Department of Biology, Lund University.

Whole-mount fluorescence in situ hybridization (WM-FISH)

Whole-mount fluorescence in situ hybridization was performed to monitor the antennal expression pattern of the IR8a and genes in IR75p/q expansions on both female and male *A. segetum* antennae [76, 77]. Digoxigenin (DIG)- and biotin-labelled antisense probes of target IRs were transcribed from linearized pCS2+ vectors with the coding region of the corresponding receptors using the T7 RNA transcription kit (Roche, Mannheim, Germany). Labelled probes were subsequently fragmented to an average length of about 800 bp by incubation in sodium

carbonate buffer (80 mM NaHCO₃, 120 mM Na₂CO₃, pH 10.2). DIG-labelled probes were visualized using anti-DIG AP-conjugated antibodies and the Vector Red Alkaline Phosphatase (AP) Substrate Kit (Vector Laboratories, Burlingame, CA, USA); biotin-labelled probes were visualized by streptavidin-horseradish peroxidase (HRP) conjugates with Cyanine 5 Plus Amplification Reagent in a TSA kit as the substrate (PerkinElmer, Boston, MA, USA). Sense probes labelled with DIG and biotin by SP6 RNA transcription kit were used as control. Antennae were dissected from the head and cut into pieces of about 4–5 mm length and then transferred to fixation solution (4% paraformaldehyde in 0.1 M Na₂CO₃, pH 9.5, 0.03% Triton X-100). The antennae were gently squeezed by fine tweezers to improve tissue penetration. The following fixation, hybridization, blocking, antibody, and substrate incubation steps were performed according to previously described protocols [76]. Finally, the tissues were mounted in Mowiol mounting medium. A confocal microscope (Leica SP8 DLS, Germany) was used to analyze the sections [77].

Functional characterization in *Xenopus* oocytes

First-strand cDNA of female antennae was synthesized from 500 ng total RNA using SuperScript™ IV First-Strand Synthesis System (Invitrogen, Carlsbad, CA, USA) following the manufacturer's instructions. Primers containing Kozak sequence ("GCCACC") and restriction sites (Additional file 1: Table S3) were designed to amplify the full-length IR genes using Platinum Pfu Polymerase (Thermo Fisher Scientific). The amplified genes were then sub-cloned into the pCS2+ expression vectors. All DNA sequences were verified by Sanger sequencing, using a capillary 3130xL Genetic Analyzer (Thermo Fisher Scientific, Waltham, MA, USA) at the sequencing facility at the Department of Biology, Lund University. The sequences of the co-receptor AsegIR8a and genes in the IR75p/q clade that were cloned and functionally tested (AsegIR75p, AsegIR75p.1, AsegIR75p.2, AsegIR75q.1, and AsegIR75q.2) have been deposited in GenBank under the accession numbers MW659848–MW659853. Large quantities of purified plasmids containing AsegIRs and AsegIR8a with verified sequences were obtained using the PureLink™ HiPure Plasmid Filter Midiprep Kit (Thermo Fisher Scientific). cRNAs of AsegIRs and AsegIR8a were synthesized from NotI (Promega) linearized recombinant pCS2+ plasmids using the mMESSAGE mMACHINE SP6 transcription kit (Thermo Fisher Scientific).

Each of the five AsegIRs from the IR75p and IR75q lineages were co-expressed with the putative co-receptor AsegIR8a in *Xenopus* oocytes, and two-electrode voltage clamp recordings were performed following

previously described protocols [78–80]. In brief, oocytes were collected from ovarian tissue of female *X. laevis* (frogs purchased from European *Xenopus* Resource Centre, University of Portsmouth, UK) and treated with 1.5 mg/mL collagenase (Sigma-Aldrich Co., St. Louis, MO, USA) in Ca^{2+} -free Oocyte Ringer 2 buffer (containing 82.5 mM NaCl, 2 mM KCl, 1 mM MgCl_2 , 5 mM HEPES, pH 7.5) at room temperature for 15–18 min. Mature healthy oocytes (stage V–VII) were isolated and co-injected with 50 ng IR75 cRNA and 50 ng IR8a cRNA and incubated for 4–7 days at 18 °C in Ringer's buffer containing sodium pyruvate (550 mg/L) and gentamicin (100 mg/L). Two-electrode voltage clamp recordings were performed to measure the ligand-induced whole-cell inward currents from oocytes in good condition at a holding potential of –80 mV, with a TEC-03BF signal amplifier. The test compounds and Ringer's buffer were applied to the oocytes through a computer-controlled perfusion system with exposure to stimuli at a rate of 2 mL/min for 20 s and extensive washing in Ringer's buffer at a rate of 4 mL/min between stimulations. Data were collected and analyzed by Cellworks software (npi electronic GmbH, Tamm, Germany).

GC-EAD

The antennal electrophysiological responses of male and female *A. segetum* to the synthetic mixture of C_6 – C_{10} straight chain acids were recorded on an Agilent 7890 gas chromatograph equipped with a flame ionization detector (FID) (Agilent Technologies, USA) and an electroantennographic detector (EAD) (Syntech, Germany). An Agilent J&W HP-5 column (30 m \times 0.25 mm i.d., 0.25 μm film thickness; Agilent Technologies, USA) was used in the GC, where the inlet temperature was set at 250 °C, the transfer line was heated at 255 °C, and the detector was set at 280 °C. Hydrogen was used as the carrier gas at a constant flow of 1.8 mL/min, and a 1:1 division of the GC effluent was directed to the FID and EAD, respectively. A PRG-2 EAG (10x gain) probe (Syntech, Germany) was used in the recording. After cutting off the tips, both antennae associated with the head of a 1–2-day-old adult were mounted on the probe using conductive gel (Blågel, CefarCompex, Sweden). Charcoal-filtered and humidified air passed over the antennal preparation. The GC oven was programmed from initial hold at 100 °C for 2 min, then increased to 160 °C at a rate of 10 °C/min and then held for 2 min at the final temperature. For each recording, a synthetic mixture containing 500 ng of each acid was injected in split mode (ratio of 40:1). Data were collected with the software GC-EAD Pro Version 4.1 (Syntech, Germany).

Two-choice Y-tube experiments

A Y-tube olfactometer was used to investigate the behavioral responses to acids. The olfactometer consisted of a 14 cm main arm with an insect introduction chamber attached and two 14-cm-long arms with an interior angle of 120° for the treatment and control odors. The inner diameter of the Y-tube was 1.6 cm. The tests were always conducted in a dark room under red light illuminating the olfactometer from above (~26 Lux). Tests were performed during the first 2 h of the scotophase at the following experimental conditions: 24 °C air temperature, 55% relative humidity, and 0.8 m/s airflow.

To test whether the honey solution is attractive to the adult moths, two cotton balls (loaded 1 mL 10% honey solution with 10 μL acetone vs 1 mL water with 10 μL acetone) were placed at inside of each Y-arm. Individual adults of *A. segetum* were released from the induce chamber at the end of main arm (29 females and 22 males). Another control experiment (cotton balls loaded 1 mL 10% honey solution with 10 μL acetone were placed at both sides of Y-tube) was performed to check whether there is a side-bias.

In this bioassay, five fatty acids were tested individually at the dose of 100 μg , including hexanoic acid, heptanoic acid, octanoic acid, nonanoic acid, and decanoic acid ($N \geq 26$ for each acid and each sex). Stock solutions of these fatty acids were prepared by diluting the neat compounds to 10 mg/mL in acetone. Before each experiment, 10 μL stock solution was mixed in 1 mL 10% honey solution, then the mixture was loaded on a cotton ball. Then, 10% honey solution with 10 μL acetone was used as control. The cotton balls were left for 10 min for the evaporation of acetone before tests and placed at inside of the ends of the Y-arms. The Y-tube and cotton balls were renewed after every 10 tests. Then, 16–20 h starved naïve individual males/females, 2–5 days old, were used in the experiment. After the screening-assay, octanoic acid was tested at a series of doses in 10-fold increments from 100 ng to 100 μg . Insects were considered to have made a choice only when they entered either of the arms and fed on the cotton within 5 min. By contrast, during the observation period (5 min), if the insect kept moving back and forth or stayed in the main arm without movement, it was considered as no choice had been made. The proportions of no choice for each treatment and each sex were lower than 5%, so we excluded the no choice data when we did the chi-square test for statistical significance.

Statistical analysis

All analyses were performed using IBM SPSS Statistics 28.0. The normality of the datasets was determined by the Shapiro-Wilk test and the homogeneity of variances

was analyzed by Levene's test. Independent samples *t*-tests were used to test for differences between two groups, i.e., the differences between males and females in the gene expression levels and in GC-EAD responses to specific acids, as well as the response of AseglR75q.1 in the oocytes to the two primary ligands at given doses. One-way ANOVA followed by an LSD test was used to compare the differences among multiple groups, including the expression levels of different genes in the antennae and of one gene among different tissues, the response of AseglR75p.1 in the oocytes to the four primary ligands at given doses, and the GC-EAD responses of the antennae to different acids. In the behavioral experiments, chi-square tests were used to assess the differences between the expected frequencies (50%) and the observed frequencies of responding individuals.

Abbreviations

A-IRs: Antennal IRs; AMPA: α -Amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid; AP: Alkaline phosphatase; DIG: Digoxigenin; D-IRs: Divergent IRs; DMSO: Dimethyl sulfoxide; FID: Flame ionization detector; FISH: Fluorescence in situ hybridization; GADPH: Glyceraldehyde 3-phosphate dehydrogenase; GC-EAD: Gas chromatography coupled to electroantennographic detection; HRP: Streptavidin-horseradish peroxidase; iGluR: Ionotropic glutamate receptor; IRs: Ionotropic receptors; LS-IRs: Lepidoptera-specific IRs; NMDA: N-methyl-D-aspartate; Orco: Odorant receptor co-receptor; ORs: Odorant receptors; OSNs: Olfactory sensory neurons; RNA-Seq: mRNA sequencing; RPS3: Ribosomal protein S3; SEM: Scanning electron microscopy; TPM: Transcripts per million; WM-FISH: Whole-mount fluorescence in situ hybridization.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12915-022-01235-0>.

Additional file 1: Table S1. Compounds used for characterization of *Agrotis segetum* ionotropic receptors (IRs), including their purities and source information. **Table S2.** Primers used in qPCR. **Table S3.** Primers used in gene cloning. **Fig. S1.** Additional representative FISH images.

Additional file 2. Amino acid sequences of AseglRs.

Additional file 3. Individual data values for Fig. 6 B and C.

Acknowledgements

We are grateful to Ola Gustafsson at the Microscopy Facility, Department of Biology, Lund University for access to the scanning electron microscope and confocal microscope. We thank Prof. Staffan Bensch's lab for providing the qPCR facility and Erling V. Jirle for his help with the technical assistance and maintenance of the insect culture.

Authors' contributions

XQH, DDZ, and CL conceived the study. XQH, DDZ, MNA, DP, and CL contributed to experimental design. XQH and DP performed transcriptomic data analysis. XQH performed the molecular work, oocyte recordings, scanning electron microscopy imaging, phylogenetic analysis, and behavioral assay. DDZ performed qPCR experiment. DDZ and XQH did in situ hybridization. HLW performed the GC-EAD. XQH and DDZ analyzed the data. XQH drafted the manuscript with contributions from HLW and DP. DDZ revised the manuscript according to reviewers' comments. All authors read and approved the final manuscript.

Funding

Open access funding provided by Lund University. We thank the Chinese Scholarship Council (CSC) for financial support of Xiao-Qing Hou's PhD study. This work was supported by grants from the Swedish Research Council (#VR-621-2013-4355 and VR-2017-03804 to CL), the Swedish Foundation for International Cooperation in Research and Higher Education STINT (#201315483 and 201315256 to CL), and FORMAS (grant #2018-01444 to MNA) and resources provided by the Swedish National Infrastructure for Computing (SNIC) through Uppsala Multidisciplinary Center for Advanced Computational Science (UPPMAX) under Project SNIC 2019/8-83.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request. The sequence reads have been deposited in the SRA database at NCBI (BioProject accession number PRJNA707654). The amino acid sequences of the IRs identified from *Agrotis segetum* antennal and ovipositor transcriptomes are available in Additional file 2; sequences of the co-receptor AseglR8a and IRs in the 75p/q clade (AseglR75p, AseglR75p.1, AseglR75p.2, AseglR75q.1, and AseglR75q.2) have been deposited in GenBank (accession numbers MW659848-MW659853).

Declarations

Ethics approval and consent to participate

The care and use of *Xenopus laevis* frogs in this study were approved by the Swedish Board of Agriculture, and the methods were carried out in accordance with the approved guidelines.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Department of Biology, Lund University, Sölvegatan 37, 223 62 Lund, Sweden.

²Present address: Agricultural Genomics Institute at Shenzhen, Chinese Academy of Agricultural Sciences, Shenzhen, China. ³Present address: Global Change Ecology Research Group, School of Science and Engineering, University of the Sunshine Coast, QLD, Sunshine Coast, Australia.

Received: 22 March 2021 Accepted: 19 January 2022

Published online: 07 February 2022

References

- Sachse S, Krieger J. Olfaction in insects. The primary processes of odor recognition and coding. *e-Neuroforum*. 2011;2(3):49–60. <https://doi.org/10.1007/s13295-011-0020-7>.
- Leal WS. Odorant reception in insects: roles of receptors, binding proteins, and degrading enzymes. *Annu. Rev. Entomol.* 2013;58:373–91. <https://doi.org/10.1146/annurev-ento-120811-153635>.
- Hansson BS, Stensmyr MC. Evolution of insect olfaction. *Neuron*. 2011;72(5):698–711. <https://doi.org/10.1016/j.neuron.2011.11.003>.
- Fleischer J, Pregitzer P, Breer H, Krieger J. Access to the odor world: olfactory receptors and their role for signal transduction in insects. *Cell. Mol. Life Sci.* 2018;75(3):485–508. <https://doi.org/10.1007/s00018-017-2627-5>.
- Clyne PJ, Warr CG, Freeman MR, Lessing D, Kim JH, Carlson JR. A novel family of divergent seven-transmembrane proteins: candidate odorant receptors in *Drosophila*. *Neuron*. 1999;22(2):327–38. [https://doi.org/10.1016/s0896-6273\(00\)81093-4](https://doi.org/10.1016/s0896-6273(00)81093-4).
- Benton R, Vannice KS, Gomez-Diaz C, Vossell LB. Variant ionotropic glutamate receptors as chemosensory receptors in *Drosophila*. *Cell*. 2009;136(1):149–62. <https://doi.org/10.1016/j.cell.2008.12.001>.
- Yao CA, Ignell R, Carlson JR. Chemosensory coding by neurons in the coeloconic sensilla of the *Drosophila* antenna. *J. Neurosci.* 2005;25(37):8359–67. <https://doi.org/10.1523/JNEUROSCI.2432-05.2005>.

8. Abuin L, Bargeton B, Ulbrich MH, Isacoff EY, Kellenberger S, Benton R. Functional architecture of olfactory ionotropic glutamate receptors. *Neuron*. 2011;69(1):44–60. <https://doi.org/10.1016/j.neuron.2010.11.042>.
9. Zhang J, Bisch-Knaden S, Fandino RA, Yan S, Obiero GF, Grosse-Wilde E, et al. The olfactory coreceptor IR8a governs larval feces-mediated competition avoidance in a hawkmoth. *Proc. Natl. Acad. Sci. U.S.A.* 2019;116(43):21828–33. <https://doi.org/10.1073/pnas.1913485116>.
10. Koh TW, He Z, Gorur-Shandilya S, Menzies K, Larter NK, Stewart S, et al. The *Drosophila* IR20a clade of ionotropic receptors are candidate taste and pheromone receptors. *Neuron*. 2014;83(4):850–65. <https://doi.org/10.1016/j.neuron.2014.07.012>.
11. Stewart S, Koh TW, Ghosh AC, Carlson JR. Candidate ionotropic taste receptors in the *Drosophila* larva. *Proc. Natl. Acad. Sci. U.S.A.* 2015;112(14):4195–201. <https://doi.org/10.1073/pnas.1503292112>.
12. Ahn JE, Chen Y, Amrein H. Molecular basis of fatty acid taste in *Drosophila*. *eLife*. 2017;6:e30115. <https://doi.org/10.7554/eLife.30115>.
13. Chen Y, Amrein H. Ionotropic receptors mediate *Drosophila* oviposition preference through sour gustatory receptor neurons. *Curr. Biol.* 2017;27(18):2741–50. <https://doi.org/10.1016/j.cub.2017.08.003>.
14. Chen C, Buhl E, Xu M, Croset V, Rees JS, Lilley KS, et al. *Drosophila* ionotropic receptor 25a mediates circadian clock resetting by temperature. *Nature*. 2015;527(7579):516–20. <https://doi.org/10.1038/nature16148>.
15. Ni L, Klein M, Svec KV, Budelli G, Chang EC, Ferrer AJ, et al. The ionotropic receptors IR21a and IR25a mediate cool sensing in *Drosophila*. *eLife*. 2016;5:e13254. <https://doi.org/10.7554/eLife.13254>.
16. Enjin A, Zaharieva EE, Frank DD, Mansourian S, Suh GS, Gallio M, et al. Humidity sensing in *Drosophila*. *Curr. Biol.* 2016;26(10):1352–8. <https://doi.org/10.1016/j.cub.2016.03.049>.
17. Knecht ZA, Silbering AF, Ni L, Klein M, Budelli G, Bell R, et al. Distinct combinations of variant ionotropic glutamate receptors mediate thermosensation and hygro-sensation in *Drosophila*. *eLife*. 2016;5:e17879. <https://doi.org/10.7554/eLife.17879>.
18. Knecht ZA, Silbering AF, Cruz J, Yang L, Croset V, Benton R, et al. Ionotropic receptor-dependent moist and dry cells control hygro-sensation in *Drosophila*. *eLife*. 2017;6:e26654. <https://doi.org/10.7554/eLife.26654>.
19. Senthilan PR, Piepenbrock D, Ovezmyradov G, Nadrowski B, Bechstedt S, Pauls S, et al. *Drosophila* auditory organ genes and genetic hearing defects. *Cell*. 2012;150(5):1042–54. <https://doi.org/10.1016/j.cell.2012.06.043>.
20. Larsson MC, Domingos AI, Jones WD, Chiappe ME, Amrein H, Vosshall LB. Or83b encodes a broadly expressed odorant receptor essential for *Drosophila* olfaction. *Neuron*. 2004;43(5):703–14. <https://doi.org/10.1016/j.neuron.2004.08.019>.
21. Sato K, Pellegrino M, Nakagawa T, Nakagawa T, Vosshall LB, Touhara K. Insect olfactory receptors are heteromeric ligand-gated ion channels. *Nature*. 2008;452(7190):1007–11. <https://doi.org/10.1038/nature06850>.
22. Wicher D, Schäfer R, Bauernfeind R, Stensmyr MC, Heller R, Heinemann SH, et al. *Drosophila* odorant receptors are both ligand-gated and cyclic-nucleotide-activated cation channels. *Nature*. 2008;452(7190):1007–11. <https://doi.org/10.1038/nature06861>.
23. Grosjean Y, Rytz R, Farine JP, Abuin L, Cortot J, Jefferis GS, et al. An olfactory receptor for food-derived odors promotes male courtship in *Drosophila*. *Nature*. 2011;478(7368):236–40. <https://doi.org/10.1038/nature10428>.
24. Silbering AF, Rytz R, Grosjean Y, Abuin L, Ramdya P, Jefferis GS, et al. Complementary function and integrated wiring of the evolutionarily distinct *Drosophila* olfactory subsystems. *J. Neurosci.* 2011;31(38):13357–75. <https://doi.org/10.1523/JNEUROSCI.2360-11.2011>.
25. Rytz R, Croset V, Benton R. Ionotropic receptors (IRs): chemosensory ionotropic glutamate receptors in *Drosophila* and beyond. *Insect Biochem. Mol. Biol.* 2013;43(9):888–97. <https://doi.org/10.1016/j.ibmb.2013.02.007>.
26. Ai M, Blais S, Park JY, Min S, Neubert TA, Suh GS. Ionotropic glutamate receptors IR64a and IR8a form a functional odorant receptor complex *in vivo* in *Drosophila*. *J. Neurosci.* 2013;33(26):10741–9. <https://doi.org/10.1523/JNEUROSCI.5419-12.2013>.
27. Hussain A, Zhang M, Üçpınar HK, Svensson T, Quillery E, Gompel N, et al. Ionotropic chemosensory receptors mediate the taste and smell of polyamines. *PLoS Biol.* 2018;14(5):e1002454. <https://doi.org/10.1371/journal.pbio.1002454>.
28. Ganguly A, Pang L, Duong VK, Lee A, Schoniger H, Varady E, et al. A molecular and cellular context-dependent role for Ir76b in detection of amino acid taste. *Cell Rep.* 2017;18(3):737–50. <https://doi.org/10.1016/j.celrep.2016.12.071>.
29. Croset V, Rytz R, Cummins SF, Budd A, Brawand D, Kaessmann H, et al. Ancient protostome origin of chemosensory ionotropic glutamate receptors and the evolution of insect taste and olfaction. *PLoS Genet.* 2010;6(8):e1001064. <https://doi.org/10.1371/journal.pgen.1001064>.
30. Andersson MN, Keeling CI, Mitchell RF. Genomic content of chemosensory genes correlates with host range in wood-boring beetles (*Dendroctonus ponderosae*, *Agrilus planipennis*, and *Anoplophora glabripennis*). *BMC Genom.* 2019;20(1):690. <https://doi.org/10.1186/s12864-019-6054-x>.
31. He Z, Luo Y, Shang X, Sun JS, Carlson JR. Chemosensory sensilla of the *Drosophila* wing express a candidate ionotropic pheromone receptor. *PLoS Biol.* 2019;17(5):e2006619. <https://doi.org/10.1371/journal.pbio.2006619>.
32. Olivier V, Monsempes C, Francois MC, Poivet E, Jacquin-Joly E. Candidate chemosensory ionotropic receptors in a Lepidoptera. *Insect Mol. Biol.* 2011;20(2):189–99. <https://doi.org/10.1111/j.1365-2583.2010.01057.x>.
33. Bengtsson JM, Trona F, Montagné N, Anfora G, Ignell R, Witzgall P, et al. Putative chemosensory receptors of the codling moth, *Cydia pomonella*, identified by antennal transcriptome analysis. *PLoS One*. 2012;7(2):e31620. <https://doi.org/10.1371/journal.pone.0031620>.
34. Poivet E, Gallot A, Montagné N, Glaser N, Legeai F, Jacquin-Joly E. A comparison of the olfactory gene repertoires of adults and larvae in the noctuid moth *Spodoptera littoralis*. *PLoS One*. 2013;8(4):e60263. <https://doi.org/10.1371/journal.pone.0060263>.
35. van Schooten B, Jiggins CD, Briscoe AD, Papa R. Genome-wide analysis of ionotropic receptors provides insight into their evolution in Heliconius butterflies. *BMC Genom.* 2016;17(1):254. <https://doi.org/10.1186/s12864-016-2572-y>.
36. Liu NY, Xu W, Dong SL, Zhu JY, Xu YX, Anderson A. Genome-wide analysis of ionotropic receptor gene repertoire in Lepidoptera with an emphasis on its functions of *Helicoverpa armigera*. *Insect Biochem. Mol. Biol.* 2018;99:37–53. <https://doi.org/10.1016/j.ibmb.2018.05.005>.
37. Zhu JY, Xu ZW, Zhang XM, Liu NY. Genome-based identification and analysis of ionotropic receptors in *Spodoptera litura*. *Sci. Nat.* 2018;105(5):38. <https://doi.org/10.1007/s00114-018-1563-z>.
38. Yuvaraj JK, Andersson MN, Zhang DD, Löfstedt C. Antennal transcriptome analysis of the chemosensory gene families from Trichoptera and basal Lepidoptera. *Front. Physiol.* 2018;9:1365. <https://doi.org/10.3389/fphys.2018.01365>.
39. Yin NN, Nuo SM, Xiao HY, Zhao YJ, Zhu JY, Liu NY. The ionotropic receptor gene family in Lepidoptera and Trichoptera: annotation, evolutionary and functional perspectives. *Genomics*. 2021;113(1):601–12. <https://doi.org/10.1016/j.jygeno.2020.09.056>.
40. Eyun SI, Soh HY, Posavi M, Munro JB, Hughes DS, Murali SC, et al. Evolutionary history of chemosensory-related gene families across the Arthropoda. *Mol. Biol. Evol.* 2017;34(8):1838–62. <https://doi.org/10.1093/molbev/msx147>.
41. Shan S, Wang SN, Song X, Khashaveha A, Lu ZY, Dhiloof KH, et al. Antennal ionotropic receptors IR64a1 and IR64a2 of the parasitoid wasp *Microplitis mediator* (Hymenoptera: Braconidae) collaboratively perceive habitat and host cues. *Insect Biochem. Mol. Biol.* 2019;114:103204. <https://doi.org/10.1016/j.ibmb.2019.103204>.
42. Tang R, Jiang NJ, Ning C, Li GC, Huang LQ, Wang CZ. The olfactory receptor of acetic acid and ionotropic receptors in the Oriental armyworm, *Mythimna separata* Walker. *Insect Biochem. Mol. Biol.* 2020;118:103312. <https://doi.org/10.1016/j.ibmb.2019.103312>.
43. Klinner CF, König MC, Werckenthin A, Daly KC, Bisch-Knaden S, et al. Functional olfactory sensory neurons housed in olfactory sensilla on the ovipositor of the hawkmoth *Manduca sexta*. *Front. Ecol. Evol.* 2016;4:130. <https://doi.org/10.3389/fevo.2016.00130>.
44. Hou X-Q, Zhang D-D, Powell D, Wang H-L, Andersson MN, Löfstedt C. Transcriptome sequencing of the turnip moth, *Agrotis segetum*. NCBI GenBank <https://www.ncbi.nlm.nih.gov/bioproject/PRJNA707654>. 2021.
45. Liu C, Pitts RJ, Bohbot JD, Jones PL, Wang G, Zwiebel LJ. Distinct olfactory signaling mechanisms in the malaria vector mosquito *Anopheles gambiae*. *PLoS Biol.* 2010;8(8):e1000467. <https://doi.org/10.1371/journal.pbio.1000467>.
46. Li B, Dewey CN. RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. *BMC Bioinform.* 2011;12:323. <https://doi.org/10.1186/1471-2105-12-323>.

47. Liu Y, Gu S, Zhang Y, Guo Y, Wang G. Candidate olfaction genes identified within the *Helicoverpa armigera* antennal transcriptome. *PLoS One*. 2012;7(10):e48260. <https://doi.org/10.1371/journal.pone.0048260>.
48. Menuz K, Larter NK, Park J, Carlson JR. An RNA-seq screen of the *Drosophila* antenna identifies a transporter necessary for ammonia detection. *PLoS Genet*. 2014;10(11):e1004810. <https://doi.org/10.1371/journal.pgen.1004810>.
49. Pitts RJ, Derryberry SL, Zhang ZW, Zwiebel LJ. Variant ionotropic receptors in the malaria vector mosquito *Anopheles gambiae* tuned to amines and carboxylic acids. *Sci. Rep*. 2017;7(1):40297. <https://doi.org/10.1038/srep40297>.
50. Mullens BA, Reifenrath WG, Butler SM. Laboratory trials of fatty acids as repellents or antifeedants against houseflies, horn flies and stable flies (Diptera: Muscidae). *Pest. Manage. Sci*. 2009;65(12):1360–6. <https://doi.org/10.1002/ps.1823>.
51. Venter GJ, Labuschagne K, Boikanyo SN, Morey L, Snyman MG. The repellent effect of organic fatty acids on Culicoides midges as determined with suction light traps in South Africa. *Vet. Parasitol*. 2011;181(2–4):365–9. <https://doi.org/10.1016/j.vetpar.2011.04.034>.
52. Legal L, Plawewski M. Comparative sensitivity of various insects to toxic compounds from *Morinda citrifolia* (L.). *Entomol. Probl*. 1995;26(2):155–9.
53. Legal L, Moulin B, Jallon JM. The relation between structures and toxicity of oxygenated aliphatic compounds homologous to the insecticide octanoic acid and the chemotaxis of two species of *Drosophila*. *Pestic Biochem Physiol*. 1999;65(2):90–101. <https://doi.org/10.1006/pest.1999.2430>.
54. Zhu JJ, Cermak SC, Kenar JA, Brewer G, Haynes KF, Boxler D, et al. Better than DEET repellent compounds derived from coconut oil. *Sci. rep*. 2018;8(1):1–12. <https://doi.org/10.1038/s41598-018-32373-7>.
55. Ramadan GRM, Abdelgaleil SAM, Shawir MS, El-bakary AS, Zhu KY, Phillips TW. Terpenoids, DEET and short chain fatty acids as toxicants and repellents for *Rhyzopertha dominica* (coleoptera: Bostrichidae) and *Lasioderma serricorne* (Coleoptera: Ptinidae). *J. Stored Prod. Res*. 2020;87:101610. <https://doi.org/10.1016/j.jspr.2020.101610>.
56. Matsuo T, Sugaya S, Yasukawa J, Aigaki T, Fuyama Y. Odorant-binding proteins OBP57d and OBP57e affect taste perception and host-plant preference in *Drosophila sechellia*. *PLoS Biol*. 2007;5(5):e118. <https://doi.org/10.1371/journal.pbio.0050118>.
57. Andrade López JM, Lanno SM, Auerbach JM, Moskowicz EC, Sligar LA, Wittkopp PJ, et al. Genetic basis of octanoic acid resistance in *Drosophila sechellia*: functional analysis of a fine-mapped region. *Mol. Ecol*. 2017;26(4):1148–60. <https://doi.org/10.1111/mec.14001>.
58. Skinner WA, Tong HC, Maibach HI, Skidmore D. Human skin surface lipid fatty acids – mosquito repellents. *Experientia*. 1970;26(7):728–30. <https://doi.org/10.1007/BF02232510>.
59. Hwang YS, Schultz GW, Axelrod H, Kramer WL, Mulla MS. Ovipositional repellency of fatty acids and their derivatives against *Culex* and *Aedes* mosquitoes. *Ecol. Entomol*. 1982;11(1):223–6. <https://doi.org/10.1093/ee/11.1.223>.
60. Guo M, Krieger J, Große-Wilde E, Mißbach C, Zhang L, Breer H. Variant ionotropic receptors are expressed in olfactory sensory neurons of coelomic sensilla on the antenna of the desert locust (*Schistocerca gregaria*). *Int. J. Biol. Sci*. 2014;10(1):1–14. <https://doi.org/10.7150/ijbs.7624>.
61. Qiu YT, Van Loon JJ, Takken W, Meijerink J, Smid HM. Olfactory coding in antennal neurons of the malaria mosquito, *Anopheles gambiae*. *Chem. Senses*. 2006;31(9):845–63. <https://doi.org/10.1093/chemse/bjl027>.
62. Missbach C, Dweck HKM, Vogel H, Vilcinskis A, Stensmyr MC, Hansson BS, et al. Evolution of insect olfactory receptors. *eLife*. 2014;3:e02115. <https://doi.org/10.7554/eLife.02115>.
63. Pophof B. Olfactory responses from sensilla coeloconica of the silkworm *Bombyx mori*. *Physiol. Entomol*. 1997;22(3):239–48. <https://doi.org/10.1111/j.1365-3032.1997.tb01164.x>.
64. Crow KD, Wagner GP. What is the role of genome duplication in the evolution of complexity and diversity? *Mol. Biol. Evol*. 2006;23(5):887–92. <https://doi.org/10.1093/molbev/msj083>.
65. Moleirinho A, Carneiro J, Matthiesen R, Silva RM, Amorim A, Azevedo L. Gains, losses and changes of function after gene duplication: study of the metallothionein family. *PLoS One*. 2011;6(4):e18487. <https://doi.org/10.1371/journal.pone.0018487>.
66. Andersson MN, Löfstedt C, Newcomb RD. Insect olfaction and the evolution of receptor tuning. *Front. Ecol. Evol*. 2015;3:53. <https://doi.org/10.3389/fevo.2015.00053>.
67. Benton R. Multigene family evolution: perspectives from insect chemoreceptors. *Trends Ecol. Evol*. 2015;30(10):590–600. <https://doi.org/10.1016/j.tree.2015.07.009>.
68. Force A, Lynch M, Pickett FB, Amores A, Yan YL, Postlethwait J. Preservation of duplicate genes by complementary, degenerative mutations. *Genetics*. 1999;151(4):1531–45. <https://doi.org/10.1093/genetics/151.4.1531>.
69. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics*. 2014;30(15):2114–20. <https://doi.org/10.1093/bioinformatics/btu170>.
70. Li W, Godzik A. Cd-hit: a fast program for clustering and comparing large sets of protein or nucleotide sequences. *Bioinformatics*. 2006;22(13):1658–9. <https://doi.org/10.1093/bioinformatics/btl158>.
71. Finn RD, Bateman A, Clements J, Coggill P, Eberhardt RY, Eddy SR, et al. Pfam: the protein families database. *Nucleic Acids Res*. 2014;42:D222–30. <https://doi.org/10.1093/nar/gkt1223>.
72. Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics*. 2015;31(19):3210–2. <https://doi.org/10.1093/bioinformatics/btv351>.
73. Huber W, Carey VJ, Gentleman R, Anders S, Carlson M, Carvalho BS, et al. Orchestrating high-throughput genomic analysis with Bioconductor. *Nat. Methods*. 2015;12(2):115–21. <https://doi.org/10.1038/nmeth.3252>.
74. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol*. 2014;15(12):550. <https://doi.org/10.1186/s13059-014-0550-8>.
75. Schmittgen TD, Livak KJ. Analyzing real-time PCR data by the comparative CT method. *Nat. Protoc*. 2008;3(6):1101–8. <https://doi.org/10.1038/nprot.2008.73>.
76. Karner T, Kellner I, Schultze A, Berre H, Krieger J. Co-expression of six tightly clustered odorant receptor genes in the antenna of the malaria mosquito *Anopheles gambiae*. *Front. Ecol. Evol*. 2015;3:26. <https://doi.org/10.3389/fevo.2015.00026>.
77. Zhang DD, Wang HL, Schultze A, Froß H, Francke W, Krieger J, et al. Receptor for detection of a type II sex pheromone in the winter moth *Operophtera brumata*. *Sci. Rep*. 2016;6(1):18576. <https://doi.org/10.1038/srep18576>.
78. Zhang DD, Zhu KY, Wang CZ. Sequencing and characterization of six cDNAs putatively encoding three pairs of pheromone receptors in two sibling species, *Helicoverpa armigera* and *Helicoverpa assulta*. *J. Insect Physiol*. 2010;56(6):586–93. <https://doi.org/10.1016/j.jinsphys.2009.12.002>.
79. Zhang DD, Löfstedt C. Functional evolution of a multigene family: orthologous and paralogous pheromone receptor genes in the turnip moth, *Agrotis segetum*. *PLoS One*. 2013;8(10):e77345. <https://doi.org/10.1371/journal.pone.0077345>.
80. Hou X, Zhang DD, Yuvaraj JK, Corcoran JA, Andersson MN, Löfstedt C. Functional characterization of odorant receptors from the moth *Eriocrania semipurpurella*: a comparison of results in the *Xenopus* oocyte and HEK cell systems. *Insect Biochem. Mol. Biol*. 2020;117:103289. <https://doi.org/10.1016/j.ibmb.2019.103289>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

