



# LUND UNIVERSITY

## Identification of two abundant *Aerococcus urinae* cell wall-anchored proteins

Senneby, Erik; Sunnerhagen, Torgny; Hallström, Björn; Lood, Rolf; Malmström, Johan; Karlsson, Christofer; Rasmussen, Magnus

*Published in:*  
International Journal of Medical Microbiology

*DOI:*  
[10.1016/j.ijmm.2019.06.005](https://doi.org/10.1016/j.ijmm.2019.06.005)

2019

*Document Version:*  
Peer reviewed version (aka post-print)

[Link to publication](#)

*Citation for published version (APA):*  
Senneby, E., Sunnerhagen, T., Hallström, B., Lood, R., Malmström, J., Karlsson, C., & Rasmussen, M. (2019). Identification of two abundant *Aerococcus urinae* cell wall-anchored proteins. *International Journal of Medical Microbiology*, 309(7), Article 151325. <https://doi.org/10.1016/j.ijmm.2019.06.005>

*Total number of authors:*  
7

*Creative Commons License:*  
CC BY-NC-ND

### General rights

Unless other specific re-use rights are stated the following general rights apply:  
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117  
221 00 Lund  
+46 46-222 00 00

**Identification of two abundant *Aerococcus urinae* cell wall-  
anchored proteins.**

Erik Senneby<sup>a\*</sup>, Torgny Sunnerhagen<sup>a</sup>, Björn Hallström<sup>b</sup>, Rolf Lood<sup>a</sup>, Johan  
Malmström<sup>a</sup>, Christofer Karlsson<sup>a</sup> and Magnus Rasmussen<sup>a</sup>.

<sup>a</sup>Division of Infection Medicine, Department of Clinical Sciences, BMC B14, 221 85,  
Lund University, Lund, Sweden.

<sup>b</sup>Centre for Translational Genomics, Division of Clinical Genetics, BMC B10, 221 85,  
Lund University, Lund, Sweden.

\*Corresponding author: [erik.senneby@med.lu.se](mailto:erik.senneby@med.lu.se)

E-mail addresses: [torgny.sunnerhagen@med.lu.se](mailto:torgny.sunnerhagen@med.lu.se), [bjorn.hallstrom@skane.se](mailto:bjorn.hallstrom@skane.se),  
[rolf.lood@med.lu.se](mailto:rolf.lood@med.lu.se), [johan.malmstrom@med.lu.se](mailto:johan.malmstrom@med.lu.se), [christofer.karlsson@med.lu.se](mailto:christofer.karlsson@med.lu.se),  
[magnus.rasmussen@med.lu.se](mailto:magnus.rasmussen@med.lu.se)

Running title: The surface proteome of *A. urinae*

Declaration of interest: none.

## Abstract

*Aerococcus urinae* is an emerging pathogen that causes urinary tract infections, bacteremia and infective endocarditis. The mechanisms through which *A. urinae* cause infection are largely unknown. The aims of this study were to describe the surface proteome of *A. urinae* and to analyse *A. urinae* genomes in search for genes encoding surface proteins. Two proteins, denoted Aerococcal surface protein (Asp) 1 and 2, were through the use of mass spectrometry based proteomics found to quantitatively dominate the aerococcal surface. The presence of these proteins on the surface was also shown using ELISA with serum from rabbits immunized with the recombinant Asp. These proteins had a signal sequence in the amino-terminal end and a cell wall-sorting region in the carboxy-terminal end, which contained an LPATG-motif, a hydrophobic domain and a positively charged tail. Twenty-three additional *A. urinae* genomes were sequenced using Illumina HiSeq technology. Six different variants of *asp* genes were found (denoted *asp1-6*). All isolates had either one or two of these *asp*-genes located in a conserved locus, designated Locus encoding Aerococcal Surface Proteins (LASP). The 25 genomes had in median 13 genes encoding LPXTG-proteins (range 6-24). For other Gram-positive bacteria, cell wall-anchored surface proteins with an LPXTG-motif play a key role for virulence. Thus, it will be of great interest to explore the function of the Asp proteins of *A. urinae* to establish a better understanding of the molecular mechanisms by which *A. urinae* cause disease.

**Key Words:** *Aerococcus urinae*, surface proteome, LPXTG-motif, genes encoding surface proteins.

## Introduction

*Aerococcus urinae* is a Gram-positive coccus that belongs to the phylum of Firmicutes. *A. urinae* is known to cause urinary tract infection (Christensen et al., 1989; Schuur et al., 1997; Senneby et al., 2015), bacteremia (Christensen et al., 1995; Senneby et al., 2016) and infective endocarditis (de Jong et al., 2010; Ebnother et al., 2002; Kristensen and Nielsen, 1995; Sunnerhagen et al., 2016), mostly in the elderly male population. Since the introduction of MALDI-TOF MS as a diagnostic tool in clinical microbiology laboratories, *A. urinae* is more frequently identified in clinical samples, especially in urine cultures, and has attracted attention as an important emerging pathogen (Rasmussen, 2016). A few studies have reported that *A. urinae* is part of a urinary tract microbiome (Hilt et al., 2014; Pearce et al., 2014). The mechanisms through which *A. urinae* establishes colonization and causes infection are largely unknown. Only one study has previously targeted the question of *A. urinae* virulence mechanisms (Shannon et al., 2010), demonstrating that it produces biofilm and is able to induce human platelet activation and aggregation. These are properties of potential importance in the process of establishing infections in the human host (Donlan and Costerton, 2002).

At the molecular level, little is known regarding aerococcal virulence factors (Carkaci et al., 2017). For other pathogenic Firmicutes, such as *Streptococcus pyogenes* and *Staphylococcus aureus*, cell wall-attached proteins with an LPXTG-motif play key roles for virulence. For instance, the M protein of *S. pyogenes* is an abundant surface protein, which possesses a broad spectrum of functions, such as the inhibition of phagocytosis and binding of several plasma proteins (Fischetti, 1989; Smeesters et al., 2010). The amino (NH<sub>2</sub>)-terminal part of the protein displays hypervariability, resulting in antigenic variation (Cunningham, 2000). The LPXTG-proteins share common features, such as a signal sequence in the NH<sub>2</sub>-terminal end and three characteristics in the carboxy (COOH)-terminal end; a cell wall-sorting region containing the LPXTG-motif, a hydrophobic membrane-spanning domain and a positively charged tail positioned in the cytoplasm. The LPXTG-motif is recognized by the membrane bound enzyme sortase and is, after cleavage between the threonine and glycine residues, covalently attached to the cell wall (Schneewind and Missiakas,

2014). It is at present unclear if *A. urinae* express proteins attached to the cell wall through the LPXTG-motif. Mass spectrometry (MS) based proteomics has previously been shown to be a useful method to detect bacterial surface-associated proteins with protein copies per cell accuracy (Malmstrom et al., 2009) and to determine the surface protein composition of gram-positive bacteria (Kilsgard et al., 2016; Rodriguez-Ortega et al., 2006; Severin et al., 2007). However, a presumption to utilize this technique is the genomic sequence for the analysed bacterial species or strain. Hence, the combination of next generation sequencing and MS-based proteomics constitutes a powerful strategy to search for potential novel virulence factors. The aims with this study were to describe the surface proteome of *A. urinae* and to describe the genes encoding surface proteins.

## Material and Methods

### Bacteria and culturing conditions

The strain ACS-120-V-Col10a (Col10a) was retrieved from the Culture collection of Gothenburg. The AU3 strain was collected at the Clinical Microbiology laboratory, Lund and originated from a blood culture as described previously (Senneby et al., 2012). Forty-six *A. urinae* isolates from blood cultures had been described previously (Senneby et al., 2016) and were designated as AuB followed by a number. *A. urinae* isolates were cultivated in Tryptic Soy Broth with 0.25% glucose (TSBG) for approximately 24 hours at 35° C in 5 % CO<sub>2</sub>.

### Bacterial surface digestion and MS sample preparation

Surface proteins were released from AU3 and Col10 stationary phase cells in triplicates with a modified protocol as previously described (Rodriguez-Ortega et al., 2006; Severin et al., 2007). The cells (~8 x10<sup>8</sup> colony forming units (CFU)) were washed with 20 mM Tris-HCl, 150 mM NaCl, pH 7.6 (TBS) and resuspended in 1 M d-arabinose, 10 mM CaCl<sub>2</sub> in TBS and 5 µg sequencing grade trypsin (Promega) and incubated at 37 °C with 500 rpm shaking for 20 min. The digested mixture was centrifuged with swing-out rotor at 2000 x g for 15 min at 4 °C. ProteaseMAX

(Promega) was added to the supernatants (surface fraction) to a final concentration of 0.01 % following by heating to 80 °C for denaturation. Cysteine residues were reduced with 25 mM tris(2-carboxyethyl)phosphine (Sigma-Aldrich, TCEP) and alkylated with 25 mM 2-Iodoacetamide (Sigma-Aldrich, IAA). Surface fraction samples were fully digested with 2 µg sequencing grade trypsin (Promega) for 10 h at 37 °C and the sample acidified with 0.5 % Trifluoroacetic acid. The digested cells (cellular fraction) were washed with TBS and resuspended in water and homogenized using a Fastprep-96 beadbeater (MPBio) with Lysing Matrix B tubes (MPBio). The cell lysates were denatured with 8 M Urea in 100 mM ammonium bicarbonate (ABC) and then reduced with 25 mM TCEP for 1 h at 37°C, and alkylated with 25 mM iodoacetamide for 45 min before diluting the sample with 100 mM ABC to a final urea concentration below 1.5 M. Proteins were digested by incubation with trypsin (trypsin:protein ratio of 1:100 (w:w)) for 10 h at 37°C. The peptides from both fractions were desalted and cleaned-up with reversed-phase spin columns (Vydac UltraMicroSpin Silica C18 300Å Columns, Harvard Apparatus) according to the manufacturer's instructions.

#### **MS data acquisition**

Peptide analyses were performed on a Q Exactive Plus mass spectrometer (Thermo Scientific) connected to an EASY-nLC 1000 ultra-high-performance liquid chromatography system (Thermo Scientific). Peptides were separated on an EASY-Spray ES802 columns (Thermo Scientific) using a linear gradient from 3 to 35% acetonitrile in aqueous 0.1% formic acid during 2 h. Data-dependent acquisition mode (DDA) and Data-independent acquisition mode (DIA) instrument settings were identical to as described in (Malmstrom et al., 2016). From the DDA data, spectral libraries were built using the TPP Fragg workflow (Teleman et al., 2017) using NCBI fasta files GCA\_000193205.1\_ASM19320 and GCA\_001649715.1\_ASM164971 respectively concatenated with iRT peptides, contaminants and decoys. The generated spectral libraries were used to extract the DIA data with DIANA v2.0.0 (Teleman et al., 2015) using a 1% peptide false discovery rate.

## **MS data analysis**

Data was processed with custom R-scripts using the tidyverse package collection together with broom package for statistical functions. For the quantification of proteins, all integrated peptide ion intensities extracted from the MS2 spectra was summed up by protein and then divided by protein length. Peptides matching more than one protein were not included in the analysis. Protein quantification data was normalized based on the sample total intensity. Proteins from Col10a and AU3 were classified into orthologous pairs using ProteinOrtho.pl v5.16b (Lechner et al., 2011) and protein domains predicted using standalone InterProScan v5.11-50 (Jones et al., 2014).

## **Genome sequencing**

Bacteria were pelleted through centrifugation and stored at -70° C until shipment on dry ice to GATC Biotech (Konstanz, Germany). Library preparation was performed using an optimized protocol and standard Illumina adapter sequences were used. Sequencing was performed with Illumina HiSeq 2500 (Illumina, Inc., San Diego, US) with paired-end reads, 2 x 125 bp. The genomes of AU3 (ASM164971v1, GenBank assembly accession GCA\_001649715.1) and ACS-120-V-Col10a (ASM19320v1, GenBank assembly accession GCA\_00193205.1) had been previously published. For assembly of the genomes, SPAdes 3.9 with the careful-mode on, was used through services on the Center for Genomic Epidemiology's website (Nurk et al., 2013). The computations were performed on resources provided by the Swedish National Infrastructure for Computing (SNIC) at Uppsala Multidisciplinary Center for Advanced Computational Science (UPPMAX). Assembly quality was evaluated using QUAST (Gurevich et al., 2013). Sample contamination was evaluated with Kraken (cross-species) (Wood and Salzberg, 2014) and by mapping the raw sequence data to a reference genome using BWA MEM (Li and Durbin, 2009) and evaluating the presence of minority bases with samtools mpileup (within-species) (Li et al., 2009). In cases with within-species contamination the contaminant gene were eliminated by coverage analysis, i.e. genes with low coverage were excluded. The BLAST software (Zhang et al., 2000) was used for searches in the genomes. Sequence alignment was performed with Clustal Omega. The maximum likelihood method was used for building phylogenetic trees through the use of the MEGA software (version 7.0.26).

Annotation of the genomes was performed with RASTtk on the PATRIC platform (Wattam et al., 2017). SignalP 4.1 was used to locate signal peptide cleavage sites (Petersen et al., 2011). Hydrophobic plots were performed using services on the Expasy website (Kyte and Doolittle, 1982). The ClustalW multiple sequence alignment was used to produce identity scores (%) for the amino acid sequences with the following parameters: similarity matrix: gonnet, open gap penalty 10, extend gap penalty 0.1-0.2, gap distance 4, delay divergent 30%. The signal sequences and the N-terminal end (starting with the LPXTG-motive) were deleted prior to analysis. Searches were also performed in 26 *A. urinae* genomes belonging to Bioproject PRJNA315093, that were accessible on the NCBI webpage. The bacterial isolates originated from the female urinary tract.

### **Expression cloning and protein production**

The MacVector software (v. 14.5.3) was used to analyse the bacterial genome for suitable primer sites for amplification of *asp* genes. Primers were ordered from Eurofins and can be found in Table 1. Chromosomal DNA from AU3 was extracted using the innuPRerp Bacterial DNA System (Analytik Jena AG). PCR products were inserted into the pGEX-6P-1 expression vector. One Shot TOP10 *E. coli* (Thermo Fisher Scientific) were transformed and used for plasmid amplification. Plasmids were extracted using Qiagen Plasmid Plus kits. The plasmids were sent to GATC GmbH for sequencing to ensure that the sequence was correct. BL21 cells with pLysS were then transformed according to the manufacturer's protocol, with selection for pGEX-6-p-1 (100 µg/ ml ampicillin) and for pLysS (32 µg/ml chloramphenicol).

For protein expression, an overnight *E. coli* culture carrying the construct was diluted 1:25 in fresh TSBG supplemented with antibiotics and grown for 3 h. IPTG (1 mM) was then added and incubation continued for 3 h at 37° C. The bacteria were pelleted and the supernatant discarded. Pellet was dissolved in BugBuster (Merck Millipore) suspended in 20 mM TRIS (pH 7.4) supplemented with 1:100 (v/v) benzonase. After 10 minutes of incubation at room temperature bacterial cell debris was collected by centrifugation and the supernatant was loaded onto an equilibrated GST-column, washed, and finally eluted (elution buffer). The sequence of the affinity-purified protein was confirmed by mass-spectrometry at the SciBlu core facility at Lund University.



## **Antiserum generation and ELISA**

To generate antiserum against the affinity-purified recombinant proteins, the proteins were sent to Biogenes GmbH for immunization. To assess binding of antibodies to the surface proteins an ELISA was performed. Nunc MaxiSorp (ThermoFisher Scientific) 96-well plates were coated with either a suspension of bacteria (AU3, Col10a or *A. viridans*), or a solution of Asp-proteins in coating buffer (1.69 g Na<sub>2</sub>CO<sub>3</sub> and 2.94 g NaHCO<sub>3</sub> in 1 L H<sub>2</sub>O). As a control, bacteria with reduced surface protein expression were generated by incubating a bacterial solution with trypsin (0.0004% w/v) at 37° C for 20 minutes after which the reaction was stopped by adding trypsin inhibitor at a molar ratio of 8:1. The trypsinated bacteria and control bacteria (where trypsin inhibitor was added before the trypsin) were then pelleted by centrifugation, washed (PBS), and finally resuspended in coating buffer. The bacteria were then heat-killed at 85° C for 30 minutes before adding the samples to the Nunc plate for coating over night at 4° C. The plates were washed (PBST), rabbit serum in different dilutions added, followed by further washes and addition of protein G coupled HRP (1:3000). As an HRP substrate, ABTS dissolved 1:20 in a substrate solution (21 g citric acid monohydrate and 17.8 g Na<sub>2</sub>HPO<sub>4</sub> 2 H<sub>2</sub>O in 1 L H<sub>2</sub>O, pH 4.5) was used in combination with H<sub>2</sub>O<sub>2</sub>; incubated in room temperature for 15 minutes. The plates were then analyzed with the iMark Microplate reader (Bio-Rad) at 415 nm.

## **Sequence similarity network (SNN) of LPXTG-containing ORFs.**

In total, 106 285 ORFs from the 25 *A. urinae* genomes were analyzed with standalone InterProScan v5.11-50 (Jones et al., 2014) and included the following analyses: Pfam, PANTHER, Gene3D, CDD, TIGRFAM, ProSitePatterns, SUPERFAMILY, Hamap, ProSiteProfiles, Coils, SMART, MobiDBLite, PRINTS, PIRSF, SignalP, SFLD and ProDom. LPXTG-motifs were predicted with InterPro signature accessions PS50847 and/or TIGR01167 and/or PF00746 and sequences identified through this procedure were analyzed by all-by-all BLAST comparisons using blastp (v2.7.1+ with default settings). The blast result was filtered with the following thresholds: *E*-value < 1×10<sup>-10</sup>, >50 % sequence identity and >75 alignment length.

The SNN was visualized using Cytoscape (v3.5.1) (Shannon et al., 2003), where each node represents a protein/ORF and an edge or line between the nodes denotes a

similarity relationship between the proteins. The “organic” layout was used whereby nodes are clustered more tightly if they are more highly interconnected and all edges were bundled.

## Results

### Quantitative profiling of *A. urinae* surface proteomes.

To search for putative novel surface-associated proteins, we released surface exposed proteins from hypotonically swelled *A. urinae* cells of the strains Col10a and AU3 with a short trypsin digestion step. This produced a surface protein fraction and a cell protein fraction. This method has previously been used for releasing surface exposed proteins from cells of other gram-positive bacteria (Rodriguez-Ortega et al., 2006; Solis et al., 2010). The trypsinated cells (the cell fraction) were homogenized and used as control samples hypothetically containing mostly intracellular and membrane proteins. Both surface and cell fractions were then prepared for MS analysis. All derived peptide samples (n = 12) were analysed using data-independent-acquisition (DIA) followed by targeted data extraction (SWATH-MS) (Fig. 1A). In this analysis we quantified a total of 1118 and 1168 proteins of which 765 and 682 proteins were detected in the surface fractions from strains Col10a and AU3 respectively (Fig. 1B). Essentially all surface fraction proteins (99.8 %) were also detected in the cellular fractions (Fig. 1C-D), but with large differences of abundances between fractions. The majority of proteins in the surface fraction were much less abundant in the cellular fraction (Fig. 2A). We defined surface associated proteins based on two criteria; 10-fold enriched in the surface associated fractions and a Hochberg adjusted p-value < 0.01 compared to the cellular fraction (Col10a, n= 24; AU3, n = 30) (Fig. 2A and B). The stringent cut-off values for the surface associated protein groups were selected to distinctly separate proteins identified in the two fractions. The defined surface associated proteins included the most abundant proteins in the surface fraction (Fig. 2C) and 70% of these proteins had a predicted signal peptide domain as determined by InterProScan analysis (Fig. 2D). Of the proteins identified in the surface fraction but not defined as surface associated, most were ribosomal proteins,

metabolic enzymes and transcriptional regulators and only ~9 % had a predicted signal peptide (Fig. 2D).

Next we compared the surface associated protein orthology between the two strains as defined by the bidirectional best-hit (Lechner et al., 2011). Of the total 54 surface associated proteins from both strains, 17 pairs were determined to be orthologous by the Proteinortho software (Lechner et al., 2011) (Fig. 2E). Based on this analysis we identified two orthologous protein pairs with predicted LPXTG-sequences and signal peptide protein domains that were abundantly produced on the surface (Fig. 2E, red nodes with dotted edges). The proteins were denoted Aerococcal surface protein (Asp) 1 and 2 and the corresponding genes were accordingly named (*asp1* and *asp2*). The Asp1 was the most abundant protein on the surface, with a relative abundance of approximately 73 % and 26 % of the total proteins on the surface (for Col10a and AU3 respectively). No proteins similar to the Asp proteins were found in the InterPro database that includes high-level structure-based classifications and sub-family classifications. Furthermore, InterPro analysis of motifs and/or domains only identified LPXTG and signal peptide sequences. Additionally, four other surface associated proteins were classified as LPXTG-anchored. However, these proteins lacked an orthologous counterpart or the orthologous counterpart did not harbour a predicated LPXTG sequence (see Fig. 2E). The total number of predicted LPXTG proteins in the genomes was 8 for AU3 and 10 for Col10a (not shown). In summary, the quantitative profiling of *A. urinae* surface proteomes revealed that less than half all theoretical LPXTG proteins are expressed during the experimental conditions used and that both surfaces are dominated by an orthologous pair of proteins (Asp1).

### **Genomic and protein sequence analysis of Aerococcal surface protein 1 and 2.**

Based on the outcome from the MS-analysis, we continued with acquiring 23 additional novel whole genome sequences of *A. urinae* isolates to investigate the presence of genes encoding Asp proteins and also other LPXTG-containing proteins. The genomes of the AU3 and the Col10a isolates were included in the analysis. Thus, 25 *A. urinae* genomes were studied. Variants of the *asp*-genes were found in all isolates sequenced. The genes were denoted *asp1-6*, based on a phylogenetic analysis (Figure 3), and the corresponding gene products were denoted Asp1-6. The *asp*-genes were located adjacent to each other in a chromosomal locus of the aerococcal

genome, in this study designated as Locus encoding Aerococcal Surface Proteins (LASP). In all isolates it was constituted by, in a 5' – 3' order, a pyruvate carboxylase gene, a gene designated as a “hypothetical cytosolic protein”, the *asp*-gene(s) and a 16S rRNA methyltransferase gene. The LASP is schematically depicted in Figure 4. The primary structure of Asp1 contained 284-298 amino acids and Asp2 contained 307-382 amino acids. The Asp3 contained 376-407 amino acids, all Asp4 had 425 amino acids and Asp5 and Asp6 had 483 amino acids. The comparison of the amino acid sequences for Asp1-6 is presented in Table 2 with identity scores (%). The amino acid sequence of the LPXTG-motif was LPATG in all isolates. The aerococcal isolates had combinations of several different *asp*-genes in their LASP. Four variants were found, designated LASP<sub>1-4</sub>. In LASP<sub>1</sub> only the *asp1* was present, LASP<sub>2</sub> had both *asp1* and *asp2*, LASP<sub>3</sub> had *asp3* and *asp6* whereas LASP<sub>4</sub> contained *asp4* and *asp5*. The 26 *A. urinae* genomes in the bio project PRJNA315093 also included *asp* genes and these isolates had either LASP<sub>1</sub>, LASP<sub>2</sub> or LASP<sub>4</sub>.

### Expression cloning and immunological assays

The predicted mature forms of Asp1 and Asp2 were recombinantly expressed in *E. coli* and were used for production of two polyclonal antisera in rabbits. ELISA demonstrated that the post-immune antisera from both rabbits reacted with Asp1 and Asp2, whereas there was no reaction between the pre-immune sera (data not shown). To confirm that Asp1 and Asp2 were present on the surface of *A. urinae*, ELISA using anti-Asp1 and anti-Asp2 antisera against whole bacteria immobilized in the wells was performed. This showed that both the antisera reacted with *A. urinae* AU3 and Col10a but not with the control *A. viridans* that lacked *asp* homologs in the genome (Figure 5A and data not shown). The signal was reduced when the bacteria were treated with trypsin, further indicating that the proteins are located on the surface (Figure 5B).

### Genome-wide associations of LPXTG-containing proteins in *A. urinae*

To expand the information on other potential surface anchored proteins in addition to the Asp-proteins we analysed all ORFs (n=106 285) from the 25 genomes for LPXTG motifs. From this analysis we identified 289 ORFs that contained LPXTG-motifs. In order to cluster these ORFs into groups we compared the translated sequences of the

289 ORFs using all-by-all BLAST comparisons, which in turn yielded 83 521 comparisons. After applying a BLAST filter as previously described (Mashiyama et al., 2014), 9 647 sequence comparisons remained and these were selected for sequence similarity network (SNN) visualization (Figure 6A). Eleven apparent clusters were identified and these clusters were numbered and colored (Figure 6B). The functional domains that were predicted in the analysis are presented in Figure 6C. Our analysis of the 25 genomes revealed that these isolates had in median 13 genes encoding LPXTG-proteins (range 6-24). Predicted domains of these proteins included the G5 domain, which is widely found in Gram-positive bacteria such as streptococcal species (Lin et al., 2012) and has been proposed to be involved in biofilm formation of *S. aureus* (Bateman et al., 2005). Also, collagen binding domains and domains with collagen triple helix repeats were predicted. Furthermore, Mucin-binding protein domains, which play a role in adhesion of *Lactobacillus* species to mucin (Chatterjee et al., 2018) and domains with Rib/alpha-like repeats were predicted. Proteins with Rib and alpha repeats can be found in surface proteins of group B *Streptococcus* (Wastfelt et al., 1996).

## Discussion

In this study, we aimed to describe the surface proteome of *A. urinae* and also to describe aerococcal genes encoding surface proteins. We selected the two strains Col10a and AU3 for the proteomic surface profiling using previously published protocols (Rodriguez-Ortega et al., 2006; Severin et al., 2007). A major issue with studying bacterial surface exposed proteins is contamination of intracellular proteins (Solis et al., 2010). Here we performed a subtraction of the contaminating proteins by comparing the protein abundances between the surface fraction and cellular fraction resulting in 54 surface associated proteins from both strains of which 17 were orthologues. A total of 18 genes encoding potential proteins with an LPXTG-motif were identified in the Col10a and AU3 genomes. However, only two orthologous pairs of them were detected in the surface fractions. These two proteins, denoted Asp1 and Asp2 in this study, were among the most abundant proteins on the aerococcal surface. The Asp1 and Asp2 share features with LPXTG-proteins from other bacterial

species (for instance streptococcal and staphylococcal species), such as a signal sequence in the NH<sub>2</sub>-terminal end, a hydrophobic membrane-spanning domain and a positively charged tail in the COOH-terminal end.

In a recent publication by Carkaci *et al.* (Carkaci et al., 2017), the authors described and compared 40 *A. urinae* and eight *A. sanguinicola* genomes of strains isolated from blood, urine and heart valves. In that study, virulence genes encoding LPXTG-containing proteins were not reported. The virulence-associated genes were however identified using a virulence factor database that only included known virulence factors. Thus, aerococcal genes with low similarity to genes in this collection, such as the *asp* genes, were prone to be missed. Prior to our investigation, it was therefore unclear whether all *A. urinae* isolates had genes encoding LPXTG-proteins.

Our results show that *A. urinae* genomes of isolates originating from both blood and urine cultures, contain genes encoding LPXTG-proteins and that some of the predicted functional domains of these proteins are of potential importance for virulence. Our results also show that *A. urinae* has a conserved locus, LASP, which comprise one or two of the *asp* genes. The organization of the LASP has a high degree of similarity to the organization of the so-called *mga* regulon of *S. pyogenes*. The genes in the *mga* regulon encode important streptococcal virulence factors (M and M-like proteins and the C5a peptidase (Fischetti, 1989; O'Connor and Cleary, 1987)), which are regulated by the transcription factor Mga (Berge et al., 1998; Caparon and Scott, 1987; McIver et al., 1995). Both the aerococcal and the streptococcal loci comprise LPXTG-containing genes, arranged in a conserved fashion, but with a variation between bacterial isolates in the number of genes present and in sequence similarities. The aerococcal protein HCP, with the corresponding gene positioned in the LASP upstream from the *asp* gene(s), was detected in the intracellular fraction in the MS analysis. It is possible that the HCP has a similar regulatory role for the LASP as Mga has for the *mga* regulon of *S. pyogenes*.

The function of the aerococcal Asp proteins is unknown. However, LPXTG-proteins in other species are involved in processes such as bacterial adhesion, immune evasion, internalization, iron acquisition and biofilm formation. Since most *A. urinae* isolates are found in urine cultures, it is possible that the Asp proteins act as adhesion molecules to epithelial cells in the human urinary tract or are involved in the process of biofilm formation, which, mentioned previously, has been described as a feature of *A. urinae* (Shannon et al., 2010). A few isolates in this study came from patients

435 diagnosed with infective endocarditis but no common LASP variant was observed for  
436 these isolates (data not shown). Conclusion regarding correlation between LASP  
437 variants and clinical features could not be made due to the small sample size.

## 439 **Conclusion**

440 In this study, we show for the first time that *A. urinae* has genes encoding surface  
441 proteins with an LPXTG-motif and that two of these proteins quantitatively dominate  
442 the surface. To establish a better understanding of how *A. urinae* colonize humans and  
443 cause disease, it is of great interest to explore the function of the Asp proteins of *A.*  
444 *urinae* further.

## 448 **Acknowledgements**

449 We acknowledge Ann-Cathrine Petersson and Gisela Hovold for important help.

## 451 **Funding:**

452 This study was supported by the Foundations of Knut and Alice Wallenberg  
453 (2016.0023), the Swedish Research Council (project 2015-02481), the Wallenberg  
454 Academy Fellow program KAW (2012.0178 and 2017.0271), the Swedish  
455 Government Funds for Clinical Research, the Royal Physiographic Society in Lund,  
456 and the foundations of Marianne and Marcus Wallenberg, Crafoord, Österlund,  
457 Lundgren and Skåne University hospital.

## References

- Bateman, A., Holden, M.T., Yeats, C., 2005. The G5 domain: a potential N-acetylglucosamine recognition domain involved in biofilm formation. *Bioinformatics* 21, 1301-1303.
- Berge, A., Rasmussen, M., Björck, L., 1998. Identification of an insertion sequence located in a region encoding virulence factors of *Streptococcus pyogenes*. *Infect Immun* 66, 3449-3453.
- Caparon, M.G., Scott, J.R., 1987. Identification of a gene that regulates expression of M protein, the major virulence determinant of group A streptococci. *Proc Natl Acad Sci U S A* 84, 8677-8681.
- Carkaci, D., Hojholt, K., Nielsen, X.C., Dargis, R., Rasmussen, S., Skovgaard, O., Fuursted, K., Andersen, P.S., Stegger, M., Christensen, J.J., 2017. Genomic characterization, phylogenetic analysis, and identification of virulence factors in *Aerococcus sanguinicola* and *Aerococcus urinae* strains isolated from infection episodes. *Microb Pathog*.
- Chatterjee, M., Pushkaran, A.C., Vasudevan, A.K., Menon, K.K.N., Biswas, R., Mohan, C.G., 2018. Understanding the adhesion mechanism of a mucin binding domain from *Lactobacillus fermentum* and its role in enteropathogen exclusion. *Int J Biol Macromol* 110, 598-607.
- Christensen, J.J., Jensen, I.P., Faerk, J., Kristensen, B., Skov, R., Korner, B., 1995. Bacteremia/septicemia due to *Aerococcus*-like organisms: report of seventeen cases. Danish ALO Study Group. *Clin Infect Dis* 21, 943-947.
- Christensen, J.J., Korner, B., Kjaergaard, H., 1989. *Aerococcus*-like organism--an unnoticed urinary tract pathogen. *APMIS* 97, 539-546.
- Cunningham, M.W., 2000. Pathogenesis of group A streptococcal infections. *Clin Microbiol Rev* 13, 470-511.
- de Jong, M.F., Soetekouw, R., ten Kate, R.W., Veenendaal, D., 2010. *Aerococcus urinae*: severe and fatal bloodstream infections and endocarditis. *J Clin Microbiol* 48, 3445-3447.
- Donlan, R.M., Costerton, J.W., 2002. Biofilms: survival mechanisms of clinically relevant microorganisms. *Clin Microbiol Rev* 15, 167-193.



501 Ebnother, C., Altwegg, M., Gottschalk, J., Seebach, J.D., Kronenberg, A., 2002.  
 502 *Aerococcus urinae* endocarditis: case report and review of the literature.  
 503 *Infection* 30, 310-313.  
 504 Fischetti, V.A., 1989. Streptococcal M protein: molecular design and biological  
 505 behavior. *Clin Microbiol Rev* 2, 285-314.  
 506 Gurevich, A., Saveliev, V., Vyahhi, N., Tesler, G., 2013. QUAST: quality assessment  
 507 tool for genome assemblies. *Bioinformatics* 29, 1072-1075.  
 508 Hilt, E.E., McKinley, K., Pearce, M.M., Rosenfeld, A.B., Zilliox, M.J., Mueller, E.R.,  
 509 Brubaker, L., Gai, X., Wolfe, A.J., Schreckenberger, P.C., 2014. Urine is not sterile:  
 510 use of enhanced urine culture techniques to detect resident bacterial flora in the  
 511 adult female bladder. *J Clin Microbiol* 52, 871-876.  
 512 Jones, P., Binns, D., Chang, H.Y., Fraser, M., Li, W., McAnulla, C., McWilliam, H.,  
 513 Maslen, J., Mitchell, A., Nuka, G., Pesseat, S., Quinn, A.F., Sangrador-Vegas, A.,  
 514 Scheremetjew, M., Yong, S.Y., Lopez, R., Hunter, S., 2014. InterProScan 5: genome-  
 515 scale protein function classification. *Bioinformatics* 30, 1236-1240.  
 516 Kilsgard, O., Karlsson, C., Malmstrom, E., Malmstrom, J., 2016. Differential  
 517 compartmentalization of *Streptococcus pyogenes* virulence factors and host  
 518 protein binding properties as a mechanism for host adaptation. *Int J Med*  
 519 *Microbiol* 306, 504-516.  
 520 Kristensen, B., Nielsen, G., 1995. Endocarditis caused by *Aerococcus urinae*, a  
 521 newly recognized pathogen. *Eur J Clin Microbiol Infect Dis* 14, 49-51.  
 522 Kyte, J., Doolittle, R.F., 1982. A simple method for displaying the hydropathic  
 523 character of a protein. *J Mol Biol* 157, 105-132.  
 524 Lechner, M., Findeiss, S., Steiner, L., Marz, M., Stadler, P.F., Prohaska, S.J., 2011.  
 525 Proteinortho: detection of (co-)orthologs in large-scale analysis. *BMC*  
 526 *Bioinformatics* 12, 124.  
 527 Li, H., Durbin, R., 2009. Fast and accurate short read alignment with Burrows-  
 528 Wheeler transform. *Bioinformatics* 25, 1754-1760.  
 529 Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G.,  
 530 Abecasis, G., Durbin, R., Genome Project Data Processing, S., 2009. The Sequence  
 531 Alignment/Map format and SAMtools. *Bioinformatics* 25, 2078-2079.  
 532 Lin, I.H., Hsu, M.T., Chang, C.H., 2012. Protein domain repetition is enriched in  
 533 *Streptococcal* cell-surface proteins. *Genomics* 100, 370-379.

534 Malmstrom, E., Kilsgard, O., Hauri, S., Smeds, E., Herwald, H., Malmstrom, L.,  
 535 Malmstrom, J., 2016. Large-scale inference of protein tissue origin in gram-  
 536 positive sepsis plasma using quantitative targeted proteomics. *Nat Commun* 7,  
 537 10261.  
 538 Malmstrom, J., Beck, M., Schmidt, A., Lange, V., Deutsch, E.W., Aebersold, R., 2009.  
 539 Proteome-wide cellular protein concentrations of the human pathogen  
 540 *Leptospira interrogans*. *Nature* 460, 762-765.  
 541 Mashiyama, S.T., Malabanan, M.M., Akiva, E., Bhosle, R., Branch, M.C., Hillerich, B.,  
 542 Jagessar, K., Kim, J., Patskovsky, Y., Seidel, R.D., Stead, M., Toro, R., Vetting, M.W.,  
 543 Almo, S.C., Armstrong, R.N., Babbitt, P.C., 2014. Large-scale determination of  
 544 sequence, structure, and function relationships in cytosolic glutathione  
 545 transferases across the biosphere. *PLoS Biol* 12, e1001843.  
 546 McIver, K.S., Heath, A.S., Green, B.D., Scott, J.R., 1995. Specific binding of the  
 547 activator Mga to promoter sequences of the *emm* and *scpA* genes in the group A  
 548 streptococcus. *J Bacteriol* 177, 6619-6624.  
 549 Nurk, S., Bankevich, A., Antipov, D., Gurevich, A.A., Korobeynikov, A., Lapidus, A.,  
 550 Prjibelski, A.D., Pyshkin, A., Sirotkin, A., Sirotkin, Y., Stepanauskas, R., Clingenpeel,  
 551 S.R., Woyke, T., McLean, J.S., Lasken, R., Tesler, G., Alekseyev, M.A., Pevzner, P.A.,  
 552 2013. Assembling single-cell genomes and mini-metagenomes from chimeric  
 553 MDA products. *J Comput Biol* 20, 714-737.  
 554 O'Connor, S.P., Cleary, P.P., 1987. In vivo *Streptococcus pyogenes* C5a peptidase  
 555 activity: analysis using transposon- and nitrosoguanidine-induced mutants. *J*  
 556 *Infect Dis* 156, 495-504.  
 557 Pearce, M.M., Hilt, E.E., Rosenfeld, A.B., Zilliox, M.J., Thomas-White, K., Fok, C.,  
 558 Kliethermes, S., Schreckenberger, P.C., Brubaker, L., Gai, X., Wolfe, A.J., 2014. The  
 559 female urinary microbiome: a comparison of women with and without urgency  
 560 urinary incontinence. *MBio* 5, e01283-01214.  
 561 Petersen, T.N., Brunak, S., von Heijne, G., Nielsen, H., 2011. SignalP 4.0:  
 562 discriminating signal peptides from transmembrane regions. *Nat Methods* 8,  
 563 785-786.  
 564 Rasmussen, M., 2016. *Aerococcus*: an increasingly acknowledged human  
 565 pathogen. *Clin Microbiol Infect* 22, 22-27.

566 Rodriguez-Ortega, M.J., Norais, N., Bensi, G., Liberatori, S., Capo, S., Mora, M.,  
 567 Scarselli, M., Doro, F., Ferrari, G., Garaguso, I., Maggi, T., Neumann, A., Covre, A.,  
 568 Telford, J.L., Grandi, G., 2006. Characterization and identification of vaccine  
 569 candidate proteins through analysis of the group A Streptococcus surface  
 570 proteome. *Nat Biotechnol* 24, 191-197.  
 571 Schneewind, O., Missiakas, D., 2014. Sec-secretion and sortase-mediated  
 572 anchoring of proteins in Gram-positive bacteria. *Biochim Biophys Acta* 1843,  
 573 1687-1697.  
 574 Schuur, P.M., Kasteren, M.E., Sabbe, L., Vos, M.C., Janssens, M.M., Buiting, A.G.,  
 575 1997. Urinary tract infections with *Aerococcus urinae* in the south of The  
 576 Netherlands. *Eur J Clin Microbiol Infect Dis* 16, 871-875.  
 577 Senneby, E., Goransson, L., Weiber, S., Rasmussen, M., 2016. A population-based  
 578 study of aerococcal bacteraemia in the MALDI-TOF MS-era. *Eur J Clin Microbiol*  
 579 *Infect Dis* 35, 755-762.  
 580 Senneby, E., Petersson, A.C., Rasmussen, M., 2012. Clinical and microbiological  
 581 features of bacteraemia with *Aerococcus urinae*. *Clin Microbiol Infect* 18, 546-  
 582 550.  
 583 Senneby, E., Petersson, A.C., Rasmussen, M., 2015. Epidemiology and antibiotic  
 584 susceptibility of aerococci in urinary cultures. *Diagn Microbiol Infect Dis* 81, 149-  
 585 151.  
 586 Severin, A., Nickbarg, E., Wooters, J., Quazi, S.A., Matsuka, Y.V., Murphy, E.,  
 587 Moutsatsos, I.K., Zagursky, R.J., Olmsted, S.B., 2007. Proteomic analysis and  
 588 identification of *Streptococcus pyogenes* surface-associated proteins. *J Bacteriol*  
 589 189, 1514-1522.  
 590 Shannon, O., Morgelin, M., Rasmussen, M., 2010. Platelet activation and biofilm  
 591 formation by *Aerococcus urinae*, an endocarditis-causing pathogen. *Infect*  
 592 *Immun* 78, 4268-4275.  
 593 Shannon, P., Markiel, A., Ozier, O., Baliga, N.S., Wang, J.T., Ramage, D., Amin, N.,  
 594 Schwikowski, B., Ideker, T., 2003. Cytoscape: a software environment for  
 595 integrated models of biomolecular interaction networks. *Genome Res* 13, 2498-  
 596 2504.  
 597 Smeesters, P.R., McMillan, D.J., Sriprakash, K.S., 2010. The streptococcal M  
 598 protein: a highly versatile molecule. *Trends Microbiol* 18, 275-282.

Solis, N., Larsen, M.R., Cordwell, S.J., 2010. Improved accuracy of cell surface shaving proteomics in *Staphylococcus aureus* using a false-positive control. *Proteomics* 10, 2037-2049.

Sunnerhagen, T., Nilsson, B., Olaison, L., Rasmussen, M., 2016. Clinical and microbiological features of infective endocarditis caused by aerococci. *Infection* 44, 167-173.

Teleman, J., Hauri, S., Malmstrom, J., 2017. Improvements in Mass Spectrometry Assay Library Generation for Targeted Proteomics. *J Proteome Res* 16, 2384-2392.


Teleman, J., Rost, H.L., Rosenberger, G., Schmitt, U., Malmstrom, L., Malmstrom, J., Levander, F., 2015. DIANA--algorithmic improvements for analysis of data-independent acquisition MS data. *Bioinformatics* 31, 555-562.

Wastfelt, M., Stalhammar-Carlemalm, M., Delisse, A.M., Cabezon, T., Lindahl, G., 1996. Identification of a family of streptococcal surface proteins with extremely repetitive structure. *J Biol Chem* 271, 18892-18897.


Wattam, A.R., Davis, J.J., Assaf, R., Boisvert, S., Brettin, T., Bun, C., Conrad, N., Dietrich, E.M., Disz, T., Gabbard, J.L., Gerdes, S., Henry, C.S., Kenyon, R.W., Machi, D., Mao, C., Nordberg, E.K., Olsen, G.J., Murphy-Olson, D.E., Olson, R., Overbeek, R., Parrello, B., Pusch, G.D., Shukla, M., Vonstein, V., Warren, A., Xia, F., Yoo, H., Stevens, R.L., 2017. Improvements to PATRIC, the all-bacterial Bioinformatics Database and Analysis Resource Center. *Nucleic Acids Res* 45, D535-D542.

Wood, D.E., Salzberg, S.L., 2014. Kraken: ultrafast metagenomic sequence classification using exact alignments. *Genome Biol* 15, R46.


Zhang, Z., Schwartz, S., Wagner, L., Miller, W., 2000. A greedy algorithm for aligning DNA sequences. *J Comput Biol* 7, 203-214.



**Figure 1. Proteomics of *A. urine* cellular and surface fractions.** (a) Summary of the proteomics workflow for *A. urine* cellular and surface fractions. Surface proteins were released from two different *A. urine* strains using a short incubation with trypsin and the fractions separated by centrifugation. Following the sample preparation, the derived peptide samples were analysed with LC-MS/MS using both shotgun- and SWATH-MS. (b) Average numbers  $\pm$  standard deviations of quantified proteins per respective strain and fractions using three replicates per condition. (c, d) Number of proteins common between fractions and per respective strain.



**Figure 2. Proteomic analysis reveals *A. urinae* surface associated proteins.** (a) Volcano plots showing Hochberg adjusted  $-\log_{10}$  p-values correlated to  $\log_2$  fold changes in surface vs cell fractions of respective strain. Proteins in red represent the surface associated proteins with a fold-change  $\geq 10$  (vertical dotted lines) and adjusted p-value  $< 0.01$  (horizontal dotted lines). (b) Heat map and unsupervised hierarchical clustering of surface associated proteins (red dots in (a)) across individual replicates and intensities normalized on total ion current (TIC) normalization (c). Average abundance distribution of surface fraction proteins of respective strain with the surface associated proteins marked with red. (d) The proportion of proteins with a predicted signal peptide of Surface Fraction proteins divided into Surface Associated and other proteins groups. The number (n) of proteins per protein group and strain are indicated in the plot. (e) Shows the ortholog pairs indicated as dotted lines (edges) between the surface associated proteins (nodes named with GenBank protein accession number) of the two strains. The node size is proportional to relative protein abundance per strain. Node colour represents predicted presence of protein domains SP: signal peptide and LPxTG: Gram-positive LPxTG cell wall anchor.




661

662 **Figure 3.**

663 Mid-point rooted maximum likelihood tree visualizing the clustering of the *asp* genes  
664 in the 25 isolates. The *asp* genes of Col10a (brown) and AU3 (grey) are highlighted in  
665 the tree. The scale length indicates 0.1 base substitutions per base.

666







**Figure 4. Schematic figure of different types of the Locus encoding Aerococcal Surface Proteins (LASP).**

The LASP includes three genes enclosing the *asp* gene(s). Four different types of the LASP were identified in the *A. urinae* isolates, denoted by us as LASP<sub>1-4</sub>.

PC; Pyruvate carboxylase gene, HCP; Hypothetical cytosolic protein gene, *asp*; aerococcal surface protein gene, *16S gm*; 16S rRNA guanine-methyltransferase gene.


The distance between genes and length of genes are not according to scale.



**Figure 5.**

Data shown as median with whiskers showing range, experiments performed in triplicate. A) Binding of serum from rabbits immunized with Asp1 was studied with ELISA. Serum from the same rabbit taken from before the immunization was used as

686 a negative control. The plate was coated with AU3, Col10a, or *A. viridans*. Dilutions  
687 of rabbit serum are shown on the x axis and range from 1:400 to no serum at all. B)  
688 AU3 and Col10a were treated with trypsin to reduce the amount of Asp1 on the  
689 surface. AU3 and Col10a with trypsin inhibitor added were used as controls.  
690



**Figure 6. Sequence similarity network of *A. urinae* LPXTG proteins.**

A) All *A. urinae* ORFs (n=106,285) from 25 genomes were analyzed with InterProScan to obtain predictions of functional domains. ORFs containing LPXTG motifs (n=289) were selected with the criteria of InterPro signature accessions PS50847 and/or TIGR01167 and/or PF00746. For sequence similarity analysis the

697 289 selected ORFs were analyzed all-by-all BLAST comparisons. B) Sequence  
698 similarity network of the 289 LPXTG ORFs with 9,647 edges each representing a  
699 BLAST hit. Eleven clusters of LPXTG ORFs are colored accordingly. Proteins that  
700 were previously defined as being surface associated in Col10a or AU3 were outlined  
701 in black. C) Predicted functional domains of the LPXTG-ORFs and in which clusters  
702 they were detected in.  
703

704

705

Direction	Gene	
	<i>asp1</i>	<i>asp2</i>
Forward	GAGGAATAATAAATGGCAGACGCA TTTGTAACACCAGTA	GAFFAATAATAATAAATGGCAGTT GCTAAAGCTGAAATGTTT
Reverse	AGTAGCTGGTAATTTAGCGTTAGCG TTAGCTTTTTGAGCTTTGC	TAATTTAGCATTAGCTTTTTCAGC TTAGCTTCTTTATC

706 **Table 1. Primers used for expression cloning**707 The primers used to construct the pGEX-6p-1 plasmids containing the *asp1* and *asp2*

708 genes were designed with an RBS.

709

Asp	1	2	3	4	5	6
1	72-100 %	40-53%	20-21%	12-13%	10-11%	11-12%
2	-	90-100 %	16-17%	14-15%	13%	13%
3	-	-	100 %	16%	14%	15%
4	-	-	-	100 %	12%	13%
5	-	-	-	-	100 %	52%
6	-	-	-	-	-	100 %

**Table 2.**  
Identity scores (%) for Asp1-6 amino acid sequences.