

LUND UNIVERSITY

Placental molecular mechanisms as pathways linking prenatal exposure to ambient air pollution to preeclampsia and fetal growth

Mandakh, Yumjirmaa

2023

Document Version: Publisher's PDF, also known as Version of record

Link to publication

Citation for published version (APA):

Mandakh, Y. (2023). *Placental molecular mechanisms as pathways linking prenatal exposure to ambient air pollution to preeclampsia and fetal growth.* [Doctoral Thesis (compilation), Department of Laboratory Medicine]. Lund University, Faculty of Medicine.

Total number of authors: 1

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights. • Users may download and print one copy of any publication from the public portal for the purpose of private study

or research.

- You may not further distribute the material or use it for any profit-making activity or commercial gain
 You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: https://creativecommons.org/licenses/

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117 221 00 Lund +46 46-222 00 00 Placental molecular mechanisms as pathways linking prenatal exposure to ambient air pollution to preeclampsia and fetal growth

YUMJIRMAA MANDAKH DEPARTMENT OF LABORATORY MEDICINE | FACULTY OF MEDICINE | LUND UNIVERSITY



Placental molecular mechanisms as pathways linking prenatal exposure to ambient air pollution to preeclampsia and fetal growth

Placental molecular mechanisms as pathways linking prenatal exposure to ambient air pollution to preeclampsia and fetal growth

Yumjirmaa Mandakh



DOCTORAL DISSERTATION

Doctoral dissertation for the degree of Doctor of Philosophy (PhD) at the Faculty of Medicine at Lund University to be publicly defended on January 27th, 2023 at 08.00 in Segerfalksalen Hall, BMC Hus A, Sölvegatan 17, 223 62 Lund, Sweden

Faculty opponent Marie Pedersen

Organization: LUND UNIVERSITY

Document name: DOCTORAL DISSERTATION

Author: YUMJIRMAA MANDAKH

Date of issue

Sponsoring organization:

Title and subtitle: Placental molecular mechanisms as pathways linking prenatal exposure to ambient air pollution to preeclampsia and fetal growth

Abstract:

Background: PE is a "multifactorial syndrome" in which ambient air pollution may contribute to the etiology of PE. However, the underlying mechanism of this association is not clearly elucidated. This dissertation aims at understanding the placental molecular mechanism underlying the relationship of prenatal exposure to ambient air pollution with preeclampsia and fetal growth by characterizing and evaluating the mediator role of placental aging using mitochondrial DNA content, telomere length, DNA methylation and gene expression analyses.

Methods: Scania in Sweden and Barcelona in Spain are the two main study areas. Maternal Air Pollution in Southern Sweden (MAPSS) is a population-based cohort consisting of 43,688 singleton pregnancies recorded from 2000 to 2009 in Scania, Sweden. For placental molecular studies, we selected a subcohort of 361 mother-child pairs from the Barcelona Life Study Cohort (BISC), a prospective cohort consisting of 1080 mother-child pairs recruited between 2018 and 2021 as well as 137 participants from unique biobank designed to study PE at Lund University. Each individual-level air pollutant concentrations were estimated either by a high-resolution dispersion model or by a passive sampling methods. Placental mtDNAcn and TL were measured using qPCR. Placental DNAm and gene expression were analyzed with EPIC array and RNA-seq, respectively. Linear, logistic regression and linear mixed effects models were used on SPSS and R statistical programs.

Results: In a setting with fair air quality in Scania, the increased risk of PE and SGA were associated with both linear exposure trend and quartile-specific exposure of ambient particles and NO_x . In a setting with moderate air quality in Barcelona, early pregnancy exposure to ambient NO_2 measured by homeoutdoor and personal sensors were associated with lower birth weight and increased odds of having SGA neonate. Although the mediatory role of placental mtDNAcn and telomere length were not confirmed, prenatal exposure to ambient NO_x and NO_2 affect the placental aging process differently depending on the exposure concentration and exposure period during pregnancy. We observed reduced mtDNAcn and shorter TL in placenta in relation to exposure during first trimester, whereas increased mtDNAcn in relation to NO_2 exposure during third trimester. Prenatal exposure to ambient NO_x during first trimester affected differential placental DNA methylation in PE and NO_x combined groups, resulting in placental age deceleration and showing sexual dimorphism.

Conclusion: Placental senescence and aging may be the underlying molecular mechanism linking the association between prenatal exposure to ambient air pollution and pregnancy complications including preeclampsia and small-for-gestational age foetuses and neonates.

Key words: Ambient air pollution, Placenta, Preeclampsia, Small-for-gestational age, Mitochondrial DNA content, Telomere length, Global DNA methylation, Gene expression, Epigenetic age

| Classification system and/or index terms (if any) | | Supplementary bibliographical information | |
|---|--|---|--|
| Language: English | | ISSN and key title: 1652-8220 | |
| SBN: 978-91-8021-348-6 Lund University, Fa | | culty of Medicine Doctoral Dissertation Series 2023:9 | |
| Recipient's notes | | Number of pages: 56 | |
| Price | | Security classification | |

I, the undersigned, being the copyright owner of the abstract of the above-mentioned dissertation, hereby grant to all reference sources permission to publish and disseminate the abstract of the above-mentioned dissertation.

Signature

М. Нитсириа

Date 2022-12-20

Placental molecular mechanisms as pathways linking prenatal exposure to ambient air pollution to preeclampsia and fetal growth

Yumjirmaa Mandakh



Coverphoto by Louise Larsson

Copyright pp 1-56 Yumjirmaa Mandakh

Paper 1 © 2020 by the authors. International Journal of Environmental Research and Public Health. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https://creativecommons.org/ licenses/by/4.0/)

Paper 2 © 2021 Mandakh, Oudin, Erlandsson, Isaxon, Hansson, Broberg and Malmqvist. Frontiers in Toxicology. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forms is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Paper 3 © 2021 by the authors. Toxics. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https://creativecommons.org/ licenses/by/4.0/)

Paper 4 © by the Authors (Manuscript unpublished)

Faculty of Medicine

Department of Laboratory Medicine

ISBN 978-91-8021-348-6

ISSN 1652-8220

Printed in Sweden by Media-Tryck, Lund University

Lund 2022



Media-Tryck is a Nordic Swan Ecolabel certified provider of printed material. Read more about our environmental work at www.mediatryck.lu.se

MADE IN SWEDEN 📲

To Grigorios

and our Anu, Vasileios and Maximos

Table of Contents

| .11 |
|-----|
| .12 |
| .13 |
| .15 |
| .15 |
| .15 |
| .15 |
| .16 |
| .16 |
| 10 |
| .18 |
| .20 |
| .21 |
| .21 |
| .25 |
| .28 |
| .28 |
| .28 |
| .29 |
| .33 |
| .38 |
| .42 |
| .44 |
| .46 |
| |
| .46 |
| |

Abstract

PE is a "multifactorial syndrome" in which ambient air pollution may contribute to the etiology of PE. However, the underlying mechanism of this association is not clearly elucidated. This dissertation aims at understanding the placental molecular mechanism underlying the relationship of prenatal exposure to ambient air pollution with preeclampsia and fetal growth by characterizing and evaluating the mediator role of placental aging using mitochondrial DNA content, telomere length, DNA methylation and gene expression analyses.

Scania in Sweden and Barcelona in Spain are the two main study areas. Maternal Air Pollution in Southern Sweden (MAPSS) is a population-based cohort consisting of 43,688 singleton pregnancies recorded from 2000 to 2009 in Scania, Sweden. For placental molecular studies, we selected a sub-cohort of 361 mother-child pairs from the Barcelona Life Study Cohort (BiSC), a prospective cohort consisting of 1080 mother-child pairs recruited between 2018 and 2021 as well as 137 participants from unique biobank designed to study PE at Lund University. Each individual-level air pollutant concentrations were estimated either by a high-resolution dispersion model or passive sampling methods. Placental mtDNAcn and TL were measured using qPCR. Placental DNAm and gene expression were analyzed with EPIC array and by RNA-seq, respectively. Linear, logistic regression and linear mixed effects models were used on SPSS and R statistical programs.

In a setting with fair air quality in Scania, the increased risk of PE and SGA were associated with both linear exposure trend and quartile-specific exposure of ambient particles and NO_x. In a setting with moderate air quality in Barcelona, early pregnancy exposure to ambient NO₂ measured by home-outdoor and personal sensors were associated with lower birth weight and increased odds of having SGA neonate, respectively. Although the mediatory role of placental mtDNAcn and telomere length were not confirmed, prenatal exposure to ambient NO_x and NO₂ affect the placental aging process differently depending on the exposure concentration and exposure period during pregnancy. We observed reduced mtDNAcn and shorter TL in placenta in relation to exposure during first trimester, whereas increased mtDNAcn in relation to NO₂ exposure during third trimester. Prenatal exposure to ambient NO_x during first trimester affected differential placental DNA methylation in PE and NO_x combined groups, resulting in placental age deceleration and showing sexual dimorphism.

Placental senescence and aging may be the underlying molecular mechanism linking the association between prenatal exposure to ambient air pollution and pregnancy complications including preeclampsia and small-for-gestational age foetuses and neonates.

Populärvetenskaplig sammanfattning

Luftföroreningar är ett allvarligt miljöproblem som bidrar till förtida dödsfall över hela världen. Luftföroreningar delas in i grova, fina och ultrafina partiklar baserat på deras storlek i förhållande till aerodynamiken. Detta är en liknande process som det fallande äpplet från trädet som inspirerade Newton att upptäcka gravitationen. Storleken på partiklarna har störst betydelse när det gäller hälsoeffekter. Ju mindre partikelstorlek (dvs ultrafina partiklar), desto mer signifikant blir den negativa effekten på hälsan. Tidigare studier har funnit att ultrafina partiklar kan färdas djupt ner till alveolerna i lungan där gasutbytet av inandat syre och utandad koldioxid äger rum.

Dessutom kan det transporteras till det systemiska blodflödet och orsaka inflammation i olika organsystem i vår kropp. Vid det här laget har du förmodligen upptäckt att ultrafina partiklar kan leta sig in i livmodern genom moderkakan. Det har bevisats i djurexperimentella studier, men dess effekt på människors hälsa visas av stora epidemiologiska studier där barn föds för tidigt eller med låg födelsevikt när deras mödrar bodde i de mycket förorenade områdena under graviditeten. Men innan de påverkar det växande barnet i livmodern, påverkar luftföroreningar hälsan hos en gravid kvinna som orsakar graviditetsinducerad hypertoni.

Ungefär en av sex mödrar dör av graviditetsinducerad hypertoni på ett år världen över. Särskilt tonårsmödrar (<20 år gamla) löper större risk än äldre kvinnor i lågoch medelinkomstländer. I min doktorsavhandling undersökte jag risken för havandeskapsförgiftning och småhet hos bebisar i relation till luftföroreningar i Sverige och Spanien. Vi utförde också placenta molekylära studier för att förstå de underliggande biologiska effekterna av denna association.

I en miljö med rättvis luftkvalitet i Skåne var prenatal exponering för luftföroreningar associerad med den ökade risken för havandeskapsförgiftning och spädbarn som är små för graviditetsåldern. I en miljö med måttlig luftkvalitet i Barcelona var tidig graviditetsexponering för omgivande NO₂ uppmätt med sensorer i hemmet och utomhus och personliga sensorer associerad med lägre födelsevikt respektive ökade chanser att få barn små för graviditetsåldern. Prenatal exponering för luftföroreningar påverkar också placentans åldrandeprocess genom DNA-metylering, mitokondriell stress och telomerförkortning.

Sammantaget har resultaten av dessa studier ett antal viktiga implikationer för framtida praktik. De reproduktiva hälsoeffekterna som behandlas i båda studiemiljöerna med rättvis och måttlig luftkvalitet stöder starkt WHO:s rekommendationer för uppdaterade globala gränsvärden för luftkvalitet.

List of Papers

Paper I

Mandakh Y, Rittner R, Flanagan E, Oudin A, Isaxon C, Familari M, Hansson SR, Malmqvist E. (2020) Maternal Exposure to Ambient Air Pollution and Risk of Preeclampsia: A Population-Based Cohort Study in Scania, Sweden. International Journal of Environmental Research and Public Health. 17(5):1744. https://doi.org/10.3390/ijerph17051744

Paper II

Mandakh Y, Oudin A, Erlandsson L, Isaxon C, Hansson SR, Broberg K and Malmqvist E (2021) Association of Prenatal Ambient Air Pollution Exposure With Placental Mitochondrial DNA Copy Number, Telomere Length and Preeclampsia. Front. Toxicol. 3:659407. doi: 10.3389/ftox.2021.659407

Paper III

Engström K, Mandakh Y, Garmire L, Masoumi Z, Isaxon C, Malmqvist E, Erlandsson L, Hansson SR. (2021) Early Pregnancy Exposure to Ambient Air Pollution among Late-Onset Preeclamptic Cases Is Associated with Placental DNA Hypomethylation of Specific Genes and Slower Placental Maturation. Toxics. 9(12):338. https://doi.org/10.3390/toxics9120338

Paper IV

Laura C.Gómez Herrera, Yumjirmaa Mandakh, Dries S. Martens, Tim S. Nawrot, Ebba Malmqvist, Mariona Bustamante, Payam Dadvand, María Dolores Gómez-Roig and Jordi Sunyer. (2022) The mediatory role of placental stress, ageing, and function in the association between air pollution exposure and fetal growth. (in manuscript)

Author's contribution to the papers

Paper I

Yumjirmaa is a sole first author. I contributed to conceptualization, methodology, validation, formal analysis, investigation, writing of original draft preparation, review and editing.

Paper II

Yumjirmaa is a sole first author. I contributed to conceptualization, methodology (laboratory work: placental DNA/RNA extraction and qPCR of relative mitochondrial DNA copy number and telomere length), formal analysis, investigation, writing of original draft, review and editing.

Paper III

Yumjirmaa is a shared first author. I contributed to formal analysis, investigation, methodology (laboratory work: placental DNA/RNA extraction), and writing of original draft, review and editing.

Paper IV

Yumjirmaa is a shared first author. I contributed to formal analysis, investigation, methodology, visualization, writing of original draft, review and editing.

Abbreviations

| BC | Black carbon |
|----------|--|
| BiSC | Barcelona Life Study Cohort |
| BMI | Body Mass Index |
| CEU | Utah Residents with Northern and Western European Ancestry |
| CI | Confidence Interval |
| COVID-19 | Coronavirus Disease |
| CPC | Control Placental Clock |
| CPR | Cerebroplacental Ratio |
| DAG | Directed Acyclic Graph |
| DEGs | Differentially Expressed Genes |
| DMPs | Differentially Methylated Positions |
| DMRs | Differentially Methylated Regions |
| DNAm | DNA Methylation |
| FC | Fold Change |
| 2logFC | Binary Logarithmic Fold Change |
| FDR | False Discovery Rate |
| FGR | Fetal Growth Restriction |
| GA | Gestational Age |
| GO | Gene Ontology |
| IQR | Interquartile Range |
| ISGlobal | Barcelona Institute for Global Health |
| ISSHP | International Society for the Study of Hypertension in Pregnancy |
| KEGG | Kyoto Encyclopedia of Genes and Genomes |
| LBW | Low Birth Weight |
| MAF | Minor Allele Frequency |
| MAPSS | Maternal Air Pollution Southern Sweden |
| mtDNAcn | Mitochondrial DNA Copy Number |
| NO2 | Nitrogen Dioxide |

| NO | Nitric Oxide |
|------------|---|
| NOx | Nitrogen Oxides |
| O3 | Ozone |
| OR | Odds Ratio |
| PC | Principal Component |
| PE | Preeclampsia |
| PM2.5 | Particles with aerodynamic diameters smaller than 2.5 m |
| PM10 | Particles with aerodynamic diameters smaller than 10 m |
| qPCR | Quantitative Real-Time Polymerase Chain Reaction Assay |
| RIN | RNA integrity number |
| RNA-seq | RNA Sequencing |
| ROS | Reactive Oxygen Species |
| Rp | Pearson Correlation |
| Rs | Spearman Correlation |
| R2 | R-Squared (coefficient of determination) |
| SARS-CoV-2 | Severe Acute Respiratory Syndrome Coronavirus 2 |
| SES | Socioeconomic Status |
| SGA | Small-for-gestational age |
| SNP | Single-nucleotide Polymorphism |
| STB | Syncytiotrophoblast |
| SVD | Singular Value Decomposition |
| TF | Transcription Factor |
| TL | Telomere Length |
| UtA | Uterine Artery |
| UA | Umbilical Artery |
| WHO | World Health Organization |

Introduction

Definitions

Preeclampsia

Preeclampsia (PE) is a persistent de novo hypertension occurring at or after 20 weeks of gestation accompanied either with proteinuria or fetal growth restriction or maternal organ dysfunction including HELLP syndrome (Hemolysis, Elevated Liver enzymes, Low Platelets) (Brown et al., 2018). Although it affects 3% to 8% of all pregnancies, PE is one of the three leading causes of direct maternal mortality worldwide ("Geographic variation in the incidence of hypertension in pregnancy. World Health Organization International Collaborative Study of Hypertensive Disorders of Pregnancy," 1988; Ghulmiyyah & Sibai, 2012; Rath & Tsikouras, 2018) It is classified into early-onset and late-onset PE based on its development before or after 34 weeks of gestation (Tranquilli et al., 2013). Although PE was recorded as moderate and severe PE in the Swedish medical birth register and mild and severe PE in the Spanish medical birth register, in this dissertation, total PE as a combination of these classifications were used as an outcome variable. In addition, PE is often associated with adverse birth outcome, such as preterm birth, small-forgestational age neonates and fetal growth restriction.

Small fetus

Low birth weight

Babies born weighing less than 2500 grams is termed as low birth weight (LBW) (WHO, 2004).

Small-for-gestational age

Smallness of fetus and neonates are defined as small-for-gestational-age (SGA) if their birth weight fell below the 10th centile (Figueras et al., 2008).

Fetal growth restriction

Small neonates can be classified into low-risk SGA and true fetal growth restriction (FGR) based on estimated birthweight centile, doppler velocimetry of uterine artery (UtA) and cerebroplacental ratio (CPR) measured at routine third trimester fetal echosonography. The CPR is calculated as the ratio of middle cerebral artery (MCA) to umbilical artery (UA) pulsatility index (PI) values, measured by Doppler ultrasound (Baschat & Gembruch, 2003). Low-risk SGA is diagnosed when small babies present any of the following: SGA in between 3rd-9th centile or CPR doppler index more than 5th centile or UtA pulsatility index less than 95th centile. Those presenting any of the following: SGA less than 3rd centile or CPR doppler index less than 5th centile or UtA pulsatility index more than 95th centile, were classified as true FGR (Figueras & Gratacós, 2014).

The two-stage model of preeclampsia

The early-onset and late-onset PE evolve in different pathways in the first stage and converges on syncytiotrophoblast stress with placental dysfunction at the onset of the maternal syndrome of PE caused by generalized vascular inflammation in the second stage (Redman et al., 2022). The incomplete remodelling of the spiral arteries along with focal oxidative stress in the vicinity of the spiral artery causes malplacentation and later on placental malperfusion in the first stage of early-onset PE with concomitant FGR. In the first stage of late-onset PE, with normal placentation, progressive placental hypoxia and fetal stress along with massive growth of chorionic villous tree persists from the beginning of the second half of pregnancy without any known well-established cause (Redman et al., 2022). The second stage of PE is syncytiotrophoblast stress, apoptosis, autophagy, necrosis and syncytial knots as well as producing large amount of vascular endothelial growth factor-1 (VEGF-1), also known as the soluble fms-like tyrosine kinase-1 (sFlt-1), and placental growth factor (PIGF).

Ambient air pollution

Ambient particulate air pollution is the fourth leading risk factor for attributable deaths for both males and females accounting for 2.92 million and 3.75 million deaths of all female and male deaths in 2019, respectively (Murray et al., 2020).

Nitrogen oxides (NO_x) is a toxic gas in the atmosphere. It includes nitric oxide (NO), a colorless and odorless gas, and nitrogen dioxide (NO_2) , a reddish-brown gas with a strong odor. Total NO₂ concentrations are composed of primary NO₂ emission

from diesel exhaust (up to 25%), secondary NO₂ formed by the reactions of NO and ozone (O3) (70%) and regional background (Casquero-Vera et al., 2019). Particulate matter is a mixture of solid particles and aerosols varying in its aerodynamic diameter: being less than 10 μ m (PM₁₀) and 2.5 μ m (PM_{2.5}). PM_{2.5} is one of the three major and rapidly increasing risk exposures globally in 2019 (Murray et al., 2020). 90% of PM_{2.5} pollution is generated from diesel engines in cities (Bernasconi et al., 2022). Other emission sources include the combustion of coal, oil, and gasoline. WHO 2021 update on global air quality guidelines recommends a maximum level of 5 μ g/m³ for PM_{2.5}, 15 μ g/m³ for PM₁₀ and 10 μ g/m³ for NO₂ for preventing the global community from health effects related to long-term exposure (World Health Organization, 2021).

Oxidative stress has been proposed to be the major mechanism of the adverse health effect of air pollution exposure. Each interquartile range increase in prenatal exposure to ambient PM_{2.5} during the first trimester was statistically significantly associated with increased maternal serum malondialdehyde (MDA), a biomarker of systemic lipid peroxidation, and total antioxidant capacity (Zhang et al., 2020). On the other hand, higher plasma levels of haematocrit, MDA and nitrates during first trimester were statistically significantly associated with the later development of PE in twin pregnancies (Ramiro-Cortijo et al., 2020). In a mouse experimental study, chronic PM_{2.5} exposure induce systemic cellular inflammatory response through Toll-like receptor (TLR)4/NADPH oxidase-dependent mechanism (Kampfrath et al., 2011). Monocytes in the peripheral blood from early-onset PE had higher TLR4 expression when compared with late-onset PE (Romão-Veiga et al., 2020). A recent meta-analysis showed that short-term exposure to ambient PM_{2.5} was associated with increased serum levels of MDA (Li et al., 2020).

PE is a "multifactorial syndrome" in which ambient air pollution may contribute to the etiology of PE (Gyselaers, 2020; Pedersen et al., 2014). Systematic reviews suggest that early pregnancy exposure to ambient air pollution is associated with increased risk of PE (Bai et al., 2020; Pedersen et al., 2014; Yu et al., 2020). However, the underlying mechanism of this association is not clearly elucidated, numerous studies have attempted to explain how external and internal factors influence the risk of hypertensive disorder of pregnancy by using the pregnancy exposome approach (Hu et al., 2020; Robinson & Vrijheid, 2015).

Placental molecular mechanisms underlying the effects of long-term exposure to ambient air pollutants

Placental mitochondrial DNA content

Mitochondria are cell organelles where oxidative phosphorylation to produce adenosine triphosphate occurs. Due to high energy electron transfer during the energy metabolism, free radicals capable of oxidation is contained within mitochondrial double membrane. When mitochondria are under stress, these free radicals are released to the cytoplasm and damage cell molecules causing oxidative stress. Mitochondrial DNA (mtDNA) is susceptible to oxidative stress damage due to its structure of two strands without histone shield (Lee & Wei, 2000). MtDNA copy number (mtDNAcn) vary by each mitochondrion, cell type, tissue and organ. MtDNA is inherited only from mother. Prenatal exposure to ambient air pollution has been found to be associated with reduced placental mtDNAcn, whereas placentas from pregnancies complicated with early-onset PE had increased mtDNAcn when compared with normotensive controls (Clemente et al., 2017; Janssen et al., 2012; Vishnyakova et al., 2016).

Placental telomere length

Human telomeres consists of tandem repeats of "TTAGGG" and are located at the end of each chromosome (Blackburn, 1991). Telomere-driven replicative cell senescence is a critical end point of a cell going to senescence and apoptosis due to telomere shortening from successive cell replication process (Hayflick & Moorhead, 1961; Pańczyszyn et al., 2020). Such high guanine content renders telomeres sensitive to reactive oxygen species (ROS) and thus leads to accelerated telomere shortening (Pańczyszyn et al., 2020; Richter & Zglinicki, 2007). Previous studies investigating the association between air pollution exposure and telomere length showed inconsistent findings. Increase in annual mean exposure to NO_x, NO₂, PM_{2.5} and PM₁₀ were associated with longer telomere (Walton et al., 2016). Maternal exposure to traffic-related air pollution and lack of green environment surrounding the residence were associated with shorter placental telomere length (Bijnens et al., 2015). Telomere was shorter in placentas from pregnancies complicated with PE, FGR and PE with FGR (Tal Biron-Shental et al., 2010).

Placental DNA methylation

DNA methylation (DNAm) is defined as an addition of methyl group at the 5' position of cytosine in CpG dinucleotides (CpGs) in CpG islands located abundantly in gene promoters and repetitive DNA elements in human genome (Rakyan et al., 2011). CpGs in gene-promoters are mostly unmethylated whereas in gene bodies and in repetitive DNA elements are mostly methylated (Suzuki & Bird, 2008). The main roles of DNAm is to regulate gene expression and to protect genome integrity by suppressing transposable element transcription which could otherwise integrate

into new chromosome by translocations (Levin & Moran, 2011). DNAm is the most stable epigenetic marker which is heritable during cell division. It also plays a critical role in cell differentiation and epigenetic reprogramming during embryonic development during first trimester (Collas et al., 2007). Environmental exposures during early pregnancy can potentially induce additional DNAm and alter gene expression resulting in increased susceptibility to disease (Jirtle & Skinner, 2007; Saenen et al., 2019).

The villous syncytiotrophoblast is the major cell type in the placenta (Gude et al., 2004) and has been suggested to explain the hypomethylation of the placental genome across gestation (Bianco-Miotto et al., 2016) due to its low levels of methylated cytosine residues (5'mC) and *de novo* methylation enzymes, namely DNMT 3a and 3b (Fogarty et al., 2015). Furthermore, the underlying differentiating cytotrophoblast cells with methylated cytosine residues (5'mC) constantly fuse to syncytiotrophoblast (Mayhew et al., 1999). Therefore placental DNA methylation has been reported to be largely dependent on the methylation profile of cytotrophoblast cells (Grigoriu et al., 2011). Moreover, number of Hofbauer cells and especially dendritic cell-specific ICAM-grabbing non-integrin (DC-SIGN) expressing Hofbauer cells in the placental villous core were found to be decreased in PE placenta (Tang et al., 2013; Yang et al., 2017). DC-SIGN positive Hofbauer cells are suggested to be involved in the immune tolerance during pregnancy (Reyes & Golos, 2018). Placental DNA methylation may be one of the underlying mechanistic links connecting prenatal exposure to ambient air pollution and PE.

Placental gene expression

DNA methylation along with other epigenetic modifications regulate gene expression (Bianco-Miotto et al., 2016), however, the findings are inconsistent among the following few multi-omics studies that have analyzed both methylome and transcriptome of the same placentas from preeclamptic and normotensive pregnant women. Xuan et al. (2016), Sundrani et al. (2013) and (Blair et al., 2013) have identified negative correlation, whereas Leavey et al. (2018) have found no association between DNA methylation and gene expression. Moreover, inhibin was found to be the only biomarker for preeclampsia by multi-omics approach (Benny et al., 2020). Nevertheless, there are only few studies investigating integrated multi-omics concerning the exposure to ambient air pollution exposure (Alfano et al., 2018). Long term exposure to ambient NO_x was associated with hypomethylation and 9 enriched pathways relating to immune regulation (Plusquin et al., 2017).

Aim

This dissertation aims at understanding the placental molecular mechanism underlying the relationship of prenatal exposure to ambient air pollution with preeclampsia and fetal growth by characterizing and evaluating the mediator role of placental aging using mitochondrial DNA content, telomere length, DNA methylation and gene expression analyses.

Paper I

The major objective of this study was to investigate the association between prenatal exposure to ambient air pollutants (black carbon, NO_x , $PM_{2.5}$, PM_{10}) and preeclampsia in Scania, Sweden

Paper II

The aim of this study was to identify how prenatal exposure to ambient NO_x was associated with preeclampsia by analysing the possible mediation effects of placental relative telomere length and mitochondrial DNA content.

Paper III

This paper investigates placental DNA methylation pattern and gene expression in relation to prenatal exposure to ambient NO_x among preeclamptic cases and normotensive controls.

Paper IV

The main aim of this investigation was to assess the several aspects of fetal growth indicators associated with prenatal exposure to ambient NO_2 measured by personal, home-outdoor and home-indoor sensors as well as to determine whether placental senescence measured by telomere length and mitochondrial DNA content explains the underlying mechanism of the association.

Material and methods

Study population

Scania in Sweden and Barcelona in Spain are the two main study areas. Scania is the southernmost and the second most densely populated province of Sweden with 1.41 million inhabitants in total and 127 people per square kilometre (km²) (Regionfakta, 2022). As can be seen from Figure 1, Scania's largest city, Malmö, ranked at 59 with fair air quality based on its $PM_{2.5}$ annual mean concentration of 8.2 µg/m³ in the past two calendar years (European Environment Agency, 2022).



Air in European cities - from the cleanest to the most polluted

| City name | Country | Rank | Fine particulate matter in ug/m3 | Population in the city | |
|--------------------------|---------|------|----------------------------------|------------------------|---|
| Umeå | Sweden | 1 | 3.1 | 125080 | • |
| Stockholm (greater city) | Sweden | 6 | 4.2 | 1745766 | • |
| Uppsala | Sweden | 7 | 4.2 | 219914 | • |
| Norrköping | Sweden | 11 | 4.9 | 140927 | ٠ |
| Göteborg | Sweden | 19 | 5.8 | 564039 | • |
| Malmö | Sweden | 59 | 8.2 | 333633 | • |

Figure 1. Air quality in Sweden (European Environment Agency, 2022)

Barcelona city is the capital of the autonomous community of Cataluña and the second most densely populated metropolitan city in Spain, with approximately 1.64 million inhabitants in total and 16094 people per km² (Instituto Nacional de Estadística, 2022). Out of 344 European cities, Barcelona ranked at 244 with

moderate air quality based on its $PM_{2.5}$ annual mean concentration of 13.3 μ g/m³ in 2020 and 2021 (Figure 2) (European Environment Agency, 2022).



Air in European cities - from the cleanest to the most polluted

| City name | Country | Rank | Fine particulate matter in ug/m3 | Population in the city | |
|-------------------|---------|------|----------------------------------|------------------------|---|
| Sevilla | Spain | 148 | 9.8 | 884899 | • |
| Jaén | Spain | 151 | 9.9 | 112757 | • |
| Gijón | Spain | 155 | 9.9 | 271717 | • |
| Valencia | Spain | 163 | 10.2 | 1417464 | • |
| Zaragoza | Spain | 176 | 10.5 | 681877 | • |
| Bilbao | Spain | 200 | 11.2 | 797590 | • |
| Valdemoro | Spain | 212 | 11.7 | 77270 | • |
| Toledo | Spain | 215 | 11.8 | 85811 | • |
| Córdoba | Spain | 223 | 12.3 | 326039 | • |
| Alicante/Alacant | Spain | 224 | 12.3 | 337482 | • |
| Torrejón de Ardoz | Spain | 230 | 12.5 | 132853 | • |
| Barcelona | Spain | 244 | 13.3 | 3755512 | • |
| Granada | Spain | 253 | 13.6 | 404809 | • |
| A Coruña | Spain | 259 | 14.0 | 247604 | • |

Figure 2. Air quality in Spain (European Environment Agency, 2022)

All studies followed the ethics conduct and data protection according to the national rules. Ethics approval was obtained from Lund University Ethical Review Board LU803-2 (2015/14) and (Dnr 696/2014) in Paper I, II and III. The Barcelona Life Study Cohort (BiSC) study was approved by the Ethical Review Board of Parc de Salut Mar (CEIm - Parc de Salut Mar) under the project reference number 2018/8050/I ("Pre-natal exposure to AIR pollution and pre- and post-Natal Brain Development") in Paper 4. All participants gave written informed consent.

Paper I

Maternal Air Pollution in Southern Sweden (MAPSS) is a population-based cohort consisting of 43,688 singleton pregnancies recorded from 2000 to 2009 in Scania, Sweden. MAPSS is based on the local birth register, Perinatal Revision Syd, with 98% coverage of all births as well as individual air pollution exposure concentrations modelled at each maternal residential address within the catchment

area of hospitals in Malmö, Lund and Trelleborg (Figure 3). Additional sociodemographic and socioeconomic factors were obtained from Statistics Sweden and linked to MAPSS based on unique personal identification number.



Figure 3. A map of Scania, Sweden

SUS – Skåne University Hospital in Lund and Malmö are marked in red and air quality monitoring stations in Scania are marked in black. Figure created by Kristoffer Mattisson.

Paper II and III

The study population of placental molecular studies, Paper II and III, in Scania were based on the unique biobank data designed to study preeclampsia at the Department of Obstetrics and Gynaecology, Lund University. The study participants were recruited from the catchment areas of Skåne University Hospital in Lund and Malmö (Figure 4).

Paper II was a cross-sectional study of 42 preeclamptic and 95 arbitrarily selected normotensive pregnant women whose prenatal exposure to ambient NO_x were modelled at their residential addresses using the Gaussian plume dispersion modelling in Scania, Sweden.

Study III involved 111 participants in total. For the DNA methylation analysis, between-subjects 2x2 factorial design was employed and therefore subjects were divided into four combined groups based on PE status and prenatal exposure to ambient NO_x: PE cases with low NO_x exposure, PE cases with high NO_x exposure, normotensive controls with low NO_x exposure, and normotensive controls with high NO_x exposure. A total of 17 subjects were included in the analyses of differential gene expression, and therefore we compared PE cases with normotensive controls.



Figure 4. A study area of placental molecular studies in relation to the association between ambient NO_x exposure during pregnancy and preeclampsia in Scania, Sweden Figure created by Kristoffer Mattisson

Paper IV

The Barcelona Life Study Cohort (BiSC) is a prospective cohort consisting of 1080 mother-child pairs recruited between 2018 and 2021 from three collaborating hospitals (Hospital Sant Joan de Déu, Hospital Clínic-La Maternitat and Hospital de Sant Pau i la Santa Creu) (Figure 5). Criteria for selecting the subjects were as follows: aged between 18 and 45 years, had singleton pregnancy and spoke either Catalan, Spanish or English. Eligible women who matched this inclusion criteria were identified by midwives during the first prenatal visit (between 8-14 weeks of gestation). Exclusion criteria consisted of having high-risk pregnancy, complex chronic conditions or mental illness and permanent residence located outside the boundary of Barcelona city. Of the initial cohort of 1080 mother-child pairs, 361 were included in the analysis based on its complete set of data on ambient NO₂ exposure, placental biomarkers, clinical and covariate data.



Figure 5. A map of Barcelona city, Cataluña, Spain Figure created by Kristoffer Mattisson

Air pollution exposure assessment

Unlike the brief description of air pollutants in the introduction, it is necessary to clarify how exposure to air pollution differs from the definition of air pollution. Exposure refers to how much quantity of air pollutant that an individual experience over certain study period. Each individual-level air pollutant concentrations were estimated either by a dispersion model or passive sampling methods in all studies. A Gaussian plume air dispersion model on ENVIMAN software was employed for estimating the individual modelled exposure levels of air pollutants (BC, locally emitted PM_{2.5} and PM₁₀, total PM_{2.5} and PM₁₀, NO_x) of MAPSS birth cohort in Paper I as well as in Paper II and III. Personal exposure to NO₂, home-outdoor and home-indoor NO₂ concentrations were measured for 24 hours and seven days during first and third trimesters by a passive diffusion tube by Gradko Environmental passive dosimeters (Gradko International Ltd., UK) in Paper IV.

To get a glimpse of how air pollution differs between the study settings, Screening for High Emission Reduction Potential on Air (SHERPA) tool allows to understand how the air quality in Malmö and Barcelona are influenced by different emission sources. As shown in Figure 6, a large proportion of total mass of urban PM_{2.5} in Malmö is a transboundary air pollution by spatial allocation as well as traffic and agriculture sectors by sectoral allocation (European Commission Joint Research Centre et al., 2021). In contrast in Barcelona, almost half of urban PM_{2.5} is emitted

in the core city mostly from the industrial and residential sources (Figure 7) (European Commission Joint Research Centre et al., 2021).



Figure 6. Urban PM2.5 spatial and sectoral allocation of Malmö, Sweden (SHERPA v.2.2.0) (European Commission Joint Research Centre et al., 2021)

Emission sources: T – transport, I – industry, A – agriculture, R – Residential, S – shipping, O – other, N – Natural, E – External.



Figure 7. Urban PM_{2.5} spatial and sectoral allocation of Barcelona, Spain (SHERPA v2.2.0) (European Commission Joint Research Centre et al., 2021) Emission sources: T – transport, I – industry, A – agriculture, R – Residential, S – shipping, O – other, N – Natural, E – External.

26

Paper I, II and III

ENVIMAN is a software with several modules. Data management module consists of emission sources data (traffic count, car models, car exhaust, type of tires etc) and meteorological data (wind speed, wind direction, temperature and global solar radiation). Dispersion modelling module estimates concentrations of NO_x , $PM_{2.5}$ and PM_{10} over the 100m x 100m grid in two years 2000 and 2011. Hourly concentrations of NO_x were calculated according to pregnancy length from the date of birth and backwards. Hourly concentrations of $PM_{2.5}$ and PM_{10} were estimated approximately by calendar months, from January till December, and for the one month prior to the conception date. Using R software, the hourly concentrations were aggregated into monthly mean concentrations for the months of 2000 and 2011.

Furthermore, in ArcGIS program, the monthly mean concentrations of 2000 and 2011 were inserted and by interpolation with atmospheric ventilation index, the individual-level pollutant concentrations of years in between (2001-2009) were estimated at each participant's residential address. Atmospheric ventilation index was calculated at the Department of Environment at City of Malmö for each month of this time period. It estimates how well the atmosphere disperses the smoke from its source in the atmospheric mixed layer on any given day. It is a product of the average wind velocity in the mixed layer and the height of the mixed layer.

Detailed description of the Gaussian plume air dispersion model can be found in Rittner et al. (2020). In brief, $PM_{2.5}$ and PM_{10} concentrations were modelled at seven monitoring stations in Scania, as shown in Figure 3 above. For validation, the measured concentrations at the monitoring stations at Malmö City Hall and Vavihill were compared with the modelled concentrations using R^2 Pearson correlation method. Moderate to strong correlation were found with a variable R^2 values between 0.46 and 0.83 for PM_{10} and 0.44 and 0.86 for $PM_{2.5}$. BC was measured for shorter period and it was unable to validate measured values with the modelled values (Rittner et al., 2020).

 NO_x model was adopted from Stroh et al. (2012). There was a good agreement between the measured and modelled concentrations of NO₂ at individual exposure levels ($r_s = 0.8$, p > 0.001).

Paper IV

We measured home-outdoor, home-indoor, and personal NO₂ concentrations for 24 hours and seven days a week around 12th week of gestation during first trimester as well as around 32^{nd} week of gestation during third trimester with Gradko Environmental passive dosimeters (Gradko International Ltd., UK). The home-outdoor measuring tube was installed on the most exposed façade in the window or a balcony. Home-indoor tube was placed near the bedside at participant's home. The personal exposure to NO₂ were measured using passive diffusion tubes installed on

the shoulder strap of a backpack. The study participants were asked to carry the backpack during the week-long measurement period. They were allowed to leave the tube hanging at a minimum height of 1.5 m while they are at home, but instructed to put on the backpack whenever they were going outside. All instructions for proper installation (during COVID-19 lockdown) and usage were given by the trained field workers.

Quality control measures were taken using at least two laboratory blank tubes in each batch when samples were sent to the laboratory. As a reference NO_2 monitor, two simultaneous tubes were installed at the urban background reference station of Palau Reial for a week every month. There was a good agreement between the reference NO_2 monitor and passive tubes ($R^2=0.88$).

Statistical methods

Clinical outcome

Outcome variables including PE and SGA were created as secondary variables in MAPSS on Paper I as well as in Paper II and III in accordance with their definitions.

In Paper IV, the customized birth weight standard for a Spanish population was adopted for estimating birthweight centile (Figueras et al., 2008). For FGR, CPR centiles were estimated using normative reference values developed by Baschat and Gembruch (2003). To following three criteria was adopted to subclassify small babies to low-risk SGA or true FGR: (1) SGA in between 3rd-9th centile or SGA less than 3rd centile; (2) CPR doppler index more than 5th centile or less than 5th centile; and (3) UtA pulsatility index less than 95th centile or more than 95th centile, respectively (Figueras & Gratacós, 2014).

Placental outcome

Paper II and IV

Placental senescence was assessed based on the changes in placental relative mtDNAcn and telomere length. In Study II (N=137 placental samples) and Study IV (N=591 placental samples), placental DNA was extracted from the collected biopsy of placental villious parenchyma using Allprep® DNA/RNA/miRNA Universal kit from Qiagen, following the manufacturer's instructions. Using a real-time quantitative polymerase chain reaction (PCR), the target gene used to determine mtDNAcn was MT-TL1 encoding the Mitochondrially Encoded TRNA-Leu (UUA/G) 1 in Paper II and MT-ND1 encoding the Mitochondrial encoded NADH dehydrogenase subunit 1 in Paper IV. For relative telomere length

quantification, *TEL1* and *TEL2* primers were used for Paper II, whereas in Paper IV, *TelC* and *TelG* primers were used. The ratios of both mtDNAcn and telomere native sequences copy numbers to nuclear single-copy reference genes (haemoglobin beta (*HBB*) in Paper II and haemoglobin subunit gamma gene (*HBG1* and *HBG2* primers) in Paper IV) were determined as relative mtDNAcn and telomere length, respectively.

Paper III

Bisulfite treatment was performed on 500 ng DNA extracted from 111 placental villous parenchyma using EZ-96 DNA Methylation kit v 1.1 (Zymo Research, Irvine, CA, USA). Genome-wide DNA methylation analysis was conducted at the Center for Translational Genomics (CTG), Lund University using Infinium MethylationEPIC BeadChip (Illumina, San Diego, CA, USA) targeting over 850,000 CpG sites. Placental DNA methylation preprocessing and data analysis were performed using the statistical software R and R packages ChAMP, minfi, Planet (placental cell type and epigenetic clock), limma, and missMethyl.

Only 17 extracted RNAs of all placental samples had RIN value over 7. At CTG, total RNA library was prepared using the TruSeq® Stranded mRNA Library Prep (20020594, Illumina, San Diego, CA, USA) and quality control was performed using LabChip DNA High Sensitivity Reagent kit (CLS138948, Perkin Elmer, Waltham, MA, USA) and DNA 1K/12K/Hi Sensitivity Assay LabChip (760517, Perkin Elmer). Using the QuantIT® dsDNA HS Assay Kit (Q33120, Thermo Fischer, Waltham, MA, USA), the library pool was quantified and sequenced as paired-end, 75-bp reads on a NovaSeq 6000 (Illumina). Raw data were processed using the FastQC tool on the bcl2fastq software (Illumina, San Diego, CA, USA) and the reference genome sequence from the Ensembl database, the Human GRCh38 as well as the StringTie software.

Data analysis

IBM SPSS Statistics software (IBM, Chicago, IL, USA), version 25 was used for the statistical analyses of Paper I and II. The statistical software R, version 4.0.208 and 4.2.2 (The R Foundation for Statistical Computing) were used for Paper III and IV.

The continuous dependent variables (mtDNAcn, TL and birthweight) were tested for normality first and in case of skewed distribution, it was log2-transformed. Complete case analysis was applied in all studies by excluding all observations with missing values in the categorical predictors. Based on a priori knowledge, maternal age, maternal pre-pregnancy body mass index (BMI), season of birth and fetal sex were the covariates adjusted for all studies (Bartsch et al., 2016; Casquero-Vera et al., 2019; Iodice et al., 2018; Martens et al., 2017; Pedersen et al., 2014; Phillips et al., 2004; Rudra & Williams, 2005; Silva et al., 2020; Wacker et al., 1998). Directed acyclic graph was employed to select the covariates for adjustment in Paper I (Figure 8). Mothers who had a history of smoking were excluded from the placental molecular studies in Scania due to its known influence on DNA methylation (Lee & Pausova, 2013) as well as its protective effect on PE (Cudmore et al., 2007; Wei et al., 2015). In Barcelona study, we added "smoking anytime during pregnancy" as a covariate for the analysis of association between prenatal NO₂ exposure and fetal growth indicators. Other covariates adjusted for the models, but not exclusively in all studies, include parity, gestational age, maternal country of birth, education level, annual household income, child ethnicity, and maternal comorbidity (diabetes mellitus, gestational diabetes, essential hypertension, and maternal alcohol consumption during pregnancy.



Figure 8. Directed Acyclic Graphs to guide the selection of potential confounders for the association between the ambient air pollution and preeclampsia: (yellow) exposure; (blue with I) outcome; (red dot) common cause; (blue dot) ancestor (Mandakh et al., 2020).

In Paper I, binary logistic regression models were applied in univariate and multivariable analyses for all pollutants during each trimester exposure period as well as entire pregnancy. The effect size was reported in adjusted odds ratio (OR) with 95% confidence interval (CI) for both linear-exposure increase and quartile-specific exposure for all pollutants. In Paper II and III, for the association between prenatal NO_x exposure and risk of PE, adjusted OR with 95% CI were estimated

using logistic regression. For the analysis of prenatal NO_x exposure and placental mtDNAcn or TL, linear regression model with same set of covariates were used to report the regression coefficient with 95% CI as an effect estimate.

In Paper III, to predict gestational age (GA) acceleration or deceleration, "control placental clock" (CPC) was applied using the R package Planet by regressing the predicted GA to the chronological GA (Horvath, 2013). If the residual value is positive, the placenta ages faster than the normal physiological process and therefore accelerating the placental epigenetic clock. In contrast, if it is negative residual value, the placenta ages slower than expected and termed as GA deceleration. Linear regression model was used to determine the GA acceleration/deceleration differences amongst the PE and NO_x combined groups.

Moreover, singular value decomposition (SVD) analysis was performed to identify the covariates, including technical and biological variables, that were associated with the DNAm. ComBat adjustments in ChAMP was applied to remove the technical variations presented in the first and second principal components. From the estimated placental cell type proportions, syncytiotrophoblast and Hofbauer cells were included in the final regression model. In addition, pre-pregnancy BMI, gestational age, fetal sex, and DNA concentration were included as covariates for the final model. Differentially Methylated Positions (DMPs) were reported in Mvalues (log₂ ratio of methylated and unmethylated probes) using linear regression model adjusted with aforementioned covariates. The q-value determined by Benjamin-Hochberg false discovery rate (FDR) method was used for assessing the statistical significance level. In addition, fetal-sex stratified analyses were performed. Furthermore, Interaction network and Gene ontology analyses were performed for genes that had CpGs with q < 0.1 in the DMP analyses using ConsensusPathDB and gometh function in the R package missMethyl. The final gene expression analysis consisted of 18,524 genes. The R packages, limma and Limma voom were used for differentially expressed genes analysis.

In Paper IV, the associations between nitrogen dioxide (NO₂), birthweight, SGA and FGR were estimated using linear and logistic regression models adjusted for covariates. For analysis of association between prenatal NO₂ exposure and placental mtDNAcn and TL, linear mixed effects models were employed by considering the random effects, including hospital of delivery and qPCR plate number.

To assess the mediatory role of placental relative mtDNA content or telomere length in the association between prenatal NO_x exposure and PE or prenatal NO₂ exposure and fetal growth indicators in Paper II and IV, the standard Baron and Kenny (1986) method of mediation analysis was employed. The mediation is tested through three step regressions including, (1) the prenatal NO_x or NO₂ exposure predicting the odds of PE or fetal growth indicators, repectively (2) the prenatal NO_x or NO₂ exposure predicting the mediators, namely, placental relative mtDNAcn or telomere length, and (3) prenatal NO_x or NO₂ exposure and the mediators predicting the clinical outcome, including PE and fetal growth indicators. At first and second steps, the associations should be statistically significant to conduct the third step. If there is complete mediation, prenatal NO_x or NO_2 exposure should not be statistically significant whereas mediators are statistically significant. On the other hand, if there is a partial mediation, the coefficient for prenatal NO_x or NO_2 exposure should be reduced after the mediator is controlled as compared to the coefficient in the first step (Baron & Kenny, 1986).

Summary of results

Paper I

The incidence of PE was 1034 (2.9%) of CCA study population (N = 35,570). Preeclampsia was more prevalent among pregnant women with the following risk factors: obese (BMI \ge 30), gestational diabetes, and both essential and gestational hypertension. Higher proportion of PE cases were observed for Nordic-born women when compared with mothers born in other countries.

As shown in Table 1, each 1 μ g/m³ increase in locally emitted BC during entire pregnancy was associated with increased risk of PE (AOR 2.14; 95% CI: 1.48, 3.09, p < 0.01). Similar patterns were observed for all trimesters as well as for each 5 μ g/m³ increment in locally emitted PM_{2.5} and PM₁₀ at all exposure windows. Early pregnancy exposure seems to be more prominent for the association with PE, as can be seen from all statistically significant associations for all pollutants, including long-range transboundary air pollution shown in the total PM_{2.5} and PM₁₀ results. **Table 1.** Adjusted odds ratios (AOR) from the complete case analysis (CCA) on the effects of maternal exposure to ambient particles on risk for preeclampsia associated with a 1 μ g/m³ increase in the concentration of BC and 5 μ g/m³ increase in the concentration of local and total PM_{2.5} and PM₁₀ (with the intermediate variables^a derived from DAGs) (Mandakh et al., 2020).

| Exposure | Entire pregnancy AOR [§] (95% CI) | First trimester AOR [§] (95% CI) | Second trimester AOR [§] (95% CI) | Third trimester AOR [§] (95% CI) |
|------------------------------------|---|--|---|--|
| Black carbon [†] (n/N) | 935/32341 2.14** (1.48, 3.09) | 970/33544 1.89** (1.34, 2.65) | 976/33475 2.08** (1.45, 2.98) | 982/33606 1.77** (1.28, 2.45) |
| Local PM2.5 [‡] (n/N) | 893/30892 2.74** (1.68, 4.47) | 962/33086 2.01** (1.34, 3.02) | 946/32547 2.11** (1.42, 3.13) | 979/33573 1.78** (1.22, 2.61) |
| Local PM10 [‡] (n/N) | 908/31033 1.50** (1.19, 1.89) | 973/33400 1.31** (1.09, 1.58) | 954/32610 1.32** (1.11, 1.58) | 972/33240 1.32** (1.11, 1.57) |
| Total PM2.5 [‡] (n/N) | 742/25050 1.98** (1.27, 3.09) | 909/31103 1.31* (1.06, 1.61) | 844/28468 1.14 (0.92, 1.42) | 827/27659 1.27* (1.03, 1.56) |
| Total PM10 [‡] (n/N) | 909/31039 1.40** (1.13, 1.72) | 974/33411 1.24** (1.09, 1.41) | 954/32605 1.10 (0.97, 1.26) | 972/33240 1.11 (0.98, 1.24) |

^aIntermediate variables are gestational diabetes, essential hypertension, gestational hypertension.

[†]Estimates are for each 1 µg/m³ increase of black carbon particles. [‡]Estimates are for each 5 µg/m³ increase of local and total PM_{2.5} and PM₁₀. [§]Adjusted for maternal age, body mass index, parity, smoking, diabetes mellitus, gestational diabetes, essential hypertension, gestational hypertension, maternal country of birth, education level, annual household income, fetal sex, year and season of birth. *p-value < 0.05 as compared to the healthy controls. **p-value < 0.01 as compared to the healthy controls.

Moreover, the association between both linear exposure trend and quartile-specific exposure of BC, local $PM_{2.5}$ and PM_{10} , the risk of PE was pronounced when PE was accompanied with SGA (Table 2).

Table 2. Adjusted analysis for PE with SGA vs. PE without SGA in relation to the exposure quartiles of local $PM_{2.5}$ particles ($\mu g/m^3$) during each window of exposure (Mandakh et al., 2020).

| | Adjusted OR [§] (95% CI) | |
|-------------------|-----------------------------------|---------------------|
| Range, µg/m³ | PE with SGA | PE without SGA |
| Entire pregnancy | 193/30192 | 700/27747 |
| Linear (5 µg/m³) | 4.33 (1.56, 12.04)** | 2.56 (1.48, 4.43)** |
| Q1 (0.13 – 0.96) | Reference | Reference |
| Q2 (0.96 – 1.50) | 1.09 (0.69, 1.73) | 1.18 (0.94, 1.49) |
| Q3 (1.50 – 2.08) | 1.42 (0.92, 2.21) | 1.24 (0.98, 1.57) |
| Q4 (2.08 – 7.52) | 1.65 (1.06, 2.58)* | 1.50 (1.19, 1.90)** |
| First trimester | 209/32783 | 761/30148 |
| Linear (5 µg/m³) | 2.57 (1.13, 5.82)* | 1.96 (1.24, 3.10)** |
| Q1 (0.09 – 0.89) | Reference | Reference |
| Q2 (0.89 – 1.42) | 1.19 (0.77, 1.83) | 1.01 (0.81, 1.25) |
| Q3 (1.42 – 2.11) | 1.36 (0.89, 2.09) | 1.21 (0.97, 1.50) |
| Q4 (2.11 – 10.68) | 1.59 (1.03, 2.45)* | 1.32 (1.05, 1.65)* |
| Second trimester | 199/31800 | 747/29230 |
| Linear (5 µg/m³) | 2.87 (1.26, 6.55)* | 2.00 (1.28, 3.11)** |
| Q1 (0.08 – 0.88) | Reference | Reference |
| Q2 (0.88 – 1.40) | 0.77 (0.49, 1.21) | 1.03 (0.82, 1.28) |
| Q3 (1.40 – 2.09) | 1.20 (0.80 1.81) | 1.02 (0.82, 1.28) |
| Q4 (2.09 – 10.5) | 1.29 (0.84, 1.99) | 1.31 (1.04, 1.65)* |
| Third trimester | 206/32800 | 773/30149 |
| Linear (5 µg/m³) | 2.65 (1.19, 5.92)* | 1.65 (1.07, 2.53)* |
| Q1 (0.10 – 0.83) | Reference | Reference |
| Q2 (0.83 – 1.35) | 1.71 (1.10, 2.65)* | 1.20 (0.97, 1.49) |
| Q3 (1.35 – 2.05) | 1.72 (1.10, 2.68)* | 1.26 (1.01, 1.57)* |
| Q4 (2.05 – 9.64) | 1.92 (1.21, 3.05)** | 1.25 (0.99, 1.58) |

Local PM_{2.5} Adjusted OR[§] (95% C

[§]Adjusted for maternal age, body mass index, parity, smoking, diabetes mellitus, gestational diabetes, essential hypertension, gestational hypertension, maternal country of birth, education level, annual household income, fetal sex, year and season of birth. *p-value < 0.05 as compared to the healthy controls. **p-value < 0.01 as compared to the healthy controls.

As shown in Table 3 in Paper I, the association between the exposure to locallyemitted $PM_{2.5}$ during the third trimester and the risk of PE showed a clear doseresponse relationship between quartiles with AOR (95% CI) being 1.27 (1.05, 1.55) at second quartile, 1.32 (1.08, 1.61) at third quartile and 1.36 (1.10, 1.67) at fourth quartile. Similar pattern was observed with local PM_{10} . When comparing the lowest quartile of exposure to the highest quartile of exposure to ambient NO_x , BC, local $PM_{2.5}$ and PM_{10} , the significant association with increased risk of PE were found for all exposure windows.

Paper II

Those exposed to higher NO_x concentrations during the first trimester (> 11.9 μ g/m³) had increased odds of developing PE (OR 4.0; 95% CI: 1.4, 11.1; *p* = 0.008). Also, higher NO_x exposure during entire pregnancy (> 12.0 μ g/m³) was associated with increased odds of PE (OR 3.7; 95% CI: 1.3, 10.4; *p* = 0.012). Those exposed to higher NO_x concentrations during the first trimester (> 11.9 μ g/m³) had reduced placental relative mtDNAcn (-0.20; 95% CI: -0.36, -0.04; *p* = 0.01). Placental relative mtDNAcn did not appear to mediate the association between prenatal exposure to ambient NO_x and PE during first trimester. No association was observed with placental relative TL with prenatal exposure to ambient NO_x and PE.

Paper III

Six DMPs (q < 0.05) were identified when comparing controls with low NO_x exposure to cases with high NO_x exposure. 14 DMPs were identified when comparing PE cases to controls with high NO_x exposure. No differentially expressed genes (q < 0.05) were found.

When comparing PE cases with high NO_x exposure to normotensive controls with low NO_x exposure, hypomethylation of *YAP1* was found. It may play a role in trophoblast fusion and proliferation in PE, thereby potentially also contributing to defective villous maturation and impaired placental function.

For those exposed to higher NO_x concentrations (> 12.1 µg/m³) during early pregnancy, PE cases had more hypomethylated DMP cg26672098, PEBP1 (cancer), MICAL1 (cytokinesis), CD163 (Hofbauer cell in placenta) than controls. This finding is consistent with the previous study in which hypomethylated DMP cg26672098 was found to be associated with early-onset PE.

Placentas of pregnancies with female fetuses presented more DMPs (N = 309) than male-derived placentas (N = 1). Female placentas had more hypomethylated positions than male placentas.

Placentas from PE cases with high NO_x exposure showed significant GA deceleration compared to placentas from controls with low NO_x exposure (beta = -0.61, p = 0.035, model R² = 0.03).



Combination PE status and NOx exposure group

Figure 8. Differences in GA acceleration between combined PE status and NOx exposure groups (Engström et al., 2021)

Paper IV

Each 1 µg/m³ of home-outdoor NO₂ increase was associated with a decrease in birth weight of 6.05 grams (95% CI: -10.4, -1.7, p=0.006) during entire pregnancy and of 5.3 grams (95% CI: -8.9, -1.6, p=0.005) during the first trimester. Moreover, personal NO₂ exposure during first trimester was associated with increased odds of SGA neonate (adjusted OR 1.033; 95% CI: 1.004, 1.062; p=0.02). Home-outdoor NO₂ exposure during third trimester was associated with increased placental relative mtDNAcn (1.003 fold-change, 95% CI: 1.0002, 1.006, p=0.038). Each 1 µg/m³ of home-indoor NO₂ increase was associated with placental relative telomere shortening (0.997 fold-change, 95% CI: 0.995, 0.999, p=0.014). The odds of SGA was associated neither with placental relative mtDNAcn (AOR 1.02; 95% CI: 0.52, 2.03; p = 0.94) nor placental relative TL (AOR 1.43; 95% CI: 0.54, 3.91; p = 0.47).

Discussion

In a setting with fair air quality in Scania, the increased risk of PE and SGA were associated with both linear exposure trend and quartile-specific exposure of ambient particles and NO_x. In a setting with moderate air quality in Barcelona, early pregnancy exposure to ambient NO₂ measured by home-outdoor and personal sensors were associated with lower birth weight and increased odds of having SGA neonate, respectively. Although the mediatory role of placental mtDNAcn and telomere length were not confirmed, prenatal exposure to ambient NO_x and NO₂ affect the placental aging process differently depending on the exposure concentration and exposure period during pregnancy. We observed reduced mtDNAcn and shorter TL in placenta in relation to exposure during first trimester, whereas increased mtDNAcn in relation to NO₂ exposure during third trimester. Prenatal exposure to ambient NO_x during first trimester affected differential placental DNA methylation in PE and NO_x combined groups, resulting in placental age deceleration and showing sexual dimorphism.

Our finding of gestational age deceleration in placentas from PE cases with high NO_x exposure indicated the effect of exposure rather than PE on placental pathology. Previous studies have reported that placentas from pregnancies complicated with PE have been found to be associated with accelerated placental aging (Girchenko et al., 2017; Mayne et al., 2017). In contrast, the exposure to ambient nitrogen dioxide was associated inversely with gestational age acceleration (White et al., 2019), whereas $PM_{2.5}$ exposure was positively associated with circadian pathway genes (Nawrot et al., 2018). Growing body of literature on molecular study of placenta found differential DNAm in PE placentas (Cirkovic et al., 2020) as well as in relation to the exposure to ambient NO₂ during first and second trimesters was associated with hypomethylation of two CpGs located at *ADORA2B* (A_{2B} adenosine receptor), a gene suggested to be involved in trophoblast proliferation and invasion in PE (Abraham et al., 2018; Darashchonak et al., 2014).

Placental DNAm findings display sexual dimorphism in the occurrence of PE in relation to early pregnancy exposure to ambient NO_2 in which placentas of pregnancies with a female fetus had more hypomethylated positions than male placentas. In singleton pregnancies, pregnancies with female fetus had higher incidence of PE when compared with those carrying male fetus (Shiozaki et al., 2011). Sexual dimorphic differences in PE have been confirmed by another meta-

analysis, suggesting that preterm PE being more common among pregnancies carrying female fetus than pregnancies with male fetus (Collaboration: et al., 2016).

Returning to the hypothesis posed at the beginning of this dissertation, it is now possible to state that placental aging may be the underlying molecular mechanism linking the association between prenatal exposure to ambient air pollution and pregnancy complications including preeclampsia and small-for-gestational age foetuses and neonates. Due to oxidative stress induced by ambient air pollution, placental DNA differential methylation may lead to syncytiotrophoblast stress whereas changes in placental mitochondrial DNA content and short telomeres trigger placental senescence (Redman et al., 2022; Sultana et al., 2018). Trophoblast senescence is a normal physiologic phenomenon and is expected to advance gradually over the course of pregnancy until full term. However, under prolonged stressors including mitochondrial stress and telomere shortening, increased premature trophoblast senescence were observed for PE and FGR (T. Biron-Shental et al., 2010; Sukenik-Halevy et al., 2016). Taken together, these mechanisms lead to perturbance in placental aging that may contribute to the etiology of preeclampsia and impaired fetal growth.

Several assumptions on confounding must be addressed when linking prenatal exposure to ambient air pollution to preeclampsia and fetal growth. Systematic reviews reported that cigarette smoking during pregnancy is associated with lower incidence of preeclampsia compared to non-smokers (Conde-Agudelo et al., 1999; Wei et al., 2015). One experimental study indicated that nicotine restored the function of human umbilical vein endothelial cells treated with soluble fms-like tyrosine kinase 1 (sFlt1) and soluble endoglin (sEng), the anti-angiogenic factors that induce endothelial dysfunction in PE (Armaly et al., 2018; Mimura et al., 2010). Another possible explanation is that carbon monoxide (CO) in the cigarette smoke lowers sFlt1 and sEng in endothelial cells through heme oxygenase-1 and its metabolite CO pathway (Cudmore et al., 2007). Prenatal chronic exposure to cigarette smoke has been associated with altered DNAm in placental genes encoding hypoxia and oxidative stress pathways (Suter et al., 2011).

Season of birth seems to be another important confounder for the association between air pollution exposure and placental molecular mechanisms as well as preeclampsia and birth weight. In Spain, NO₂ and NO_x ambient concentrations present clear seasonal trends with lower levels during summer compared to winter, partially due to tropospheric ozone formation based on warmer temperatures, extended sunlight and higher emissions of NO_x from primary emission sources including diesel vehicles (Casquero-Vera et al., 2019; Silva et al., 2020). Full-term human placental gene expression analysis found increased expression of circadian clock, vitamin-D receptor and hypoxia-related genes during winter births (Clarkson-Townsend et al., 2020). Birth weight is significantly associated with season of birth indicating higher mean birth weight in the summer months and low birth weight in February (Day et al., 2015). Seasonal variation is also observed in the incidence

rates of hypertensive disorders of pregnancy, particularly increased sunlight before delivery was associated with decreased rates of pregnancy hypertension (Algert et al., 2010).

To study the association between preeclampsia and prenatal exposure to air pollution, central monitoring data have been used to model each individual exposure using the dispersion model. The problem with using the central monitoring data is high spatial variation of air pollutants over a short distance. Within the European Study of Cohorts for Air Pollution Effects (ESCAPE), dispersion model estimates of ambient NO₂ at residence level had a good agreement with land use regression model estimates, probably due to NO₂ traffic sources in the urban areas contributing to small-scale spatial variation (de Hoogh et al., 2014). However, dispersion model estimates for ambient PM_{2.5} and PM₁₀ had less agreement with land use regression estimates, suggesting difference in effect size in epidemiological studies employing these models (de Hoogh et al., 2014). Considering this aspect, modelled NO_x concentrations are robust, while the results of prenatal PM_{2.5} and PM₁₀ therefore need to be interpreted with caution in Paper I.

Gradkosampler tend to overestimate ambient NO_2 concentrations by 10% to 30% when compared with chemiluminescence analyzers due to interconversion between gas-phase NO_x species in the diffusion tube body (Bush et al., 2001; Heal et al., 1999; Yu et al., 2008). In addition to personal sensor measurement by passive sampling tube, time-activity data was collected with follow-up questionnaire as well as GPS tracking of movement collected with ExpoApp. The limitations concerning this aspect was that it was prone to recall bias and the participants did not answer the follow-up questionnaire completely.

Further research needs to investigate more closely the links between exogenous NO_x exposure and endogenous nitric oxide (NO) generation in pregnancies complicated with PE and FGR. NO is synthesized physiologically by the nitric oxide synthases (NOS) that catalyse the oxidation of L-arginine to L-citrulline and NO (Michel & Vanhoutte, 2010; Palmer et al., 1987). Endothelial NOS (eNOS) is expressed excessively on placental villous vessel endothelium as well as in syncytiotrophoblast of placentas of pregnancies complicated with PE and FGR (Myatt et al., 1997). Moreover, as a marker of eNOS, the total nitrite (NO₂) and nitrate (NO₃) concentrations were significantly increased in umbilical venous serum of PE group when compared with normotensive controls, but not in maternal circulation (Lyall et al., 1995). Oxidative stress and smoking among other factors can down-regulate the release of NO by the endothelium (Michel & Vanhoutte, 2010). In state of increased levels of ROS, NO reacts with superoxide (O_2) and generates NO-derived free radicals, such as peroxynitrite (ONOO), leading to further degradation on bioavailable NO and endothelial dysfunction by nitrosative stress (Leiva et al., 2016). In vitro study on metabolic fate of exogenous NO_x on human placental homogenates found higher NO_x concentrations and the formation of iron nitrosyl (FeNO) species in placental villous tissue collected from pregnancies complicated with FGR (Mukosera et al., 2020). It would be interesting to assess the effects of ambient NO_x or NO_2 exposure on both oxidative and nitrosative stress in normal and pathologic pregnancies complicated with PE and FGR.

Taken together, the findings of these studies have a number of important implications for future practice. The reproductive health effects addressed at both study settings with fair and moderate air quality strongly support the WHO recommendations on updated global air quality limit values.

Conclusions

In a setting with fair air quality in Scania, Sweden, prenatal exposure to ambient air pollution was associated with increased risk of developing PE with or without SGA;

- The increased risk of PE associated with the linear exposure trend of ambient particles indicates the importance of reducing PM levels.
- Ambient NO_x exposure during the first trimester was associated with reduced placental relative mtDNAcn and an increased risk of PE. Placental relative mtDNAcn did not appear to mediate the association between prenatal exposure to ambient NO_x and PE.
- Early pregnancy exposure to ambient NO_x affected differential placental DNA methylation in PE and NO_x combined groups, resulting in placental immaturity and showing sexual dimorphism.

In contrast, in a setting with moderate air quality in Barcelona, Spain, prenatal exposure to ambient NO_2 was associated with changes in fetal growth indicators as well as in placental biomarkers;

- For home-outdoor sensor, each $1 \ \mu g/m^3$ of ambient NO₂ increase was associated with a decrease in birth weight of 6.05 grams during entire pregnancy and of 5.3 grams during the first trimester.
- Personal NO₂ exposure during first trimester was associated with increased odds of SGA neonate.
- Placental relative mtDNAcn was positively associated with home-outdoor NO₂ exposure during third trimester.
- placental relative telomere length was inversely associated with homeindoor NO₂ exposure during first trimester.

Premature placental senescence and placental aging may be the underlying molecular mechanism linking the association between prenatal exposure to ambient air pollution and pregnancy complications including preeclampsia and small-forgestational age foetuses and neonates. Due to oxidative stress induced by ambient air pollution, placental DNA differential methylation may lead to syncytiotrophoblast stress whereas changes in placental mitochondrial DNA content and short telomeres trigger placental senescence. Taken together, these mechanisms may contribute to the etiology of preeclampsia and impaired fetal growth.

Considerably more work needs to be done to determine the dose-response relationship of exposure concentrations and the risk of PE and adverse birth outcome to further explore the significant differences in molecular dynamics of placental epigenetic and toxicity markers. Also, more information on both oxidative and nitrosative stress in the placenta would help us to establish a greater degree of accuracy on the association between ambient NO_x and PE or SGA.

Acknowledgments

First and foremost, I would like to express my sincere gratitude to Erasmus Mundus Joint Doctorate in FetalMed-PhD Programme for the opportunity to be a part of this fellowship. An immense thank you to EM FetalMed-PhD Management Team: Dr. Eduard Gratacós, Dr. Stefan Hansson, Anette Saltin, and Maite Aguilera.

It has been an honour to be the recipient of EMJD FetalMed PhD Program Fellowship with support of the Erasmus + Programme of the European Union (Framework Agreement number: 2013-0040) as well as co-fund by ReproUnion collaborative study co-financed by the European Union, Interreg V ÖKS. I am indebted to the funders in providing me complete academic freedom in this research.

I want to give my deepest appreciation to my main supervisor Ebba Malmqvist for providing valuable guidance and feedback, reviewing my progress constantly, and challenging me to grow as a scientist during my PhD journey. I would also like to extend my sincere gratitude to my co-supervisors at Lund University, Dr. Stefan Hansson, Anna Oudin, and Christina Isaxon for all the guidance, support and outstanding feedback. A special thanks to Dr. Lola Gómez, my host supervisor at Hospital Sant Joan de Déu, University of Barcelona, and Mariona Bustamante, my mentor at Barcelona Institute for Global Health (ISGlobal) for your unlimited support and unconditional guidance during my 6-month mobility period of EMJD FetalMed PhD Programme in Barcelona, Spain.

It has been an amazing experience working with many different research groups across different departments and countries. I am grateful to all at Ergonomics and Aerosol Technology as well as Division of Occupational and Environmental Medicine for your inclusivity and support when I started my PhD study at Lund University. Thanks to Lena and Eva at Department of Obstetrics and Gyneacology for all your support on placental DNA/RNA extraction. I would also like to extend my thanks to Daniella and Helena at Genotoxicity lab, Division of Occupational and Environmental Medicine for your unlimited guidance and support on conducting qPCR of mitochondrial DNA content and telomere length as well as being a great company in and out of the lab. In addition, I want to thank Cecilia and Srinivas at Center for Translational Genomics for your support and guidance throughout the placental DNA methylation project. A special thanks to all my co-authors for all your support and input, in particular Karin Broberg and Karin Engström from whom I have learned so much. Thank you to my research group: Emilie, Kristoffer, Ralf and Erin, for always being there when I needed your support and for being a great company in and out of the office.

I am grateful to everyone I have collaborated with during my mobility in Barcelona: BCNatal Fetal Medicine Research Center and Barcelona Life Study Cohort (BiSC) project team at ISGlobal: your friendship and support have added a lot to my life. A special thanks to Lina Youssef, Laura C. Gómez Herrera, Marta Cosin and Mireia Gascon for being so supportive at all times when I needed your help.

Finally, I am always thankful to my parents, Mandakh and Tsolmon, and my sisters, Tserenlkham and Urjinlkham, for their love and support. Most importantly, I would like to thank my incredible husband, Grigorios, and our Anu Marina, Vasileios Khanui and Maximos Khulegu. You are my everything.

List of Tables

| Table 1. Adjusted odds ratios (AOR) from the complete case analysis | 34 |
|---|----|
| Table 2. Adjusted analysis for PE with SGA vs.). | 35 |

List of figures

| Figure 1. Air quality in Sweden (European Environment Agency, 2022) | 21 |
|---|----|
| Figure 2. Air quality in Spain (European Environment Agency, 2022) | 22 |
| Figure 3. A map of Scania, Sweden | 23 |
| Figure 4. A study area of placental molecular studies in relation to the association between | |
| ambient NO _x exposure during pregnancy and preeclampsia in Scania, Sweden | 24 |
| Figure 5. A map of Barcelona city, Cataluña, Spain | 25 |
| Figure 6. Urban PM2.5 spatial and sectoral allocation of Malmö, Sweden (SHERPA v.2.2.0) | |
| (European Commission Joint Research Centre et al., 2021) | 26 |
| Figure 7. Urban PM _{2.5} spatial and sectoral allocation of Barcelona, Spain (SHERPA v2.2.0) | |
| (European Commission Joint Research Centre et al., 2021) | 26 |
| Figure 8. Directed Acyclic Graphs to guide the selection of potential confounders for the | |
| association between the ambient air pollution and preeclampsia: (yellow) exposure; (blue with I) | |
| outcome; (red dot) common cause; (blue dot) ancestor (Mandakh et al., 2020) | 30 |

References

- Abraham, E., Rousseaux, S., Agier, L., Giorgis-Allemand, L., Tost, J., Galineau, J., Hulin, A., Siroux, V., Vaiman, D., Charles, M. A., Heude, B., Forhan, A., Schwartz, J., Chuffart, F., Bourova-Flin, E., Khochbin, S., Slama, R., & Lepeule, J. (2018).
 Pregnancy exposure to atmospheric pollution and meteorological conditions and placental DNA methylation. *Environ Int*, *118*, 334-347. https://doi.org/10.1016/j.envint.2018.05.007
- Alfano, R., Herceg, Z., Nawrot, T. S., Chadeau-Hyam, M., Ghantous, A., & Plusquin, M. (2018). The Impact of Air Pollution on Our Epigenome: How Far Is the Evidence? (A Systematic Review). *Curr Environ Health Rep*, 5(4), 544-578. <u>https://doi.org/10.1007/s40572-018-0218-8</u>
- Algert, C. S., Roberts, C. L., Shand, A. W., Morris, J. M., & Ford, J. B. (2010). Seasonal variation in pregnancy hypertension is correlated with sunlight intensity. *Am J Obstet Gynecol*, 203(3), 215.e211-215.e215. <u>https://doi.org/https://doi.org/10.1016/j.ajog.2010.04.020</u>
- Armaly, Z., Jadaon, J. E., Jabbour, A., & Abassi, Z. A. (2018). Preeclampsia: Novel Mechanisms and Potential Therapeutic Approaches [Review]. *Front Physiol*, 9. <u>https://doi.org/10.3389/fphys.2018.00973</u>
- Bai, W., Li, Y., Niu, Y., Ding, Y., Yu, X., Zhu, B., Duan, R., Duan, H., Kou, C., Li, Y., & Sun, Z. (2020). Association between ambient air pollution and pregnancy complications: A systematic review and meta-analysis of cohort studies. *Environmental Research*, 185, 109471. <u>https://doi.org/https://doi.org/10.1016/j.envres.2020.109471</u>
- Baron, R. M., & Kenny, D. A. (1986). The moderator-mediator variable distinction in social psychological research: conceptual, strategic, and statistical considerations. J Pers Soc Psychol, 51(6), 1173-1182. <u>https://doi.org/10.1037//0022-3514.51.6.1173</u>
- Bartsch, E., Medcalf, K. E., Park, A. L., & Ray, J. G. (2016). Clinical risk factors for preeclampsia determined in early pregnancy: systematic review and meta-analysis of large cohort studies. *Bmj*, 353, i1753. <u>https://doi.org/10.1136/bmj.i1753</u>
- Baschat, A. A., & Gembruch, U. (2003). The cerebroplacental Doppler ratio revisited. Ultrasound in Obstetrics & Gynecology, 21(2), 124-127. <u>https://doi.org/https://doi.org/10.1002/uog.20</u>
- Benny, P. A., Alakwaa, F. M., Schlueter, R. J., Lassiter, C. B., & Garmire, L. X. (2020). A review of omics approaches to study preeclampsia. *Placenta*, 92, 17-27. <u>https://doi.org/10.1016/j.placenta.2020.01.008</u>

- Bernasconi, S., Angelucci, A., & Aliverti, A. (2022). A Scoping Review on Wearable Devices for Environmental Monitoring and Their Application for Health and Wellness. Sensors (Basel), 22(16). <u>https://doi.org/10.3390/s22165994</u>
- Bianco-Miotto, T., Mayne, B. T., Buckberry, S., Breen, J., Lopez, C. M. R., & Roberts, C. T. (2016). Recent progress towards understanding the role of DNA methylation in human placental development. *Reproduction*, 152(1), R23. <u>https://doi.org/10.1530/rep-16-0014</u>
- Bijnens, E., Zeegers, M. P., Gielen, M., Kicinski, M., Hageman, G. J., Pachen, D., Derom, C., Vlietinck, R., & Nawrot, T. S. (2015). Lower placental telomere length may be attributed to maternal residential traffic exposure; a twin study. *Environ Int*, 79, 1-7. <u>https://doi.org/10.1016/j.envint.2015.02.008</u>
- Biron-Shental, T., Sukenik-Halevy, R., Sharon, Y., Goldberg-Bittman, L., Kidron, D., Fejgin, M. D., & Amiel, A. (2010). Short telomeres may play a role in placental dysfunction in preeclampsia and intrauterine growth restriction. *Am J Obstet Gynecol*, 202(4), 381.e381-387. <u>https://doi.org/10.1016/j.ajog.2010.01.036</u>
- Biron-Shental, T., Sukenik-Halevy, R., Sharon, Y., Goldberg-Bittman, L., Kidron, D., Fejgin, M. D., & Amiel, A. (2010). Short telomeres may play a role in placental dysfunction in preeclampsia and intrauterine growth restriction. *Am J Obstet Gynecol*, 202(4), 381.e381-381.e387. https://doi.org/https://doi.org/10.1016/j.ajog.2010.01.036
- Blackburn, E. H. (1991). Structure and function of telomeres. *Nature*, *350*(6319), 569-573. <u>https://doi.org/10.1038/350569a0</u>
- Blair, J. D., Yuen, R. K. C., Lim, B. K., McFadden, D. E., Von Dadelszen, P., & Robinson, W. P. (2013). Widespread DNA hypomethylation at gene enhancer regions in placentas associated with early-onset pre-eclampsia. *Molecular Human Reproduction*, 19(10), 697-708. <u>https://doi.org/10.1093/molehr/gat044</u>
- Brown, M. A., Magee, L. A., Kenny, L. C., Karumanchi, S. A., McCarthy, F. P., Saito, S., Hall, D. R., Warren, C. E., Adoyi, G., & Ishaku, S. (2018). The hypertensive disorders of pregnancy: ISSHP classification, diagnosis & management recommendations for international practice. *Pregnancy Hypertens*, 13, 291-310. <u>https://doi.org/10.1016/j.preghy.2018.05.004</u>
- Bush, T., Smith, S., Stevenson, K., & Moorcroft, S. (2001). Validation of nitrogen dioxide diffusion tube methodology in the UK. *Atmospheric Environment*, 35(2), 289-296. <u>https://doi.org/https://doi.org/10.1016/S1352-2310(00)00172-2</u>
- Casquero-Vera, J. A., Lyamani, H., Titos, G., Borrás, E., Olmo, F. J., & Alados-Arboledas, L. (2019). Impact of primary NO2 emissions at different urban sites exceeding the European NO2 standard limit. *Science of The Total Environment*, 646, 1117-1125. <u>https://doi.org/https://doi.org/10.1016/j.scitotenv.2018.07.360</u>
- Cirkovic, A., Garovic, V., Milin Lazovic, J., Milicevic, O., Savic, M., Rajovic, N., Aleksic, N., Weissgerber, T., Stefanovic, A., Stanisavljevic, D., & Milic, N. (2020).
 Systematic review supports the role of DNA methylation in the pathophysiology of preeclampsia: a call for analytical and methodological standardization. *Biol Sex Differ*, 11(1), 36. <u>https://doi.org/10.1186/s13293-020-00313-8</u>

- Clarkson-Townsend, D. A., Kennedy, E., Everson, T. M., Deyssenroth, M. A., Burt, A. A., Hao, K., Chen, J., Pardue, M. T., & Marsit, C. J. (2020). Seasonally variant gene expression in full-term human placenta. *Faseb j*, 34(8), 10431-10442. <u>https://doi.org/10.1096/fj.202000291R</u>
- Clemente, D., Casas, M., Janssen, B., Lertxundi, A., Santa Marina, L., Iñiguez, C., Llop, S., Sunyer, J., Guxens, M., Nawrot, T., & Vrijheid, M. (2017). Prenatal ambient air pollution exposure, infant growth and placental mitochondrial DNA content in the INMA birth cohort. *Environmental Research*, 157, 96-102. <u>https://doi.org/10.1016/j.envres.2017.05.018</u>
- Collaboration:, O. b. o. t. G. P., Schalekamp-Timmermans, S., Arends, L. R., Alsaker, E., Chappell, L., Hansson, S., Harsem, N. K., Jälmby, M., Jeyabalan, A., Laivuori, H., Lawlor, D. A., Macdonald-Wallis, C., Magnus, P., Myers, J., Olsen, J., Poston, L., Redman, C. W., Staff, A. C., Villa, P., . . . Steegers, E. A. (2016). Fetal sex-specific differences in gestational age at delivery in pre-eclampsia: a meta-analysis. *International Journal of Epidemiology*, 46(2), 632-642. <u>https://doi.org/10.1093/ije/dyw178</u>
- Collas, P., Noer, A., & Timoskainen, S. (2007). Programming the genome in embryonic and somatic stem cells [https://doi.org/10.1111/j.1582-4934.2007.00079.x]. Journal of Cellular and Molecular Medicine, 11(4), 602-620. https://doi.org/https://doi.org/10.1111/j.1582-4934.2007.00079.x
- Conde-Agudelo, A., Althabe, F., Belizán, J. M., & Kafury-Goeta, A. C. (1999). Cigarette smoking during pregnancy and risk of preeclampsia: A systematic review. Am J Obstet Gynecol, 181(4), 1026-1035. <u>https://doi.org/https://doi.org/10.1016/S0002-9378(99)70341-8</u>
- Cudmore, M., Ahmad, S., Al-Ani, B., Fujisawa, T., Coxall, H., Chudasama, K., Devey, L. R., Wigmore, S. J., Abbas, A., Hewett, P. W., & Ahmed, A. (2007). Negative regulation of soluble Flt-1 and soluble endoglin release by heme oxygenase-1. *Circulation*, 115(13), 1789-1797. <u>https://doi.org/10.1161/circulationaha.106.660134</u>
- Darashchonak, N., Koepsell, B., Bogdanova, N., & von Versen-Höynck, F. (2014). Adenosine A2Breceptors induce proliferation, invasion and activation of cAMP response element binding protein (CREB) in trophoblast cells. *BMC Pregnancy Childbirth*, 14(1), 2. <u>https://doi.org/10.1186/1471-2393-14-2</u>
- Day, F. R., Forouhi, N. G., Ong, K. K., & Perry, J. R. B. (2015). Season of birth is associated with birth weight, pubertal timing, adult body size and educational attainment: a UK Biobank study. *Heliyon*, 1(2), e00031. <u>https://doi.org/https://doi.org/10.1016/j.heliyon.2015.e00031</u>
- de Hoogh, K., Korek, M., Vienneau, D., Keuken, M., Kukkonen, J., Nieuwenhuijsen, M. J., Badaloni, C., Beelen, R., Bolignano, A., Cesaroni, G., Pradas, M. C., Cyrys, J., Douros, J., Eeftens, M., Forastiere, F., Forsberg, B., Fuks, K., Gehring, U., Gryparis, A., . . . Bellander, T. (2014). Comparing land use regression and dispersion modelling to assess residential exposure to ambient air pollution for epidemiological studies. *Environment International*, *73*, 382-392. https://doi.org/https://doi.org/10.1016/j.envint.2014.08.011

- Engström, K., Mandakh, Y., Garmire, L., Masoumi, Z., Isaxon, C., Malmqvist, E., Erlandsson, L., & Hansson, S. R. (2021). Early Pregnancy Exposure to Ambient Air Pollution among Late-Onset Preeclamptic Cases Is Associated with Placental DNA Hypomethylation of Specific Genes and Slower Placental Maturation. *Toxics*, 9(12), 338. <u>https://www.mdpi.com/2305-6304/9/12/338</u>
- European Commission Joint Research Centre, Wilson, J., Bessagnet, B., Pisoni, E., Thunis, P., De Meij, A., Vignati, E., & Mascherpa, A. (2021). Urban PM2.5 atlas : air quality in European cities : 2021 report. Publications Office of the European Union. https://doi.org/doi/10.2760/356670
- European Environment Agency, E. (2022). *European city air quality viewer*. Retrieved 2022/11/27 from <u>https://www.eea.europa.eu/themes/air/urban-air-quality/european-city-air-quality-viewer</u>
- Figueras, F., & Gratacós, E. (2014). Update on the Diagnosis and Classification of Fetal Growth Restriction and Proposal of a Stage-Based Management Protocol. *Fetal Diagn Ther*, 36(2), 86-98. <u>https://doi.org/10.1159/000357592</u>
- Figueras, F., Meler, E., Iraola, A., Eixarch, E., Coll, O., Figueras, J., Francis, A., Gratacos, E., & Gardosi, J. (2008). Customized birthweight standards for a Spanish population. *Eur J Obstet Gynecol Reprod Biol*, 136(1), 20-24. https://doi.org/10.1016/j.ejogrb.2006.12.015
- Fogarty, N., Burton, G., & Ferguson-Smith, A. (2015). Different epigenetic states define syncytiotrophoblast and cytotrophoblast nuclei in the trophoblast of the human placenta. *Placenta*, *36*, 796-802.
- Geographic variation in the incidence of hypertension in pregnancy. World Health Organization International Collaborative Study of Hypertensive Disorders of Pregnancy. (1988). *Am J Obstet Gynecol*, *158*(1), 80-83.
- Ghulmiyyah, L., & Sibai, B. (2012). Maternal mortality from preeclampsia/eclampsia. *Semin Perinatol*, *36*(1), 56-59. <u>https://doi.org/10.1053/j.semperi.2011.09.011</u>
- Girchenko, P., Lahti, J., Czamara, D., Knight, A. K., Jones, M. J., Suarez, A., Hämäläinen, E., Kajantie, E., Laivuori, H., Villa, P. M., Reynolds, R. M., Kobor, M. S., Smith, A. K., Binder, E. B., & Räikkönen, K. (2017). Associations between maternal risk factors of adverse pregnancy and birth outcomes and the offspring epigenetic clock of gestational age at birth. *Clin Epigenetics*, *9*, 49. <u>https://doi.org/10.1186/s13148-017-0349-z</u>
- Grigoriu, A., Ferreira, J. C., Choufani, S., Baczyk, D., Kingdom, J., & Weksberg, R. (2011). Cell specific patterns of methylation in the human placenta. *Epigenetics*, 6(3), 368-379. <u>https://doi.org/10.4161/epi.6.3.14196</u>
- Gude, N., Roberts, C., Kalionis, B., & King, R. (2004). Growth and function of the normal human placenta. *Thrombosis Research*, 114, 397-407. <u>https://www.thrombosisresearch.com/article/S0049-3848(04)00342-1/fulltext</u>
- Gyselaers, W. (2020). Preeclampsia Is a Syndrome with a Cascade of Pathophysiologic Events. *Journal of clinical medicine*, 9(7), 2245. <u>https://doi.org/10.3390/jcm9072245</u>
- Hayflick, L., & Moorhead, P. S. (1961). The serial cultivation of human diploid cell strains. *Exp Cell Res*, 25, 585-621. <u>https://doi.org/10.1016/0014-4827(61)90192-6</u>

- Heal, M. R., O'Donoghue, M. A., & Cape, J. N. (1999). Overestimation of urban nitrogen dioxide by passive diffusion tubes: a comparative exposure and model study. *Atmospheric Environment*, 33(4), 513-524. https://doi.org/https://doi.org/10.1016/S1352-2310(98)00290-8
- Horvath, S. (2013). DNA methylation age of human tissues and cell types. *Genome Biology*, 14(10), 3156. <u>https://doi.org/10.1186/gb-2013-14-10-r115</u>
- Hu, H., Zhao, J., Savitz, D. A., Prosperi, M., Zheng, Y., & Pearson, T. A. (2020). An external exposome-wide association study of hypertensive disorders of pregnancy. *Environ Int*, 141, 105797. <u>https://doi.org/10.1016/j.envint.2020.105797</u>
- Instituto Nacional de Estadística, I.-S. (2022). *Continuous Register Statistics* Retrieved 2022/11/24 from <u>https://www.ine.es/dyngs/INEbase/en/operacion.htm?c=Estadistica_C&cid=1254736</u> <u>177012&menu=ultiDatos&idp=1254734710990</u>
- Iodice, S., Hoxha, M., Ferrari, L., Carbone, I. F., Anceschi, C., Miragoli, M., Pesatori, A. C., Persico, N., & Bollati, V. (2018). Particulate Air Pollution, Blood Mitochondrial DNA Copy Number, and Telomere Length in Mothers in the First Trimester of Pregnancy: Effects on Fetal Growth. Oxid Med Cell Longev, 2018, 5162905. <u>https://doi.org/10.1155/2018/5162905</u>
- Janssen, B. G., Munters, E., Pieters, N., Smeets, K., Cox, B., Cuypers, A., Fierens, F., Penders, J., Vangronsveld, J., Gyselaers, W., & Nawrot, T. S. (2012). Placental mitochondrial DNA content and particulate air pollution during in utero life. *Environ Health Perspect*, 120(9), 1346-1352. <u>https://doi.org/10.1289/ehp.1104458</u>
- Jirtle, R. L., & Skinner, M. K. (2007). Environmental epigenomics and disease susceptibility. *Nature reviews. Genetics*, 8(4), 253-262. <u>https://doi.org/10.1038/nrg2045</u>
- Kampfrath, T., Maiseyeu, A., Ying, Z., Shah, Z., Deiuliis, J. A., Xu, X., Kherada, N., Brook, R. D., Reddy, K. M., Padture, N. P., Parthasarathy, S., Chen, L. C., Moffatt-Bruce, S., Sun, Q., Morawietz, H., & Rajagopalan, S. (2011). Chronic Fine Particulate Matter Exposure Induces Systemic Vascular Dysfunction via NADPH Oxidase and TLR4 Pathways. *Circulation Research*, 108(6), 716-726. <u>https://doi.org/doi:10.1161/CIRCRESAHA.110.237560</u>
- Leavey, K., Wilson, S. L., Bainbridge, S. A., Robinson, W. P., & Cox, B. J. (2018). Epigenetic regulation of placental gene expression in transcriptional subtypes of preeclampsia. *Clinical Epigenetics*, 10(1). <u>https://doi.org/10.1186/s13148-018-0463-6</u>
- Lee, H. C., & Wei, Y. H. (2000). Mitochondrial role in life and death of the cell. *J Biomed Sci*, 7(1), 2-15. <u>https://doi.org/10.1007/bf02255913</u>
- Lee, K., & Pausova, Z. (2013). Cigarette smoking and DNA methylation [Review]. *Frontiers in Genetics*, 4. <u>https://doi.org/10.3389/fgene.2013.00132</u>
- Leiva, A., Fuenzalida, B., Barros, E., Sobrevia, B., Salsoso, R., Sáez, T., Villalobos, R., Silva, L., Chiarello, I., Toledo, F., Gutiérrez, J., Sanhueza, C., Pardo, F., & Sobrevia, L. (2016). Nitric Oxide is a Central Common Metabolite in Vascular Dysfunction Associated with Diseases of Human Pregnancy. *Curr Vasc Pharmacol*, 14(3), 237-259. <u>https://doi.org/10.2174/1570161114666160222115158</u>

- Levin, H. L., & Moran, J. V. (2011). Dynamic interactions between transposable elements and their hosts. *Nature Reviews Genetics*, 12(9), 615-627. <u>https://doi.org/10.1038/nrg3030</u>
- Li, Z., Liu, Q., Xu, Z., Guo, X., & Wu, S. (2020). Association between short-term exposure to ambient particulate air pollution and biomarkers of oxidative stress: A meta-analysis. *Environmental Research*, 191, 110105. https://doi.org/https://doi.org/10.1016/j.envres.2020.110105
- Lyall, F., Young, A., & Greer, I. A. (1995). Nitric oxide concentrations are increased in the fetoplacental circulation in preeclampsia. *Am J Obstet Gynecol*, 173(3, Part 1), 714-718. <u>https://doi.org/https://doi.org/10.1016/0002-9378(95)90328-3</u>
- Mandakh, Y., Rittner, R., Flanagan, E., Oudin, A., Isaxon, C., Familari, M., Hansson, S. R., & Malmqvist, E. (2020). Maternal Exposure to Ambient Air Pollution and Risk of Preeclampsia: A Population-Based Cohort Study in Scania, Sweden. *Int J Environ Res Public Health*, 17(5). <u>https://doi.org/10.3390/ijerph17051744</u>
- Martens, D. S., Cox, B., Janssen, B. G., Clemente, D. B. P., Gasparrini, A., Vanpoucke, C., Lefebvre, W., Roels, H. A., Plusquin, M., & Nawrot, T. S. (2017). Prenatal air pollution and newborns' predisposition to accelerated biological aging [Article]. *JAMA Pediatr*, 171(12), 1160-1167. https://doi.org/10.1001/jamapediatrics.2017.3024
- Mayhew, T. M., Leach, L., McGee, R., Wan Ismail, W., Myklebust, R., & Lammiman, M. J. (1999). Proliferation, Differentiation and Apoptosis in Villous Trophoblast at 13–41 Weeks of Gestation (Including Observations on Annulate Lamellae and Nuclear Pore Complexes). *Placenta*, 20(5-6), 407-422. https://doi.org/10.1053/plac.1999.0399
- Mayne, B. T., Leemaqz, S. Y., Smith, A. K., Breen, J., Roberts, C. T., & Bianco-Miotto, T. (2017). Accelerated placental aging in early onset preeclampsia pregnancies identified by DNA methylation. *Epigenomics*, 9(3), 279-289. <u>https://doi.org/10.2217/epi-2016-0103</u>
- Michel, T., & Vanhoutte, P. M. (2010). Cellular signaling and NO production. *Pflugers* Arch, 459(6), 807-816. <u>https://doi.org/10.1007/s00424-009-0765-9</u>
- Mimura, K., Tomimatsu, T., Sharentuya, N., Tskitishvili, E., Kinugasa-Taniguchi, Y., Kanagawa, T., & Kimura, T. (2010). Nicotine restores endothelial dysfunction caused by excess sFlt1 and sEng in an in vitro model of preeclamptic vascular endothelium: a possible therapeutic role of nicotinic acetylcholine receptor (nAChR) agonists for preeclampsia. *Am J Obstet Gynecol*, 202(5), 464.e461-466. <u>https://doi.org/10.1016/j.ajog.2010.01.037</u>
- Mukosera, G. T., Clark, T. C., Ngo, L., Liu, T., Schroeder, H., Power, G. G., Yellon, S. M., Parast, M. M., & Blood, A. B. (2020). Nitric oxide metabolism in the human placenta during aberrant maternal inflammation. *J Physiol*, 598(11), 2223-2241. <u>https://doi.org/10.1113/jp279057</u>

- Murray, C. J. L., Aravkin, A. Y., Zheng, P., Abbafati, C., Abbas, K. M., Abbasi-Kangevari, M., Abd-Allah, F., Abdelalim, A., Abdollahi, M., Abdollahpour, I., Abegaz, K. H., Abolhassani, H., Aboyans, V., Abreu, L. G., Abrigo, M. R. M., Abualhasan, A., Abu-Raddad, L. J., Abushouk, A. I., Adabi, M., . . . Lim, S. S. (2020). Global burden of 87 risk factors in 204 countries and territories, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019. *The Lancet*, 396(10258), 1223-1249. <u>https://doi.org/10.1016/S0140-6736(20)30752-2</u>
- Myatt, L., Eis, A. L., Brockman, D. E., Greer, I. A., & Lyall, F. (1997). Endothelial nitric oxide synthase in placental villous tissue from normal, pre-eclamptic and intrauterine growth restricted pregnancies. *Hum Reprod*, *12*(1), 167-172. https://doi.org/10.1093/humrep/12.1.167
- Nawrot, T. S., Saenen, N. D., Schenk, J., Janssen, B. G., Motta, V., Tarantini, L., Cox, B., Lefebvre, W., Vanpoucke, C., Maggioni, C., & Bollati, V. (2018). Placental circadian pathway methylation and in utero exposure to fine particle air pollution. *Environ Int*, 114, 231-241. <u>https://doi.org/10.1016/j.envint.2018.02.034</u>
- Neven, K. Y., Saenen, N. D., Tarantini, L., Janssen, B. G., Lefebvre, W., Vanpoucke, C., Bollati, V., & Nawrot, T. S. (2018). Placental promoter methylation of DNA repair genes and prenatal exposure to particulate air pollution: an ENVIRONAGE cohort study. *The Lancet Planetary Health*, 2(4), e174-e183. https://doi.org/https://doi.org/10.1016/S2542-5196(18)30049-4
- Palmer, R. M. J., Ferrige, A. G., & Moncada, S. (1987). Nitric oxide release accounts for the biological activity of endothelium-derived relaxing factor. *Nature*, 327(6122), 524-526. <u>https://doi.org/10.1038/327524a0</u>
- Pańczyszyn, A., Boniewska-Bernacka, E., & Goc, A. (2020). The role of telomeres and telomerase in the senescence of postmitotic cells. DNA Repair (Amst), 95, 102956. <u>https://doi.org/10.1016/j.dnarep.2020.102956</u>
- Pedersen, M., Stayner, L., Slama, R., Sorensen, M., Figueras, F., Nieuwenhuijsen, M. J., Raaschou-Nielsen, O., & Dadvand, P. (2014). Ambient air pollution and pregnancyinduced hypertensive disorders: a systematic review and meta-analysis. *Hypertension*, 64(3), 494-500. <u>https://doi.org/10.1161/hypertensionaha.114.03545</u>
- Phillips, J. K., Bernstein, I. M., Mongeon, J. A., & Badger, G. J. (2004). Seasonal Variation in Preeclampsia Based on Timing of Conception. Obstetrics & Gynecology, 104(5). <u>https://journals.lww.com/greenjournal/Fulltext/2004/11000/Seasonal_Variation_in_P</u> reeclampsia Based on Timing.21.aspx
- Plusquin, M., Guida, F., Polidoro, S., Vermeulen, R., Raaschou-Nielsen, O., Campanella, G., Hoek, G., Kyrtopoulos, S. A., Georgiadis, P., Naccarati, A., Sacerdote, C., Krogh, V., Bas Bueno-de-Mesquita, H., Monique Verschuren, W. M., Sayols-Baixeras, S., Panni, T., Peters, A., Hebels, D. G. A. J., Kleinjans, J., . . . Chadeau-Hyam, M. (2017). DNA methylation and exposure to ambient air pollution in two prospective cohorts. *Environment International*, *108*, 127-136. https://doi.org/https://doi.org/10.1016/j.envint.2017.08.006

- Rakyan, V. K., Down, T. A., Balding, D. J., & Beck, S. (2011). Epigenome-wide association studies for common human diseases. *Nature Reviews Genetics*, 12(8), 529-541. <u>https://doi.org/10.1038/nrg3000</u>
- Ramiro-Cortijo, D., de la Calle, M., Rodriguez-Rodriguez, P., Phuthong, S., López de Pablo, Á. L., Martín-Cabrejas, M. A., & Arribas, S. M. (2020). First trimester elevations of hematocrit, lipid peroxidation and nitrates in women with twin pregnancies who develop preeclampsia. *Pregnancy Hypertens*, 22, 132-135. <u>https://doi.org/https://doi.org/10.1016/j.preghy.2020.09.005</u>
- Rath, W., & Tsikouras, P. (2018). [Maternal Deaths Worldwide Falling But Commonly Preventable]. Z Geburtshilfe Neonatol, 222(4), 143-151. <u>https://doi.org/10.1055/a-0607-2816</u> (Müttersterbefälle weltweit rückläufig, aber häufig vermeidbar!)

Redman, C. W. G., Staff, A. C., & Roberts, J. M. (2022). Syncytiotrophoblast stress in preeclampsia: the convergence point for multiple pathways. *Am J Obstet Gynecol*, 226(2, Supplement), S907-S927. <u>https://doi.org/https://doi.org/10.1016/j.ajog.2020.09.047</u>

Regionfakta. (2022). *Skåne län*. Retrieved 2022/11/27 from <u>https://www.regionfakta.com/skane-lan/in-english/population1/population/</u>

- Reyes, L., & Golos, T. G. (2018). Hofbauer Cells: Their Role in Healthy and Complicated Pregnancy [Review]. *Frontiers in Immunology*, 9(2628). <u>https://doi.org/10.3389/fimmu.2018.02628</u>
- Richter, T., & Zglinicki, T. v. (2007). A continuous correlation between oxidative stress and telomere shortening in fibroblasts. *Experimental Gerontology*, 42(11), 1039-1042. <u>https://doi.org/https://doi.org/10.1016/j.exger.2007.08.005</u>
- Rittner, R., Gustafsson, S., Spanne, M., & Malmqvist, E. (2020). Particle concentrations, dispersion modelling and evaluation in southern Sweden. SN Applied Sciences, 2(6), 1013. <u>https://doi.org/10.1007/s42452-020-2769-1</u>
- Robinson, O., & Vrijheid, M. (2015). The Pregnancy Exposome. *Curr Environ Health Rep*, 2(2), 204-213. <u>https://doi.org/10.1007/s40572-015-0043-2</u>
- Romão-Veiga, M., Bannwart-Castro, C. F., Borges, V. T. M., Golim, M. A., Peraçoli, J. C., & Peraçoli, M. T. S. (2020). Increased TLR4 pathway activation and cytokine imbalance led to lipopolysaccharide tolerance in monocytes from preeclamptic women. *Pregnancy Hypertens*, 21, 159-165. <u>https://doi.org/10.1016/j.preghy.2020.06.002</u>
- Rudra, C. B., & Williams, M. A. (2005). Monthly variation in preeclampsia prevalence: Washington state, 1987–2001. *The Journal of Maternal-Fetal & Neonatal Medicine*, 18(5), 319-324. <u>https://doi.org/10.1080/14767050500275838</u>
- Saenen, N. D., Martens, D. S., Neven, K. Y., Alfano, R., Bové, H., Janssen, B. G., Roels, H. A., Plusquin, M., Vrijens, K., & Nawrot, T. S. (2019). Air pollution-induced placental alterations: an interplay of oxidative stress, epigenetics, and the aging phenotype? *Clinical Epigenetics*, 11(1), 124. <u>https://doi.org/10.1186/s13148-019-0688-z</u>
- Shiozaki, A., Matsuda, Y., Satoh, S., & Saito, S. (2011). Impact of fetal sex in pregnancyinduced hypertension and preeclampsia in Japan. J Reprod Immunol, 89(2), 133-139. <u>https://doi.org/10.1016/j.jri.2010.12.011</u>

- Silva, A. V., Oliveira, C. M., Canha, N., Miranda, A. I., & Almeida, S. M. (2020). Long-Term Assessment of Air Quality and Identification of Aerosol Sources at Setúbal, Portugal. *International Journal of Environmental Research and Public Health*, 17(15), 5447. <u>https://www.mdpi.com/1660-4601/17/15/5447</u>
- Stroh, E., Rittner, R., Oudin, A., Ardö, J., Jakobsson, K., Björk, J., & Tinnerberg, H. (2012). Measured and modeled personal and environmental NO2 exposure. *Popul Health Metr*, 10(1), 10. <u>https://doi.org/10.1186/1478-7954-10-10</u>
- Sukenik-Halevy, R., Amiel, A., Kidron, D., Liberman, M., Ganor-Paz, Y., & Biron-Shental, T. (2016). Telomere homeostasis in trophoblasts and in cord blood cells from pregnancies complicated with preeclampsia. *American Journal of Obstetrics & Gynecology*, 214(2), 283.e281-283.e287. <u>https://doi.org/10.1016/j.ajog.2015.08.050</u>
- Sultana, Z., Maiti, K., Dedman, L., & Smith, R. (2018). Is there a role for placental senescence in the genesis of obstetric complications and fetal growth restriction? Am J Obstet Gynecol, 218(2, Supplement), S762-S773. https://doi.org/https://doi.org/10.1016/j.ajog.2017.11.567
- Sundrani, D. P., Reddy, U. S., Joshi, A. A., Mehendale, S. S., Chavan-Gautam, P. M., Hardikar, A. A., Chandak, G. R., & Joshi, S. R. (2013). Differential placental methylation and expression of VEGF, FLT-1 and KDR genes in human term and preterm preeclampsia. *Clinical Epigenetics*, 5(1), 6. <u>https://doi.org/10.1186/1868-7083-5-6</u>
- Suter, M., Ma, J., Harris, A., Patterson, L., Brown, K. A., Shope, C., Showalter, L., Abramovici, A., & Aagaard-Tillery, K. M. (2011). Maternal tobacco use modestly alters correlated epigenome-wide placental DNA methylation and gene expression. *Epigenetics*, 6(11), 1284-1294. <u>https://doi.org/10.4161/epi.6.11.17819</u>
- Suzuki, M. M., & Bird, A. (2008). DNA methylation landscapes: provocative insights from epigenomics. *Nature Reviews Genetics*, 9(6), 465-476. <u>https://doi.org/10.1038/nrg2341</u>
- Tang, Z., Buhimschi, I. A., Buhimschi, C. S., Tadesse, S., Norwitz, E., Niven-Fairchild, T., Huang, S.-T. J., & Guller, S. (2013). Decreased Levels of Folate Receptor-β and Reduced Numbers of Fetal Macrophages (Hofbauer Cells) in Placentas from Pregnancies with Severe Pre-Eclampsia. *American Journal of Reproductive Immunology*, 70(2), 104-115. <u>https://doi.org/https://doi.org/10.1111/aji.12112</u>
- Tranquilli, A. L., Brown, M. A., Zeeman, G. G., Dekker, G., & Sibai, B. M. (2013). The definition of severe and early-onset preeclampsia. Statements from the International Society for the Study of Hypertension in Pregnancy (ISSHP). *Pregnancy Hypertension: An International Journal of Women's Cardiovascular Health*, 3(1), 44-47. https://doi.org/https://doi.org/10.1016/j.preghy.2012.11.001
- Vishnyakova, P. A., Volodina, M. A., Tarasova, N. V., Marey, M. V., Tsvirkun, D. V., Vavina, O. V., Khodzhaeva, Z. S., Kan, N. E., Menon, R., Vysokikh, M. Y., & Sukhikh, G. T. (2016). Mitochondrial role in adaptive response to stress conditions in preeclampsia. *Sci Rep*, 6, 32410. <u>https://doi.org/10.1038/srep32410</u>
- Wacker, J., Schulz, M., Frühauf, J., Chiwora, F., Solomayer, E., & Bastert, G. (1998). Seasonal change in the incidence of preeclampsia in Zimbabwe. Acta Obstetricia et Gynecologica Scandinavica, 77(7), 712-716. <u>https://doi.org/10.1080/j.1600-0412.1998.770703.x</u>

- Walton, R. T., Mudway, I. S., Dundas, I., Marlin, N., Koh, L. C., Aitlhadj, L., Vulliamy, T., Jamaludin, J. B., Wood, H. E., Barratt, B. M., Beevers, S., Dajnak, D., Sheikh, A., Kelly, F. J., Griffiths, C. J., & Grigg, J. (2016). Air pollution, ethnicity and telomere length in east London schoolchildren: An observational study. *Environ Int*, 96, 41-47. <u>https://doi.org/10.1016/j.envint.2016.08.021</u>
- Wei, J., Liu, C. X., Gong, T. T., Wu, Q. J., & Wu, L. (2015). Cigarette smoking during pregnancy and preeclampsia risk: a systematic review and meta-analysis of prospective studies. *Oncotarget*, 6(41), 43667-43678. <u>https://doi.org/10.18632/oncotarget.6190</u>
- White, A. J., Kresovich, J. K., Keller, J. P., Xu, Z., Kaufman, J. D., Weinberg, C. R., Taylor, J. A., & Sandler, D. P. (2019). Air pollution, particulate matter composition and methylation-based biologic age. *Environ Int*, 132, 105071. <u>https://doi.org/10.1016/j.envint.2019.105071</u>
- WHO. (2004). International statistical classification of diseases and related health problems. In (Second edition ed.). WHO.
- World Health Organization, W. (2021). WHO global air quality guidelines: particulate matter (PM2.5 and PM10), ozone, nitrogen dioxide, sulfur dioxide and carbon monoxide. World Health Organization. <u>https://apps.who.int/iris/handle/10665/345329</u>
- Xuan, J., Jing, Z., Yuanfang, Z., Xiaoju, H., Pei, L., Guiyin, J., & Yu, Z. (2016). Comprehensive analysis of DNA methylation and gene expression of placental tissue in preeclampsia patients. *Hypertens Pregnancy*, 35(1), 129-138. <u>https://doi.org/10.3109/10641955.2015.1117099</u>
- Yang, S. W., Cho, E. H., Choi, S. Y., Lee, Y. K., Park, J. H., Kim, M. K., Park, J. Y., Choi, H. J., Lee, J. I., Ko, H. M., Park, S. H., Hwang, H. S., & Kang, Y. S. (2017). DC-SIGN expression in Hofbauer cells may play an important role in immune tolerance in fetal chorionic villi during the development of preeclampsia. *Journal of Reproductive Immunology*, 124, 30-37. https://doi.org/https://doi.org/10.1016/j.jri.2017.09.012
- Yu, C. H., Morandi, M. T., & Weisel, C. P. (2008). Passive dosimeters for nitrogen dioxide in personal/indoor air sampling: a review. J Expo Sci Environ Epidemiol, 18(5), 441-451. <u>https://doi.org/10.1038/jes.2008.22</u>
- Yu, H., Yin, Y., Zhang, J., & Zhou, R. (2020). The impact of particulate matter 2.5 on the risk of preeclampsia: an updated systematic review and meta-analysis. *Environmental Science and Pollution Research*, 27(30), 37527-37539. <u>https://doi.org/10.1007/s11356-020-10112-8</u>
- Zhang, Y., Wang, J., Gong, X., Chen, L., Zhang, B., Wang, Q., Han, B., Zhang, N., Xue, F., Vedal, S., & Bai, Z. (2020). Ambient PM(2.5) exposures and systemic biomarkers of lipid peroxidation and total antioxidant capacity in early pregnancy. *Environ Pollut*, 266(Pt 2), 115301. <u>https://doi.org/10.1016/j.envpol.2020.115301</u>





Department of Laboratory Medicine

Lund University, Faculty of Medicine Doctoral Dissertation Series 2023:9 ISBN 978-91-8021-348-6 ISSN 1652-8220

