LUND UNIVERSITY

(18)F-choline PET/CT for early detection of metastases in biochemical recurrence following radical prostatectomy.

Kjölhede, Henrik; Ahlgren, Göran; Almquist, Helen; Liedberg, Fredrik; Lyttkens, Kerstin; Ohlsson, Thomas; Bratt, Ola

Published in:
World Journal of Urology

DOI:
[10.1007/s00345-015-1547-y](https://doi.org/10.1007/s00345-015-1547-y)

2015

[Link to publication](#)

Citation for published version (APA):

Kjölhede, H., Ahlgren, G., Almquist, H., Liedberg, F., Lyttkens, K., Ohlsson, T., & Bratt, O. (2015). (18)F-choline PET/CT for early detection of metastases in biochemical recurrence following radical prostatectomy. *World Journal of Urology*, 33(11), 1749-1752. <https://doi.org/10.1007/s00345-015-1547-y>

Total number of authors:
7

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

¹⁸F-choline PET/CT for early detection of metastases in biochemical recurrence following radical prostatectomy

Kjölhede H¹, Ahlgren G², Almquist H³, Liedberg F², Lyttkens K³, Ohlsson T⁴, Bratt O⁵

1 Section of Urology, Department of Surgery, Växjö Hospital, Lund University, Sweden

2 Department of Urology, Skåne University Hospital, Lund University, Sweden

3 Center for Medical Imaging and Physiology, Skåne University Hospital, Lund University, Sweden

4 Radiation Physics, Skåne University Hospital, Lund University, Sweden

5 Department of Urology, Helsingborg Hospital, Lund University, Sweden

Corresponding author:

Henrik Kjölhede, Department of Surgery, Växjö Hospital, 351 85 Växjö, Sweden

Phone: +46-702-988922

Email: henrik.kjolhede@med.lu.se

Manuscript includes

1451 words and abstract 185 words, 1 table

Key words: Prostate Cancer, Recurrence, Metastasis, Positron Emission Tomography, Computed X Ray Tomography

Abstract

Purpose: Salvage radiotherapy (SRT) for biochemical recurrence (BCR) following radical prostatectomy (RP) should if possible be added at a prostate-specific antigen (PSA) level of less than 1-2 ng/mL. The value of positron emission tomography combined with computed tomography (PET/CT) at such low PSA values is not defined. The purpose was to determine what proportion of a well-defined cohort of hormone-naïve patients who were candidates for early salvage radiotherapy had ¹⁸F-choline PET/CT findings suggesting metastases.

Materials and methods: Patients with untreated BCR following RP, PSA < 2 ng/mL, and Gleason score ≥ 7 or PSA doubling time ≤ 6 months underwent ¹⁸F-choline PET/CT. Focal choline uptake in lymph nodes or skeletal sites were recorded.

Results: PET/CT indicated metastases in 16 (28%) of 58 patients. In five (9%) patients the scans suggested bone metastases and in 11 (19%) the scans suggested regional lymph node metastases only. For patients with PSA levels < 1.0 ng/mL, the PET/CT scans indicated metastatic recurrence in 25%.

Conclusions: ¹⁸F-choline PET/CT may be valuable for selecting patients with BCR following RP for SRT or experimental treatment of oligometastases, even at low PSA values.

Introduction

Radical prostatectomy (RP) is one of the cornerstones in curative treatment of prostate cancer [1], significantly reducing cancer-specific mortality compared to watchful waiting [2]. Still, a large proportion of patients, up to 60% depending on tumour characteristics [3,4], will develop biochemical recurrence (BCR) and may need additional treatment. BCR is usually treated with either radiotherapy (RT) or hormonal therapy (HT), depending on the likelihood that the recurrence is local only [5]. Salvage RT, which is the curative option, is associated with significant risk of morbidity [6] and should therefore only be given when there is a reasonable chance for cure. A high prostate-specific antigen (PSA) value, short time to relapse, short PSA doubling time, Gleason score 8-10 and lymph node metastases at lymphadenectomy are associated with increased risk of metastatic recurrence, and reduced chance of complete remission after salvage therapy [7,8]. Especially PSA > 2.0 ng/mL has been associated with a low probability of complete remission [8–10]. However, even in the group of patients with PSA < 2.0 ng/mL, some have metastatic disease and require systemic therapy, rather than local RT.

¹⁸F-choline positron emission tomography combined with computed tomography (PET/CT) has been studied as a modality for detecting regional and distant metastases in patients with newly detected prostate cancer, but the role of PET/CT in restaging after post-surgical recurrence is still uncertain [11,12]. Few previous studies have reported specifically on patients with low PSA values, i.e. when the choice of secondary treatment usually has to be made [13–18]. Most of the studies included a large proportion of patients on HT, for whom the association between PSA values and tumour burden is different than for hormone-naïve patients, and for whom treatment with additional RT is not evidence based. The primary aim of this study was to determine how large a proportion of hormone-naïve patients with BCR after RP would have positive ¹⁸F-choline PET/CT findings at PSA values < 2 ng/mL.

Materials and methods

Patients

Between June 2008 and November 2012, urologists in the Southern health-care region of Sweden were invited to refer patients with BCR after RP for a whole-torso ^{18}F -choline PET/CT scan according to a specific protocol. The inclusion criteria were: a rising PSA ≥ 0.2 ng/mL below 2 ng/mL, eligibility for salvage RT, and either Gleason Score ≥ 7 at prostatectomy or a PSA doubling time ≤ 6 months. Patients with ongoing or previous androgen deprivation therapy or anti-androgens were excluded. Enrollment was prospective according to the protocol, but the acquisition of the outcome data was performed retrospectively at the end of the study. The study aimed at including 50 patients. It was approved by the Research Ethical Review Board at the University of Lund (EPN Dnr 552/2007).

Imaging

^{18}F -choline was synthesized according to the method described by Kryza et al [19] using the TracerLab MXFDG module (GE Healthcare, Stockholm, Sweden). PET/CT images were obtained using an integrated PET/CT system (Philips Gemini TF, Philips Medical Systems, Ohio, USA) at the Center for Medical Imaging and Physiology, Skåne University Hospital in Malmö and Lund. All patients were fasting for four hours before tracer injection. Whole-torso PET was performed 1-1½ after intravenous injection of 4 MBq/kg of ^{18}F -choline, using 2 minutes per bed position. A CT scan was performed immediately prior to the PET scan with a multi-detector CT scanner using 5 mm reconstructed slice thickness. 1000 ml oral contrast (Omnipaque, GE Healthcare, Stockholm, Sweden) was given 1 hour before the scan. The CT scan was performed in three phases; the liver was scanned without contrast, the thorax was scanned with intravenous contrast in arterial phase during breath-holding, and then a femur to skull base scan was obtained in portal contrast phase during normal breathing [20].

All PET/CT scans were interpreted by a nuclear medicine physician as well as by a radiologist. Focal tracer uptakes in regional or retroperitoneal lymph nodes, or in skeletal sites without evidence

of degenerative causes or fractures, were considered positive findings. Focal uptake in the prostatic bed was noted, but was not included in further analyses.

Statistics

Statistical analysis was performed using SPSS Statistics 22 (IBM, Chicago, Ill, USA). PSA levels at the time of PET/CT, PSA doubling time, PSA nadir after RP and time to BCR were categorized according to Table 1. PSA doubling time was calculated by the log/slope method [21], using all available PSA levels after prostatectomy.

Results

A total of 58 patients were included. The clinical characteristics of all the patients are presented in Table 1. The PET/CT scans indicated metastases in 16 patients (28%): regional lymph node metastases in 11 (19%) and distant metastases in 5 (9%). At PSA cutoff levels of 1.5, 1.0 and 0.5 ng/mL, metastatic recurrence was indicated in 26%, 25% and 24%, respectively. Local PET tracer uptake in the prostatic bed was noted in four patients (7%), none of which had evidence of metastases.

Discussion

We investigated the utility of ¹⁸F-fluorocholine PET/CT in men with BCR after RP and PSA values in the low range (< 2 ng/mL), when the decision whether to add salvage RT usually has to be made. The PET/CT scans suggested metastatic recurrence in 28% of the patients, who were selected by having either Gleason score ≥ 7 cancers or a PSA doubling time shorter than 6 months. In the patients with PSA < 1.5 ng/mL 26% of the scans suggested metastases, and for patients with PSA < 1.0 ng/mL the corresponding figure was 25%.

There are only few other reports on PET/CT in patients with untreated BCR after RP and PSA < 2.0 ng/mL. Castelluci et al reported ¹¹C-choline PET/CT scans indicating metastases in 23% of 605 patients with PSA < 2 ng/mL, and in 18% of 291 patients with PSA < 1.0 ng/mL [15]. This is slightly less than in our study, but almost 25% of the patients in their study received HT at the time of the PET/CT investigation, which is known to affect the sensitivity of choline PET/CT [22]. Also, their study population included more patients with Gleason Score 5-6, with a lower likelihood of metastatic recurrence [7]. From a similar study, Giovacchini and associates reported 29% positive PET/CT scans in 358 patients with PSA < 2 ng/mL, but 43% of the patients received androgen deprivation therapy and it is unclear how many of the positive findings were metastases and how many were local recurrences [14]. These results are in contrast to those from a study including patients with very low PSA levels only (≤ 0.76 ng/mL); PET/CT scans indicated metastases in only 1 of 22 patients [13]. However, taken together, these studies suggest that a significant proportion of patients with BCR after RP and PSA levels < 2.0 ng/mL have metastatic recurrence, rather than local recurrence only. However, a limitation in our study, as well as in the others, is the lack of histopathological verification of the suspected metastases. Tilki et al reported a positive predictive value of only 50% for choline PET/CT compared to a secondary lymphadenectomy following BCR after RP and PSA < 2.0 ng/mL [23]. However, their study included only 12 patients. In contrast, Jilg and associates found a positive predictive value of 89% in a similar study, but with much more heterogeneous clinical characteristics [24]. Further validation is therefore required before choline PET/CT should be recommended outside clinical trials in the setting of BCR.

PET tracer uptake suggesting local recurrence was noted in only four patients. However, since all patients were planned to receive salvage RT if the scan did not reveal convincing evidence of metastases, we do not consider the absence or presence of uptake in the prostatic fossa as clinically relevant.

The most common purpose of using PET/CT for patients with BCR is to exclude those with metastases from local salvage RT. However, it is possible that aggressive treatment of oligo-metastatic disease, as determined by PET/CT, could lead to improved survival. Suardi recently reported 59% complete biochemical response and 17% recurrence-free 5-year survival following salvage lymphadenectomy in patients with nodal metastases only on choline PET/CT [25]. Schick et al reported similar results of RT targeted against oligometastases for BCR after RP or RT [26]. Further prospective, randomized trials in this setting could possibly lead to an improved management of patients with BCR.

Conclusions

In this study of hormone-naïve patients with BCR after RP, ^{18}F -choline PET/CT scans indicated metastatic recurrence in 28% of patients with PSA < 2 ng/mL, and in 25% of patients with PSA < 1 ng/mL. These results suggest that ^{18}F -choline PET/CT may be valuable in selecting patients for local salvage RT or experimental treatment of regional oligometastases.

Acknowledgements

This study has been funded by The Swedish Cancer Society (2012/475), Cancerstiftelsen Kronoberg, Region Skånes FoU-enhet, FoU Kronoberg and Philips.

Conflict of interest

The authors declare that they have no conflict of interest.

Authors' contribution

H Kjölhede: Data collection, data analysis, manuscript writing.

G Ahlgren: Protocol, manuscript editing

H Almquist: Protocol, data collection, manuscript editing

F Liedberg: Data analysis, manuscript editing

K Lyttkens: Protocol, data collection, manuscript editing

T Ohlsson: Data collection, manuscript editing

O Bratt: Protocol, data analysis, manuscript writing

References

1. Heidenreich A, Bastian PJ, Bellmunt J, Bolla M, Joniau S, van der Kwast T, et al. EAU guidelines on prostate cancer. part 1: screening, diagnosis, and local treatment with curative intent-update 2013. *Eur Urol.* 2014; 65(1):124–37.
2. Bill-Axelsson A, Holmberg L, Garmo H, Rider JR, Taari K, Busch C, et al. Radical prostatectomy or watchful waiting in early prostate cancer. *N Engl J Med.* 2014; 370(10):932–42.
3. Han M, Partin AW, Zahurak M, Piantadosi S, Epstein JI, Walsh PC. Biochemical (prostate specific antigen) recurrence probability following radical prostatectomy for clinically localized prostate cancer. *J Urol.* 2003; 169(2):517–23.
4. Sooriakumaran P, Haendler L, Nyberg T, Gronberg H, Nilsson A, Carlsson S, et al. Biochemical recurrence after robot-assisted radical prostatectomy in a European single-centre cohort with a minimum follow-up time of 5 years. *Eur Urol.* 2012; 62(5):768–74.
5. Heidenreich A, Bastian PJ, Bellmunt J, Bolla M, Joniau S, van der Kwast T, et al. EAU guidelines on prostate cancer. Part II: Treatment of advanced, relapsing, and castration-resistant prostate cancer. *Eur Urol.* 2014; 65(2):467–79.
6. Bolla M, van Poppel H, Collette L, van Cangh P, Vekemans K, Da Pozzo L, et al. Postoperative radiotherapy after radical prostatectomy: a randomised controlled trial (EORTC trial 22911). *Lancet.* 2005; 366(9485):572–8.
7. Slovin SF, Wilton AS, Heller G, Scher HI. Time to detectable metastatic disease in patients with rising prostate-specific antigen values following surgery or radiation therapy. *Clin Cancer Res.* 2005; 11(24 Pt 1):8669–73.
8. Stephenson AJ, Scardino PT, Kattan MW, Pisansky TM, Slawin KM, Klein E a, et al. Predicting the outcome of salvage radiation therapy for recurrent prostate cancer after radical prostatectomy. *J Clin Oncol.* 2007; 25(15):2035–41.
9. King CR. The timing of salvage radiotherapy after radical prostatectomy: a systematic review. *Int J Radiat Oncol Biol Phys.* 2012; 84(1):104–11.
10. Pfister D, Bolla M, Briganti A, Carroll P, Cozzarini C, Joniau S, et al. Early salvage radiotherapy following radical prostatectomy. *Eur Urol.* 2014; 65(6):1034–43.
11. Kjölhede H, Ahlgren G, Almquist H, Liedberg F, Lyttkens K, Ohlsson T, et al. 18F-fluorocholine PET/CT compared with extended pelvic lymph node dissection in high-risk prostate cancer. *World J Urol.* 2014; 32(4):965–70.
12. Evangelista L, Guttilla A, Zattoni F, Muzzio PC, Zattoni F. Utility of choline positron emission tomography/computed tomography for lymph node involvement identification in intermediate- to high-risk prostate cancer: a systematic literature review and meta-analysis. *Eur Urol.* 2013; 63(6):1040–8.
13. Veas H, Buchegger F, Albrecht S, Khan H, Husarik D, Zaidi H, et al. 18F-choline and/or 11C-acetate positron emission tomography: detection of residual or progressive subclinical disease

at very low prostate-specific antigen values (<1 ng/mL) after radical prostatectomy. *BJU Int.* 2007; 99(6):1415–20.

14. Giovacchini G, Picchio M, Coradeschi E, Bettinardi V, Gianolli L, Scattoni V, et al. Predictive factors of [(11)C]choline PET/CT in patients with biochemical failure after radical prostatectomy. *Eur J Nucl Med Mol Imaging.* 2010; 37(2):301–9.
15. Castellucci P, Ceci F, Graziani T, Schiavina R, Brunocilla E, Mazzarotto R, et al. Early Biochemical Relapse After Radical Prostatectomy: Which Prostate Cancer Patients May Benefit from a Restaging 11C-Choline PET/CT Scan Before Salvage Radiation Therapy? *J Nucl Med.* 2014; 55(9):1424–9.
16. Krause BJ, Souvatzoglou M, Tuncel M, Herrmann K, Buck AK, Praus C, et al. The detection rate of [11C]choline-PET/CT depends on the serum PSA-value in patients with biochemical recurrence of prostate cancer. *Eur J Nucl Med Mol Imaging.* 2008; 35(1):18–23.
17. Rinnab L, Mottaghy FM, Blumstein NM, Reske SN, Hautmann RE, Hohl K, et al. Evaluation of [11C]-choline positron-emission/computed tomography in patients with increasing prostate-specific antigen levels after primary treatment for prostate cancer. *BJU Int.* 2007; 100(4):786–93.
18. Scattoni V, Picchio M, Suardi N, Messa C, Freschi M, Roscigno M, et al. Detection of lymph-node metastases with integrated [11C]choline PET/CT in patients with PSA failure after radical retropubic prostatectomy: results confirmed by open pelvic-retroperitoneal lymphadenectomy. *Eur Urol.* 2007; 52(2):423–9.
19. Kryza D, Tadino V, Filannino MA, Villeret G, Lemoucheux L. Fully automated [18F]fluorocholine synthesis in the TracerLab MX FDG Coincidence synthesizer. *Nucl Med Biol.* 2008; 35(2):255–60.
20. Kjölhede H, Ahlgren G, Almquist H, Liedberg F, Lyttkens K, Ohlsson T, et al. Combined 18F-fluorocholine and 18F-fluoride positron emission tomography/computed tomography imaging for staging of high-risk prostate cancer. *BJU Int.* 2012; 110(10):1501–6.
21. Pound CR, Partin AW, Eisenberger MA, Chan DW, Pearson JD, Walsh PC. Natural history of progression after PSA elevation following radical prostatectomy. *JAMA.* 1999; 281(17):1591–7.
22. Beheshti M, Haim S, Zakavi R, Steinmair M, Waldenberger P, Kunit T, et al. Impact of 18F-choline PET/CT in prostate cancer patients with biochemical recurrence: influence of androgen deprivation therapy and correlation with PSA kinetics. *J Nucl Med.* 2013; 54(6):833–40.
23. Tilki D, Reich O, Graser A, Hacker M, Silchinger J, Becker AJ, et al. 18F-Fluoroethylcholine PET/CT identifies lymph node metastasis in patients with prostate-specific antigen failure after radical prostatectomy but underestimates its extent. *Eur Urol.* 2013; 63(5):792–6.
24. Jilg C a, Schultze-Seemann W, Drendel V, Vach W, Wieser G, Krauss T, et al. Detection of lymph node metastasis in patients with nodal prostate cancer relapse using (18)f/(11)c-choline positron emission tomography/computerized tomography. *J Urol.* 2014; 192(1):103–11.

25. Suardi N, Gandaglia G, Gallina A, Di Trapani E, Scattoni V, Vizziello D, et al. Long-term Outcomes of Salvage Lymph Node Dissection for Clinically Recurrent Prostate Cancer: Results of a Single-institution Series with a Minimum Follow-up of 5 Years. *Eur Urol.* (2014), In press <http://dx.doi.org/10.1016/j.eururo.2014.02.011>.
26. Schick U, Jorcano S, Nouet P, Rouzaud M, Veas H, Zilli T, et al. Androgen deprivation and high-dose radiotherapy for oligometastatic prostate cancer patients with less than five regional and/or distant metastases. *Acta Oncol (Madr).* 2013; 52(8):1622–8.