



LUND UNIVERSITY

Plasticity of mitochondrial function safeguards phosphorylating respiration during in vitro simulation of rest-phase hypothermia

García-Díaz, Carmen C.; Chamkha, Imen; Elmer, Eskil; Nord, Andreas

Published in:
FASEB Journal

DOI:
[10.1096/fj.202201613R](https://doi.org/10.1096/fj.202201613R)

2023

Document Version:
Publisher's PDF, also known as Version of record

[Link to publication](#)

Citation for published version (APA):
García-Díaz, C. C., Chamkha, I., Elmer, E., & Nord, A. (2023). Plasticity of mitochondrial function safeguards phosphorylating respiration during in vitro simulation of rest-phase hypothermia. *FASEB Journal*, 37(4), Article e22854. <https://doi.org/10.1096/fj.202201613R>

Total number of authors:
4

Creative Commons License:
CC BY-NC-ND

General rights

Unless other specific re-use rights are stated the following general rights apply:
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

RESEARCH ARTICLE

Plasticity of mitochondrial function safeguards phosphorylating respiration during in vitro simulation of rest-phase hypothermia

Carmen C. García-Díaz¹  | Imen Chamkha²  | Eskil Elmér²  | Andreas Nord¹ 

¹Department of Biology, Section for Evolutionary Ecology, Lund University, Lund, Sweden

²Department of Clinical Sciences, Mitochondrial Medicine, Lund University, Lund, Sweden

Correspondence

Andreas Nord, Department of Biology, Section for Evolutionary Ecology, Lund University, Lund, Sweden.
Email: andreas.nord@biol.lu.se

Funding information

Birgit and Hellmuth Hertz Foundation, Grant/Award Number: 2017-39034; Royal Physiographic Society in Lund, Grant/Award Number: 2019-41011; Swedish Research Council, Grant/Award Number: 2020-04686; Stiftelsen Lunds Djurskyddsfond

Abstract

Many animals downregulate body temperature to save energy when resting (rest-phase hypothermia). Small birds that winter at high latitudes have comparatively limited capacity for hypothermia and so pay large energy costs for thermoregulation during cold nights. Available evidence suggests this process is fueled by adenosine triphosphate (ATP)-dependent mechanisms. Most ATP is produced by oxidative phosphorylation in the mitochondria, but mitochondrial respiration may be lower during hypothermia because of the temperature dependence of biological processes. This can create conflict between increased organismal ATP demand and a lower mitochondrial capacity to provide it. We studied this in blood cell mitochondria of wild great tits (*Parus major*) by simulating rest-phase hypothermia via a 6°C reduction in assay temperature in vitro. The birds had spent the night preceding the experiment in thermoneutrality or in temperatures representing mild or very cold winter nights, but night temperatures never affected mitochondrial respiration. However, across temperature groups, endogenous respiration was 14% lower in hypothermia. This did not reflect general thermal suppression of mitochondrial function because phosphorylating respiration was unaffected by thermal state. Instead, hypothermia was associated with a threefold reduction of leak respiration, from 17% in normothermia to 4%

Abbreviations: ATP, Adenosine triphosphate; BAT, Brown adipose tissue, i.e., brown fat; EGTA, Egtazic acid; E-L coupling efficiency, Flow control ratio indicating how tightly electron transport is coupled to ATP production when the mitochondria are maximally stimulated by an artificial uncoupler; E-R control efficiency, Flow control ratio indicative of surplus respiratory capacity; that is, the proportion of maximum working capacity remaining during endogenous respiration; ESM, Electronic supplementary material to accompany the main text; ETS, Electron transport system; here in the context of its maximum respiration capacity when uncoupled from ATP synthase; FCCP, Cyanide-p-trifluoro-methoxyphenyl-hydrazone; a mitochondrial uncoupler; FCR, Flow control ratio; that is, a ratio between two different mitochondrial respiration states; Hepes, Hydroxyethyl piperazineethanesulfonic acid; ID, Subject identification number; K2-EDTA, Dipotassium ethylenediaminetetraacetic acid; LEAK, Mitochondrial respiration to offset proton leak across the inner mitochondrial membrane when no ATP is being produced; lm, Linear model; lmer, Linear mixed effects regression model; MiR05, Mitochondrial respiration medium 05; OXPHOS, Mitochondrial respiration driving oxidative phosphorylation during ATP synthesis; PIT tag, Passive integrative transponder; here used for non-contact body temperature measurement; R-L control efficiency, Flow control ratio indicating the proportion of endogenous respiration channeled toward ATP production via oxidative phosphorylation; RMR, resting metabolic rate; ROS, Reactive oxygen species; ROUTINE, Baseline mitochondrial respiration on endogenous substrates; SERCA, Sarco/endoplasmic reticulum Ca²⁺-ATPase; i.e., an ion pump that moves Ca²⁺ from the cytosol into the sarcoplasmic reticulum.

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2023 The Authors. *The FASEB Journal* published by Wiley Periodicals LLC on behalf of Federation of American Societies for Experimental Biology.

in hypothermia. Thus, the coupling of total respiration to ATP production was 96% in hypothermia, compared to 83% in normothermia. Our study shows that the thermal insensitivity of phosphorylation combined with short-term plasticity of leak respiration may safeguard ATP production when endogenous respiration is suppressed. This casts new light on the process by which small birds endure harsh winter cold and warrants future tests across tissues *in vivo*.

KEYWORDS

blood cells, body temperature, cell respiration, great tit, mitochondria, thermoregulation

1 | INTRODUCTION

Endothermic animals, such as mammals and birds, use considerable amounts of energy to regulate a high and relatively stable body temperature across a wide range of air temperatures. The associated energy cost of thermoregulation can sometimes become prohibitively high, for example, when food is scarce or when weather conditions deteriorate. Many small to medium-sized endotherms use rest-phase hypothermia, a regulated reduction of resting body temperature below normal levels, to mitigate such energetic challenges. This may reduce metabolic demands by 50% or more.¹ It follows that most animals using rest-phase hypothermia have high metabolic intensity, lose heat quickly, and rely on high-energy or ephemeral food sources.² This fits the description of “the little bird in winter”,³ i.e., small (<25g) songbirds that are year-round residents at high latitudes where they routinely meet environmental temperatures 50–60°C below body temperature during the cold season. Yet, most birds in northern forests, such as tits and chickadees, typically show daily variation in body temperature within 5 °C (e.g., Refs. [4–6]) and generally do not exceed 7–8°C even in severe cold,^{(7–9} but see Ref. [10] for an exception). Theoretical models predict that even such moderate body temperature reduction can carry survival benefits.^{11,12} However, the modest extent of hypothermia means that the little bird in winter can never forego a significant increase in energetically costly heat production to counter winter cold, even when maximally hypothermic (e.g., Ref. [13]).

Most of the energy used by the working body is in the form of adenosine triphosphate (ATP) which is produced mainly by oxidative phosphorylation in the mitochondria.¹⁴ Since more energy is needed to stay warm in winter, it is not surprising that many endotherms upregulate mitochondrial content and/or respiration rate in key metabolic tissues when exposed to cold, such as in the wintertime.^{15–19} However, little is known about mitochondrial adjustment in response to shorter-term variation in air temperature, for example, during a sudden drop in night-time temperature coincident with a clearing of an overcast sky. Moreover, a potential problem for a hypothermic animal is that the activity of the respiratory chain could be slower in a colder

body (e.g., Ref. [20]), causing a conflict between increased organismal demand for fuel but reduced mitochondrial capacity to provide it. However, not all the proton motive force is used to produce ATP since some protons backflow into the mitochondrial matrix bypassing ATP synthase. This is mediated by three principal mechanisms. Physiological properties maintain a basal level of proton conductance of the mitochondrial membrane that can be further up or downregulated under the influence of uncoupling proteins (reviewed by Ref. [21]). A functionally similar phenomenon occurs when protons spontaneously “slip” back through the proton pumps of the electron transport chain, though the contribution of slippage to total uncoupling is probably low.²¹ Compensation for these processes makes up what is known as leak respiration. This constitutes some 20%–25% of resting metabolic rate in endotherms,^{22,23} meaning there is scope to adjust the coupling of electron transport to ATP production according to the animal's need. For example, increased mitochondrial uncoupling in brown adipose tissue (BAT) is the basis for non-shivering heat production in mammals.¹⁴ Birds, by contrast, lack BAT and so are thought to produce heat mostly using ATP-dependent shivering in skeletal muscles, that is, without non-shivering contributions such as mitochondrial uncoupling or futile pumping of Ca²⁺ in the SERCA pump^{24,25} (but see Refs. [26,27] for possible exceptions). Thus, in birds, it can be speculated that acute cold exposure could be associated with improved mitochondrial coupling, to make ATP production more effective for a given respiration rate, which has been suggested for other energy-demanding processes such as fasting (e.g., Refs. [28–31]). This could secure adequate fuel delivery even when cell respiration is reduced, such as when heat production must remain high, but rest-phase hypothermia causes the mitochondria to work at a slower pace. There is support for this compensation hypothesis from studies on ectotherms^{32–34} (but see Refs. [35,36] for exceptions). Similar studies on endotherms suggest that while phosphorylating respiration is often suppressed in hypometabolic animals, non-phosphorylating (i.e., leak) respiration is sometimes reduced, sometimes unaltered, and sometimes increased (reviewed by Ref. [37]).

Previous endotherm studies have largely been performed in mammals during hibernation or in deep torpor; thermal states characterized by substantial metabolic downregulation¹ beyond what is feasible in most birds (but see Ref. [38]). Thus, previous data are not easily extrapolated to the problems faced by the little bird in winter that must increase energy expenditure sharply to counter cold temperature even when hypothermic, and must do so using ATP-dependent thermogenesis. Hence, we studied mitochondrial responses to both cold environments and rest-phase hypothermia in the great tit (*Parus major* L.)—a small (16–20 g) songbird that winters at high latitudes and shows pronounced diel variation in body temperature.^{4,39} We exploited the fact that birds contain functional mitochondria in their red blood cells. Previous studies have found that avian blood cell mitochondria respond in the short-term to variation in energetic demands,^{40–42} over longer time periods in line with age-related metabolic senescence⁴³ and according to environmental context.¹⁸ There is also evidence suggesting blood cell respiration in mammals and birds is correlated to respiration in tissues that are important for heat production (skeletal muscle) and energy management (heart, kidney, brain),^{44–46} all of which show pronounced seasonal acclimatization in small, northern birds.⁴⁷ This emphasizes the need for further studies exploring the utility of blood cell respirometry as a minimally invasive method for assessing mitochondrial responses to environmental stressors. On this basis, we, first, tested if the immediate thermal environment of the bird impacted mitochondrial function by manipulating the night preceding measurements to be representative of mild or very cold winter temperature, or thermoneutrality. Then, we performed an *in vitro* simulation of rest-phase hypothermia, to test if lower body temperature caused a reduction in mitochondrial respiration and if this was associated with any changes in the efficiency of ATP production. We expected that low night-time temperature would be associated with increased mitochondrial respiration rate to facilitate ATP-dependent heat production. In contrast, we predicted that lower assay temperature would cause thermal suppression of respiration traits. However, if better-coupled mitochondria serve as an adaptation to safeguard ATP production when cold ambient temperatures increase organismal-level demand for ATP, or when a low body temperature reduces overall energy expenditure of the cell, we expected tighter coupling of electron transport to ATP production both in response to lower air temperature at night and during *in vitro* hypothermia.

2 | MATERIALS AND METHODS

The experimental design and timeline are outlined in Figure 1.




FIGURE 1 Overview of the experimental design and timeline of events when studying the effects of night temperature and an *in vitro* simulation of thermal state (normothermia or hypothermia) on functional properties of mitochondria in avian blood cells. Studies were performed using wild great tits (*Parus major*) that wintered outside the city of Lund in southernmost Sweden.

2.1 | Study site and animals

Great tits were caught at two sites in the vicinity of Lund in southernmost Sweden – Räfteå (n = 37) and Vomb (n = 21)—between January and February 2021. Räfteå (55° 43'N, 13° 17' E'') is an oak (*Quercus robur*) dominated 32 ha woodlot surrounded by arable fields. The Vomb site (55° 39'N, 13° 33' E) is part of a pine (*Pinus sylvestris*) plantation with a rich deciduous understory composed mainly of willow (*Salix* spp.), birch (*Betula pendula*), and rowan (*Sorbus aucuparia*). Birds in Räfteå were caught in mist nets close to feeders baited with sunflower seeds that were erected 1.5 m above ground in January 2021. The birds in Vomb were caught shortly before sunrise when roosting in nest boxes. Four to eight birds were caught each day of the study and were brought to Lund University within 2 h of capture, where they were alternately allocated to temperature treatment groups in batches of four. We housed the birds in cages measuring 77×44×77 cm (length×width×height) at 5°C under simulated natural photoperiod and provided sunflower

seeds, peanuts, and water ad libitum. One-two days after capture, the birds were implanted with a temperature-sensitive PIT tag (BioTherm13, Biomark, Boise, ID, USA) into the intraperitoneal cavity under local anesthesia (5% lidocaine) as part of a different study. Eight to 32 h later, the birds were put in a 1 L hermetically sealed respirometry glass container placed inside a climate chamber (Weiss Umwelttechnik C180, Reiskirchen, Germany). The respirometry chamber was ventilated with dry, pressurized, air at 400–500 mL min⁻¹ (recorded using a FB-8 mass flow meter, Sable Systems, Las Vegas, NV, USA) for measurement of resting metabolic rate (RMR; calculated from oxygen consumption recorded in carbon dioxide-free air using a FC-10 oxygen analyzer, Sable Systems) and body temperature (recorded using a custom-built multiplexed antennae system from BioMark). These measurements were performed at thermoneutrality (25°C; *n* = 20) [13], during a simulated mild winter night (5°C; *n* = 19) and during a simulated very cold winter night (−15°C; *n* = 19) for 13–15 h. Four birds were measured each night. Mean RMR was 0.36 ± 0.01 W (mean ± 1 standard error.), 0.51 ± 0.01 W, and 0.68 ± 0.02 W in +25, +5, and −15°C, respectively. Mean nightly body temperature was 37.9 ± 0.3 °C (range of individual means: 35.2–40.1°C; minimum: 33.8°C), 39.3 ± 0.1°C (range of individual means: 37.6–40.2°C; minimum: 37.0°C), and 39.2 ± 0.1°C (range of individual means: 38.0–40.3°C; minimum: 37.6°C), in the cold, mild, and thermoneutral night conditions.

Air temperature in Lund (10–40 km from the study sites) ranged from −12.5 to +4.6°C during the study period and photoperiod ranged from 8:20 to 10:20 h.⁴⁸

2.2 | Blood sampling and mitochondrial measurement in whole blood

In the morning after the metabolic measurements, a 100–150 µL blood sample (i.e., ≤10% of the estimated total blood volume) was obtained from the jugular vein using a heparinized syringe within 5 min of removing the bird from the metabolic chamber. The sample was stored at 10–12°C in 2 mL K2-EDTA (dipotassium ethylenediaminetetraacetic acid) tubes (BD Vacutainer®, Franklin Lakes, NJ, USA) without agitation until analyzed 0.5–2 h later. All birds were released at the site of capture within 2 h of blood sampling.

We measured mitochondrial respiration in whole blood using an Oxygraph O2k high-resolution respirometer (Oroboros Instruments, Innsbruck, Austria), following the protocol described in Nord, Chamkha & Elmér.⁴⁹ To test how rest-phase hypothermia affected mitochondrial function, we ran duplicate samples from the same individual simultaneously in each of two thermal states: a

representative normothermic daytime bird body temperature (41°C; henceforth “normothermia”) and a hypothermic night-time body temperature in the lower range of the measured natural variation (35°C; henceforth “hypothermia”)⁵⁰ using two respirometers. These were calibrated at assay temperature (i.e., 35 or 41°C) daily, and we changed instrument/assay temperature combinations between days to avoid confounding experimental results by any variation pertaining to the specific instruments. The main purpose of our study was to demonstrate the presence of a putative mitochondrial response to acute body temperature change. To achieve this goal, we purposely selected a low but biologically relevant hypothermic assay temperature, under the assumption that any phenotypic response would be linear over the range of in vitro temperatures considered.

The blood samples were allowed 5–10 min on a rotating mixer at room temperature before the start of the assay. Then, we added a 50 µL sample to 1.95 mL MiRO5 respiration medium (0.5 mmol L⁻¹ EGTA, 3 mmol L⁻¹ MgCl₂, 60 mmol L⁻¹ K-lactobionate, 20 mmol L⁻¹ taurine, 10 mmol L⁻¹ KH₂PO₄, 20 mmol L⁻¹ HEPES, 110 mmol L⁻¹ sucrose, free fatty acid bovine serum albumin (1 g L⁻¹), pH 7.1) that was contained within the respirometer chambers and so was pre-equilibrated at assay temperature. The final volume contained in the respirometer chamber was, thus, 2 mL.

Mitochondrial respiration rate was measured as the rate of decline in O₂ concentration in the chamber. Once the O₂ consumption had stabilized upon closing the chamber, baseline O₂ consumption on endogenous substrates (“ROUTINE”) was recorded for 2–3 min. Then, we assessed mitochondrial function by sequential addition of three mitochondrial agonists and antagonists. First, we added the ATP-synthase inhibitor oligomycin at a final concentration of 5 µmol L⁻¹, which blocks the process of oxidative phosphorylation (“OXPHOS”). Thus, the remaining O₂ after oligomycin addition is attributed to proton leak (“LEAK”). Next, we added the mitochondrial uncoupler FCCP (carbonyl cyanide-p-trifluoro-methoxyphenyl-hydrazone), which abolishes the proton gradient across the inner mitochondrial membrane, causing the electron transport system to work at maximum capacity to restore it (“ETS”). Because an excessive concentration of FCCP inhibits mitochondrial respiration and can lead to acidification of the cell cytoplasm [14], we first added 1.5 µL of 1 mmol L⁻¹ FCCP and then titrated FCCP in 0.5 µL increments until maximum O₂ consumption was reached. The final concentration ranged 1–1.75 µmol L⁻¹. Finally, we added 5 µmol L⁻¹ of antimycin A, which blocks Complex III of the electron transport chain. Therefore, the remaining O₂ consumption after antimycin A addition is due to non-mitochondrial oxygen utilization.

2.3 | Data handling and statistical analyses

We excluded five of the 57 duplicate samples measured at 41°C that showed low endogenous respiration that was not inhibited by the addition of oligomycin, most likely because of unknown, aberrant, consequences of sample processing before the start of the experiment. Thus, the final data set consisted of 57 observations at 35°C and 52 observations at 41°C. We then calculated mitochondrial respiration states and flux control ratios (FCR; i.e., ratios between the respiratory states) following Gnaiger.⁵¹ We considered three FCRs: (1) E-R control efficiency ($1 - \text{ROUTINE}/\text{ETS}$), which measures how much excess capacity there is in the electron transport system until maximum respiration is reached; (2) R-L control efficiency ($(\text{ROUTINE} - \text{LEAK})/\text{ROUTINE}$), which measures the proportion of endogenous respiration that is channeled toward oxidative phosphorylation; (3) E-L coupling efficiency ($1 - \text{LEAK}/\text{ETS}$) which is indicative of mitochondrial efficiency in the context of coupling between electron transport and ATP production in a stimulated cellular state. The derivation and definition of all respiration traits and FCRs are detailed in Table 1.

All statistical analyses were performed using R version 4.1.2. To test how a sudden change in night temperature

affected mitochondrial respiration the morning after, we used linear models (lm in R base) to test how each respiration state and FCR assayed at normothermic temperature (i.e., 41°C) was affected by the preceding night temperature. We used linear mixed models (lmer in the lme4 package)⁵² to test how respiration traits and FCRs were affected by thermal state (i.e., hypo- or normothermia) using bird ID as a random factor to account for individual variation in the Y-intercept and non-independence of repeated measurements on the same birds. To test if the phenotypic response to a change in thermal state differed depending on the depth of hypothermia during the preceding night, we included mean nightly body temperature and body temperature \times thermal state in these models. Significances were assessed using *F*-tests with degrees of freedom based on the Kenward-Roger approximation, implemented using the KRmodcomp function in the pbkrtest package.⁵³ The significance of the random term (i.e., bird ID) in final models was assessed using log-likelihood tests using the drop1 function in base R. The interaction was removed when non-significant ($p > 0.05$), but all main effects were retained. Predicted values and standard errors for the lmer models were calculated using the emmeans function in the emmeans package.⁵⁴ For the tests of night temperature effects, estimates and their standard errors were taken from the linear model output.

TABLE 1 Calculation method and definition of mitochondrial respiration states and flux control ratios.

Parameter	Calculation method	Definition
ROUTINE	$R_{\text{Baseline}} - R_{\text{Antimycin A}}$	Baseline O ₂ consumption during oxidative phosphorylation under endogenous cellular conditions
OXPHOS	$R_{\text{Baseline}} - R_{\text{Oligomycin}}$	O ₂ consumption used to drive oxidative phosphorylation where ATP is produced
ETS	$R_{\text{FCCP}} - R_{\text{Antimycin A}}$	O ₂ consumption when the electron transport system is operating at its maximum capacity
LEAK	$R_{\text{Oligomycin}} - R_{\text{Antimycin A}}$	O ₂ consumption needed to compensate for proton leakiness of the mitochondria
E-R control efficiency	$1 - (\text{ROUTINE}/\text{ETS})$	Proportion of ETS maximum working capacity remaining under endogenous cellular conditions
R-L control efficiency	$(\text{ROUTINE} - \text{LEAK})/\text{ROUTINE}$	Proportion of endogenous respiration used for ATP production via oxidative phosphorylation
E-L coupling efficiency	$1 - (\text{LEAK}/\text{ETS})$	Tightness of electron transport during ATP production in a stimulated cellular state

Note: All traits were derived based on changes in O₂ consumption rate in response to the addition of mitochondrial agonists and antagonists during the assay as described in the main text.

3 | RESULTS

A representative normothermic experiment is presented in [Figure S1](#). All details of the statistical tests pertaining to the effects of night temperature are presented in [Table 2](#), and those pertaining to the effects of thermal state and body temperature are presented in [Table 3](#).

3.1 | Effects of preceding night temperature

All but one of the mitochondrial respiration metrics were unaffected by the temperature of the preceding night ($p \geq 0.16$; [Table 2](#)). However, there was a tendency for R-L control efficiency, which measures how much of

Model and Variable	Estimate \pm SEM	<i>n</i>	df	<i>F</i>	<i>p</i>
ROUTINE (pmol O ₂ s ⁻¹ μL ⁻¹)		52			
Night temp (-15, 5 or 25°C)			2, 49	0.51	0.607
Night temp = -15°C	0.565 \pm 0.041	17			
Night temp = 5°C	0.546 \pm 0.058	17			
Night temp = 25°C	0.602 \pm 0.057	18			
OXPPOS (pmol O ₂ s ⁻¹ μL ⁻¹)		52			
Night temp (-15, 5 or 25°C)			2, 49	0.79	0.461
Night temp = -15°C	0.487 \pm 0.034	17			
Night temp = 5°C	0.442 \pm 0.047	17			
Night temp = 25°C	0.498 \pm 0.047	18			
ETS (pmol O ₂ s ⁻¹ μL ⁻¹)		52			
Night temp (-15, 5 or 25°C)			2, 49	0.99	0.376
Night temp = -15°C	0.981 \pm 0.090	17			
Night temp = 5°C	0.993 \pm 0.127	17			
Night temp = 25°C	0.934 \pm 0.125	18			
LEAK (pmol O ₂ s ⁻¹ μL ⁻¹)		52			
Night temp (-15, 5 or 25°C)			2, 49	1.56	0.220
Night temp = -15°C	0.078 \pm 0.012	17			
Night temp = 5°C	0.103 \pm 0.016	17			
Night temp = 25°C	0.104 \pm 0.016	18			
E-R control efficiency ('mitochondrial reserve capacity')		52			
Night temp (-15, 5 or 25°C)			2, 49	1.60	0.213
Night temp = -15°C	0.408 \pm 0.023	17			
Night temp = 5°C	0.419 \pm 0.033	17			
Night temp = 25°C	0.462 \pm 0.032	18			
R-L control efficiency ('phosphorylating capacity')		52			
Night temp (-15, 5 or 25°C)			2, 49	2.92	0.063
Night temp = -15°C	0.858 \pm 0.014	17			
Night temp = 5°C	0.811 \pm 0.020	17			
Night temp = 25°C	0.834 \pm 0.019	18			
E-L coupling efficiency ('tightness of electron transport')		52			
Night temp (-15, 5 or 25°C)			2, 49	1.90	0.160
Night temp = -15°C	0.915 \pm 0.009	17			
Night temp = 5°C	0.892 \pm 0.013	17			
Night temp = 25°C	0.911 \pm 0.012	18			

TABLE 2 Parameter estimates, sample sizes (*n*), test statistics (*F*), degrees of freedom (df), and resultant *p*-values from linear models testing how a sudden change in night temperature affected mitochondrial respiration parameters and flux control ratios in great tit whole blood the morning after.

Note: Samples were assayed at a normothermic bird body temperature (41°C). Model-estimated means are presented \pm 1 standard error (SEM). Effects for which $0.05 > p < 0.10$ are given in italics.

TABLE 3 Parameter estimates, sample sizes (n), degrees of freedom (df), test statistics (F/LRT), and resultant p -values from linear mixed-effects models used to investigate how mitochondrial respiration parameters and flux control ratios in great tit whole blood were affected by a change from a normothermic (41°C) to a hypothermic (35°C) thermal state in vitro.

Model and Variable	Estimate \pm SEM	$n_{\text{hyp}}/n_{\text{norm}}$	df	F/LRT	p	$\sigma_{\text{Bird}} \sigma_{\text{Residual}}$
ROUTINE (pmol O ₂ s ⁻¹ μ L ⁻¹)		57/52				
Body temperature \times thermal state			1, 53.0	0.04	0.850	
Body temperature (°C)	-0.002 \pm 0.019		1, 54.5	0.02	0.903	
Thermal state			1, 53.1	22.16	<0.0001	
Hypothermia (35°C)	0.499 \pm 0.019					
Normothermia (41°C)	0.567 \pm 0.019					
Bird ID (random)			1	41.23	<0.0001	0.124 0.074
OXPHOS (pmol O ₂ s ⁻¹ μ L ⁻¹)		57/52				
Body temperature \times thermal state			1, 53.3	0.29	0.592	
Body temperature (°C)	-0.002 \pm 0.016		1, 54.5	0.02	0.879	
Thermal state			1, 53.3	0.11	0.744	
Hypothermia (35°C)	0.476 \pm 0.017					
Normothermia (41°C)	0.472 \pm 0.017					
Bird ID (random)			1	34.74	<0.0001	0.104 0.068
ETS (pmol O ₂ s ⁻¹ μ L ⁻¹)		57/52				
Body temperature \times thermal state			1, 51.7	0.10	0.748	
Body temperature (°C)	0.054 \pm 0.048		1, 54.2	1.25	0.268	
Thermal state			1, 52.4	0.33	0.570	
Hypothermia (35°C)	1.050 \pm 0.047					
Normothermia (41°C)	1.040 \pm 0.048					
Bird ID (random)			1	93.27	<0.0001	0.337 0.104
LEAK (pmol O ₂ s ⁻¹ μ L ⁻¹)		57/52				
Body temperature \times thermal state			1, 54.7	3.16	0.081	
Body temperature (°C)	0.000 \pm 0.005		1, 54.3	0.00	0.971	
Thermal state			1, 54.2	173.24	<0.0001	
Hypothermia (35°C)	0.023 \pm 0.005					
Normothermia (41°C)	0.096 \pm 0.005					
Bird ID (random)			1	14.28	0.0002	0.027 0.028
E-R control efficiency ('mitochondrial reserve capacity')		57/52				
Body temperature \times thermal state			1, 54.1	0.02	0.882	
Body temperature (°C)	0.041 \pm 0.010		1, 54.5	16.62	0.0001	
Thermal state			1, 53.7	51.34	<0.0001	
Hypothermia (35°C)	0.505 \pm 0.011					
Normothermia (41°C)	0.434 \pm 0.011					
Bird ID (random)			1	21.79	<0.0001	0.062 0.051
R-L control efficiency ('phosphorylating capacity')		57/52				

(Continues)

TABLE 3 (Continued)

Model and Variable	Estimate ± SEM	$n_{\text{hyp}}/n_{\text{norm}}$	df	F/LRT	<i>p</i>	$\sigma_{\text{Bird}} \sigma_{\text{Residual}}$
Body temperature × thermal state			1, 55.5	4.57	0.037	
Hypothermia (35°C)	0.006 ± 0.007					
Normothermia (41°C)	−0.012 ± 0.008					
Body temperature (°C)						
Thermal state						
Hypothermia (35°C)	0.956 ± 0.007 ^a					
Normothermia (41°C)	0.834 ± 0.007 ^a					
Bird ID (random)			1	7.44	0.006	0.033 0.042
E-L coupling efficiency ('tightness of electron transport')		57/52				
Body temperature × thermal state			1, 56.0	0.83	0.365	
Body temperature (°C)	0.003 ± 0.004		1, 53.0	0.72	0.400	
Thermal state			1, 55.0	194.57	<0.0001	
Hypothermia (35°C)	0.978 ± 0.004					
Normothermia (41°C)	0.907 ± 0.004					
Bird ID (random)			1	4.02	0.045	0.016 0.027

Note: Model-estimated means are presented ±1 standard error (SEM). Significant effects (i.e., $p \leq 0.05$) are presented in bold font, and effects for which $0.05 > p > 0.10$ are given in italics. Degrees of freedom were calculated using the Kenward-Roger approximation. The test statistic is *F* for all fixed effects. For the random term, the test statistic is the likelihood ratio (LRT). Note that *p*-values for main effects were not provided when the interaction was significant.

Abbreviations: σ_{Bird} , standard deviation of the random intercept; σ_{Residual} , residual standard deviation.

^aMain effect to be interpreted with caution due to involvement in an interaction with mean nightly body temperature.

the ROUTINE respiration that is channeled toward ATP production, to vary with night temperature ($p = 0.063$). Specifically, 86% of endogenous respiration was used to drive oxidative phosphorylation after a night at -15°C , which was higher than after a night at $+5^{\circ}\text{C}$ (81%) but similar to a night at $+25^{\circ}\text{C}$ (83%) (Table 2).

3.2 | Effects of thermal state and nightly body temperature

There were profound changes to mitochondrial respiration when the assay temperature changed from a normothermic to a hypothermic thermal state. ROUTINE respiration was 14% higher in normothermia ($0.567 \pm 0.019 \text{ pmol O}_2 \text{ s}^{-1} \mu\text{L}^{-1}$) than in hypothermia ($0.499 \pm 0.019 \text{ pmol O}_2 \text{ s}^{-1} \mu\text{L}^{-1}$) ($p < 0.0001$) (Figure 2A), that is, endogenous respiration was lower during in vitro hypothermia. However, neither OXPHOS nor ETS were affected by the thermal state ($p = 0.7$ and 0.6 , respectively) (Figures 2B,C). Instead, the drop in ROUTINE was mediated by a pronounced reduction in LEAK respiration ($p < 0.0001$), which was more than three times higher in normothermia ($0.096 \pm 0.004 \text{ pmol O}_2 \text{ s}^{-1} \mu\text{L}^{-1}$) compared to in hypothermia ($0.023 \pm 0.004 \text{ pmol O}_2 \text{ s}^{-1} \mu\text{L}^{-1}$) (Figure 2D).

E-R control efficiency was lower during hypothermia ($p < 0.0001$), meaning that a hypothermic bird had more mitochondrial reserve capacity compared to a normothermic bird (0.505 ± 0.012 and 0.434 ± 0.012 , of 1, respectively) (Figure 3A). The efficiency of oxidative phosphorylation was higher in the hypothermic state (i.e., R-L control efficiency increased) such that nearly all (96%) of endogenous respiration was used to produce ATP in hypothermia, compared to 83% in normothermia (Figure 3B). In other words, LEAK accounted for nearly 17% of endogenous respiration during normothermia, but in hypothermia, LEAK contributed only 4% to ROUTINE. E-L coupling efficiency (i.e., coupling of the electron transport system when maximal respiration was stimulated using FCCP) increased from 91% (0.907 ± 0.004) in normothermia to 98% (0.978 ± 0.004) in hypothermia (Figure 3C). Thus, the electron transport chain was tighter in the hypothermic state.

Most mitochondrial respiration traits were unaffected by nightly mean body temperature, both as a main effect and in interaction with thermal state (Table 3; Figure 4). However, E-R control efficiency increased with increasing body temperature (by 0.041 ± 0.010 per $^{\circ}\text{C}$) across thermal states (Figure 4D), suggesting that birds that were more hypothermic during the preceding night had mitochondria that respired at a greater proportion of maximum. There




FIGURE 2 Mitochondrial respiration traits in whole blood of winter-adapted, wild, great tits (*Parus major*) at a representative daytime normothermic body temperature (41°C) and a representative night-time hypothermic body temperature (35°C). The same individuals were measured in both thermal states. (A) baseline (“ROUTINE”) oxygen consumption during oxidative phosphorylation on endogenous substrates. (B) oxygen consumption devoted to ATP production alone (“OXPHOS”). (C) maximum working capacity of the electron transport system (“ETS”). (D) respiration used to offset proton leak across the inner mitochondrial membrane (“LEAK”). Solid plotting symbols with error bars represent model-predicted means ± 1 standard error. Gradient lines are plotted to show individual responses, but should not be taken as unequivocal evidence for linearity of change in the absence of measurement at intermediate temperatures. Sample sizes and statistics are reported in Table 3.

was also a thermal-state-dependent relationship between R-L control efficiency and body temperature, whereby oxidative efficiency decreased by 0.012 ± 0.008 per °C in the normothermic state but was unaffected by body temperature in the hypothermic state. However, when tested in assay temperature-specific models, the linear regression between R-L control and mean night-time body temperature was not significant (hypothermia: $p = 0.39$; normothermia: $p = 0.13$).

4 | DISCUSSION

We tested whether the plasticity of mitochondrial function could compensate for any reduced cellular respiration rate

during rest-phase hypothermia in great tit blood cells. As predicted, endogenous respiration (ROUTINE) was lower in the hypothermic thermal state (Figure 2A). Surprisingly, this was not a reflection of a general downregulation of cellular respiration during hypothermia, because there was no temperature-dependence of phosphorylating respiration (OXPHOS; in which ATP is produced) (Figure 2B). Instead, lower ROUTINE was primarily caused by a threefold reduction in LEAK (Figure 2D). Lack of temperature effects on OXPHOS contrasts previous studies on mammals, where a lowering of assay temperature by 10–12°C to simulate a deep hypothermic state was associated with a 50%–70% decrease of both phosphorylating and non-phosphorylating respiration.^{55,56} It has been proposed that, in hibernating mammals, active suppression of phosphorylating respiration only becomes noticeable at body temperatures below 30°C,³⁷ which could explain part of the thermal insensitivity of OXPHOS recorded here for a 6°C temperature reduction. If the thermal threshold concept applies also to birds, a logical conclusion is that passive thermal effects do not influence phosphorylating respiration strongly over the range of temperatures in our study. By contrast, leak respiration showed thermal sensitivity at, or beyond, that recorded in other studies,^{55,56} but our experimental design permits neither analysis of the contribution of passive thermal and actively regulated effects, nor firm conclusions related to the linearity of this response. Mechanisms aside, the combined actions of thermal insensitivity of OXPHOS and pronounced thermal sensitivity of LEAK meant that the hypothermic thermal state was associated with significant increases in both phosphorylating efficiency (i.e., R-L control efficiency was higher; Figure 3B) and coupling efficiency of electron transport when the cells were stimulated to work at a maximal rate (i.e., E-L coupling efficiency was higher; Figure 3C), broadly in keeping with results from mammal studies that manipulated assay temperature.^{55,56} In fact, nearly all respiration (96%) was directed toward ATP production during hypothermia, and the electron transport system was almost perfectly coupled (98%) in this thermal state. It is also interesting to note that we found no effect of simulated hypothermia on maximally stimulated mitochondrial respiration (i.e., ETS; Figure 2C), suggesting a lack of noticeable thermal sensitivity of the electron transport system. As a result, hypothermic cells had higher reserve capacity (Figure 3A), which could have functional significance in face of a challenge to energy metabolism. Collectively, our study indicates that the plasticity of mitochondrial leak respiration combined with thermal insensitivity of phosphorylating and maximal respiration may allow the little bird in winter to secure sufficient ATP production to meet thermoregulatory demands during rest-phase hypothermia or in other energetically stressful situations. This notion now needs confirmation in other tissues and by in vivo studies.




FIGURE 3 Flux control ratios for mitochondrial respiration in great tit whole blood in a normothermic (41°C) and a hypothermic (35°C) thermal state in vitro. (A) E-R control efficiency indicates how intensively the mitochondria were respiring under endogenous conditions (i.e., ROUTINE) relative to their maximum working capacity (i.e., ETS). For example, a value of 0.4 indicates that mitochondrial respiration can increase by 40% before reaching maximum. (B) The fraction of endogenous respiration that is used for oxidative phosphorylation where ATP is formed. (C) Coupling efficiency of electron transport when the ETS is stimulated to work at maximum capacity, where a value of 0 indicates a fully uncoupled system and a value of 1 a fully coupled system (i.e., where all respiration is channeled towards phosphorylation). Samples were collected from wild great tits and the same individual was measured in both thermal states. Solid plotting symbols with error bars represent model-predicted means \pm 1 standard error. Gradient lines show individual responses, but they are not necessarily meant to be suggestive of linearity of change. Sample sizes and statistics are reported in [Table 3](#).

Reduced LEAK and improved phosphorylating and coupling efficiency during hypothermia are in keeping with the hypothesis of biochemical compensation for thermal suppression of cell respiration at low body temperature. However, the tighter coupling can potentially result in increased production of reactive oxygen species (ROS) as a by-product of electron transport,^{21,57} provided that the resultant increase in protonmotive force is not dissipated through ATP synthase. Unless quenched, ROS causes oxidative stress with resultant macromolecular damage (e.g.,⁵⁸), which is related to many medical disorders.⁵⁹ It is interesting to speculate that a hypothermic bird might have to accept an oxidative cost for improved oxidative and coupling efficiencies. In this context, our study could provide proximate insights into why oxidative stress has been found to be higher in birds that are more hypothermic at night.⁶⁰ However, the role of mitochondria as sources of cellular ROS has been contended^{61–63} and empirical evidence suggests that the relationship between body temperature, ATP production, and ROS generation is not straightforward.^{31,64,65} In line with this, even though (pharmacological) uncoupling protects birds from cold-induced oxidative stress,⁶⁶ the evidence for clear fitness effects thereof is equivocal.^{67,68}

We found no significant effects of temperature of the preceding night on mitochondrial respiration traits and flux control ratios, though phosphorylating capacity tended to be the highest after the coldest night-time temperature ([Table 2](#)). This contrasts what has been recorded

as part of seasonal acclimatization of blood cell respiration in the same model system¹⁸ and in response to cold exposure in other tissue types.^{15–17} Responses in other studies were likely driven by seasonal upregulation of mitochondrial content (e.g.,¹⁸) and so it is possible that the period of exposure (one night) and acute nature of the air temperature manipulation were too short or too stochastic to serve as a reliable environmental cue prompting a change to cellular respiratory capacity. It is also possible that seasonal changes in mitochondrial content or respiration rate in the studies above were not causally related to cold temperature. In line with this, cold acclimation did not impact the respiration of liver mitochondria in the lesser hedgehog tenrec (*Echinops telfairi*).⁵⁵ Even so, the difference in short- and long-term responses of mitochondrial respiration coincident with colder environmental temperature suggests it would be interesting to study the temporal resolution of mitochondrial thermal acclimatization in some detail. For example, it is possible that, in the short term, mitochondrial function responds primarily to a change of tissue temperature in endothermic animals. This could be beneficial since endotherm body temperature is determined by the balance between heat production and heat loss rates, and so may be partly or fully independent of air temperature.

On average, there were few effects of night-time body temperature on mitochondrial respiration traits the day after ([Figure 4](#)). Similar results were obtained for liver mitochondria in the golden spiny mouse (*Acomys russatus*)




FIGURE 4 The relationship between mitochondrial respiration traits measured in great tit whole blood and mean body temperature of the individual during the preceding night. Panels (A–C) show these relationships for ROUTINE, ETS, and LEAK respiration, neither of which were statistically significant as denoted by dotted lines. Panel (D) shows the significant relationship with E-R control efficiency, an index of mitochondrial reserve capacity, denoted by the solid lines. Data were collected from wild great tits and respiration was assayed at each of a normothermic (41°C; red triangles) and a hypothermic (35°C; blue circles) thermal state *in vitro*. Sample sizes and statistics are reported in Table 3.

measured at a constant assay temperature.⁶⁹ Thus, for most of the respiration traits investigated here, between-individual variation in nighttime body temperature did not impact the thermal plasticity of blood cell mitochondria. One notable exception was the mitochondrial reserve capacity (i.e., E-R control efficiency), which decreased significantly with decreasing body temperature in both assay temperatures (Figure 4D). Thus, *in vitro*, a hypothermic thermal state was associated with *higher* surplus capacity (Figure 3A) whereas *in vivo*, more hypothermic birds had mitochondria with *lower* surplus capacity. This invites the speculation that the scope for plastic upregulation of mitochondrial work rate, e.g., in response to stochastic air temperature changes, may be limited by individual differences in the underlying mitochondrial phenotype. It remains to be tested if such constraints on plasticity hinder adaptation to the environment over short- and long-temporal scales. Finally, it

is interesting to note that the reduction in surplus mitochondrial capacity with decreasing body temperature is well in line with the observation that more hypothermic birds are typically of lower phenotypic quality.^{5,6} This raises the question of whether rest-phase hypothermia in the little bird in winter is an adaptive response to save energy or if it instead is an emergent phenotypic expression enforced by the inability to stay warm.

This study provides new perspectives to our understanding of how cellular and whole animal metabolism interact when the little bird in winter adapts to low temperatures. We conclude that *in vitro* hypothermia was associated with an immediate reduction in proton leak and a resultant increase in coupling and oxidative efficiency, which can explain why blood cell mitochondria could maintain phosphorylating respiration despite acute temperature reduction. Somewhat surprisingly, low assay temperature did not suppress OXPHOS or maximum working capacity of the electron transport system (i.e., ETS), leading to increased phosphorylating and respiratory scope during hypothermia (Figure 3). However, it is possible that such mitochondrial plasticity during rest-phase hypothermia may come at the cost of increased oxidative stress.⁶⁰ Future work should explore this hypothesis by direct measurement of ATP and ROS production, and membrane potential, to conclude on the consequences of changes to the efficiency of phosphorylating respiration for redox balance in normo- and hypothermic birds. Several endotherm studies show that mitochondrial function in blood cells correlates to that in tissues and organs with clearly defined roles in heat production and seasonal acclimatization (see Introduction), suggesting that data presented here may be representative of cellular responses to hypothermia at organismal levels. More broadly, our work aligns with the body of evidence highlighting the utility of blood cell respirometry to gain minimally invasive insight into functional ecology, evolutionary physiology, and gerontology in birds.^{18,40–43,70} Even so, it is important that future studies formally address the uniformity of the thermal plasticity of mitochondrial function across tissues and organs. Likewise, future investigations should aim to increasingly integrate tissue-based studies with an *in vivo* approach to understand mitochondrial regulation during hypothermia at the organismal level. More broadly, it would be relevant to increase taxonomic and phenotypic coverage of these studies, spanning cold-hardy, non-hypothermic, birds (e.g., Ref. [71]) to those that routinely use shallow (e.g., [72]) or deep (e.g., Refs. [73,74]) torpor.

AUTHOR CONTRIBUTIONS

AN conceived the idea and designed the protocol with input from EE. AN performed the fieldwork, and CCGD, IM, and AN performed the laboratory work. All authors interpreted the results. CCGD and AN analyzed the data

and wrote the first draft. CCGD and AN produced the graphic materials. All authors critically revised and edited the manuscript. AN procured funding.

ACKNOWLEDGMENTS

The authors thank Stefan Nord for managing the Räfte feeding stations by which birds for the study were caught. SN also assisted with fieldwork and provided miscellaneous technical support during the study. An anonymous reviewer provided critical comments that improved the manuscript. We thank Antoine Stier for helpful discussions.

FUNDING INFORMATION

This study was supported by the Birgit and Hellmuth Hertz Foundation (grant no. 2017-39034), The Royal Physiographic Society in Lund (grant no. 2019-41011), the Swedish Research Council (grant no. 2020-04686) and Stiftelsen Lunds Djurskyddsfond (all to AN).

DATA AVAILABILITY STATEMENT

Data are deposited in Figshare: <https://doi.org/10.6084/m9.figshare.19902277>.

DISCLOSURES

The authors declare that they have no competing or financial interests.

ETHICS STATEMENT

All procedures were approved by the Malmö/Lund Animal Ethics Committee, acting under the jurisdiction of the Swedish Board of Agriculture (permit no. 9246-19). Bird ringing was permitted by the Swedish Bird Ringing Centre at the Natural History Museum of Sweden (license no. 723, to AN).

ORCID

Carmen C. García-Díaz  <https://orcid.org/0000-0002-7827-8104>

Imen Chamkha  <https://orcid.org/0000-0002-6919-585X>

Eskil Elmér  <https://orcid.org/0000-0001-7901-1826>

Andreas Nord  <https://orcid.org/0000-0001-6170-689X>

REFERENCES

- Ruf T, Geiser F. Daily torpor and hibernation in birds and mammals. *Biol Rev*. 2015;90(3):891-926. doi:10.1111/brv.12137
- Geiser F. *Ecological physiology of daily torpor and hibernation*. Springer Nature Switzerland AG; 2021:317.
- Brodin A. Theoretical models of adaptive energy management in small wintering birds. *Phil Trans R Soc*. 2007;B362(1486):185-187.
- Nilsson JF, Nilsson J-Å, Broggi J, Watson H. Predictability of food supply modulates nocturnal hypothermia in a small passerine. *Biol Lett*. 2020;16(6):20200133. doi:10.1098/rsbl.2020.0133
- Nord A, Nilsson JF, Nilsson J-Å. Nocturnal body temperature in wintering blue tits is affected by roost-site temperature and body reserves. *Oecologia*. 2011;167(1):21-25.
- Nord A, Nilsson JF, Sandell MI, Nilsson J-Å. Patterns and dynamics of rest-phase hypothermia in wild and captive blue tits during winter. *J Comp Physiol B*. 2009;179(6):737-745. doi:10.1007/s00360-009-0357-1
- Reinertsen RE, Haftorn S. The effect of short-time fasting on metabolism and nocturnal hypothermia in the willow tit *Parus montanus*. *J Comp Physiol*. 1984;154(1):23-28.
- Reinertsen RE, Haftorn S. Nocturnal hypothermia and metabolism in the willow tit *Parus montanus* at 63 degrees N. *J Comp Physiol*. 1983;151(2):109-118.
- Haftorn S. Hypothermia of tits in the Arctic winter. *Orn Scand*. 1972;3(2):153-166.
- Cooper SJ, Gessaman JA. Nocturnal hypothermia in seasonally acclimatized mountain chickadees and juniper titmice. *Condor*. 2005;107(1):151-155.
- Brodin A, Nilsson J-Å, Nord A. Adaptive temperature regulation in the little bird in winter: predictions from a stochastic dynamic programming model. *Oecologia*. 2017;185(1):43-54. doi:10.1007/s00442-017-3923-3
- Welton NJ, Houston AI, Ekman J, McNamara JM. A dynamic model of hypothermia as an adaptive response by small birds to winter conditions. *Acta Biotheor*. 2002;50(1):39-56.
- Andreasson F, Nord A, Nilsson J-Å. Age differences in nighttime metabolic rate and body temperature in a small passerine. *J Comp Physiol B*. 2020;190(3):349-359. doi:10.1007/s00360-020-01266-5
- Nicholls DG, Ferguson SJ. *Bioenergetics*. 4th ed. Academic Press; 2013.
- Liknes ET, Swanson DL. Phenotypic flexibility in passerine birds: seasonal variation of aerobic enzyme activities in skeletal muscle. *J Therm Biol*. 2011;36(7):430-436. doi:10.1016/j.jtherbio.2011.07.011
- Wang Y, Shan S, Zhang H, Dong B, Zheng W, Liu J. Physiological and biochemical thermoregulatory responses in male Chinese hwameis to seasonal acclimatization: phenotypic flexibility in a small passerine. *Zool Stud*. 2019;58:e6. doi:10.6620/ZS.2019.58-06
- Zheng WH, Li M, Liu J-S, Shao S-L, Xu X-J. Seasonal variation of metabolic thermogenesis in Eurasian tree Sparrows (*Passer montanus*) over a latitudinal gradient. *Physiol Biochem Zool*. 2014;87(5):704-718. doi:10.1086/676832
- Nord A, Metcalfe NB, Page JL, Huxtable A, McCafferty DJ, Dawson NJ. Avian red blood cell mitochondria produce more heat in winter than in autumn. *FASEB J*. 2021;35(5):e21490. doi:10.1096/fj.202100107R
- Schaeffer PJ, Villarin JJ, Lindstedt SL. Chronic cold exposure increases skeletal muscle oxidative structure and function in *Monodelphis domestica*, a marsupial lacking Brown adipose tissue. *Physiol Biochem Zool*. 2003;76(6):877-887. doi:10.1086/378916
- Willis JS. Hibernation: cellular aspects. *Annu Rev Physiol*. 1979;41:275-286. doi:10.1146/annurev.ph.41.030179.001423
- Divakaruni AS, Brand MD. The regulation and physiology of mitochondrial proton leak. *Phys Ther*. 2011;26(3):192-205. doi:10.1152/physiol.00046.2010
- Porter RK, Brand MD. Causes of differences in respiration rate of hepatocytes from mammals of different body mass.

- Am J Physiol.* 1995;269(5):R1213-R1224. doi:10.1152/ajpregu.1995.269.5.R1213
23. Rolfe DF, Brand MD. Contribution of mitochondrial proton leak to skeletal muscle respiration and to standard metabolic rate. *Am J Physiol.* 1996;271(4):C1380-C1389. doi:10.1152/ajpce.1996.271.4.C1380
 24. Stager M, Cheviron ZA. Is there a role for sarcolipin in avian facultative thermogenesis in extreme cold? *Biol Lett.* 2020;16:20200078.
 25. Hohtola E. Shivering thermogenesis in birds and mammals. In: Barnes BM, Carey HV, eds. *Life in the Cold—Mechanisms, Adaptation and Application.* University of Alaska; 2004:241-252.
 26. Teulier L, Rouanet JL, Letexier D, et al. Cold-acclimation-induced non-shivering thermogenesis in birds is associated with upregulation of avian UCP but not with innate uncoupling or altered ATP efficiency. *J Exp Biol.* 2010;14:2476-2482.
 27. Nowack J, Giroud S, Arnold W, Ruf T. Muscle non-shivering thermogenesis and its role in the evolution of endothermy. *Front Physiol.* 2017;8:889.
 28. Roussel D, Marmillot V, Monternier P-A, et al. Skeletal muscle metabolism in sea-acclimatized king penguins. II. Improved efficiency of mitochondrial bioenergetics. *J Exp Biol.* 2020;223(21):jeb233684. doi:10.1242/jeb.233684
 29. Bourguignon A, Rameau A, Toullec G, Romestaing C, Roussel D. Increased mitochondrial energy efficiency in skeletal muscle after long-term fasting: its relevance to animal performance. *J Exp Biol.* 2017;220(13):2445-2451. doi:10.1242/jeb.159087
 30. Monternier PA, Marmillot V, Rouanet JL, Roussel D. Mitochondrial phenotypic flexibility enhances energy savings during winter fast in king penguin chicks. *J Exp Biol.* 2014;217(15):2691-2697. doi:10.1242/jeb.104505
 31. Boël M, Romestaing C, Duchamp C, et al. Improved mitochondrial coupling as a response to high mass-specific metabolic rate in extremely small mammals. *J Exp Biol.* 2020;223(5):jeb215558. doi:10.1242/jeb.215558
 32. Glanville EJ, Seebacher F. Compensation for environmental change by complementary shifts of thermal sensitivity and thermoregulatory behaviour in an ectotherm. *J Exp Biol.* 2006;209(24):4869-4877. doi:10.1242/jeb.02585
 33. Seebacher F. A review of thermoregulation and physiological performance in reptiles: what is the role of phenotypic flexibility? *J Comp Physiol B.* 2005;175(7):453-461. doi:10.1007/s00360-005-0010-6
 34. Bouchard P, Guderley H. Time course of the response of mitochondria from oxidative muscle during thermal acclimation of rainbow trout, *Oncorhynchus mykiss.* *J Exp Biol.* 2003;206(Pt 19):3455-3465. doi:10.1242/jeb.00578
 35. St-Pierre J, Brand MD, Boutilier RG. The effect of metabolic depression on proton leak rate in mitochondria from hibernating frogs. *J Exp Biol.* 2000;203(9):1469-1476.
 36. Bishop T, St-Pierre J, Brand MD. Primary causes of decreased mitochondrial oxygen consumption during metabolic depression in snail cells. *Am J Physiol.* 2002;282(2):R372-R382. doi:10.1152/ajpregu.00401.2001
 37. Staples JF, Brown JCL. Mitochondrial metabolism in hibernation and daily torpor: a review. *J Comp Physiol B.* 2008;178(7):811-827.
 38. McKechnie AE, Lovegrove BG. Avian facultative hypothermic responses: a review. *Condor.* 2002;104(4):705-724.
 39. Nord A, Chiriac S, Hasselquist D, Nilsson J-Å. Endotoxin injection attenuates rest-phase hypothermia in wintering great tits through the onset of fever. *Funct Ecol.* 2013;27(1):236-244. doi:10.1111/1365-2435.12003
 40. Casagrande S, Stier A, Monaghan P, et al. Increased glucocorticoid concentrations in early life cause mitochondrial inefficiency and short telomeres. *J Exp Biol.* 2020;223(15):jeb222513. doi:10.1242/jeb.222513
 41. Stier A, Bize P, Hsu B-Y, Ruuskanen S. Plastic but repeatable: rapid adjustments of mitochondrial function and density during reproduction in a wild bird species. *Biol Lett.* 2019;15(11):20190536. doi:10.1098/rsbl.2019.0536
 42. Udino E, George JM, McKenzie M, et al. Prenatal acoustic programming of mitochondrial function for high temperatures in an arid-adapted bird. *Proc R Soc Lond B.* 2021;288:20211893. doi:10.1098/rspb.2021.1893
 43. Dawson NJ, Salmón P. Age-related increase in mitochondrial quantity may mitigate a decline in mitochondrial quality in red blood cells from zebra finches (*Taeniopygia guttata*). *Exp Gerontol.* 2020;133:110883. doi:10.1016/j.exger.2020.110883
 44. Stier A, Romestaing C, Schull Q, et al. How to measure mitochondrial function in birds using red blood cells: a case study in the king penguin and perspectives in ecology and evolution. *Meth Ecol Evol.* 2017;8(10):1172-1182. doi:10.1111/2041-210x.12724
 45. Stier A, Monaghan P, Metcalfe NB. Experimental demonstration of prenatal programming of mitochondrial aerobic metabolism lasting until adulthood. *Proc R Soc Lond B.* 2022;289(1970):20212679. doi:10.1098/rspb.2021.2679
 46. Karamercan MA, Weiss SL, Villarroel JP, et al. Can peripheral blood mononuclear cells be used as a proxy for mitochondrial dysfunction in vital organs during hemorrhagic shock and resuscitation? *Shock.* 2013;40(6):476-484. doi:10.1097/shk.000000000000026
 47. Petit M, Lewden A, Vézina F. How does flexibility in body composition relate to seasonal changes in metabolic performance in a small passerine wintering at northern latitude? *Physiol Biochem Zool.* 2014;87(4):539-549. doi:10.1086/676669
 48. Swedish Meteorological and Hydrological Institute (SMHI). Unpublished data. Accessed February 01, 2022 <http://www.smhi.se/>
 49. Nord A, Chamkha I, Elmér E. A whole blood approach improves speed and accuracy when measuring mitochondrial respiration in intact avian blood cells. *FASEB J.* 2023;37:e22766. doi:10.1096/fj.202201749R
 50. Prinzing R, Pressmar A, Schleucher E. Body temperature in birds. *Comp Biochem Physiol A.* 1991;99(4):499-506.
 51. Gnaiger E. Mitochondrial pathways and respiratory control: an Introduction to OXPHOS analysis. 5th ed. 2020.
 52. Bates D, Maechler M, Bolker B, Walker S. Fitting linear mixed-effects models using lme4. *J Stat Softw.* 2015;67:1-48.
 53. Halekoh U, Hojsgaard S. A Kenward-Roger approximation and parametric bootstrap methods for tests in linear mixed models—the R package pbrtest. *J Stat Softw.* 2014;59(9):1-32.
 54. Lenth RV. emmeans: estimated marginal means, aka least-squares means. R package version 1.3.3. 2019 <https://CRAN.R-project.org/package=emmeans>
 55. Polymeropoulos ET, Oelkrug R, Jastroch M. Mitochondrial proton leak compensates for reduced oxidative power during frequent hypothermic events in a protoendothermic mammal, *Echinops telfairi.* *Front Physiol.* 2017;8:909.

56. Dufour S, Rouse N, Canioni P, Diolez P. Top-down control analysis of temperature effect on oxidative phosphorylation. *Biochem J*. 1996;314:743-751.
57. Demine S, Renard P, Arnould T. Mitochondrial uncoupling: a key controller of biological processes in physiology and diseases. *Cell*. 2019;8(8):795. doi:10.3390/cells8080795
58. Brand MD, Buckingham JA, Esteves TC, et al. Mitochondrial superoxide and aging: uncoupling-protein activity and superoxide production. *Biochem Soc Symp*. 2004;71:203-213. doi:10.1042/bss0710203
59. Grasemann H, Holguin F. Oxidative stress and obesity-related asthma. *Paed Resp Rev*. 2021;37:18-21. doi:10.1016/j.prrv.2020.05.004
60. Zagkle E, Grosiak M, Bauchinger U, Sadowska ET. Rest-phase hypothermia reveals a link between aging and oxidative stress: a novel hypothesis. *Front Physiol*. 2020;11:575060. doi:10.3389/fphys.2020.575060
61. Zhang Y, Wong HS. Are mitochondria the main contributor of reactive oxygen species in cells? *J Exp Biol*. 2021;224(5):eb221606. doi:10.1242/jeb.221606
62. Mookerjee SA, Divakaruni AS, Jastroch M, Brand MD. Mitochondrial uncoupling and lifespan. *Mech Ageing Dev*. 2010;131(7-8):463-472. doi:10.1016/j.mad.2010.03.010
63. Brown GC, Borutaite V. There is no evidence that mitochondria are the main source of reactive oxygen species in mammalian cells. *Mitochondrion*. 2012;12(1):1-4. doi:10.1016/j.mito.2011.02.001
64. Roussel D, Voituron Y. Mitochondrial costs of being hot: effects of acute thermal change on liver bioenergetics in toads (*Bufo bufo*). *Front Physiol*. 2020;11:153. doi:10.3389/fphys.2020.00153
65. Boël M, Veyrunes F, Durieux A-C, Freyssenet D, Voituron Y, Roussel D. Does high mitochondrial efficiency carry an oxidative cost? The case of the African pygmy mouse (*Mus mattheyi*). *Comp Biochem Physiol A*. 2022;264:111111. doi:10.1016/j.cbpa.2021.111111
66. Stier A, Massemin S, Criscuolo F. Chronic mitochondrial uncoupling treatment prevents acute cold-induced oxidative stress in birds. *J Comp Physiol B*. 2014;184(8):1021-1029. doi:10.1007/s00360-014-0856-6
67. Stier A, Bize P, Roussel D, Schull Q, Massemin S, Criscuolo F. Mitochondrial uncoupling as a regulator of life-history trajectories in birds: an experimental study in the zebra finch. *J Exp Biol*. 2014;217(19):3579-3589. doi:10.1242/jeb.103945
68. Stier A, Bize P, Massemin S, Criscuolo F. Long-term intake of the illegal diet pill DNP reduces lifespan in a captive bird model. *Comp Biochem Physiol C*. 2021;242:108944. doi:10.1016/j.cbpc.2020.108944
69. Grimpo K, Kutschke M, Kastl A, et al. Metabolic depression during warm torpor in the Golden spiny mouse (*Acomys russatus*) does not affect mitochondrial respiration and hydrogen peroxide release. *Comp Biochem Physiol A*. 2014;167:7-14.
70. Stier A, Bize P, Schull Q, et al. Avian erythrocytes have functional mitochondria, opening novel perspectives for birds as animal models in the study of ageing. *Front Zool*. 2013;10(1):33. doi:10.1186/1742-9994-10-33
71. Nord A, Folkow LP. Seasonal variation in the thermal responses to changing environmental temperature in the world's northernmost land bird. *J Exp Biol*. 2018;221(1):jeb171124. doi:10.1242/jeb.171124
72. Romano AB, Hunt A, Welbergen JA, Turbill C. Nocturnal torpor by superb fairy-wrens: a key mechanism for reducing winter daily energy expenditure. *Biol Lett*. 2019;15(6):20190211. doi:10.1098/rsbl.2019.0211
73. Hiebert SM. Energy costs and temporal organization of torpor in the rufous hummingbird (*Selasphorus rufus*). *Physiol Zool*. 1990;63:1082-1097.
74. Brigham RM. Daily torpor in a free-ranging goatsucker, the common poorwill (*Phalaenoptilus nuttallii*). *Physiol Zool*. 1992;65(2):457-472. doi:10.2307/30158263

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: García-Díaz CC, Chamkha I, Elmér E, Nord A. Plasticity of mitochondrial function safeguards phosphorylating respiration during in vitro simulation of rest-phase hypothermia. *The FASEB Journal*. 2023;37:e22854. doi:10.1096/fj.202201613R