



# LUND UNIVERSITY

## Deaths caused by Single Stab Injuries to the Trunk - Homicide or Suicide?

Berg von Linde, Maria; Wingren, Carl Johan

2021

*Document Version:*

Publisher's PDF, also known as Version of record

[Link to publication](#)

*Citation for published version (APA):*

Berg von Linde, M., & Wingren, C. J. (2021). *Deaths caused by Single Stab Injuries to the Trunk - Homicide or Suicide?*. Poster session presented at Deutschen Gesellschaft für rechtsmedizin, München, Germany.

*Total number of authors:*

2

*Creative Commons License:*

Unspecified

**General rights**

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

**Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117  
221 00 Lund  
+46 46-222 00 00

# Deaths caused by Single Stab Injuries to the Trunk - Homicide or Suicide?

Maria Berg von Linde, MD, Department of Forensic Medicine in Lund, Swedish National Board of Forensic Medicine, Sweden, email: [maria.bergvonlinde@rmv.se](mailto:maria.bergvonlinde@rmv.se)  
Carl Johan Wingren, MD, PhD, Department of Forensic Medicine in Lund, Swedish National Board of Forensic Medicine and Lund University, Sweden

## Introduction

To identify the manner of death in single stab injuries is a challenge for forensic pathologists. Previous comparative studies (1)(2) are limited.

We aimed to define variables to differentiate between homicidal and suicidal deaths in single stab injuries to the trunk.

## Method

Using the Swedish forensic autopsy register, we identified 75 homicides and 38 suicides that died between 2010 and 2019 following a single stab injury to the trunk.

Logistic regression models were performed.



## Results

We identified variables with conclusive associations to homicide, namely victim aged 20-44 years, scene of death located outdoors, injury affecting bone/cartilage of the ribcage, injury involving vasculature and positive alcohol results (table 1). The majority of all the stab injuries were located to the anterior thorax, from which a large proportion were located at the left side (figure 1 and 2). There were no suicidal stab injuries located axillary or posterior.



Table 1. Variables with conclusive associations to homicide

Variable (reference)	Homicide OR (95% CI)
20-44 years (45-65 years)	2.77 (1.09-7.04)
Found outdoors (victim's home)	15.74 (5.09-48.72)
Injury affecting the ribcage (intercostal space)	4.34 (1.71-10.99)
Injury involving vessel (not involving vessel)	8.28 (2.67-25.64)
Positive alcohol results (negative alcohol)	8.14 (2.85-23.7)



## Discussion and conclusion

We were able to reproduce several findings from previous research on single stab injuries (1)(2). The variables with conclusive associations to homicide could be of value in determining the manner of death in cases of single stab injury to the trunk.

We also attempted to use variables to frame a victim profile. If the following criteria were met, the victim showed a 58.7% sensitivity and 92.1% specificity of being subjected to homicide:

- ✓ male
- ✓ <=65 years
- ✓ positive ethanol/narcotics
- ✓ not found in victim's home

A weakness of the study is the small population and thereby low statistical power of some analyses. Therefore, further research based on larger case series is needed.