



# LUND UNIVERSITY

## Blockade of CD27/CD70 pathway to reduce the generation of memory T cells and markedly prolong the survival of heart allografts in presensitized mice

Dai, Helong; Chen, Jibing; Shao, Wei; Wang, Feng; Xu, Shuo; Peng, Yuanzheng; Lin, Yingying; Xia, Junjie; Ekberg, Henrik; Wang, Xiaomin; Qi, Zhongquan

*Published in:*  
Transplant Immunology

*DOI:*  
[10.1016/j.trim.2011.02.002](https://doi.org/10.1016/j.trim.2011.02.002)

2011

[Link to publication](#)

### *Citation for published version (APA):*

Dai, H., Chen, J., Shao, W., Wang, F., Xu, S., Peng, Y., Lin, Y., Xia, J., Ekberg, H., Wang, X., & Qi, Z. (2011). Blockade of CD27/CD70 pathway to reduce the generation of memory T cells and markedly prolong the survival of heart allografts in presensitized mice. *Transplant Immunology*, 24(4), 195-202. <https://doi.org/10.1016/j.trim.2011.02.002>

*Total number of authors:*  
11

### **General rights**

Unless other specific re-use rights are stated the following general rights apply: Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117  
221 00 Lund  
+46 46-222 00 00

## **Title page**

### **Blockade of CD27/CD70 pathway to reduce the generation of memory T cells and markedly prolong the survival of heart allografts in presensitized mice**

Helong Dai<sup>a\*</sup>, Jibing Chen<sup>b\*</sup>, Wei Shao<sup>a</sup>, Feng Wang<sup>a</sup>, Shuo Xu<sup>a</sup>, Yuanzheng Peng<sup>a</sup>, Yingying Lin<sup>a</sup>, Junjie Xia<sup>a</sup>, Henrik Ekberg<sup>c</sup>, Xiaomin Wang<sup>d#</sup> and Zhongquan Qi<sup>a#</sup>

<sup>a</sup>Organ Transplantation Institute, Xiamen University, Fujian Province, P. R. China

<sup>b</sup>Basic Medical Department of Medical College, Xiamen University, Fujian Province, P. R. China

<sup>c</sup>Department of Nephrology and Transplantation, Skane University Hospital, Malmö, Sweden

<sup>d</sup> Department of Hepato-Biliary Surgery, Affiliated Zhongshan Hospital of Xiamen University, Xiamen City, Fujian Province, P. R. China

\*Both authors contributed equally to the study and share first authorship.

**Corresponding author:** <sup>#</sup>Zhongquan Qi, Organ Transplantation Institute, Xiamen University, Fujian Province 361005, PR China. Phone 86-0592-2180126, Email: oti@xmu.edu.cn or <sup>#</sup>Xiaomin Wang, Department of Hepato-Biliary surgery, Affiliated Zhongshan Hospital of Xiamen University, Xiamen City, Fujian Province 361005, PR China. Email: wxm@xmzsh.com.

**Helong Dai**, performed research and wrote the paper

**Jibing Chen**, designed research and modified the paper

**Wei Shao**, performed qRT-PCR

**Feng Wang**, performed heart transplantation

**Shuo Xu**, performed mixed lymphocyte reaction (MLR) assays

**Yuanzheng Peng**, performed flow cytometry

**Yingying Lin**, performed ELISA

**Junjie Xia**, modified the paper

**Henrik Ekberg**, contributed important reagents

**Xiaomin Wang**, contributed supporting funds

**Zhongquan Qi**, contributed important reagents and supporting funds

**Running title:** Blockade of CD27/CD70 pathway to achieve long-term survival

## **Abstract**

**Background.** Alloreactive memory T cells are a major obstacle to transplantation acceptance due to their capacity for accelerated rejection.

**Methods.** C57BL/6 mice that had rejected BALB/c skin grafts 4 weeks earlier were used as recipients. The recipient mice were treated with anti-CD154/LFA-1 with or without anti-CD70 during the primary skin transplantation and anti-CD154/LFA-1 or not during the secondary transplantation of BALB/c heart. We evaluated the impact of combinations of antibody-mediated blockade on the generation of memory T cells and graft survival after fully MHC-mismatched transplantations.

**Results.** One month after the primary skin transplantation, the proportions of CD4<sup>+</sup> memory T cells/CD4<sup>+</sup> T cells and CD8<sup>+</sup> memory T cells/CD8<sup>+</sup> T cells in the anti-CD154/LFA-1 combination group were 47.32±4.28% and 23.18±2.77%, respectively. In the group that included anti-CD70 treatment, the proportions were reduced to 34.10±2.71% and 12.19±3.52% (P<0.05 when comparing the proportion of memory T cells between the two groups). The addition of anti-CD70 to the treatment regimen prolonged the mean survival time following secondary heart transplantation from 10 days to more than 90 days (P<0.001). Furthermore, allogenic proliferation of recipient splenic T cells and graft-infiltrating lymphocytes were significantly decreased. Meanwhile, the proportion of regulatory T cells were increased to 9.46±1.48% on day 100 post-transplantation (P<0.05).

**Conclusions.** The addition of anti-CD70 to the anti-CD154/LFA-1 combination given during the primary transplantation reduced the generation of memory T cells. This therapy regimen provided a potential means to alleviate the accelerated rejection mediated by memory T cells during secondary heart transplantation and **markedly prolong the survival of heart allografts.**

**Keywords.** costimulatory blockade; anti-CD70;memory T cells; heart transplantation.

## **Introduction**

Transplant patients may develop alloreactive memory T cells after exposure to alloantigens during previous transplantations, blood transfusions, pregnancies, or due to continuous exposure to bacterial and viral pathogens[1,2]. In adult humans, 40–50% of T cells circulating in the peripheral blood have memory phenotypes[3,4]. Memory T cells, including CD4<sup>+</sup> and CD8<sup>+</sup> subsets, play a key role in accelerated rejection. Furthermore, there is a growing body of evidence that B-cell production of alloantibodies is also an important element in the accelerated rejection[5]. Therefore, inhibiting the generation of memory T cells and alloantibodies is a critical step in the prevention of rejection.

A variety of methods to induce memory cells tolerance in mouse models have been identified, such as the use of antilymphocyte serum and antithymocyte globulin for general clearance of T and B lymphocytes [6], anti-CD122 to deplete CD8<sup>+</sup>CD122<sup>+</sup> memory T cells [7,8], and anti-CD20 to deplete B-lymphocytes during treatment of autoimmune diseases [9,10]. Here, we developed a protocol designed to minimize the generation of memory T cells and production of alloantibodies during primary transplantation in order to significantly prolong the secondary allograft survival.

The CD40/CD154 and LFA-1/ICAM-1 pathways have been shown to play an important role in the activation of T cells [11-13]. The CD40/CD154 pathway also provides important signals regulating B lymphocytes clonal expansion, antibody production and isotype switching, as well as the development of humoral memory[14,15]. CD70 (CD27 ligand) is a type II transmembrane glycoprotein belonging to the TNF family that is induced upon T and B cell activation [16]. CD27/CD70 engagement has been implicated in T-cell development, T-cell activation, and T-cell-dependent antibody production by B cells [17,18]. CD70 blockade can inhibit clonal expansion of CD8<sup>+</sup> T cells and reduce the generation of memory CD8<sup>+</sup> T cells [19,20].

In this study, we attempted to reduce the generation of memory T cells and alloantibodies by treating mice with anti-CD70 added in combination with an anti-CD154/LFA-1 regimen during primary transplantation. We provided evidence that this approach could alleviate the accelerated rejection mediated by memory T cells during

secondary heart transplantation, and we explored the possible mechanism of the obviously prolonged survival.

## **MATERIALS AND METHODS**

### **Animals**

Female C57BL/6 (B6, H-2<sup>b</sup>) and BALB/c(H-2<sup>d</sup>) mice (8–12 weeks old) were purchased from Slac Laboratory Animal Co. Ltd. (Shanghai, China) and used as graft recipients and donors, respectively. All animals were maintained and bred in a pathogen-free facility, and all procedures were performed according to the Institutional Animal Care and Use Committee (IACUC) guidelines.

### **Antibodies**

All administered antibodies were produced by Bioexpress (West Lebanon, NH, USA), including anti-CD70 (FR-70), anti-CD154 (MR-1), anti-LFA-1(M17/4), and their respective isotype controls. Antibodies used for flow cytometric analysis, including FITC-anti-CD4 (GK1.5), FITC-anti-CD8 (53-6.7), PE-anti-CD44 (IM7), PECy5-anti-CD62L (MEL14), PE-anti-IgM (RMM-1), FITC-anti-IgG1 (RMG1-1), FITC-anti-IgG2a (RMG2a-62), and their isotype controls were purchased from Biolegend (San Diego, CA, USA). CD4<sup>+</sup>Foxp3<sup>+</sup> regulatory T cells (Tregs) were detected using the Mouse Regulatory T cell Staining Kit from eBioscience (San Diego, CA, USA).

### **Skin transplantation**

Full-thickness skin grafts were prepared from the lateral thoracic skin of BALB/c mice, cut into circular pieces (~1.2 cm<sup>2</sup> in area), and engrafted onto the lumbar region of B6 mice. After the fully MHC-mismatched transplantation, the B6 mice received a 2-antibody treatment regimen consisting of 0.25 mg of anti-CD154 and 0.1 mg of anti-LFA-1, or a 3-antibody treatment regimen consisting of 0.25 mg of anti-CD154, 0.1 mg of anti-LFA-1, and 0.25 mg of anti-CD70. Control group mice were treated

with isotype antibodies (Table 1). The drugs were intraperitoneally (i.p.) administered on days 0 and 2 post-transplantation.

### **Alloantigen-primed heart transplantation model**

Four weeks after skin grafting, the B6 mice were defined as alloantigen-primed mice. Vascularized heterotopic heart transplantations from BALB/c donors to B6 recipients were performed with anastomosis to the vessels of the neck using a non-suture cuff technique as described previously [21]. The mice were treated with 0.25 mg anti-CD154 and 0.1 mg of anti-LFA-1 or isotype controls on days 0, 2, 4, and 6 post-transplantation (Table 1). Graft survival was monitored by daily palpation. Rejection was defined as complete loss of palpable heart beat.

### **Mixed lymphocyte reactions (MLR)**

T lymphocytes were isolated from spleens of B6 mice using nylon wool columns (Wako, Osaka, Japan) and used as responder cells. Donor spleen cells were used as stimulator cells and treated with mitomycin(40  $\mu$ g/ml, Amresco, Solon, OH, USA) before used in the MLR assay. For proliferation assays,  $10^5$  stimulator cells were cultured with  $5 \times 10^5$  responder cells in RPMI1640 media supplemented with 10% fetal bovine serum (FBS) and 1% penicillin and streptomycin in 96-well plates. Three separate wells were dedicated to each responder-stimulator combination and each experiment was repeated three times. Cells were incubated for 72 h at 37°C in 95% humidified air mixed with 5% carbon dioxide. After 72 h of culture, cell proliferation was quantified using a BrdU kit (Roche Applied Science, Mannheim, Germany).

### **Enzyme-linked immunosorbent assay (ELISA)**

ELISAs were performed using commercially available kits (NeoBioscience Technology Limited Company, Shenzhen, China) to detect the concentrations of IL-2, IFN- $\gamma$ , IL-10, and TGF-  $\beta$  in the supernatants from MLR according to the manufacturer's instructions. A standard curve was generated using known amounts of

purified recombinant murine cytokines.

### **Graft pathological analysis**

The heart allografts were resected from the recipient mice on day 7 or day 100 post-transplantation. Tissues were fixed in 10% buffered formalin solution, embedded in paraffin, cut into 5  $\mu\text{m}$  sections and stained with hematoxylin and eosin (H&E). Graft rejection was graded on the extent of infiltration and the anatomical localization of inflammatory cells according to the International Society of Heart and Lung Transplantation (ISHLT) standard [22,23].

### **qRT-PCR**

RNA was isolated from the heart allografts using Trizol reagent (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's instructions. Reverse transcription and qRT-PCR were performed using commercially available reagents (TOYOBO, Osaka, Japan) with the StepOne Real-Time PCR System (ABI, Foster, UK). Syber Green I was used to detect amplification and  $\beta$ -actin was used as a normalizing control. Calculations were performed using the  $2^{-\Delta\Delta\text{CT}}$  method. Each reaction was carried out in triplicate. The primer sequences used for the qRT-PCR were listed in Table 2.

### **Extraction of lymphocytes from heart allografts**

The harvested heart allografts were minced with a sterile blade and incubated in 10 ml buffered saline with 2% bovine serum albumin (BSA) and 2 mg/ml collagenase at 37°C for 2 h. The cells were strained through a 70  $\mu\text{m}$  nylon cell strainer (Becton Dickinson, Franklin Lakes, NJ, USA). Lymphocytes were isolated from these cells using EZ<sup>Sep</sup><sup>TM</sup> Mouse lymphocyte separation medium (Dakewe Biotech Company, shenzhen, China) with centrifugation for 20 min at 1600 rpm. After washed twice in RPMI 1640, lymphocytes were resuspended in phosphate buffered saline (PBS) with 10% fetal bovine serum for the Flow cytometry.



## **Flow cytometry**

A total of 30  $\mu$ l of sera from different mice in each group was collected following rejection of the allografts in the control group mice. BALB/c spleen cells ( $1 \times 10^6$  cells/100  $\mu$ l) were incubated with 10  $\mu$ l of recipient sera at 4°C for 30 min. After washing twice in PBS, the cells were incubated with PE-IgM and FITC-IgG1 or IgG2a antibodies at 4°C for 30 min. Approximately  $1 \times 10^6$  T lymphocytes from recipients and heart grafts were stained using fluorescent antibodies according to the manufacturer's instructions. At last, all of the positive cells were analyzed with a FACS instrument (Partec Co, Germany). Data were analyzed with Flow Jo 7.5 software.

## **Statistical analyses**

The mean survival times (MST) of the six groups were analyzed by the Kaplan-Meier method and Log-rank test. All other data were analyzed by one-way analysis of variance (ANOVA). Because multiple comparisons were made during the analysis, a Bonferroni correction was calculated and applied. A P value < 0.05 was considered statistically significant; P < 0.01 and P < 0.001 indicated highly significant differences. All analyses were performed using GraphPad Prism<sup>®</sup> (GraphPad, Inc., La Jolla, CA, USA) software.

## **Results**

### **Addition of anti-CD70 to anti-CD154/LFA-1 combination treatment during primary skin transplantation reduced the generation of memory T cells**

In order to determine whether anti-CD70 treatment would impact the generation of alloreactive effector memory T cells ( $CD44^{\text{high}}CD62L^{\text{low}}$ ), we isolated B6 mouse splenic T lymphocytes four weeks after skin transplantation and analyzed CD4, CD8, CD44, and CD62L expression using flow cytometry (Fig. 1). The proportions of

CD4<sup>+</sup> memory T cells/CD4<sup>+</sup> T cells and CD8<sup>+</sup> memory T cells/CD8<sup>+</sup> T cells in the S0 group mice following skin transplantation were 57.71±3.25% and 31.04±1.06%, respectively. In the S2 treated group, the proportions were 47.32±4.28% and 23.18±2.77%, respectively. Finally, in the S3 group, which had anti-CD70 included in the combination treatment, the proportions were reduced to 34.10±2.71% and 12.19±3.52%, respectively (Fig. 1). P<0.001 when comparing the proportion of CD4<sup>+</sup> and CD8<sup>+</sup> memory T cells between treatment group S3 to S0 group. P<0.05 when comparing the proportion of CD4<sup>+</sup> and CD8<sup>+</sup> memory T cells between S2 group to S0 and S3 groups.

### **Evaluations of groups without treatment during secondary heart transplantation**

To determine the impact of the reduction of effector memory T cells observed in the S3 group on secondary transplantation, we selected B6 mice which came from the three groups (S0, S2, S3) as recipients for heart transplantation. Mice from all three groups were treated with isotype control antibodies during the secondary heart transplantation. The mean survival times following heart transplantation were 3.5 d for the S0H0 control group, 4.5 d for the S2H0 treated group, and 6.5 d for the S3H0 treated group. The P value for the S3H0 vs. the S0H0 and S2H0 groups was less than 0.001. The P value for the S2H0 group vs. the S0H0 group was less than 0.05. (Fig. 2A ). Splens from recipient mice were removed on day 3.5 post-transplantation following identification of rejection in the control S0H0 group. The values of MLR in the S3H0 and S2H0 group were reduced compared to the S0H0 control group (Fig. 2B, P < 0.05). There were no statistical differences in IL-2 concentrations between groups. The concentration of IFN- $\gamma$  was reduced (Fig. 2B, P < 0.001). The proportions of CD4<sup>+</sup> memory T cells/CD4<sup>+</sup> T cells and CD8<sup>+</sup> memory T cells/CD8<sup>+</sup> T cells were reduced, as was previously observed following primary skin transplantation. P<0.001 when comparing the proportion of CD4<sup>+</sup> and CD8<sup>+</sup> memory T cells between treatment groups (S2H0 and S3H0) to positive control group (S0H0). P<0.05 when comparing the proportion of CD4<sup>+</sup> and CD8<sup>+</sup> memory T cells between S2H0 group to S3H0 group (Fig. 2C). Splenic Tregs were not significantly different between groups. The

alloantibodies IgG1 and IgG2a of recipient sera were nearly 100% in the control group. However, there were no IgG1 and IgG2a in the S2H0 and S3H0 groups, and alloantibodies IgM were not different between the groups.

**Addition of anti-CD70 to combination treatment during primary skin transplantation markedly prolonged survival when anti-CD154/LFA-1 were also given during secondary heart transplantation.**

Four weeks following primary skin transplantation, B6 mice selected as recipients for heart transplantation were treated with anti-CD154/LFA-1. The MST for group S3H2 was greater than 90 days, as compared to 6.5 days for group S0H2 and 10.0 days for group S2H2. These results indicated that only mice from group S3H2 which had received primary skin transplantation with anti-CD70 treatment could markedly prolong survival. The P value between each group was less than 0.001 (Fig. 3A ).

In order to discover the mechanism of obviously prolonged allografts survival induced by the treatment including anti-CD70, we performed functional tests on the immune cells of the recipient mice. First, we examined the proliferative responses of recipient splenic T cells to donor cells. Recipient splenic T cells were isolated at day 7 after heart transplantation. MLR results showed that mice that received anti-CD70 treatment during primary skin transplantation had lower cell proliferative responses than mice that did not receive anti-CD70 treatment (Fig. 3B,  $P < 0.05$ ). Furthermore, cell proliferation in the S3H2 group mice was further reduced on day 100 post-transplantation (Fig. 3B,  $P < 0.05$ ). TGF- $\beta$  concentrations in the MLR supernatants were increased in the S3H2 mice compared to the S2H2 mice that did not receive anti-CD70 (Fig. 3B,  $P < 0.05$ ). By contrast, IL-10 expression was not different between the groups in the supernatants. IL-2 concentrations in the MLR supernatants were reduced in the S3H2 and S2H2 mice compared to the S0H2 mice (Fig. 3B,  $P < 0.05$ ). The concentrations of IFN- $\gamma$  in the MLR supernatants were lower in the S3H2 mice compared to the S2H2 mice (Fig. 3B,  $P < 0.05$ ). Flow cytometric analysis of the splenic cell populations showed that the S3H2 100d group mice had a higher proportion of Tregs than mice in the S0H2, S2H2 and S3H2 groups (Fig. 3C,  $P < 0.05$  ). Furthermore, Treg populations in the S3H2 mice were increased to  $9.46 \pm 1.48\%$  on day 100 post-transplantation.  $P < 0.001$  when comparing the proportion

of CD4<sup>+</sup> memory T cells between S3H2 group to S0H2 and S2H2 groups.  $P < 0.01$  when comparing the proportion of CD8<sup>+</sup> memory T cells between S2H2 group to S0H2 group (Fig. 3C). Finally, the levels of alloantibodies IgG1, IgG2a and IgM detected in all groups were low.

### **Addition of anti-CD70 to combination treatment during primary skin transplantation effectively prevented lymphocyte infiltration and induced high Foxp3 gene expression in the graft**

As shown in Figure 4A, heart allografts tissue from mice in the S3H2 group showed low levels of lymphocytic infiltration and fewer changes in myocardial structural integrity compared to S2H2 groups ( $P < 0.01$ ). This result was consistent with the detection of low levels of graft-infiltrating lymphocytes by flow cytometry. The proportion of CD8<sup>+</sup> T cells/graft-infiltrated cells was  $43.38 \pm 4.83\%$  in the S2H2 group that had not receive anti-CD70 treatment. In contrast, the proportion was  $23.39 \pm 2.78\%$  in the S3H2 group mice on day 7 post-transplantation and was further reduced to  $13.44 \pm 2.99\%$  by day 100 post-transplantation.  $P < 0.01$  when comparing the proportion of CD8<sup>+</sup> T cells/graft-infiltrated cells between S3H2 group to S2H2 group.  $P < 0.001$  when comparing the proportion of CD8<sup>+</sup> T cells/graft-infiltrated cells between S3H2 100d group to S2H2 group. (Fig. 4B). There were very little CD4<sup>+</sup> T cells infiltration detected in the grafts (Fig. 4B). The intragraft immune response was dominated by effector T cells (data not shown). In order to determine the mechanism of graft protection, we examined graft rejection and tolerance-related gene expression. Compared with the no treatment group, the inflammatory factors (IL-2, IFN- $\gamma$ ) in S3H2 group were significantly suppressed (Fig. 4C,  $P < 0.001$ ). Treatment with anti-CD70 induced higher Foxp3 gene expression (Fig. 4C,  $P < 0.001$ ) and TGF- $\beta$  expression in the graft, although there was no difference in IL-10 expression between groups. TGF- $\beta$  levels remained high in the S3H2 group on day 100 post-transplantation (Fig. 4C).

### **Discussion**

Although the generation of memory T cells specific for pathogens is beneficial in providing protective immunity, memory T cells specific for alloantigens can be deleterious to the recipient of a transplanted organ. Graft rejection is accelerated and

allograft acceptance is more difficult to induce in hosts whose immune repertoires contain donor-reactive memory T cells [2,24].

Nicolls et al. demonstrated that combining monoclonal antibody therapies targeting CD154 and LFA-1 induced transplantation acceptance to pancreatic islet allografts [25]. Other studies had shown that anti-CD154 or anti-CD154/CD28 treatment could prevent the occurrence of anti-donor antibodies following xenogeneic transplantation [26,27]. Here, we showed that anti-CD154/LFA-1 and anti-CD70 combination treatment during primary skin transplantation reduced the generation of memory T cells. CD8<sup>+</sup> memory T cells in particular were maintained close to the level observed in naïve mice (Fig. 1). This was consistent with previous studies showing that blocking the CD27/70 pathway effectively inhibited the function and generation of memory T cells [28,29]. Furthermore, we found that this treatment regimen abrogated the production of IgG1/IgG2a alloantibodies in recipient mice, were consistent with a prior study by Nathan et al. demonstrating that anti-CD154 treatment could abrogate the production of alloantibodies [30]. CD154 is a key molecule in the contact-mediated signaling required for B cell activation and differentiation and can induce Ig isotype switching [31,32], while the CD27/CD70 pathway is important in stimulating B cell activation, plasma cell formation, and Ab production [33]. Therefore, the anti-CD154/LFA-1 and anti-CD70 regimen following primary skin transplantation provided the benefit of abrogating alloantibodies production along with reducing the generation of memory T cells.

Treatment with anti-CD154/LFA-1 and anti-CD70 during primary skin transplantation prolonged secondary graft survival compared to the control group, but the survival time was not very long (Fig. 2A). In view of these results, we treated mice with anti-CD154/LFA-1 during secondary heart transplantation to suppress effector T cell function, markedly prolong survival in presensitized mice.

The results obtained from the S0H2 group mice are comparable to transplantation patients who have a history of previous transplantations, blood transfusions, or pregnancies, as well as continuous exposure to bacterial and viral pathogens. These patients have higher levels of memory T cells and alloreactive antibodies in vivo which are barriers to secondary transplantation. In contrast, the S2H2 group mice are comparable to primary transplant patients given conventional

treatment, and the S3H2 group mice are comparable to primary transplant patients given treatments to decrease production of allogenic memory T cells. Repeated transplantation from the same donor would not be performed in human transplantation. However, our model could research the alloreactive memory T cells conveniently. Our study was performed in pathogen free mice and therefore did not study alloreactivity by viral specific memory T-cells. Therefore, it would be particularly useful to know if this regimen could also prolong survival in pathogen exposed mice. This could be particularly useful for our future work. Based on survival curves, mice in the S3H2 group which had anti-CD70 treatment combined with anti-CD154/LFA-1 during the primary skin transplantation could markedly prolong allograft survival (Fig. 3A). Nicolls et al. found that combined anti-CD154/LFA-1 therapy induced "dominant" transplantation acceptance in naïve mice and that this combination therapy did not deplete lymphocytes or require the long-term deletion of donor-reactive T lymphocytes to maintain allograft survival [25]. Here, we explored an effective approach designed to markedly prolong allograft survival in previous transplant recipients. Our novel method used anti-CD70 treatment combined with anti-CD154/LFA-1 treatment during primary skin transplantation to reduce the generation of memory T cells, and anti-CD154/LFA-1 treatment during secondary heart transplantation to inhibit the function of effector T cells.

To explore the mechanisms of the obviously prolonged allograft survival associated with our treatment regimen, we examined the proliferative response of recipient splenic T cells to donor antigens in the presensitized mice. Mice treated with anti-CD70 during primary skin transplantation had lower cell proliferation than mice that did not receive anti-CD70. Furthermore, cell proliferation remained low in the anti-CD70 treated group at day 100 post-transplantation (Fig. 3B). These results showed that anti-CD70 treatment effectively suppressed the recipient cellular immune response *in vivo*. As shown in Figure 3C, the group treated with anti-CD70 exhibited increased memory T cells and Tregs production on day of 100 post-transplantation. Moreover, the group treated with anti-CD70 had fewer graft-infiltrating CD8<sup>+</sup> T cells (Fig. 4B). Through these results, we speculated that the majority of T cells existed in the spleen as effector memory T cells [34] and were kept in balance with the Tregs population.

As we can see from the pathological analysis of allografts, the group treated with anti-CD70 showed only a small amount of lymphocytic infiltration and few changes in myocardial structural integrity compared to mice not treated with anti-CD70 (Fig. 4A,  $P < 0.01$ ). This was consistent with flow cytometric analysis of graft-infiltrating lymphocytes. These results showed that anti-CD70 treatment could reduce T cells infiltration to the allografts. We found few graft-infiltrating  $CD4^+$  T cells, suggesting that  $CD4^+$  T cells play a key role in secondary lymphoid organs. This was consistent with previous results published by Yalai et al. [35]. The Foxp3 gene expression was higher in the group treated with anti-CD70 (Fig. 4C,  $P < 0.001$ ). Moreover, the anti-CD70 treatment group exhibited higher levels of TGF- $\beta$  expression but not IL-10 expression in the grafts that were maintained until at least day 100 post-transplantation (Fig. 4C). This result, in combination with the high ratio of Tregs in the spleen, showed that mice treated with anti-CD70 might protect allografts using Th3-type Tregs. A few studies had shown that T-cell-derived IL-10 was not essential for induction of allograft acceptance in mice treated with a combined CTLA-4Ig/anti-CD154/LFA-1 blockade, but this treatment limited T-cell expansion in the recipients. These results further indicated that allograft acceptance was maintained by intragraft Tregs [36]. Arefanian et al. found that the combination of anti-LFA-1 and anti-CD154 monoclonal antibodies induced allograft acceptance to neonatal porcine islet xenografts maintained by Tregs [37]. As with allograft acceptance induced by anti-CD154/LFA-1 in naïve mice, our obviously prolonged allograft survival in pre-sensitized mice demonstrated in this study might be also maintained by intragraft Tregs. We will further examine the differences and similarities between the two in future studies.

In summary, we had markedly prolonged allograft survival in presensitized mice by addition of anti-CD70 to the anti-CD154/LFA-1 combination treatment during primary skin transplantation. Our regimen reduced the generation of memory T cells during primary transplantation and inhibited effector T cells function during secondary heart transplantation. This obviously prolonged allograft survival induction program inhibited the proliferative response of recipient splenic T cells to donor antigens, greatly reduced the infiltration of T lymphocytes to the grafts, and might successfully protect the allografts by Tregs. The program can provide a reference for clinical organ transplantation and show that minimizing allogeneic memory T cells

and allogeneic antibodies in primary transplantation patients could allow treatment with conventional therapy during secondary transplantation.

### **Acknowledgments**

We thank Technician Zhigang Liu (School of Life Sciences, Xiamen University) for his help with the flow cytometry analysis.

### **References**

- [1] Amir AL, D'Orsogna LJ, Roelen DL, et al. Allo-HLA reactivity of virus-specific memory T cells is common. *Blood* 2010;115(15):3146-57.
- [2] Bingaman AW, Farber DL. Memory T cells in transplantation: generation, function, and potential role in rejection. *Am J Transplant* 2004;4(6):846-52.
- [3] McFarland RD, Douek DC, Koup RA, Picker LJ. Identification of a human recent thymic emigrant phenotype. *Proc Natl Acad Sci U S A* 2000;97(8):4215-20.
- [4] Douek DC, McFarland RD, Keiser PH, et al. Changes in thymic function with age and during the treatment of HIV infection. *Nature* 1998;396(6712):690-5.
- [5] Pescovitz MD. B cells: a rational target in alloantibody-mediated solid organ transplantation rejection. *Clin Transplant* 2006;20(1):48-54.
- [6] Haudebourg T, Poirier N, Vanhove B. Depleting T-cell subpopulations in organ transplantation. *Transpl Int* 2009;22(5):509-18.
- [7] Mbitikon-Kobo FM, Vocanson M, Michallet MC, et al. Characterization of a CD44/CD122int memory CD8 T cell subset generated under sterile inflammatory conditions. *J Immunol* 2009;182(6):3846-54.
- [8] Minamimura K, Sato K, Yagita H, Tanaka T, Arai S, Maki T. Strategies to induce marked prolongation of secondary skin allograft survival in alloantigen-primed mice. *Am J Transplant* 2008;8(4):761-72.
- [9] Wu GD, He Y, Chai NN, et al. Anti-CD20 antibody suppresses anti-HLA antibody formation in a HLA-A2 transgenic mouse model of sensitization. *Transpl Immunol* 2008;19(3-4):178-86.
- [10] Uchida J, Lee Y, Hasegawa M, et al. Mouse CD20 expression and function. *Int Immunol* 2004;16(1):119-29.
- [11] Corbascio M, Mahanty H, Osterholm C, et al. Anti-lymphocyte function-associated antigen-1 monoclonal antibody inhibits CD40 ligand-independent immune responses and prevents chronic vasculopathy in CD40 ligand-deficient mice. *Transplantation* 2002;74(1):35-41.
- [12] Jones ND, Van Maurik A, Hara M, et al. CD40-CD40 ligand-independent activation of CD8+ T cells can trigger allograft rejection. *J Immunol* 2000;165(2):1111-8.
- [13] Ni HT, Deeths MJ, Li W, Mueller DL, Mescher MF. Signaling pathways activated by leukocyte function-associated Ag-1-dependent costimulation. *J Immunol* 1999;162(9):5183-9.
- [14] Bishop GA, Hostager BS. The CD40-CD154 interaction in B cell-T cell liaisons. *Cytokine Growth Factor Rev* 2003;14(3-4):297-309.



- [15] Bishop GA, Hostager BS. Signaling by CD40 and its mimics in B cell activation. *Immunol Res* 2001;24(2):97-109.
- [16] Tesselaar K, Gravestain LA, van Schijndel GM, Borst J, van Lier RA. Characterization of murine CD70, the ligand of the TNF receptor family member CD27. *J Immunol* 1997;159(10):4959-65.
- [17] Jacquot S. CD27/CD70 interactions regulate T dependent B cell differentiation. *Immunol Res* 2000;21(1):23-30.
- [18] Gravestain LA, Borst J. Tumor necrosis factor receptor family members in the immune system. *Semin Immunol* 1998;10(6):423-34.
- [19] Yamada A, Salama AD, Sho M, et al. CD70 signaling is critical for CD28-independent CD8+ T cell-mediated alloimmune responses in vivo. *J Immunol* 2005;174(3):1357-64.
- [20] Taraban VY, Rowley TF, Al-Shamkhani A. Cutting edge: a critical role for CD70 in CD8 T cell priming by CD40-licensed APCs. *J Immunol* 2004;173(11):6542-6.
- [21] Matsuura A, Abe T, Yasuura K. Simplified mouse cervical heart transplantation using a cuff technique. *Transplantation* 1991;51(4):896-8.
- [22] Winters GL, Marboe CC, Billingham ME. The International Society for Heart and Lung Transplantation grading system for heart transplant biopsy specimens: clarification and commentary. *J Heart Lung Transplant* 1998;17(8):754-60.
- [23] Billingham ME, Cary NR, Hammond ME, et al. A working formulation for the standardization of nomenclature in the diagnosis of heart and lung rejection: Heart Rejection Study Group. The International Society for Heart Transplantation. *J Heart Transplant* 1990;9(6):587-93.
- [24] Sprent J, Surh CD. T cell memory. *Annu Rev Immunol* 2002;20:551-79.
- [25] Nicolls MR, Coulombe M, Beilke J, Gelhaus HC, Gill RG. CD4-dependent generation of dominant transplantation tolerance induced by simultaneous perturbation of CD154 and LFA-1 pathways. *J Immunol* 2002;169(9):4831-9.
- [26] Bucher P, Gang M, Morel P, et al. Transplantation of discordant xenogeneic islets using repeated therapy with anti-CD154. *Transplantation* 2005;79(11):1545-52.
- [27] Elwood ET, Larsen CP, Cho HR, et al. Prolonged acceptance of concordant and discordant xenografts with combined CD40 and CD28 pathway blockade. *Transplantation* 1998;65(11):1422-8.
- [28] Yamaura K, Boenisch O, Watanabe T, et al. Differential requirement of CD27 costimulatory signaling for naive versus alloantigen-primed effector/memory CD8+ T cells. *Am J Transplant* 2010;10(5):1210-20.
- [29] Croft M. Co-stimulatory members of the TNFR family: keys to effective T-cell immunity? *Nat Rev Immunol* 2003;3(8):609-20.
- [30] Nathan MJ, Yin D, Eichwald EJ, Bishop DK. The immunobiology of inductive anti-CD40L therapy in transplantation: allograft acceptance is not dependent upon the deletion of graft-reactive T cells. *Am J Transplant* 2002;2(4):323-32.
- [31] Kirk AD, Blair PJ, Tadaki DK, Xu H, Harlan DM. The role of CD154 in organ transplant rejection and acceptance. *Philos Trans R Soc Lond B Biol Sci* 2001;356(1409):691-702.
- [32] Aversa G, Punnonen J, Carballido JM, Cocks BG, de Vries JE. CD40 ligand-CD40 interaction in Ig isotype switching in mature and immature human B cells. *Semin Immunol* 1994;6(5):295-301.
- [33] Li XC, Rothstein DM, Sayegh MH. Costimulatory pathways in transplantation: challenges and new developments. *Immunol Rev* 2009;229(1):271-93.
- [34] Oberbarnscheidt MH, Ng YH, Chalasani G. The roles of CD8 central and effector memory T-cell subsets in allograft rejection. *Am J Transplant* 2008;8(9):1809-18.
- [35] Bai Y, Liu J, Wang Y, et al. L-selectin-dependent lymphoid occupancy is required to induce alloantigen-specific tolerance. *J Immunol* 2002;168(4):1579-89.

[36] Oderup C, Malm H, Ekberg H, et al. Costimulation blockade-induced cardiac allograft tolerance: inhibition of T cell expansion and accumulation of intragraft CD4(+)Foxp3(+) T cells. *Transplantation* 2006;82(11):1493-500.

[37] Arefanian H, Tredget EB, Rajotte RV, Gill RG, Korbitt GS, Rayat GR. Short-term administrations of a combination of anti-LFA-1 and anti-CD154 monoclonal antibodies induce tolerance to neonatal porcine islet xenografts in mice. *Diabetes* 2010;59(4):958-66.

Table 1. The meanings of abbreviation in this study

abbreviation	treatment	days post-transplantation
S0	Skin transplantation+isotype antibodies	at day 0 and 2
S2	Skin transplantation+anti-CD154/LFA-1	at day 0 and 2
S3	Skin transplantation+ anti-CD154/LFA-1/CD70	at day 0 and 2
H0	Heart transplantation+ isotype antibodies	at day 0, 2, 4 and 6
H2	Heart transplantation+ anti-CD154/LFA-1	at day 0, 2, 4 and 6

Table legend: mAbs were pooled and administered i.p. after the transplantation procedure as follows: 0.25 mg/dose anti-CD154, 0.1 mg/dose anti-LFA-1, and 0.25 mg/dose anti-CD70. Heart transplantations were performed one month after primary skin transplantation.

Table 2. Primer sequences used for qRT-PCR

Primer name	Sequences
$\beta$ -actin	forward 5'-CATCCGTAAGACCTCTATGCCAAC-3', reverse 5'-ATGGAGCCACCGATCCACA-3'
IFN- $\gamma$	forward 5'-CGGCACAGTCATTGAAAGCCTA-3', reverse 5'-GTTGCTGATGGCCTGATTGTC-3'

---

IL-2	forward 5'-GGAGCAGCTGTTGATGGACCTAC-3', reverse 5'-AATCCAGAACATGCCGCAGAG-3'
IL-10	forward 5'-GACCAGCTGGACAACATACTGCTAA-3', reverse 5'-GATAAGGCTTGGCAACCCAAGTAA-3'
Foxp3	forward 5'- CAGCTCTGCTGGCGAAAGTG -3', reverse 5'-TCG.TCTGAAGGCAGAGTCAGGA -3'
TGF- $\beta$	forward 5'- TGACGTCACTGGAGTTGTACGG -3', reverse 5'- GGTTTCATGTCATGGATGGTGC -3'

---


**Figure 1. Changes in the proportion of memory T cells in each treatment group following skin transplantation.** Proportions of CD4<sup>+</sup> memory T cells/CD4<sup>+</sup> T cells (top) and CD8<sup>+</sup> memory T cells/CD8<sup>+</sup> T cells (bottom) were analyzed using flow cytometry. Naïve B6 mice were used as negative controls (n=3 mice/group). The abbreviation of S0 represented mice received isotype antibodies only at the day 0 and 2 post skin transplantation. S2 represented mice received a 2-antibody treatment regimen consisting of anti-CD154/LFA-1 at the day 0 and 2 post skin transplantation, S3 represented mice received a 3-antibody treatment regimen consisting of anti-CD154/LFA-1/CD70 at the day 0 and 2 post skin transplantation. Data are represented of three separate experiments with similar results.




**Figure 2. Graft survival time, recipient splenic lymphocyte function, and serum alloantibody levels in mice that received no treatment during secondary heart transplantation.** Spleens and sera were recovered from recipient mice at day 3.5 post-transplantation for analysis. (A) Graft survival were showed by Kaplan-Meier method and compared by Log-rank test (n=6 mice/group). (B) MLR assays were used to test the proliferative responses of recipient splenic T cells to donor BALB/c antigens (top; n=3 mice/group; \*  $P < 0.05$ ). Three separate wells were dedicated to each responder-stimulator combination and each experiment was repeated three times. Concentrations of IL-2 and IFN- $\gamma$  in the supernatant from MLR were detected by EISA (bottom; n=3 mice/group; \*\*\*  $P < 0.001$ ). Each reaction was carried out in triplicate. (C) The proportions of CD4<sup>+</sup> memory T cells/CD4<sup>+</sup> T cells, CD8<sup>+</sup> memory T cells/CD8<sup>+</sup> T cells, and Tregs (CD4<sup>+</sup>Foxp3<sup>+</sup>) were analyzed by flow cytometry taking into account the proportion of splenic lymphocytes (n=3 mice/group). (D) Alloantibodies in recipient sera were detected by flow cytometry (n =3 mice/group).






Data are represented of three separate experiments with similar results.





**Figure 3. Graft survival time, recipient spleen lymphocyte function, and serum alloantibody levels in mice treated with anti-CD154/LFA-1 during secondary heart transplantation.** Splens and sera were recovered from recipient mice at day 7 post-transplantation for analysis. Mice in the S3H2 group were sacrificed at either day 7 or 100 post-transplantation. (A) Graft survival were showed by Kaplan-Meier method and compared by Log-rank test. (B) MLR assays were used to test the proliferative responses of recipient splenic T cells to donor BALB/c antigens (top; n=3 mice/group; \* $P < 0.05$ ). Three separate wells were dedicated to each responder-stimulator combination and each experiment was repeated three times. Concentrations of IL-2, IFN- $\gamma$ , IL-10, and TGF- $\beta$  in the supernatant from MLR were detected by ELISA (bottom; n=3 mice/group; \* $P < 0.05$ ; \*\* $P < 0.01$ ). Each reaction was carried out in triplicate. (C) The proportions of CD4<sup>+</sup> memory T cells/CD4<sup>+</sup>T cells, CD8<sup>+</sup> memory T cells/CD8<sup>+</sup>T cells, and Tregs (CD4<sup>+</sup>Foxp3<sup>+</sup>) were analyzed by flow cytometry taking into account the proportion of splenic lymphocytes (n=3 mice/group). (D) Alloantibodies in recipient sera were detected by flow cytometry (n=3 mice/group). Data are represented of three separate experiments with similar results.

**Figure 4. Pathological analysis of grafts, detection of graft-infiltrating lymphocytes, and determination of rejection/tolerance levels associated with cytokines in grafts from mice treated with anti-CD154/LFA-1 during secondary heart transplantation.** Recipient mice were sacrificed at day 7 post-transplantation. Mice from the S3H2 group were sacrificed at either day of 7 or 100 post-transplantation. (A) H&E staining of grafts and the ISHLT score. Each point represents the score from one slice. The horizontal line represents the mean for each group (n=3 mice/group; \*\* $P < 0.01$ ; \*\*\*  $P < 0.001$ ). (B) Flow cytometric analysis of graft-infiltrating lymphocytes. (C) Gene expression levels of IL-2, IFN- $\gamma$ , IL-10, Foxp3, and TGF- $\beta$  detected by qRT-PCR. Each reaction was carried out in triplicate. Data are representative of three separate experiments (n=3 mice/group; \* $P < 0.05$ ; \*\*\* $P < 0.001$ ).














**Supplement figure1. The character of CD8<sup>+</sup> T in the grafts after secondary transplantation.** The lymphocytes from the recipients' grafts were analyzed by flow cytometry. CD8<sup>+</sup> T cells were effector/memory T cells. Data are representative of three independent experiments.