



LUND UNIVERSITY

Acupuncture in the treatment of infantile colic

Landgren, Kajsa; Raith, Wolfgang; Schmoelzer, Georg M.; Skjeie, Holgeir; Skonnord, Trygve

Published in:
The Italian Journal of Pediatrics

DOI:
[10.1186/s13052-014-0105-3](https://doi.org/10.1186/s13052-014-0105-3)

2015

[Link to publication](#)

Citation for published version (APA):
Landgren, K., Raith, W., Schmoelzer, G. M., Skjeie, H., & Skonnord, T. (2015). Acupuncture in the treatment of infantile colic. *The Italian Journal of Pediatrics*, 41, Article 1. <https://doi.org/10.1186/s13052-014-0105-3>

Total number of authors:
5

General rights

Unless other specific re-use rights are stated the following general rights apply:
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00



LETTER TO THE EDITOR

Open Access

Acupuncture in the treatment of infantile colic

Kajsa Landgren^{1*}, Wolfgang Raith^{2,3}, Georg M Schmölzer^{3,4,5}, Holgeir Skjeie⁶ and Trygve Skonnord⁶

Abstract

Regarding the recently published review "Looking for new treatments of Infantile Colic" by Savino et al. we want to add that positive effects of acupuncture have been demonstrated to release pain and agitation and that acupuncture seems to be a safe treatment when performed by trained acupuncturists. Inconclusive results in the few published articles on the subject can be due to different acupuncture points, different insertion time, different needling methods, differences in the outcome variables, in how the crying was measured and insufficient sample sizes. Further research is needed on understanding the utility, safety, and effectiveness of acupuncture in infants with colic.

Keywords: Infantile colic, Acupuncture, Safety

To the Editor

We read with interest "Looking for new treatments of Infantile Colic" by Savino et al [1]. This is an excellent review describing the current therapies in infantile colic. Treating infantile colic using acupuncture is feasible and without serious side effects [1], however the evidence seems to be inconclusive [1-3]. Savino et al. clearly state that there is a need for a detailed evaluation of the indications of acupuncture in newborn infants with colic, particularly during early infancy when responses are difficult to evaluate.

While we agree with the authors' conclusion, we would like to add for the audience of the *Italian Journal of Pediatrics* and especially for centers performing acupuncture in newborn infants, that positive effects of acupuncture have been demonstrated to release pain and agitation [4]. In addition, both a recent and a 3 years older systematic review of acupuncture for children and newborns found that acupuncture is a safe treatment when performed by trained and licensed acupuncturists [4,5].

The current evidence includes three previous randomized trials [2,3,6,7] and a case series with 913 newborn infants included [8], investigating acupuncture treatment in infantile colic. Landgren et al [2,6] used

two seconds of unilateral needling at acupuncture point LI4 (Large Intestine 4), and Reinthal et al [7,8] used bilateral needling for 10–20 seconds at LI4. Both reported a reduction of crying frequency and intensity in the acupuncture group compared to the control group [2,7]. A third study by Skjeie et al [3] used a different acupuncture point, bilateral needling of ST36 (Stomach 36) for 30 seconds, which is in accordance with the Norwegian Society of Medical Acupuncture, and reported no statistically significant difference in reduction of crying time between the acupuncture and the control group. These three trials [2,3,6,7] used different acupuncture points, different insertion time, and different needling methods, which potentially contributed to the inconclusive results. Other contributing factors to the variable results can be differences in the outcome variables, in how the crying was measured and insufficient sample size. For example Skjeie et al. recruited 79 patients of the 120 who were requested in the power analysis. Furthermore we want to add, that everybody who is performing paediatric acupuncture should have had adequate training and experience in acupuncture of newborn infants and children. We need future studies, recruiting larger population with feasible placebo control, to better understand the utility, safety, and effectiveness of acupuncture in newborn infants with colic.

* Correspondence: kajsa.landgren@med.lu.se

¹Department of Health Sciences, Faculty of Medicine, Lund University, Box 157, 22100 Lund, Sweden

Full list of author information is available at the end of the article

Abbreviations

LI4: the acupuncture point "Large Intestine 4"; ST36: the acupuncture point "Stomach 36".

Competing interests

The authors declare that they have no competing interest.

Authors' contributions

All authors contributed in preparing this letter and approved it.

Author details

¹Department of Health Sciences, Faculty of Medicine, Lund University, Box 157, 22100 Lund, Sweden. ²Division of Neonatology, Department of Paediatrics, Medical University of Graz, Graz, Austria. ³Research Group for Paediatric Traditional Chinese Medicine, TCM Research Center Graz, Medical University of Graz, Graz, Austria. ⁴Department of Paediatrics, University of Alberta, Edmonton, Canada. ⁵Neonatal Research Unit, Royal Alexandra Hospital, Edmonton, Alberta Health Services, Edmonton, Canada. ⁶Department of General Practice, Institute of Health and Society, University of Oslo, Oslo, Norway.

Received: 15 October 2014 Accepted: 16 December 2014

Published online: 15 January 2015

References

1. Savino F, Ceratto S, De Marco A. Looking for new treatments of Infantile Colic. *Ital J Pediatr.* 2014;5:40–53.
2. Landgren K, Kvornings N, Hallström I. Acupuncture reduces crying in infants with infantile colic: a randomised, controlled, blind clinical study. *Acupunct Med.* 2010;28(4):174–9.
3. Skjeie H, Skonnord T, Fetveit A, Brekke M. Acupuncture for infantile colic: a blinding-validated, randomized controlled multicentre trial in general practice. *Scand J Prim Health Care.* 2013;31(4):190–6.
4. Raith W, Urlesberger B, Schmölzer GM. Efficacy and safety of acupuncture in preterm and term infants. *Evid Based Complement Alternat Med.* 2013;2013:739414.
5. Adams D, Cheng F, Jou H, Aung S, Yasui Y, Vohra S. The safety of pediatric acupuncture: a systematic review. *Pediatrics.* 2011;128(6):e1575–587.
6. Landgren K, Kvornings N, Hallström I. Feeding, stooling and sleeping patterns in infants with colic—a randomized controlled trial of minimal acupuncture. *BMC Complement Altern Med.* 2011;11:93. doi:10.1186/1472-6882-11-93.
7. Reinthal M, Andersson S, Gustafsson M, Plos K, Lund I, Lundeberg T. Effects of minimal acupuncture in children with infantile colic: A prospective, quasi-randomised single blind controlled trial. *Acupunct Med.* 2008;26:171–82.
8. Reinthal M, Lund I, Ullman D, Lundeberg T. Gastrointestinal symptoms of infantile colic and their change after light needling of acupuncture: a case series study of 913 infants. *Chin Med.* 2011;6:28. doi:10.1186/1749-8546-6-28.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

