



# LUND UNIVERSITY

## Airborne SARS-CoV-2 during childbirth

Thuresson, Sara; Alsved, Malin; Medstrand, Patrik; Löndahl, Jakob; Fraenkel, Carl-Johan

2023

[Link to publication](#)

*Citation for published version (APA):*

Thuresson, S., Alsved, M., Medstrand, P., Löndahl, J., & Fraenkel, C.-J. (2023). *Airborne SARS-CoV-2 during childbirth.*

*Total number of authors:*

5

*Creative Commons License:*

CC BY-NC-ND

### General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117  
221 00 Lund  
+46 46-222 00 00



LUND  
UNIVERSITY

# Airborne SARS-CoV-2 during childbirth

S. THURESSON<sup>1</sup>, M. ALSVED<sup>1</sup>, P. MEDSTRAND<sup>2</sup>, J. LÖNDAHL<sup>1</sup>, CJ FRAENKEL<sup>3,4</sup>

<sup>1</sup> Division of Ergonomics and Aerosol Technology, Department of Design Sciences, Lund University

<sup>2</sup> Department of Translational Medicine, Lund University

<sup>3</sup> Department of Infection Control, Region Skåne, Lund, Sweden

<sup>4</sup> Division of Infection Medicine, Department of Clinical Sciences, Lund University

## Background

Airborne SARS-CoV-2 is considered to play a major role in covid-19 transmission, and has been found in several hospital environments, including corridors, patient rooms and cohort rooms. A number of factors have been found to increase the levels of airborne virus, such as low ventilation, patient viral load and in some cases, certain medical procedures. However, other specific medical situations still deserve further investigation.

Childbirth is one such interesting situation, as respiratory emissions are increased due to heavy breathing. In the case of an infected mother, these emissions could contain virus. Moreover, the patient's use of face masks is limited in these situations, due to the exertion of pushing during delivery. To date, published studies on SARS-CoV-2 during childbirth only include a small number of air samples collected with both passive and active air samplers. Most of these fail to detect airborne SARS-CoV-2 RNA.


The aim of the current study was to further explore the presence of airborne SARS-CoV-2 RNA during labor and delivery.



The Coriolis  $\mu$  sampler, a liquid cyclone, was used to collect air samples. Air is sampled with  $200 \text{ L min}^{-1}$  and the collection liquid is analysed for virus.

## Methods

Air was sampled before, during and after childbirth in rooms where the women giving birth had tested positive for covid-19. Sampling was performed using a liquid cyclone (Coriolis  $\mu$ , Bertin Instruments, France), operating at  $200 \text{ L min}^{-1}$  for 10 min, with 15 mL of phosphate-buffered saline solution as collection liquid. The collection liquid was concentrated using Amicon Ultra-15 centrifugal filter units (50 kDa cutoff, Merck Millipore). SARS-CoV-2 RNA was detected from concentrated samples by real time reverse transcription polymerase chain reaction (RT-qPCR).



## Results

- ❖ 44 air samples were collected on **six** occasions.
- ❖ **Six** samples, collected during **two** different childbirths, were positive for SARS-CoV-2 in RT-qPCR.
- ❖ The concentrations of RNA were generally **low**, with cycle threshold (Ct) values between 37.7 and 40.
- ❖ Positive air samples were found both **before, during and after delivery.**

## Conclusion

There is a need to investigate the presence of airborne SARS-CoV-2 in other hospital areas than traditional infectious disease wards. The results from the present study can contribute to a better understanding of the risk of covid-19 transmission by aerosols at delivery wards. This can improve guidelines for protective equipment for healthcare personnel working with such patients.

## Contact

Sara Thuresson, PhD student  
sara.thuresson@design.lth.se

FORMAS

afa  
FÖRSÄKRING