



# LUND UNIVERSITY

## Alterations of particle deposition in the respiratory tract at increasing activity

Linell, Julia; Isaxon, Christina; Stroh, Emilie; Olsson, Bo; Löndahl, Jakob; Wollmer, Per; Rissler, Jenny

2023

[Link to publication](#)

### *Citation for published version (APA):*

Linell, J., Isaxon, C., Stroh, E., Olsson, B., Löndahl, J., Wollmer, P., & Rissler, J. (2023). *Alterations of particle deposition in the respiratory tract at increasing activity*. Poster session presented at Kick-off för LTHs Profilmråde AEROSOLS, Lund, Sweden.

*Total number of authors:*

7

### **General rights**

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117  
221 00 Lund  
+46 46-222 00 00



# Alterations of particle deposition in the respiratory tract at increasing activity

J. Linell<sup>1,2</sup>, C. Isaxon<sup>1,2</sup>, E. Stroh<sup>3</sup>, B. Olsson<sup>4</sup>, J. Löndahl<sup>1,2</sup>, P. Wollmer<sup>5,6</sup> and J. Rissler<sup>1,2,7</sup>

<sup>1</sup>Ergonomics and Aerosol Technology, Lund University, Sweden <sup>2</sup>NanoLund, Lund University, Sweden <sup>3</sup>Occupational and Environmental Medicine, Lund University, Sweden <sup>4</sup>Emmace Consulting AB, Lund, Sweden <sup>5</sup>Centre for Medical Imaging and Physiology, Skåne University Hospital, Malmö, Sweden <sup>6</sup>Translational Medicine, Lund University, Sweden <sup>7</sup>Bioeconomy and health, RISE Research Institutes of Sweden, Lund, Sweden.

## Conclusions

Deposition of  $< 0.1 \mu\text{m}$  particles increase with activity in the alveolar region while the deposition in the entire lung does not change considerably.

This increased deposition in the alveolar region is driven by a larger breath volume at increased activity.

## Motivation

The **deposited dose** of particles in the respiratory tract is an important **factor for understanding health effects of aerosol particles**. It is a combination of the exposure, volume of inhaled air and particle deposition.

At **increasing activity** the minute ventilation (volume of breath x number of breaths) increase. Little is known about if and how this **affects the regional particle deposition**.




Fig.1. Size-resolved particle deposition in the entire respiratory tract (total) and alveolar region of an adult (oral breathing). Similar patterns were seen for children.




Fig.2. Regional deposition for three particle size ranges for an adult (oral breathing). Similar patterns were seen for children.




Fig.3. Deposition of  $0.005\text{--}0.1 \mu\text{m}$  particles keeping the breath volume constant while increasing the breathing frequency (left) and keeping the breathing frequency constant while increasing the breath volume (right).

## Method

The alteration of breath volume and minute ventilation during increasing activity was parametrized using data from previous studies. The results were used to model the effect of activity on lung deposition for children and adults.

The extra-thoracic deposition model from NCRP (1997) and the thoracic deposition model and lung anatomy model from Yeh and Schum (1980) were used in the software Mimetikos Preludium™ (Olsson and Bäckman, 2018).

## Acknowledgements

This research was supported by Formas [prn 2018-00693].

## Results

Particle deposition was altered with activity for different particle sizes in the entire respiratory tract at increasing activity (Fig.1). The largest effects were seen in the alveolar region where the peak deposition was shifted from  $\sim 30 \text{ nm}$  to  $15 \text{ nm}$  (Fig.1).

Deposition in the alveolar region increased for  $< 0.1 \mu\text{m}$  particles while it decreased for  $> 0.1 \mu\text{m}$  particles at increasing activity (Fig.2). Similar patterns were seen for both adults and children.

The deposition of  $< 0.1 \mu\text{m}$  particles in the alveolar region seemed unaffected by an increased breathing frequency (Fig.3).

The increased deposition for  $< 0.1 \mu\text{m}$  particles in the alveolar region was driven by an increased volume of the inhaled air (Fig.3) and a decreased diffusional deposition in the conducting airways (Fig.2).

## Outlook

Increasing activity lead to a shift in the size distribution of particles that deposit in different regions of the respiratory tract. This can be important to consider when calculating deposited doses for health risk assessments.

## Contact

Julia Linell, Ergonomics and Aerosol Technology, LTH  
julia.linell@design.lth.se