



# LUND UNIVERSITY

## Neuropeptide Y autoantibodies in patients with long-term type 1 and type 2 diabetes and neuropathy.

Skärstrand, Hanna; Dahlin, L B; Lernmark, Åke; Vaziri Sani, Fariba

*Published in:*  
Journal of Diabetes and its Complications

*DOI:*  
[10.1016/j.jdiacomp.2013.06.007](https://doi.org/10.1016/j.jdiacomp.2013.06.007)

2013

[Link to publication](#)

*Citation for published version (APA):*  
Skärstrand, H., Dahlin, L. B., Lernmark, Å., & Vaziri Sani, F. (2013). Neuropeptide Y autoantibodies in patients with long-term type 1 and type 2 diabetes and neuropathy. *Journal of Diabetes and its Complications*, 27(6), 609-617. <https://doi.org/10.1016/j.jdiacomp.2013.06.007>

*Total number of authors:*  
4

### General rights

Unless other specific re-use rights are stated the following general rights apply:  
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117  
221 00 Lund  
+46 46-222 00 00



## **Neuropeptide Y autoantibodies in patients with long-term type 1 and type 2 diabetes and neuropathy.**

H. Skärstrand, MSc., L B. Dahlin, M.D, Ph.D., Å. Lernmark, Ph.D., F. Vaziri-Sani, Ph.D.

Department of Clinical Sciences, Malmö, Lund University, Skåne University Hospital, Sweden.

Running title: NPY autoantibodies in diabetes.

Corresponding author:

Hanna Skärstrand

Lund University/CRC

Department of Clinical Sciences, Malmö

Jan Waldenströms gata 35

Skåne University Hospital SUS

SE-20502 Malmö, Sweden

Phone +46 40 391123

Fax +46 40 391919

Email [hanna.skarstrand@med.lu.se](mailto:hanna.skarstrand@med.lu.se)

### **Grant support**

The study was supported in part by the EU 7<sup>th</sup> Framework Programme: DIAPREPP (Diabetes type 1 Prediction, Early Pathogenesis and Prevention, grant agreement 202013), the Swedish Child Diabetes Foundation (Barndiabetesfonden), the Swedish Research Council (Grant 14064 and 05188), the Swedish Diabetes Association Research Fund (Diabetesfonden), the Skåne County Council Foundation for Research and Development, SUS Fund as well as the Swedish Association of Local Authorities and Regions (SKL) and Lund University, Medical Faculty.

### **Presented at meeting**

Presented in part as a poster at the 12<sup>th</sup> Immunology of Diabetes Society (IDS) conference June 15-19, 2012 in Victoria, BC, Canada. Printed in the meeting abstract book on page 140.

## **Abstract**

**Aims:** The neurotransmitter Neuropeptide Y (NPY) was previously reported as a minor autoantigen in newly diagnosed type 1 diabetes (T1D) patients. The single nucleotide polymorphism at rs16139 (T1128C, L7P) in the NPY gene was associated with an increased risk for the development of type 2 diabetes (T2D). We aimed to develop a radiobinding assay for NPY-L (Leucine) and NPY-P (Proline) autoantibodies (A) to study the levels and the association with other islet autoantibodies and neuropathy. **Methods:** Autoantibodies against NPY-L, NPY-P, ZnT8, GAD65 and IA-2 were studied in T1D (n=48) and T2D (n=26) patients with duration up to 42 and 31 years. A subgroup of T1D (n=32) patients re-examined, 5-8 years after first visit, were tested for peripheral (Z-score) and autonomic neuropathy (E/I ratio). **Results:** NPY-LA and NPY-PA were detected in 23% and 19% in T1D ( $p<0.001$ ), and 12% and 23% in T2D patients ( $p<0.001$ ) compared to 2.5% controls (n=398). The levels of NPYA declined during follow-up in the T1D patients ( $p<0.001$ ). The neuropathy was not related to the NPYA or the other islet autoantibodies. **Conclusions:** Regardless of the absence of an association between NPYA and neuropathy, NPY may contribute to the pathogenesis of T1D and T2D as a minor autoantigen.

## **Keywords**

1. Minor autoantigen
2. NPY autoantibodies
3. Autoimmunity
4. Neuropathy
5. Long-term diabetes

## 1. Introduction

Type 1 diabetes (T1D) is strongly associated with autoantibodies (A) against zinc transporter 8 (ZnT8A), glutamic acid decarboxylase 65 (GADA), insulinoma-associated antigen-2 (IA-2A) and insulin (IAA) (reviewed in [1]). Together or alone, they are important not only for classification of diabetes at the time of clinical onset, but also for the prediction of T1D in subjects at risk [2] [3] [4]. More than 90% of patients with T1D have multiple autoantibodies, and less than 5% of T1D patients were found negative for any of these four islet autoantibodies at clinical onset [5]. However, it has been suggested that islet autoantibody negative individuals may be positive for yet other beta-cell autoantigen-specific autoantibodies [5]. Neuropeptide Y (NPY) was previously reported as a minor autoantigen in newly diagnosed T1D patients [6].

NPY is a neurotransmitter widely expressed within the central and peripheral nervous systems affecting vasoconstriction, insulin secretion and food intake [7] [8]. NPY has also been demonstrated in islet cells and pancreatic nerve fibers [9]. Furthermore, a single nucleotide polymorphism (SNP) at rs16139 (T1128C) identified an amino acid substitution from Leucine (L) to Proline (P) (L7P) in the signal peptide [10]. The minor allele (P) correlated with elevated NPY levels in plasma [11] perhaps due to an increased biosynthesis [12] [13]. It was suggested that the L7P affects NPY secretion as a consequence of a modified secondary and tertiary structure [12]. It is therefore of interest that the minor allele was associated with atherosclerosis [14], impaired glucose tolerance [15] as well as with a three- to four-fold increased risk for Type 2 diabetes (T2D) [15] [16]. More importantly, the minor allele also increased the risk for retinopathy [14] and nephropathy [17] in patients with T2D.

Autonomic neuropathy in patients with T1D was associated with impaired NPY responses to insulin-induced hypoglycaemia [18]. As T1D neuropathy was also associated with GADA [19] [20], we tested the hypothesis that NPY autoantibodies (NPYA) may be present in T1D patients with neuropathy and that NPYA would be related to the rs16139 polymorphism.

The aims of this study were to 1) develop a radiobinding assay (RBA) for NPY-L (Leucine) and NPY-P (Proline) autoantibodies (NPYA) in controls and in patients with long-term T1D and T2D, 2) relate both NPY-LA and NPY-PA with autoantibodies against ZnT8R (Arginine), ZnT8W (Tryptophan), ZnT8Q (Glutamine), GAD65 and IA-2; and 3) study these autoantibody levels during follow-up of T1D patients tested for neuropathy.

## **2. Materials and Methods**

### **2.1. Patient and control samples**

A total of 100 patients, 1.5-18.8 years of age, with T1D diagnosed in 1996-2001 were studied (Table 1). Serum samples from 48 long-term T1D and 26 long-term T2D patients were collected in 1998 at the Clinic of Diabetes at Skåne University Hospital in Malmö, Sweden (Table 2). A subgroup of 32 patients with long-term T1D, first seen in 1998, was followed-up in 2005 (Table 2). Sera for control samples were obtained from 398 healthy blood donors at 19-81 years of age (Table 2).

### **2.2. Nerve function tests during follow-up**

The long-term T1D patients underwent peripheral and cardiac autonomic nerve function tests during the follow-up in 2005 (n=31; Table 2). The peripheral nerve function (peripheral neuropathy) was evaluated by measurement of conduction velocity (CV) and amplitude (AMP) of the sural and peroneal nerves in the lower extremities and calculated as a mean expressed in age-corrected values [Z-score in standard deviations (SD)]. A composite Z-score leg, reflecting the nerve function in the leg, was calculated: (Z-score CV peroneal nerve + Z-score CV sural nerve + Z-score AMP sural nerve)/3. The summation of nerve conduction attributes has been used elsewhere [21]. The cardiac autonomic nerve test (autonomic neuropathy) of the parasympathetic vagal nerve function (E/I ratio) was also tested [22]. The E/I ratio was calculated as the mean of the longest R-R interval during expiration (E) divided by the mean of the shortest R-R interval during inspiration (I) and expressed in age-corrected values [Z-scores in SD]. E/I Z-score values less than -1.64 SD (95% confidence interval, one-sided test) below the age-related reference values were considered as abnormal [22].

### **2.3. Subcloning of pJ201prepro-NPY**

Prepro-Neuropeptide Y (NPY) (aa 1-97) cDNA was inserted in a de novo synthesised pJ201 vector (2759 bp) (DNA 2.0, Menlo Park, CA, USA) with both restriction cloning sites of Xho1 and Not1 at the 5' end and 3' end of the insert. The pJ201NPY cDNA (2 µg) was cut with two restriction enzymes, Xho1 and Not1 (FastDigest™, Fermentas, Helsingborg, Sweden). The NPY insert was subcloned into the pTnT™ vector (2871 bp) (Promega, Madison, WI, USA) and transformed using DH5α E.coli competent cells (Invitrogen AB, Stockholm, Sweden). The

pThNPY plasmid DNA was extracted using the QiaPrep Spin Miniprep Kit (QIAGEN AB, Solna, Sweden) according to manufacturer's instructions followed by sequencing using (3700/3730 BigDye Terminator 3.1 Cycle Sequencing Kit (Applied Biosystems, Foster City, CA, USA) at the Region Skånes Competence Centre (RSKC) in Malmö, Sweden.

#### **2.4. Generation of pTnTprepro-NPY SNP T1128C-Proline variant**

Both forward and reverse primers were designed according to manufacturer's instructions (DNA Technology, Risskov, Denmark). A Phusion™ site directed mutagenesis Kit (Thermo Fisher Scientific, Leicestershire, UK) was used to obtain Proline (P) in substitution for Leucine (L) (L7P) of the prepro-NPY insert. The pThNPY-P variant was transformed in DH5α E.coli competent cells prior plasmid DNA extraction and sequencing as previously described.

#### **2.5. Radiolabeling of prepro-NPY protein products**

Both pThNPY-L and pThNPY-P (1μg each) were subjected to *in vitro* coupled transcription translation to generate <sup>35</sup>S-methionine labeled human recombinant proteins (<sup>35</sup>S-met, 500 uCi, PerkinElmer, Waltham, MA, USA) using the TNT® SP6 Coupled Reticulocyte Lysate System (Promega, Madison, WI, USA) in a 100 μL reaction mix for 90 min at 30 °C degrees during shaking (300 rpm). The translation mix was separated by gel filtration on Illustra™ NAP-5 Columns (GE Healthcare Bio-Sciences AB, Uppsala, Sweden) by adding antigen buffer (150 mmol/L NaCl, 20 mmol/L Tris, pH 7.4, 0.15%, Tween20, 0.1% BSA) in various volumes from 100-1500 μL in eleven steps. Prior gel filtration, columns were washed and equilibrated with antigen buffer. The amount of <sup>35</sup>S-met incorporated in both <sup>35</sup>S-met NPY-L and <sup>35</sup>S-met NPY-P was determined (1450 MicroBeta Counter, Perkin Elmer). The <sup>35</sup>S-met NPY-L and <sup>35</sup>S-met NPY-P protein products were stored up to one month at -80 °C degrees.

#### **2.6. SDS-polyacrylamide gel electrophoresis**

<sup>35</sup>S-met NPY protein products were verified by 16% sodium dodecyl sulfate (SDS) polyacrylamide gel electrophoresis (PAGE) (ClearPage™, C.B.S Scientific, Delmar, CA, USA) described elsewhere [23].

#### **2.7. NPY autoantibody analysis**

Serum samples were analyzed for autoantibodies against NPY-L and NPY-P using a standard RBA. In duplicates, 5  $\mu$ L of serum sample was mixed with 60  $\mu$ L of either  $^{35}$ S-met NPY-L or  $^{35}$ S-met NPY-P protein products to a final concentration of  $425 \pm 25$  cpm/ $\mu$ L by dilution with antigen buffer in a 96-well plate. The reaction mix (antigen-antibody solution) was incubated over night at 4 °C with constant shaking. Cold washing buffer (150 mM NaCl, 20 mM Tris, pH 7.4, 0.15% Tween 20) of 200  $\mu$ L per plate were prepared on the first day. 200  $\mu$ L of antigen buffer was added to each well of a 96-well filter plate and incubated at room temperature over night to block unspecific binding. The following day, the filter plates were emptied and 50  $\mu$ L of the reaction mix was added to 20% v/v Protein A Sepharose (PAS, Invitrogen) in the filter plate and incubated for 1 h at 4 °C with constant shaking (Perkin Elmer, Waltham, MA, USA). The filter plate was washed eight times with washing buffer using a ELx50 Microplate Strip Washer (Biotek, Winooski, VT, USA). Antibody-bound radioactivity was analyzed in a  $\beta$ -counter (1450 MicroBeta Counter, Perkin Elmer). PAS-bound radioactivity was converted into in-house units (U) using a polyclonal rabbit IgG antibody against full length human NPY (Abcam, Cambridge Science Park, Cambridge, UK) in seven doubling dilution steps as a standard for both NPY antigens. The binding of both  $^{35}$ S-met NPY-L and  $^{35}$ S-met NPY-P to the standard IgG was displaced in a concentration-dependent manner by both cold NPY-L and NPY-P (Supplement Figure 1) generated by *in vitro* transcription translation as described [24]. Patient and control samples were analysed for autoantibodies in parallel on the same plate to reduce assay variances.

## **2.8. ZnT8, GAD65, IA-2 autoantibody analyses**

The RBA for each of individual ZnT8 autoantibody (ZnT8RA, ZnT8WA, ZnT8QA) was carried out as described [25]. Autoantibodies against GAD65 and IA-2 were similarly analyzed as described previously [26].

## **2.9. Statistical analysis**

Quantile-quantile (Q-Q) analysis was used for the quantiles of the autoantibody distribution against the quantiles of a standard distribution. Pearson Chi-square (and two-sided Fisher's exact test and Yate's correction for continuity value when applied) was used to assess differences of autoantibody frequency in un-paired groups, and McNemar test in paired groups. Mann-Whitney U-test was used for calculation of differences in autoantibody levels and Wilcoxon signed ranks



test for change in autoantibody levels over time. Spearman two-tailed correlation test was used for correlation analysis. Agreement of islet autoantibodies was calculated with Kappa statistics. A value of one indicated perfect agreement, whereas a kappa of zero was what would be expected by chance and negative values a systematic disagreement between the observers. A p-value of  $<0.05$  was considered significant. IBM SPSS Statistics 18.0 was used for the statistical calculations.

Interpretation of kappa agreement:

$<0$  Less than chance agreement

0.01-0.20 Slight agreement

0.21-0.40 Fair agreement

0.41-0.60 Moderate agreement

0.61-0.80 Substantial agreement

0.81-0.99 Almost perfect agreement

### **3. Results**

#### **3.1. NPY and islet autoantibodies in controls, type 1 and type 2 diabetes patients**

In the newly diagnosed T1D patients (n=100) NPY-LA or NPY-PA showed a frequency of 29% for both autoantigens (Table 1). NPY-LA and NPY-PA double positives were found in 20% of the newly diagnosed patients.

The long-term T1D (n=48) patients were diagnosed at 15-26 years of age compared to the T2D (n=26) patients diagnosed at 31-50 years of age (Table 2). Duration of the disease was comparable between the two groups of patients. However, while all patients with long-term T1D had low level of C-peptide, the long-term T2D patients showed a marked variability (Table 2). A total of 32 T1D patients were re-examined after 21-50 years of disease (Table 2). The  $^{35}\text{S}$ -met labeled NPY-L and NPY-P generated by *in vitro* transcription translation had the expected molecular mass of Mr 10.8 (Figure 1). NPY-LA and NPY-PA levels in serum from healthy controls were normally distributed. The cut-off levels at the 97.5<sup>th</sup> percentile were 44 U/mL for NPY-LA and 32 U/mL for NPY-PA (Figure 2, panels A-B). The Q-Q plots of the long-term 48 T1D patients for NPY-LA and NPY-PA, respectively, deviated from a normal distribution (Figure 2, panel C). The frequency at baseline in the long-term T1D patients was 23% for NPY-LA ( $p<0.001$ ) and 19% for NPY-PA ( $p<0.001$ ) compared to controls (Table 3). There were 6/48 (12.5%) NPY-LA and NPY-PA double positive T1D patients (Figure 2, panel D). The corresponding frequency in the patients with T2D were 12% for NPY-LA ( $p=0.039$ ) and 23% for

NPY-PA ( $p < 0.001$ ) as compared to the controls (Table 3). NPY-LA and NPY-PA double positive T2D were 3/26 (11.5%).

The median levels of the two NPYA were different from the controls in both the T1D (NPY-LA:  $p < 0.001$  and NPY-PA:  $p < 0.001$ , respectively) and the T2D patients (NPY-LA:  $p = 0.05$  and NPY-PA:  $p < 0.001$ ) (Figure 3, panel a). The titers of the two NPYA variants correlated in both the T1D ( $R^2 = 0.547$ ,  $p < 0.001$ ) (Figure 2, panel D) and in T2D patients ( $R^2 = 0.802$ ,  $p < 0.001$ ) (data not shown).

Compared to the controls, the frequencies of the ZnT8A variants were 40% for ZnT8RA ( $p < 0.001$ ), 40% for ZnT8WA ( $p < 0.001$ ) and 35% for ZnT8QA ( $p < 0.001$ ) in the T1D and 15% ( $p = 0.004$ ), 12% ( $p = 0.031$ ) and 12% ( $p = 0.031$ ), respectively, in T2D patients (Table 3). The median levels of all ZnT8A variants (ZnT8RA, ZnT8WA, ZnT8QA) were higher in long-term patients with both T1D (ZnT8RA,  $p < 0.001$ ; ZnT8WA,  $p < 0.001$ ; ZnT8QA,  $p < 0.001$ ) and in T2D (ZnT8RA,  $p < 0.001$ ; ZnT8WA,  $p < 0.001$ ; ZnT8QA,  $p < 0.001$ ) compared to the controls (Figure 3, panel B).

GADA and IA-2A were detected in 52% ( $p < 0.001$ ) and in 31% ( $p < 0.001$ ), respectively, of the long-term T1D patients compared to the controls (Table 3) but were negative in all T2D patients. However, the median levels including also the samples below the cut-off of both GADA and IA-2A were higher in both patient groups compared to the controls (T1D: GADA  $p < 0.001$ , IA-2A  $p < 0.001$ ; T2D: GADA  $p < 0.001$ , IA-2A  $p = 0.02$ ) (Figure 3, panel C).

### **3.2. Combinations between NPYA and islet autoantibodies at baseline**

We tested whether NPY-LA, NPY-PA, or both, occurred more often in patients with any of the ZnT8A variants, GADA or IA-2A. At baseline, neither of the single NPYA variants nor both of them taken together, appeared more often with ZnT8RA, ZnT8WA, ZnT8QA, GADA or IA-2A in patients with long-term T1D (Table 4). However, in 3/6 (50%) of T2D patients NPYA were double positive with both ZnT8RA ( $p = 0.028$ ) and ZnT8WA ( $p = 0.008$ ). NPY-PA showed a substantial agreement with ZnT8WA (kappa 0.61,  $p = 0.001$ ) and a moderate agreement with ZnT8RA (kappa 0.51,  $p = 0.001$ ) in the T2D patients (Table 4). Among the NPY-LA positive T1D patients 6/11 (55%) were also positive for NPY-PA ( $p = 0.002$ ). The two NPYA variants showed a moderate agreement (kappa 0.50,  $p = 0.001$ ) in T1D and a substantial agreement (kappa 0.61,  $p = 0.001$ ) in T2D patients (Table 4).

### 3.3. Number of islet autoantibodies at baseline

Among the autoantibody positive (41/48, 85%) patients with long-term T1D, there were 25/48 (52%) positive for at least one ZnT8A variant, and 12/48 (25%) positive for all three ZnT8A (Table 3). Most of the T1D patients, 13/48 (27%) had one autoantibody, although 9/48 (19%) were positive for four autoantibodies (Table 5). One patient with T1D had all seven autoantibodies whereas one had NPY-LA only. Most of the T2D patients (18/26, 69%) were negative for all autoantibodies.

### 3.4. NPY and islet autoantibodies during follow-up of long-term T1D patients

The subgroup of 32 patients with long-term T1D was analyzed after duration of 21-50 years and 5-8 years after baseline (Table 2). In general, 27/32 (84%) initially autoantibody positive T1D patients, 17/32 (53%) remained autoantibody positive ( $p=0.002$ ) during follow-up. The NPY-LA frequency decreased from 9/32 (28%) to 0/32 (0%) and for NPY-PA from 5/32 (16%) to 0/32 (0%) (Table 6). The median levels of NPY-LA declined from 34 U/mL to 21 U/mL ( $p<0.001$ ) (Figure 4, panel A) and of NPY-PA from 20 U/mL to 12 U/mL ( $p<0.001$ ) (Figure 4, panel B). NPY-LA and NPY-PA levels correlated ( $R^2=0.446$ ;  $p=0.011$ ) at baseline, however this correlation was essentially lost ( $R^2=0.348$ ;  $p=0.051$ ) at follow-up.

In the same group of 32 patients, the frequency of ZnT8A decreased from 14/32 (44%) to 4/32 (13%). Also, the median levels of the ZnT8A variants were lower at the time of follow-up (ZnT8RA,  $p<0.001$ ; ZnT8WA,  $p<0.001$ ; ZnT8QA,  $p<0.001$ ) compared to baseline (Figure 5, panels A-C).

The frequency of GADA ( $p=0.687$ ) and IA-2A ( $p=1.0$ ) did not change between baseline and follow-up (Table 6). The levels of GADA ( $p=0.005$ ) and IA-2A ( $p<0.001$ ) declined during follow-up (Figure 5, panels D-E).

It was also tested if NPY-LA and NPY-PA correlated to any of the islet autoantibodies in the 32 T1D patients. The levels of NPY-LA did not correlate with any of the ZnT8A variants (ZnT8RA,  $R^2=0.341$ ,  $p=0.056$ ; ZnT8WA,  $R^2=0.307$ ,  $p=0.087$ ; ZnT8QA,  $R^2=0.329$ ,  $p=0.066$ ) at baseline, although there were significant correlations at follow-up (ZnT8RA,  $R^2=0.376$ ;  $p=0.034$ ; ZnT8WA,  $R^2=0.390$ ,  $p=0.027$ ; ZnT8QA,  $R^2=0.593$ ,  $p<0.001$ ). The levels of NPY-PA correlated with the levels of ZnT8WA ( $R^2=0.546$ ,  $p=0.001$ ) and ZnT8QA ( $R^2=0.524$ ,  $p=0.002$ ), but not

with ZnT8RA ( $R^2=0.287$ ,  $p=0.111$ ) at baseline. At follow-up, the correlation between the NPY-PA levels with ZnT8WA ( $R^2=0.395$ ,  $p=0.025$ ) and ZnT8QA ( $R^2=0.364$ ,  $p=0.040$ ) variants remained (ZnT8RA,  $R^2=0.302$ ,  $p=0.093$ ). The levels of the two NPYA did not correlate with GADA or IA-2A at baseline or at follow-up.

### **3.5. NPYA and neuropathy in re-examined T1D patients**

Peripheral neuropathy was related to cardiac autonomic neuropathy ( $R^2=0.328$ ;  $p<0.001$ ) in the long-term follow-up patients with T1D (Figure 6). It was observed that males (12/14; 86%), more often had an abnormal cardiac autonomic neuropathy (cut-off at -1.64) compared to females (2/14; 14%) ( $p=0.007$ ). Taking these gender differences into account, we also addressed the question whether peripheral, autonomic, or both, neuropathies in the 32 T1D patients re-examined after 5-8 years were associated to any of the autoantibodies. Neither peripheral nor cardiac autonomic neuropathy was related to the presence or levels of the NPYA nor was there a correlation with ZnT8A, GADA or IA-2A.

## **4. Discussion**

In this study, we established an autoantibody radiobinding assay to assess the levels and frequency of two NPY autoantibody variants (NPY-LA and NPY-PA) in patients with T1D and T2D compared to controls. The major findings were as follows. First, newly diagnosed T1D patients had 29% NPY-LA alone, 29% NPY-PA alone but 20% being double positive. Second, long duration T1D patients had 23% and 19% compared to 12% and 23%, respectively, in the patients with T2D using the control 97.5<sup>th</sup> percentile as cut-off. Previously, NPY-LA was reported in 9% of T1D patients newly diagnosed below the age of 19 years [6]. It was therefore of interest that the present group of patients not only were diagnosed as adults but also had disease duration of median 26 years at the time when the first samples were obtained (referred to as baseline). More remarkably, at follow-up the levels had decreased significantly. These findings were comparable to GADA, IA-2A and the ZnT8A, which showed decreased levels during follow-up despite the lack of correlation between the levels of any of the two NPYA with GADA and IA-2A. The loss of autoantibodies over time in the follow-up samples may be due to both storage and an unknown number of freezing and thawing cycles.

Other studies of newly diagnosed patients with T1D in active follow-up [27] will be needed to determine if NPYA will decrease rapidly after diagnosis such as IA-2A and ZnT8A or slowly similar to GADA [28]. It should also be noted that the present group of long-term T1D patients had frequencies of GADA, IA-2A and ZnT8A at baseline ranging between 31-52%.

Another major finding was the comparable frequencies of NPYA among the patients with T2D. These patients were also long-term and their duration amounted to an average of 23 years. The NPYA in this group of patients may be easily explained by the possibility that at least some of these patients might have had autoimmune diabetes (LADA) as opposed to T2D. In the event that LADA would be the classification, none of the patients would qualify as neither GADA nor IA-2A were found among the T2D patients. If the ZnT8A would allow these patients to be classified as LADA, 23% of our T2D patients would be classified with LADA. The frequency of ZnT8A in LADA patients has been reported to vary between 2% [29], 34% [30] and 42% [31]. Our observation that the NPY-PA variant correlated to ZnT8RA and ZnT8WA in the T2D patients (Table 4) is of interest for two reasons. The first is that ZnT8A in addition to GADA and IA-2A may also be a marker of autoimmune diabetes or LADA [32]. The second is that the co-occurrence of NPYA and ZnT8A would further support the notion that also long-term 56-73 year old T2D patients may have autoimmune diabetes [32]. The results of our study demonstrate that NPY is not necessarily a minor autoantigen to T1D, rather that it may be related to autoimmune diabetes. Further studies are needed to clarify to what extent NPYA and ZnT8A tend to occur together in newly diagnosed diabetes patients classified with either T1D or T2D.

To our knowledge this is the first study to investigate two different NPY autoantibody variants in patients with long-term T1D and T2D. Although the appearance of islet autoantibodies prior to the clinical onset of T1D have made it possible to predict disease, it is still of interest to establish to what extent islet autoimmunity may be related to late diabetes complications. It should be noted that the T1D patients in the present study were primarily investigated with a second hypothesis in mind namely to test whether NPYA are related to neuropathy. As a response to nerve injury, NPY increases in different regions of the nervous system [33] [34] [35]. Also, it has been reported that the NPY levels [36] and the release [37] are increased in the hypothalamic nuclei of streptozotocin-induced diabetic rats. However, in response to insulin-induced hypoglycemia the NPY expression is instead lost in patients with autonomic neuropathy [18]. On

the other hand, duration [38] and type [39] of diabetes or chronic hyperglycemia [40] have been suggested to be strongly related to the risk and degree of neuropathy in the lower extremities.

The appearances of islet antibodies GADA, IA-2A and ZnT8A predict beta-cell failure in T1D, while in T2D, where they are less common, have a higher preserved beta-cell function [41]. Yet, the possible relevance of autoantibodies in neuropathy is less known. It has been reported that increased GADA levels at onset were associated with retinopathy in patients with T1D [42]. In addition, higher GADA levels were found in T1D patients with retinopathy and nephropathy compared to T1D patients without complications, which may suggest GADA as a complication predictor [43]. Also, T1D patients with higher GADA levels was reported to have a slower motor nerve conduction velocity compared to T1D patients with lower GADA levels [19]. The ability to predict peripheral and autonomic neuropathy by studying the correlation with the islet autoantibody levels and frequency in patients with T1D was also investigated in the present study. Neither of the peripheral nor the autonomic neuropathy correlated to the presence of the autoantibodies in the patients. There was a clear difference of the abnormal autonomic neuropathy between genders, but this difference was not related to any of the islet autoantibodies, including NPY-LA or NPY-PA. This lack of association may be due to the small number of T1D patients (n=32). Thus, we cannot conclude that the presently used autoantibodies can predict presence of peripheral or autonomic neuropathy in T1D. We noted that 6/26 T2D patients (5/6 were overweight or obese, data not shown) had at least one of the NPYA variants. Similarly, ZnT8A were also positive in 5/6 overweight or obese T2D. Although there was no relation between NPYA (or ZnT8A) and neuropathy in the T1D patients, it cannot be excluded that there may be such a relation in T2D. Unfortunately, we did not have any neuropathy data of the T2D patients. Further studies are warranted to investigate this possibility.

It is known that the ZnT8 transporter is targeted by three ZnT8A variants dependent on the T2D associated SNP rs13266634 at the amino acid at position 325 [44] [45] [26]. Previously, it was indicated that the structure of the mature NPY protein is dependent on the amino acid residue at SNP rs16139 [12] resulting in an altered packaging and secretion of NPY [13]. The genetic polymorphisms of the patients in this study were not known. It would be of interest to determine if autoantibodies against NPY are related to the L7P genetic polymorphism. Our kappa analysis suggests that there was substantial agreement between the two NPYA variants, which suggest that NPY L7P polymorphisms may not be as strong as the ZnT8 R325W polymorphism [24] to

determine a crucial epitope. In order to detect specific serum autoreactivity against either NPY-P or NPY-L protein, displacement assays with a cold inhibitor was used. However, specific sera were rare and further analysis of large number of sera will be needed to search for possible isoform-specific sera. It cannot be excluded that individuals may have such autoantibodies.

## **5. Conclusions**

Autoantibodies against both NPY-L and NPY-P were present in long-term T1D and T2D patients, and the NPYA levels declined during follow-up. Neither the frequency nor the titers of the NPYA variants were related to peripheral or cardiac autonomic neuropathy in the T1D patients. We conclude that NPY fulfills the criteria for a minor autoantigen in T1D. It cannot be excluded that NPY autoimmunity may contribute to the pathogenesis in long-term T1D, perhaps also in T2D.

## **Acknowledgements**

We thank Carina Törn, Anita Nilsson and Ingrid Wigheden for expert advice. We are especially thankful to Göran Sundkvist and Victoria Granberg for recruiting and collecting the patient samples. Also, Ann Radelius was invaluable to the organization and to all the patients who participated in this study.

## **Conflict of interests**

Declaration of competing interest: Nothing to declare.

## **References**

1. Winter, W.E. and D.A. Schatz, *Autoimmune markers in diabetes*. Clinical chemistry, 2011. **57**(2): p. 168-75.
2. Kulmala, P., et al., *Prediction of insulin-dependent diabetes mellitus in siblings of children with diabetes. A population-based study. The Childhood Diabetes in Finland Study Group*. The Journal of clinical investigation, 1998. **101**(2): p. 327-36.
3. Lernmark, A., *Autoimmune diseases: are markers ready for prediction?* The Journal of clinical investigation, 2001. **108**(8): p. 1091-6.

4. Plotz, P.H., *The autoantibody repertoire: searching for order*. Nature reviews. Immunology, 2003. **3**(1): p. 73-8.
5. Andersson, C., et al., *The three ZNT8 autoantibody variants together improve the diagnostic sensitivity of childhood and adolescent type 1 diabetes*. Autoimmunity, 2011.
6. Hirai, H., et al., *Selective screening of secretory vesicle-associated proteins for autoantigens in type 1 diabetes: VAMP2 and NPY are new minor autoantigens*. Clinical immunology, 2008. **127**(3): p. 366-74.
7. Zukowska-Grojec, Z., *Neuropeptide Y. A novel sympathetic stress hormone and more*. Annals of the New York Academy of Sciences, 1995. **771**: p. 219-33.
8. Gerald, C., et al., *A receptor subtype involved in neuropeptide-Y-induced food intake*. Nature, 1996. **382**(6587): p. 168-71.
9. Adeghate, E. and A. Ponery, *Pancreatic peptides, neuropeptides and neurotransmitters in diabetes mellitus: a review*. Int J Diabetes & Metabolism, 2003(11): p. 1-6.
10. Karvonen, M.K., et al., *Association of a leucine(7)-to-proline(7) polymorphism in the signal peptide of neuropeptide Y with high serum cholesterol and LDL cholesterol levels*. Nature medicine, 1998. **4**(12): p. 1434-7.
11. Jaakkola, U., et al., *The Leu7Pro polymorphism of the signal peptide of neuropeptide Y (NPY) gene is associated with increased levels of inflammatory markers preceding vascular complications in patients with type 2 diabetes*. Microvascular research, 2010. **80**(3): p. 433-9.
12. Kallio, J., et al., *Altered intracellular processing and release of neuropeptide Y due to leucine 7 to proline 7 polymorphism in the signal peptide of preproneuropeptide Y in humans*. The FASEB journal : official publication of the Federation of American Societies for Experimental Biology, 2001. **15**(7): p. 1242-4.
13. Mitchell, G.C., et al., *A common single nucleotide polymorphism alters the synthesis and secretion of neuropeptide Y*. The Journal of neuroscience : the official journal of the Society for Neuroscience, 2008. **28**(53): p. 14428-34.
14. Niskanen, L., et al., *Leucine 7 to proline 7 polymorphism in the neuropeptide y gene is associated with retinopathy in type 2 diabetes*. Experimental and clinical endocrinology & diabetes : official journal, German Society of Endocrinology [and] German Diabetes Association, 2000. **108**(3): p. 235-6.



15. Nordman, S., et al., *Leu7Pro polymorphism in the neuropeptide Y (NPY) gene is associated with impaired glucose tolerance and type 2 diabetes in Swedish men*. Experimental and clinical endocrinology & diabetes : official journal, German Society of Endocrinology [and] German Diabetes Association, 2005. **113**(5): p. 282-7.
16. Ukkola, O. and Y.A. Kesaniemi, *Leu7Pro polymorphism of PreproNPY associated with an increased risk for type II diabetes in middle-aged subjects*. European journal of clinical nutrition, 2007. **61**(9): p. 1102-5.
17. Jaakkola, U., et al., *The Leu7Pro polymorphism of neuropeptide Y is associated with younger age of onset of type 2 diabetes mellitus and increased risk for nephropathy in subjects with diabetic retinopathy*. Experimental and clinical endocrinology & diabetes : official journal, German Society of Endocrinology [and] German Diabetes Association, 2006. **114**(4): p. 147-52.
18. Bolinder, J., et al., *Autonomic neuropathy is associated with impaired pancreatic polypeptide and neuropeptide Y responses to insulin-induced hypoglycaemia in Type I diabetic patients*. Diabetologia, 2002. **45**(7): p. 1043-4.
19. Hoeldtke, R.D., et al., *Antibodies to glutamic acid decarboxylase and peripheral nerve function in type 1 diabetes*. The Journal of clinical endocrinology and metabolism, 2000. **85**(9): p. 3297-308.
20. Hoeldtke, R.D., et al., *Antibodies to GAD65 and peripheral nerve function in the DCCT*. Journal of neuroimmunology, 2007. **185**(1-2): p. 182-9.
21. Pourhamidi, K., et al., *Heat shock protein 27 is associated with better nerve function and fewer signs of neuropathy*. Diabetologia, 2011. **54**(12): p. 3143-9.
22. Forsen, A., et al., *A 14-year prospective study of autonomic nerve function in Type I diabetic patients: association with nephropathy*. Diabetic medicine : a journal of the British Diabetic Association, 2004. **21**(8): p. 852-8.
23. Kjelleras, J., F. Vaziri-Sani, and D. Agardh, *Improved efficacy by using the pTnT-rhtTG plasmid for the detection of celiac disease specific tissue transglutaminase autoantibodies in radioligand binding assays*. Scandinavian journal of clinical and laboratory investigation, 2011. **71**(8): p. 701-4.
24. Skarstrand, H., A. Lernmark, and F. Vaziri-Sani, *Antigenicity and epitope specificity of ZnT8 autoantibodies in type 1 diabetes*. Scandinavian journal of immunology, 2012.

25. Vaziri-Sani, F., et al., *A novel triple mix radiobinding assay for the three ZnT8 (ZnT8-RWQ) autoantibody variants in children with newly diagnosed diabetes*. Journal of immunological methods, 2011. **371**(1-2): p. 25-37.
26. Delli, A.J., et al., *Zinc Transporter 8 Autoantibodies and Their Association With SLC30A8 and HLA-DQ Genes Differ Between Immigrant and Swedish Patients With Newly Diagnosed Type 1 Diabetes in the Better Diabetes Diagnosis Study*. Diabetes, 2012.
27. Williams, A.J. and P.J. Bingley, *Worth the wait: type 1 diabetes prospective birth cohort studies enter adolescence*. Diabetologia, 2012. **55**(7): p. 1873-6.
28. Savola, K., et al., *Autoantibodies associated with Type I diabetes mellitus persist after diagnosis in children*. Diabetologia, 1998. **41**(11): p. 1293-7.
29. Hawa, M.I., et al., *Adult-Onset Autoimmune Diabetes in Europe Is Prevalent With a Broad Clinical Phenotype: Action LADA 7*. Diabetes care, 2012.
30. Andersen, M.K., et al., *Zinc transporter type 8 autoantibodies (ZnT8A): prevalence and phenotypic associations in latent autoimmune diabetes in adults and type 1 diabetes diagnosed >35 years*. Autoimmunity, 2013.
31. Vaziri-Sani, F., et al., *ZnT8 autoantibody titers in type 1 diabetes patients decline rapidly after clinical onset*. Autoimmunity, 2010. **43**(8): p. 598-606.
32. Rolandsson, O. and J.P. Palmer, *Latent autoimmune diabetes in adults (LADA) is dead: long live autoimmune diabetes!* Diabetologia, 2010. **53**(7): p. 1250-3.
33. Wakisaka, S., et al., *Cell size-specific appearance of neuropeptide Y in the trigeminal ganglion following peripheral axotomy of different branches of the mandibular nerve of the rat*. Brain research, 1993. **620**(2): p. 347-50.
34. Noguchi, K., et al., *Quantification of axotomy-induced alteration of neuropeptide mRNAs in dorsal root ganglion neurons with special reference to neuropeptide Y mRNA and the effects of neonatal capsaicin treatment*. Journal of neuroscience research, 1993. **35**(1): p. 54-66.
35. Ji, R.R., et al., *Expression of neuropeptide Y and neuropeptide Y (Y1) receptor mRNA in rat spinal cord and dorsal root ganglia following peripheral tissue inflammation*. The Journal of neuroscience : the official journal of the Society for Neuroscience, 1994. **14**(11 Pt 1): p. 6423-34.

36. Williams, G., et al., *Increased neuropeptide Y concentrations in specific hypothalamic regions of streptozocin-induced diabetic rats*. Diabetes, 1989. **38**(3): p. 321-7.
37. Sahu, A., et al., *Neuropeptide Y release from the paraventricular nucleus increases in association with hyperphagia in streptozotocin-induced diabetic rats*. Endocrinology, 1992. **131**(6): p. 2979-85.
38. Bergstrom, B., et al., *Autonomic neuropathy in type I diabetes: influence of duration and other diabetic complications*. Acta medica Scandinavica, 1987. **222**(2): p. 147-54.
39. Dyck, P.J., et al., *Risk factors for severity of diabetic polyneuropathy: intensive longitudinal assessment of the Rochester Diabetic Neuropathy Study cohort*. Diabetes care, 1999. **22**(9): p. 1479-86.
40. Harati, Y., *Diabetic peripheral neuropathies*. Annals of internal medicine, 1987. **107**(4): p. 546-59.
41. Ekholm, E., et al., *No signs of progressive beta cell damage during 20 years of prospective follow-up of autoantibody-negative diabetes*. Acta diabetologica, 2012. **49**(1): p. 57-62.
42. Jensen, R.A., et al., *HLA Genes, Islet Autoantibodies and Residual C-Peptide at the Clinical Onset of Type 1 Diabetes Mellitus and the Risk of Retinopathy 15 Years Later*. PloS one, 2011. **6**(3): p. e17569.
43. Khan, M.W., et al., *Detection of autoantibodies against reactive oxygen species modified glutamic acid decarboxylase-65 in type 1 diabetes associated complications*. BMC immunology, 2011. **12**: p. 19.
44. Wenzlau, J.M., et al., *Identification of a major humoral epitope in Slc30A8 (ZnT8)*. Annals of the New York Academy of Sciences, 2008. **1150**: p. 252-5.
45. Brorsson, C., et al., *Correlations between islet autoantibody specificity and the SLC30A8 genotype with HLA-DQB1 and metabolic control in new onset type 1 diabetes*. Autoimmunity, 2011. **44**(2): p. 107-14.

**Table 1.** Frequency of NPY-L (Leucine), NPY-P (Proline), ZnT8R (Arginine), ZnT8W (Tryptophan), ZnT8Q (Glutamine), GAD65 and IA-2 autoantibodies in newly diagnosed T1D (n=100) patients compared to controls (n=398).

	<b>T1D n=100 (100%)</b>	<b>Controls n=398 (%)</b>	<b><i>p</i></b>
<b>Males, n (%)</b>	58 (58)	66 (263)	
<b>Age – median (range) years</b>	N.A.	44 (19-81)	
<b>Age at diagnosis – median (range) years</b>	9.8 (1.5-18.8)	N.A.	
<b>NPY-LA</b>	29 (29)	10 (3)	p<0.001
<b>NPY-PA</b>	29 (29)	8 (2)	p<0.001
<b>NPYA (1 ≥)</b>	38 (38)	12 (3)	p<0.001
<b>NPY-LA only</b>	1 (1)	1 (0.03)	N.A.
<b>NPY-PA only</b>	0 (0)	1 (0.03)	N.A.
<b>ZnT8RA</b>	48 (48)	8 (2)	p<0.001
<b>ZnT8WA</b>	50 (50)	9 (2)	p<0.001
<b>ZnT8QA</b>	34 (34)	9 (2)	p<0.001
<b>ZnT8A (1 ≥)</b>	68 (68)	14 (4)	p<0.001
<b>GADA</b>	64 (64)	8 (2)	p<0.001
<b>IA-2A</b>	71 (71)	4 (1)	p<0.001
<b>Autoantibody negative</b>	4 (4)	367 (92)	p<0.001

Differences of autoantibody frequency between patients and controls were considered significant when p-value of <0.05 was applied.

Abbreviations: NPY-LA, Neuropeptide Y-Leucine autoantibodies; NPY-PA, Neuropeptide Y-Proline autoantibodies; N.A. is not applicable.

**Table 2.** Demographic data of patients with long-term Type 1 (n=48) diabetes (T1D) and Type 2 (n=26) diabetes (T2D) at baseline compared to healthy controls (n=398), and long-term T1D patients (n=32) at follow-up.

<b>BASELINE</b>	<b>T1D</b>	<b>T2D</b>	<b>Controls</b>
<b>Subjects, n</b>	48	26	398
<b>Males, n (%)</b>	29 (60)	24 (92)	263 (66)
<b>Age – median (range) years</b>	46 (30-67)	63 (56-73)	43 (19-81)
<b>Age at onset – median (range) years</b>	20 (15-26)	42 (31-50) <sup>a</sup>	N.A.
<b>Duration – median (range) years</b>	26 (13-42)	23 (10-31)	N.A.
<b>HbA1c – median, range mmol/mol (%)</b>	66, 42-117 (8.2, 6.0-12.9) <sup>b</sup>	N.A.	N.A.
<b>C-peptide - median (range)</b>	<0.10 (<0.10-0.18)	0.61 (0.11-1.32) <sup>a</sup>	N.A.
<b>FOLLOW-UP</b>			
<b>Subjects, n</b>	32	N.A.	398
<b>Males, n (%)</b>	19 (59)	N.A.	263 (66)
<b>Age at onset – median (range) years</b>	19 (15-26)	N.A.	N.A.
<b>Duration – median (range) years</b>	33 (21-50)	N.A.	N.A.
<b>HbA1c – median, range mmol/mol (%)</b>	67, 42-93 (8.3, 6.0-10.7)	N.A.	N.A.
<b>C-peptide - median (range)</b>	<0.1 (<0.1-0.18)	N.A.	N.A.
<b>Composite Z-score leg – median (range)</b>	-2.8 (-7.0- -0.5) <sup>c</sup>	N.A.	N.A.
<b>E/I Z-score – median (range)</b>	-1.5 (-2.5-07)	N.A.	N.A.
<b>Abnormal E/I ratio (&lt; -1.64), n (%)</b>	14 (44)	N.A.	N.A.

The Z-score reflects the peripheral nerve function in the leg (peroneal and sural nerves) and expressed in age-corrected values. The E/I Z-score reflects the parasympathetic vagal nerve function and expressed in age-corrected values. E/I Z-score values less than -1.64 SD (95% confidence interval, one-sided test) below the age-related reference values were considered abnormal.

Abbreviations: T1D, Type 1 diabetes; T2D, Type 2 diabetes; N.A. is not applicable. Number of patients is indicated as followed; a=25; b=27; c=31.

**Table 3.** Frequency of NPY-L (Leucine), NPY-P (Proline), ZnT8R (Arginine), ZnT8W (Tryptophan), ZnT8Q (Glutamine), GAD65 and IA-2 autoantibodies in long-term T1D (n=48) and T2D (n=26) patients at baseline compared to the controls (n=398).

	<b>T1D n=48 (%)</b>	<b>T2D n=26 (%)</b>	<b>Controls n=398 (%)</b>	<b><i>p</i></b>
<b>NPY-LA</b>	11 (23)	3 (12)	10 (3)	*p<0.001 **p=0.039
<b>NPY-PA</b>	9 (19)	6 (23)	8 (2)	*p<0.001 **p<0.001
<b>NPYA (1 ≥)</b>	14 (29)	6 (23)	12 (3)	*p<0.001 **p<0.001
<b>NPY-LA only</b>	1 (2)	0 (0)	1 (0.03)	*N.A. **N.A.
<b>NPY-PA only</b>	0 (0)	3 (12)	1 (0.03)	*N.A. **N.A.
<b>ZnT8RA</b>	19 (40)	4 (15)	8 (2)	*p<0.001 **p=0.004
<b>ZnT8WA</b>	19 (40)	3 (12)	9 (2)	*p<0.001 **p=0.031
<b>ZnT8QA</b>	17 (35)	3 (12)	9 (2)	*p<0.001 **p=0.031
<b>ZnT8A (1 ≥)</b>	25 (52)	6 (23)	14 (4)	*p<0.001 **p<0.001
<b>GADA</b>	25 (52)	0 (0)	8 (2)	*p<0.001 **N.A.
<b>IA-2A</b>	15 (31)	0 (0)	4 (1)	*p<0.001 **N.A.
<b>Autoantibody negative</b>	7 (15)	18 (69)	367 (92)	*p<0.001 **p<0.001

P-values are shown as \*T1D patients compared to controls and \*\*T2D patients compared to controls. Differences of autoantibody frequency between patients and controls were considered significant when p-value of <0.05 was applied.

Abbreviations: NPY-LA, Neuropeptide Y-Leucine autoantibodies; NPY-PA, Neuropeptide Y-Proline autoantibodies; N.A. is not applicable.

**Table 4.** Agreement of NPY-L (Leucine) and NPY-P (Proline) autoantibodies with ZnT8, GAD65 and IA-2 islet autoantibodies to mark autoimmunity in long-term T1D and T2D patients at base-line.

<b>T1D patients (n=48)</b>				<b>T2D patients (n=26)</b>			
<b>Antibody1-Antibody2</b>	<b>Rank<sup>a</sup></b>	<b>kappa</b>	<b>p</b>	<b>Antibody1-Antibody2</b>	<b>Rank<sup>a</sup></b>	<b>kappa</b>	<b>p</b>
NPY-LA - NPY-PA	1	0.50	0.001	NPY-LA - NPY-PA	1	0.61	0.001
NPY-PA - ZnT8RA	2	0.23	0.065	NPY-PA - ZnT8WA	2	0.61	0.001
NPY-LA - ZnT8QA	3	0.21	0.131	NPY-PA - ZnT8RA	3	0.51	0.007
NPY-PA - ZnT8QA	4	0.19	0.161	NPY-PA - ZnT8QA	4	0.34	0.057
NPY-LA - ZnT8RA	5	0.16	0.248	NPY-LA - ZnT8WA	5	0.25	0.209
NPY-LA - ZnT8WA	5	0.16	0.248	NPY-LA - ZnT8QA	5	0.25	0.209
NPY-PA - ZnT8WA	7	0.14	0.277	NPY-LA - ZnT8RA	6	0.18	0.360
NPY-LA - IA-2A	8	0.06	0.677	NPY-LA - GADA		N.A.	
NPY-PA - IA-2A	9	0.02	0.881	NPY-LA - IA-2A		N.A.	
NPY-PA - GADA	10	-0.06	0.611	NPY-PA - GADA		N.A.	
NPY-LA - GADA	11	-0.06	0.616	NPY-PA - IA-2A		N.A.	

A P-value < 0.05 was considered significant, a= ranked by agreement to mark autoimmunity (kappa).

Abbreviations: NPY-LA, Neuropeptide Y-Leucine autoantibodies; NPY-PA, Neuropeptide Y-Proline autoantibodies; N.A. is not applicable.

**Table 5.** The number of autoantibodies in sera from long-term T1D (n=48), T2D patients (n=26) and in the healthy controls (n=398) detected by radioligand binding assay at baseline, respectively.

Subjects	Number of autoantibodies, n (%)								Total
	0	1	2	3	4	5	6	7	
<b>T1D</b>	7 (15)	13 (27)	10 (21)	2 (4)	9 (19)	3 (6)	3 (6)	1 (2)	48 (100)
<b>T2D</b>	18 (69)	2 (8)	4 (15)	0 (0)	1 (4)	1 (4)	0 (0)	0 (0)	26 (100)
<b>Controls</b>	367 (92.2)	16 (4)	10 (2.5)	2 (0.5)	1 (0.3)	0 (0)	2 (0.5)	0 (0)	398 (100)



**Table 6.** The autoantibody frequency of the NPY-L (Leucine), NPY-P (Proline), ZnT8R (Arginine), ZnT8W (Tryptophan), ZnT8Q (Glutamine), GAD65 and IA-2 autoantibodies in the subgroup (n=32) of long-term T1D patients at baseline (duration range 13-42 years) and at follow-up 5-8 years after (duration range 21-50 years) baseline.

<b>T1D patients, n (%)</b>	<b>32 (100)</b>		
<b>Autoantibodies n (%)</b>	<b>Baseline</b>	<b>Follow-up</b>	<b>p</b>
<b>NPY-LA</b>	9 (28)	0 (0)	N.A.
<b>NPY-PA</b>	5 (16)	0 (0)	N.A.
<b>ZnT8RA</b>	13 (41)	3 (9)	p=0.002
<b>ZnT8WA</b>	14 (44)	3 (9)	p=0.001
<b>ZnT8QA</b>	13 (41)	1 (3)	p<0.001
<b>GADA</b>	14 (44)	12 (38)	p=0.687
<b>IA-2A</b>	9 (28)	8 (25)	p=1.0

Differences of autoantibody frequency between baseline and follow-up were considered significant when p-value of <0.05 was applied.

Abbreviations: NPY-LA, Neuropeptide Y-Leucine autoantibodies; NPY-PA, Neuropeptide Y-Proline autoantibodies; N.A. is not applicable.

## **Legends to Figures**

**Figure 1.**  $^{35}\text{S}$ -methionine labeled NPY-L (Leucine) and NPY-P (Proline) proteins (expected Mr 10.8) generated by *in vitro* transcription and translation as analyzed by sodium dodecyl sulfate (SDS) polyacrylamide gel electrophoresis (PAGE).


**Figure 2.** The healthy control (n=398) levels in quantile-quantile (Q-Q) plots showed that the cut-off at 97.5<sup>th</sup> percentile was 44 U/mL for NPY-L (Leucine) (panel A) and 32 U/mL for NPY-P (Proline) (panel B) autoantibodies. The cut-off levels (dotted line) were used to distinguish NPY-L (■) and NPY-P (●) autoantibody positive T1D patients (n=48) from negative T1D patients (panel C). At baseline, 6/48 (12.5%) long-term T1D patients were positive for both NPY autoantibody variants (filled symbol in the upper right square) in panel D). Also, the titers of the NPY autoantibodies correlated ( $R^2=0.547$ ,  $p < 0.001$ ). A p-value of  $< 0.05$  was considered significant.

**Figure 3.** Box plots of autoantibody levels of NPY-L (Leucine) and NPY-P (Proline) (panel A), ZnT8R (Arginine), ZnT8W (Tryptophan) and ZnT8Q (Glutamine) (panel B) and GAD65 and IA-2 (panel C) in long-term 48 T1D (▣) and 26 T2D (■) patients compared to 398 healthy controls (□). A p-value of  $< 0.05$  was considered significant.

**Figure 4.** In the subgroup of long-term T1D patients (n=32) analyzed during follow-up (duration 21-50 years), 5-8 years after baseline, the median levels of NPY-LA (panel A) declined from 34 U/mL to 21 U/mL ( $p < 0.001$ ) and of NPY-PA (panel B) from 20 U/mL to 12 U/mL ( $p < 0.001$ ). A p-value of  $< 0.05$  was considered significant.

**Figure 5.** In the subgroup of long-term T1D patients (n=32) analyzed during follow-up (duration 21-50 years), 5-8 years after baseline, the median levels of ZnT8A, GADA and IA-2A declined demonstrated in panel A) ZnT8RA from 33 U/mL to 17 U/mL ( $p < 0.001$ ), panel B) ZnT8WA from 25 U/mL to 14 U/mL ( $p < 0.001$ ) and panel C) ZnT8QA from 52 U/mL to 28 U/mL ( $p < 0.001$ ). The median levels of GADA (panel D) declined from 33 U/mL to 20 U/mL ( $p = 0.005$ ) and of IA-2A (panel E) from 3 U/mL to 2 U/mL ( $p < 0.001$ ). A p-value of  $< 0.05$  was considered significant.

**Figure 6.** Correlation between cardiac autonomic Z-score for E/I ratio (mean of the longest R-R interval during expiration (E) divided by the mean of the shortest R-R interval during inspiration (I) expressed in age-corrected values) and peripheral (sural and peroneal function) composite Z-score were analyzed in the subgroup (n=32) of long-term T1D patients. The dotted line represents the cut-off (−1.64) for abnormal cardiac autonomic neuropathy calculated as E/I ratio. The solid line represents the correlation ( $R^2=0.328$ ,  $p<0.001$ ) between the E/I ratio and the peripheral Z-score composite. A p-value of  $<0.05$  was considered significant.



**Figure 1.**






Figure 2.



**Figure 3.**

(A)



(B)




Figure 4.





Figure 5.





**Figure 6.**