



LUND UNIVERSITY

Depression and use of antidepressants in Swedish nursing homes: a 12-month follow-up study.

Midlöv, Patrik; Andersson, Martin L; Östgren, Carl Johan; Mölsted, Sigvard

Published in:
International Psychogeriatrics

DOI:
[10.1017/S1041610213002354](https://doi.org/10.1017/S1041610213002354)

2014

[Link to publication](#)

Citation for published version (APA):

Midlöv, P., Andersson, M. L., Östgren, C. J., & Mölsted, S. (2014). Depression and use of antidepressants in Swedish nursing homes: a 12-month follow-up study. *International Psychogeriatrics*, 26(4), 669-675.
<https://doi.org/10.1017/S1041610213002354>

Total number of authors:
4

General rights

Unless other specific re-use rights are stated the following general rights apply:
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

Depression and use of antidepressants in Swedish nursing homes: a 12-month follow-up study

Patrik Midlöv,¹ Martin Andersson,¹ Carl Johan Östgren² and Sigvard Mölstedt¹

¹Department of Clinical Sciences in Malmö, Center for Primary Health Care Research, Lund University, Malmö, Sweden

²Department of Medical and Health Sciences, Linköping University, Linköping, Sweden

ABSTRACT

Background: The prescription of antidepressants in nursing homes has increased markedly since the introduction of SSRIs, while at the same time depressive symptoms often go unrecognized and untreated. The aim of this study was to examine whether depression among residents in nursing homes is treated adequately.

Methods: A sample of 429 participants from 11 Swedish nursing homes was selected and was assessed with the Cornell Scale for Depression in Dementia (CSDD) and using medical records and drug prescription data. For 256 participants a follow-up assessment was performed after 12 months.

Results: The prevalence of depression, according to medical records, was 9.1%, and the prevalence of CSDD score of ≥ 8 was 7.5%. Depression persisted in more than 50% of cases at the 12-month follow-up. Antidepressants were prescribed to 33% of the participants without a depression diagnosis or with a CSDD score of < 8 . 46.2% of all participants were prescribed antidepressants. 14% of the participants without a depression diagnosis or with a CSDD score of < 8 had psychotropic polypharmacy. 15.2% of all participants had psychotropic polypharmacy, which persisted at the 12-month follow-up in three-quarters of cases.

Conclusion: The prescription of antidepressants in frail elderly individuals is extensive and may be without clear indication. The clinical implication is that there is a need for systematic drug reviews at nursing homes, paying special attention to the subjects which are on antidepressants.

Key words: depression, antidepressants, nursing homes, Sweden, dementia

Introduction

Depression is a treatable condition and should not be considered a part of normal aging. Residents in nursing homes with multiple illnesses are at increased risk of acquiring additional mental disorders. Studies suggest that the prevalence of depressive symptoms in nursing homes could range from 14% to 82%, with a median prevalence of 29% (Seitz *et al.*, 2010; Snowden, 2010).

Feelings of anxiety and unease are common in nursing homes, with a prevalence of 13% to 65% depending on studies, and anxiety disorder prevalence varies from 3.5% to 11% (Seitz *et al.*, 2010). Recommended treatment for anxiety is similar to treatment for depression: selective serotonin reuptake inhibitors (SSRIs), serotonin-

norepinephrine reuptake inhibitors (SNRIs) and/or cognitive behavioral therapy (CBT) (NICE, 2011).

Antidepressants

In different European studies, the prevalence of antidepressant use in nursing homes has been around 30% (Mann *et al.*, 2009; Lustenberger *et al.*, 2011). In Sweden, there has been a five-fold increase in the prescription of antidepressants in individuals aged 80+ years from 1995 to 2005 in men, and a three-fold increase in women. This could partly be explained by SSRIs having milder side-effects, particularly on cognition, than TCAs (tricyclic antidepressants), which were widely used to treat depression before the introduction of SSRIs (Weitoft *et al.*, 2012).

A Swedish study from 2004 reported the prevalence of prescribed antidepressants in nursing homes to be 41%, whereas in the general Swedish population of people aged 80+ years, the usage was 11% in men and 18% in women (Fastbom and Schmidt, 2004).

Correspondence should be addressed to: Patrik Midlöv, Center for Primary Health Care Research, Lund University, Clinical Research Centre (CRC), Jan Waldenströms gata 35, Skåne University Hospital, 205 02 Malmö, Sweden. Phone: +4640 391363; Fax: +4640 391370. Email: patrik.midlov@med.lu.se. Received 4 Sep 2013; revision requested 18 Oct 2013; revised version received 5 Nov 2013; accepted 12 Nov 2013. First published online 16 December 2013.

Few studies have been conducted on adverse outcomes of SSRIs in nursing home settings, but a previous study indicated an increased risk of all cause-mortality, stroke/transient ischemic attack, falls, fracture, epilepsy/seizures, and hyponatremia in older people who were using SSRI. Other antidepressants showed similar risks, but TCAs, produced fewer of these adverse effects (Coupland *et al.*, 2011). In a Norwegian study discontinuation of antidepressant treatment in patients with dementia led to an increase in depressive symptoms (Bergh *et al.*, 2012). A Swedish study however concluded that withdrawal of SSRI treatment is often successful in nursing home patients (Lindstrom *et al.*, 2007).

A Cochrane review indicated that TCAs and SSRI were of the same efficacy but that TCAs are associated with a higher withdrawal rate due to side effects (Mottram *et al.*, 2006).

Recommended treatment for depression

According to Swedish guidelines for the treatment of depressive disorders (similar to NICE guidelines, (NICE, 2009)), CBT, alone or in combination with antidepressants, is recommended as the first-line treatment for mild to moderate depression.

Another Cochrane review indicated that CBT could be used to treat depressed elderly individuals, although only a few studies with small sample sizes have been conducted, and more research is needed (Wilson *et al.*, 2008).

Elderly patients with dementia and concurrent moderate major depressive disorder do benefit from antidepressant treatment both in respect of depressive symptoms and behavior disturbance (Lyketsos *et al.*, 2003), although the benefit is modest and scientific evidence is limited. Furthermore, one study showed no effect of SSRIs on depression in elderly patients with concurrent dementia (Banerjee *et al.*, 2011).

Aim

The aim of this study was to examine whether symptoms of depression among residents in nursing homes are treated adequately, and whether antidepressants are used in an appropriate manner.

Material and methods

Sample

Data were collected from the Study of Health and Drugs in the Elderly (SHADES), a longitudinal cohort study of elderly people living in 11 nursing




Figure 1. Flow chart of patients in the SHADES study.

homes in three cities in the southern part of Sweden (Linköping, Jönköping and Eslöv). All residents of the 11 nursing homes were invited to participate in the study and when a participant moved or died, the next person who moved to the nursing home was asked to participate. Individuals who were at the nursing home temporarily for short-term rehabilitation or palliative care were excluded. Persons with language difficulties and persons under the age of 65 were also excluded. A total of 429 individuals were included. The inclusion and exclusion criteria are shown in Figure 1.

Data collection

Specially trained nurses examined the participants in their respective nursing homes every six months between 2008 and 2011. All nursing homes patients have a general practitioner (GP) who regularly comes to the nursing home. Information on medical diagnoses, hospital admissions, rehabilitation plans (if any), and prescription of drugs were obtained from nursing home records and the GP's medical records.

Diagnoses were defined by present state and not by history of the disease.

Psychotropic polypharmacy was defined according to the Swedish National Board of Health and Welfare (The National Board of Health and Welfare, 2010) as simultaneous treatment with three or more psychotropic drugs from ATC (Anatomic Therapeutic Chemical classification system) groups N05A (antipsychotics), N05B (sedatives), N05C

Table 1. Baseline characteristics of the study participants

	ALL PARTICIPANTS (<i>n</i> = 429)			PARTICIPANTS FOR WHOM COMPLETE CSDD DATA WERE AVAILABLE (<i>n</i> = 401)		
	DEPRESSION <i>n</i> = 39	NO DEPRESSION <i>n</i> = 390	<i>p</i> -VALUE	CSDD ≥8 <i>n</i> = 30	CSDD <8 <i>n</i> = 371	<i>p</i> -VALUE
Age, mean (SD) ^a	86.3 (6.2)	84.8 (7.0)	0.203 ^a	85.2 (8.6)	84.9 (6.8)	0.862 ^a
Female, <i>n</i> (%)	31 (79.5)	274 (70.3)	0.225 ^c	23 (76.7)	263 (70.9)	0.501 ^c
Dementia, <i>n</i> (%)	12 (30.8)	165 (42.3)	0.163 ^c	12 (40.0)	151 (40.7)	0.940 ^c
≥3 psychotropic drugs, <i>n</i> (%)	11 (28.2)	54 (13.8)	0.017 ^c	10 (33.3)	50 (13.5)	0.007 ^d
≥5 drugs, <i>n</i> (%)	35 (92.3)	294 (75.4)	0.017 ^c	23 (76.7)	285 (76.8)	0.985 ^c
≥10 drugs, <i>n</i> (%)	8 (20.5)	82 (21.0)	0.940 ^c	6 (20.0)	81 (21.8)	0.815 ^c
CSDD ≥8, <i>n</i> (%)	2 (6.5)	28 (7.6)	1.000 ^d	–	–	–
Heart disease, <i>n</i> (%)	13 (33.3)	151 (38.7)	0.509 ^c	7 (23.3)	151 (40.7)	0.061 ^c
Cerebrovascular disease, <i>n</i> (%)	16 (41.0)	79 (20.3)	0.003 ^c	4 (13.3)	85 (22.9)	0.225 ^c
Diabetes, <i>n</i> (%)	3 (7.7)	75 (19.2)	0.075 ^c	5 (16.7)	68 (18.3)	0.820 ^c
Asthma, <i>n</i> (%)	0 (0.0)	12 (3.1)	0.613 ^d	0 (0.0)	11 (3.0)	1.000 ^d
Arthritis, <i>n</i> (%)	0 (0.0)	6 (1.5)	1.000 ^d	0 (0.0)	5 (1.3)	1.000 ^d
SSRI, <i>n</i> (%)	24 (61.5)	125 (32.1)	<0.001 ^c	13 (43.3)	125 (33.7)	0.319 ^c
MAOI, <i>n</i> (%)	1 (2.6)	4 (1.0)	0.381 ^d	1 (3.3)	4 (1.1)	0.324 ^d
Other antidepressants ^b , <i>n</i> (%)	16 (41.0)	56 (14.4)	<0.001 ^c	8 (26.7)	58 (15.6)	0.126 ^d
Sedatives, <i>n</i> (%)	21 (53.8)	161 (41.2)	0.130 ^c	17 (56.7)	151 (40.7)	0.088 ^c

^aIndependent samples t-test.

^bATC code N06AX (tryptophan, mianserin, mirtazapine, bupropion, venlafaxine, reboxetine, duloxetine or agomelatine).

^cPearson's chi-squared test.

^dFisher's exact test.

(hypnotics) or N06A (antidepressants), including subgroups (WHO Collaborating Centre for Drug Statistics Methodology, 2013).

In-person testing

The Cornell Scale for Depression in Dementia (CSDD) was used to measure depressive symptoms. The CSDD uses staff members to assist residents in giving answers, which makes it suitable for participants with cognitive dysfunction. In our study the CSDD was completed solely with the assistance of staff members. The CSDD consists of 19 questions regarding mood-related signs, behavioral disturbance, physical signs, cyclic functions, and ideational disturbance. Each answer is scored from 0 to 2 (0 = absent, 1 = mild/intermittent, 2 = severe), and the maximum score for the scale is 38. The cut-off score for depression was set at 8 (Alexopoulos *et al.*, 1988; Barca *et al.*, 2010). The mini-mental state examination (MMSE) was used to evaluate cognitive functions (Folstein *et al.*, 1975).

Statistical analyses

Descriptive statistics (Pearson's chi-squared test, Fisher's exact test, and independent t-tests) were

used to analyze differences in the material. Levene's test was used for equality of variances. Data analysis was conducted with SPSS Statistics for Windows Version 21.0 (IBM Corp, Armonk, NY, USA).

Ethical considerations

SHADES was approved by the Ethical Committee in Linköping, Sweden (application number M 150–07). Informed consent was obtained from all participants. If the patient could not understand the information and give informed consent, it was obtained from next of kin.

Results

At baseline, 15.2% of participants had prescriptions for three or more psychotropic drugs. Baseline characteristics are shown in Table 1. The prevalence of depression according to medical records was 9.1% overall and the prevalence of CSDD score of ≥8 was 7.5%.

In all, 46.2% of participants were prescribed one or more antidepressant at baseline and 33.0% of participants not diagnosed with depression and with a CSDD score of <8 were prescribed one or more antidepressants.

Table 2. Persistence of depression and use of medications at 12-month follow-up in the same individuals (*n* = 256)

	BASILINE, NUMBER OF PATIENTS	12-MONTH FOLLOW-UP, NUMBER OF PATIENTS (PERSISTENCE RATE, %)
Depression	25	14 (56.0)
CSDD ≥ 8	13	7 (53.8)
Antidepressants (one or more)	117	108 (92.3)
Psychotropic polypharmacy (≥ 3 drugs)	41	31 (75.6)
Number of drugs (≥ 5)	195	181 (92.8)
Number of drugs (≥ 10)	56	42 (75.0)
CSDD score ≥ 8 , treated with antidepressants at baseline	5	3 (60.0)
Sedatives or hypnotics (one or more)	108	93 (86.1)

Table 3. Comparisons between men and women at baseline

	MEN (<i>n</i> = 124)	WOMEN (<i>n</i> = 305)	<i>p</i> -VALUE
Antidepressant use, <i>n</i> (%)	57 (46.0)	141 (46.2)	0.961 ^a
Antidepressants use and age 80+ years, <i>n</i> (%), ^b	30 (35.7)	102 (42.7)	0.264 ^a
Depression, <i>n</i> (%)	8 (6.5)	31 (10.2)	0.225 ^a
CSDD score ≥ 8 , <i>n</i> (%)	7 (5.6)	23 (7.5)	0.501 ^a
Dementia, <i>n</i> (%)	51 (41.1)	126 (41.3)	0.972 ^a
Polypharmacy ≥ 10 , <i>n</i> (%) (<i>n</i> = 90)	28 (22.6)	62 (20.3)	0.603 ^a
Polypharmacy ≥ 5 , <i>n</i> (%) (<i>n</i> = 330)	100 (80.6)	230 (75.4)	0.243 ^a
Psychotropic polypharmacy (≥ 3 drugs), <i>n</i> (%) (<i>n</i> = 65)	19 (15.3)	46 (15.1)	0.950 ^a
MMSE score < 24 , <i>n</i> (%), ^c	77 (78.6)	205 (81.7)	0.508 ^a

^aPearson's chi-squared test.

^bIn total 323 nursing home residents were 80 years or older.

^cMMSE was assessed in 349 patients. For 80 patients this was not possible.

Persistence rates for depression and prescribed drugs at the 12-month follow-up are presented in Table 2, thus more than 90% of those treated with antidepressants were still on this treatment after 12 months.

No participant was treated with CBT.

There was a significant difference in median CSDD score between baseline and 12-month follow-up (1.0 and 2.0, respectively; $p < 0.001$, Wilcoxon signed ranked test).

The mean number of psychotropic drugs was 1.2 (SD 1.2) at baseline and 1.3 (SD 1.2) at 12-month follow-up, and the mean total number of drugs was 6.9 (SD 3.1) at baseline and 7.0 (SD 3.0) at 12-month follow-up. The most frequently prescribed group of psychotropic drugs was SSRIs (34.7%). The most frequently used drugs overall were aspirin (54.1%) and paracetamol (41.7%).

There were no significant differences between men and women regarding depression or use of antidepressants (Table 3). Patients without dementia were more often treated with sedatives/hypnotics but there were no associations between dementia and antidepressant prescription or psychotropic polypharmacy (Table 4).

Discussion

The prescription of antidepressants in frail elderly individuals was extensive and sometimes without clear indication. In most cases it persisted after 12 months although many of these patients had no depression according to medical records and a CSDD score that did not indicate depression.

There was an increase in CSDD score at 12-month follow-up compared to baseline, (median score 1.0 and 2.0, respectively), but it cannot be considered clinically significant as these numbers are much lower than the cut-off for depression (set at 8). Of the thirteen participants with CSDD scores of ≥ 8 , over seven still had a CSDD score of ≥ 8 at 12-month follow-up. This could be explained by symptoms of depression not being recognized, but conclusions are hard to draw due to the very low number of cases.

There was an association between psychotropic polypharmacy and diagnosed depression, as suspected, but nearly 14% of the non-depressed participants/participants with CSDD scores of < 8 were prescribed three or more psychotropic drugs. Psychotropic drugs have been used to treat

Table 4. Comparisons between participants with and without dementia, according to medical records, at baseline

	DEMENTIA (<i>n</i> = 177)	NO DEMENTIA (<i>n</i> = 252)	<i>p</i> -VALUE
Age (years), mean (SD) ^a	84.0 (6.4)	85.6 (7.2)	0.02 ^a
Female, <i>n</i> (%)	126 (71.2)	179 (71.0)	0.972 ^b
Psychotropic polypharmacy (>3 drugs), <i>n</i> (%)	26 (14.7)	39 (15.5)	0.823 ^b
Diagnosed depression, <i>n</i> (%)	12 (6.8)	27 (10.7)	0.163 ^b
CSDD score ≥8, <i>n</i> (%)	12 (6.8)	18 (7.1)	0.940 ^b
SSRI, <i>n</i> (%)	60 (33.9)	89 (35.3)	0.761 ^b
MAOI, <i>n</i> (%)	3 (1.7)	2 (0.8)	0.407 ^c
Other antidepressants, <i>n</i> (%)	35 (19.8)	37 (14.7)	0.165 ^b
One or more antidepressants, <i>n</i> (%)	88 (49.7)	110 (43.7)	0.215 ^b
Sedatives/hypnotics, <i>n</i> (%)	61 (34.5)	121 (48.0)	0.005 ^b

^aLevene's test for equality of variances.

^bPearson's chi-squared test.

^cFisher's exact test.

Behavioral and Psychological Symptoms of Dementia (BPSD), but when comparing participants with and without dementia, there were no associations between dementia and antidepressant prescription or psychotropic polypharmacy. There was however a possible under-diagnosis of dementia since 282 participants had MMSE scores <24 but only 177 participants had a dementia diagnosis.

At 12-month follow-up, three-quarters of the participants with three or more psychotropic drugs showed continued use. This may be adequate, but it raises questions about the necessity to treat residents with multiple other illnesses and medications with psychotropic drugs for this length of time.

The low prevalence of diagnosed anxiety (0.9%) probably did not reflect the real presence of anxiety and unease, which might not have been considered sufficiently severe to require a diagnosis.

A substantial proportion of the participants in this study had continued prescriptions of antidepressants after 12 months without an obvious indication.

There was no significant association between use of sedatives/hypnotics and diagnosed depression/CSDD score ≥8 at baseline, but patients without dementia, according to the medical records, used sedatives/hypnotics more often than patients with dementia.

This study has some limitations. There was no new examination by a physician to confirm the diagnosis of depression, as defined by CSDD score, at follow-up. Scales cannot diagnose depression, but only indicate the probability of depression. We did not analyze the relationship between CSDD score and depression diagnosis. The reason is that patients diagnosed with depression and subjected to treatment may have normal CSDD score.

Also, if conducting more comparisons, the effect of multiple tests and significance levels should be considered.

The sample used was not randomly selected, but rather selected for reasons of convenience from three different areas in Sweden, with persons living in nursing homes whose staffs were interested in joining the project. For residents who chose not to participate or were excluded, we have no information about diagnoses or their current medication. Thus, we do not know if these residents differed in baseline characteristics from the included subjects in any way.

Almost all of the participants with diagnosed depression were treated with antidepressants, and 56% were still depressed after 12 months. However, we do not have data on the current dosages of antidepressants. Similarly, 53.8% of participants with CSDD scores of ≥8 at baseline had scores of ≥8 at follow-up. This is consistent with another study that showed persistence rates of over 50% for depression at 12 months in frail elderly patients (Bergdahl *et al.*, 2005).

The prevalence of diagnosed depression was 9.1%. This is in line with prevalence of major depression in nursing homes that in a literature review was 10% (Seitz *et al.*, 2010).

We do not have any data on neuropsychiatric symptoms which is a limitation since this could have been valuable in patients with depression and dementia.

The prevalence of one or more antidepressants in the participants aged 80+ years was 42.7% in women and 35.7% in men. These values are considerably higher than the figures from the National Public Health Report 2012, according to which 18% of women and 11% of men in the

same age group were using antidepressants (Weitoft *et al.*, 2012). The health report does not separate frail elderly individuals from healthier individuals, which could explain the discrepancy. One should however in patients with multimorbidity consider whether guideline recommendations are applicable and be aware that these guidelines may drive polypharmacy for patients in whom treatment burden will sometimes be overwhelming (Hughes *et al.*, 2013).

A Norwegian qualitative study, based on interviews with physicians and nurses in nursing homes, indicated that decisions to prescribe antidepressants were not always based on clinical diagnoses, but more often relied on registered nurses' opinions, and that there was a lack of follow-up (Iden *et al.*, 2011). This may well be the case in the present study, as 33% of all participants without diagnosed depression or a CSDD score of ≥ 8 were prescribed antidepressants, without an obvious indication, which is also consistent with another study (Shah *et al.*, 2012).

A report from the Swedish Council on Health Technology Assessment (SBU) did not assess CSDD due to a lack of high quality studies and urged for more studies on the subject (Swedish Council on Health Technology Assessment (SBU), 2012).

A Norwegian reliability and validity study of CSDD suggested a cut-off point of 8 based on the ICD-10 criteria for depression (Barca *et al.*, 2010), while a Japanese study suggested a cut-off score at 5 (Schreiner *et al.*, 2003). We chose to use a cut-off of 8, but there have been few validity studies of the CSDD.

Although chronic diseases and depression have been shown to be associated with each other, in this study there was only evidence of associations between cerebrovascular disease and diagnosed depression. This may be due to the low numbers of participants with a chronic disease and concomitant depression/CSDD score ≥ 8 (Moussavi *et al.*, 2007).

This study shows that despite the extensive knowledge in this field, little has been changed regarding the treatment of frail elderly individuals, regarding the use of antidepressants without documented indication or the lack of non-pharmacological treatment of depression. We conclude that the guidelines were not being followed, especially not concerning non-pharmacological treatment, which was virtually non-existent in the elderly. Non-pharmacological treatment should be considered feasible in the elderly, especially in view of the extent of polypharmacy in this group and the associated hazards.

There is a need for intervention studies comparing different kinds of pharmacological and

non-pharmacological treatments for depression in nursing home residents. Furthermore, the clinical implication from this study is that there is a need for systematic drug reviews at nursing homes, paying special attention to subjects who are on antidepressants.

Conflict of interest

None.

Description of authors' roles

PM, CJÖ, and SM designed the study. MA performed the data analysis. PM and MA drafted the paper. CJÖ and SMÖ revised the paper. All authors read and approved the final version of the paper.

Acknowledgments

This study was financially supported by the Medical Research Council of Southeast Sweden (FORSS) and the Janne Elgqvist Foundation.

We are indebted to Stephen Gilliver for his expertise and invaluable advice in editing the paper.

References

- Alexopoulos, G. S., Abrams, R. C., Young, R. C. and Shamoian, C. A. (1988). Cornell Scale for Depression in Dementia. *Biological Psychiatry*, 23, 271–284.
- Banerjee, S. *et al.* (2011). Sertraline or mirtazapine for depression in dementia (HTA-SADD): a randomised, multicentre, double-blind, placebo-controlled trial. *Lancet*, 378, 403–411.
- Barca, M. L., Engedal, K. and Selbaek, G. (2010). A reliability and validity study of the cornell scale among elderly inpatients, using various clinical criteria. *Dementia and Geriatric Cognitive Disorders*, 29, 438–447.
- Bergdahl, E. *et al.* (2005). Depression among the oldest old: the Umea 85+ study. *International Psychogeriatrics*, 17, 557–575.
- Bergh, S., Selbaek, G. and Engedal, K. (2012). Discontinuation of antidepressants in people with dementia and neuropsychiatric symptoms (DESEP study): double blind, randomised, parallel group, placebo controlled trial. *British Medical Journal*, 344, e1566.
- Coupland, C., Dhiman, P., Morriss, R., Arthur, A., Barton, G. and Hippisley-Cox, J. (2011). Antidepressant use and risk of adverse outcomes in older people: population based cohort study. *British Medical Journal*, 343, d4551.
- Fastbom, J. and Schmidt, I. (2004). Shortages in the treatment of depression of the elderly in nursing homes in Stockholm. Diagnosis is based on flimsy ground, follow up performed early. *Lakartidningen*, 101, 3683–3684, 3686–3688.

- Folstein, M. F., Folstein, S. E. and McHugh, P. R.** (1975). "Mini-mental state." A practical method for grading the cognitive state of patients for the clinician. *Journal of Psychiatric Research*, 12, 189–198.
- Hughes, L. D., McMurdo, M. E. and Guthrie, B.** (2013). Guidelines for people not for diseases: the challenges of applying UK clinical guidelines to people with multimorbidity. *Age Ageing*, 42, 62–69.
- Iden, K. R., Hjorleifsson, S. and Ruths, S.** (2011). Treatment decisions on antidepressants in nursing homes: a qualitative study. *Scandinavian Journal of Primary Health Care*, 29, 252–256.
- Lindstrom, K., Ekedahl, A., Carlsten, A., Martensson, J. and Molstad, S.** (2007). Can selective serotonin inhibitor drugs in elderly patients in nursing homes be reduced? *Scandinavian Journal of Primary Health Care*, 25, 3–8.
- Lustenberger, I., Schupbach, B., von Gunten, A. and Mosimann, U.** (2011). Psychotropic medication use in Swiss nursing homes. *Swiss Medical Weekly*, 141, w13254.
- Lyketsos, C. G. et al.** (2003). Treating depression in Alzheimer disease: efficacy and safety of sertraline therapy, and the benefits of depression reduction: the DIADS. *Archives of General Psychiatry*, 60, 737–746.
- Mann, E., Kopke, S., Haastert, B., Pitkala, K. and Meyer, G.** (2009). Psychotropic medication use among nursing home residents in Austria: a cross-sectional study. *BMC Geriatrics*, 9, 18.
- Mottram, P. G., Wilson, K. and Strobl, J. J.** (2006). Antidepressants for depressed elderly. *Cochrane Database of Systematic Reviews*, CD003491.
- Moussavi, S., Chatterji, S., Verdes, E., Tandon, A., Patel, V. and Ustun, B.** (2007). Depression, chronic diseases, and decrements in health: results from the World Health Surveys. *Lancet*, 370, 851–858.
- NICE** (2009). *Depression: The Treatment and Management of Depression in Adults*. London: National Institute for Health and Clinical Excellence (NICE).
- NICE** (2011). *Generalised Anxiety Disorder and Panic Disorder (with or Without Agoraphobia) in Adults*. London: National Institute for Health and Clinical Excellence (NICE).
- Schreiner, A. S., Hayakawa, H., Morimoto, T. and Kakuma, T.** (2003). Screening for late life depression: cut-off scores for the Geriatric Depression Scale and the Cornell Scale for Depression in Dementia among Japanese subjects. *International Journal of Geriatric Psychiatry*, 18, 498–505.
- Seitz, D., Purandare, N. and Conn, D.** (2010). Prevalence of psychiatric disorders among older adults in long-term care homes: a systematic review. *International Psychogeriatrics*, 22, 1025–1039.
- Shah, S., Schoenbachler, B., Streim, J. and Meeks, S.** (2012). Antidepressant prescribing patterns in the nursing home: second-generation issues revisited. *Journal of the American Medical Directors Association*, 13, 406 e413–e408.
- Snowdon, J.** (2010). Depression in nursing homes. *International Psychogeriatrics*, 22, 1143–1148.
- Swedish Council on Health Technology Assessment (SBU).** (2012). Diagnosis and monitoring of mood disorders – a systematic review. Stockholm.
- The National Board of Health and Welfare** (2010). Indicators for evaluation of the quality of drug use by elderly. Stockholm.
- Weitoft, G. R., Ericsson, O. and Fastbom, J.** (2012). Prescription drugs: Health in Sweden: The National Public Health Report 2012. Chapter 18. *Scandinavian Journal of Public Health*, 40, 293–304.
- WHO Collaborating Centre for Drug Statistics Methodology.** (2013). Anatomical Therapeutic Chemical (ATC) Classification Index. Oslo: World Health Organisation.
- Wilson, K., Mottram, P. G. and Vassilas, C.** (2008). Psychotherapeutic treatments for older depressed people. *Cochrane Database of Systematic Reviews*, CD004853.