LUND UNIVERSITY

IGF1 htSNPs in relation to IGF-1 levels in young women from high-risk breast cancer families: implications for early-onset breast cancer

Henningson, Maria; Hietala, Maria; Törngren, Therese; Olsson, Håkan; Jernström, Helena

Published in:
Familial Cancer

DOI:
[10.1007/s10689-010-9404-z](https://doi.org/10.1007/s10689-010-9404-z)

2011

[Link to publication](#)

Citation for published version (APA):

Henningson, M., Hietala, M., Törngren, T., Olsson, H., & Jernström, H. (2011). IGF1 htSNPs in relation to IGF-1 levels in young women from high-risk breast cancer families: implications for early-onset breast cancer. *Familial Cancer*, 10(2), 173-185. <https://doi.org/10.1007/s10689-010-9404-z>

Total number of authors:
5

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

IGF1 htSNPs in relation to IGF-1 levels in young women from high-risk breast cancer families: implications for early-onset breast cancer

Maria Henningson¹, Maria Hietala¹, Therese Törngren¹, Håkan Olsson^{1,2} and Helena Jernström¹

¹Department of Oncology, Clinical Sciences, Lund University, Sweden

² Department of Cancer Epidemiology, Clinical Sciences, Lund University, Sweden

Corresponding Author: Helena Jernström, Department of Oncology, Clinical Sciences, Lund University, Barngatan 2B, SE-221 85 Lund, Sweden, Email: helena.jernstrom@med.lu.se
Fax: + 46 46 14 73 27

Keywords: *BRCA1/2*; insulin-like growth factor-1; oral contraception; *IGF1* polymorphism; *IGF1* diplotype; breast cancer

Running head: *IGF1* polymorphisms in high-risk women

Abbreviations:

IGF1	insulin-like growth factor-1
htSNPS	haplotype tagging single nucleotide polymorphisms
BRCA	breast cancer associated gene
ECR	evolutionary conserved regions
LD	linkage disequilibrium
IGF1R	type 1 IGF receptor
IRS1	insulin receptor substrate 1

IGFBP2 IGF binding protein 2

OCs oral contraceptives

BMI body mass index

Abstract

High levels of insulin-like growth factor-1 (IGF-1) have been associated with increased risk of developing several types of cancer including breast cancer. A set of nine haplotype tagging SNPs (htSNPs) in the *IGF1* gene were associated with IGF-1 levels and prostate cancer in a Swedish population. We aimed to study the nine htSNPs in three haplotype blocks (block1: rs855211, rs35765, rs2162679; block2: rs1019731, rs7956547, rs5742632; and block3: rs2033178, rs7136446, rs6220) combined into diplotypes, and three additional SNPs (rs5742612, rs35765817, rs35455143) in relation to IGF-1 levels, *BRCA* status, the *IGF1* CA-repeat microsatellite, and breast cancer in a population of 325 Swedish women from breast cancer high-risk families. Questionnaire data and blood samples for IGF-1 and genetic analyses were obtained twice during the menstrual cycle from 269 women aged 40 years or younger. SNP analyses were also performed in 56 *BRCA1/2* mutation carriers. Women (n=14) with any rare variant block1 diplotype had higher odds to be *BRCA1* mutation carriers OR 4.1 (95% CI 1.4-12.2), to lack the common *IGF1* 19 CA-repeat allele OR 33.3 (95% CI 6.6-166.7), and were more likely to develop early-onset breast cancer (Log Rank $P < 0.001$) than women with common block1 diplotypes. In the subgroup of *BRCA1* mutation carriers, block1 rare diplotypes were associated with earlier diagnosis (Log Rank $P = 0.031$). No association was found between IGF-1 levels and individual SNPs or diplotypes. If confirmed, these rare diplotypes may identify women with particularly high risk for early-onset breast cancer and this group should be included in forthcoming studies.

Introduction

Insulin-like growth factor-1 (IGF-1) is important for normal breast development [1]. It stimulates cell proliferation and decreases apoptosis [2]. High levels of circulating IGF-1 have repeatedly been associated with several types of cancer, including prostate and premenopausal breast cancer [3-5]. The levels of circulating IGF-1 are highly influenced by genetic factors [6], and several polymorphic variants including single nucleotide polymorphisms (SNPs) and microsatellites have been associated with both IGF-1 levels and breast cancer risk [7-10]. SNPs (rs35765817, rs35455143, and rs3839984) located in evolutionary conserved regions (ECR) close to binding sites for a transcription factor that is important for IGF-1 expression were associated with circulating IGF-1 levels in a study of Caucasian women [10]. Another SNP (rs5742612) was found to be in linkage disequilibrium (LD) with variations in a microsatellite 969 bp upstream from the *IGF1* transcription start site among Singapore Chinese women [11].

Haplotype tagging SNPs (htSNPs) can be used as markers that cover the majority of genetic diversity of several individual SNPs. In the *IGF1* gene, a set of nine htSNPs (rs855211, rs35765, rs2162679, rs1019731, rs7956547, rs5742632, rs2033178, rs7136446, and rs6220) in three haplotype blocks with three htSNPs in each was studied in relation to prostate cancer risk in a Swedish population. One haplotype (TCC in block three) in the 3' region of the gene was associated with increased risk of developing prostate cancer [12].

Genetic variations in genes other than *IGF1* are likely to also influence IGF-1 levels. BRCA1 has been shown to interact with IGF signaling. In *Brc1*-deficient mice, the expression of IGF1R, IRS1, and IGFBP2 was increased and so were the serum levels of IGF-1 compared with those of control mice expressing *Brc1* [13]. BRCA1 suppresses IGF1R promoter activity

in MCF-7 breast cancer cell lines [14]. In one study, the IGF1R levels were higher in tumors from *BRCA1* mutation carriers compared with non-carriers [15]. IGF-1 is also important for intrauterine growth and *BRCA1* mutation carriers have been reported to be smaller than non-carriers from *BRCA1* mutation families [16], which further supports a link between *BRCA1* and IGF-1. Moreover, we have previously reported an association between absence of the common *IGF1* 19 CA-repeat polymorphism and *BRCA1* mutation status [17]. Absence of the 19 CA-repeat has been associated with increased premenopausal breast cancer risk [7, 8]. *BRCA1* and *BRCA2* mutation carriers have a substantially increased risk of developing breast cancer. However, the lifetime risk estimates range from 20 to 80% [18], a range that suggests that risk modifying genetic and non-genetic factors play an important role in carcinogenesis. We hypothesized that other genetic variations in the *IGF1* gene may, by altering IGF-1 levels, contribute to the variation in breast cancer susceptibility between and within families with a high risk of developing breast cancer.

The first aim of our study was to determine the frequencies of the SNP in LD with the *IGF1* microsatellite (rs5742612), two of the SNPs in the ECR (rs35765817 and rs35455143), and the nine individual htSNPs in the *IGF1* gene identified in the Swedish prostate cancer study within a population of women from Swedish breast cancer high-risk families. The second aim was to determine whether circulating IGF-1 levels were associated with the SNPs and diplotypes. The third aim was to determine diplotype frequencies in relation to *BRCA* status, *IGF1* CA-repeat genotypes and breast cancer risk.

Materials and Methods

Study Population

The study population consisted of 325 ethnic Swedish women from 194 breast cancer high-risk families, from the Lund Oncogenetic Clinic. The original population included 269 young healthy women with varying *BRCA* mutation status from 161 families. These women were 40 years or younger at the time of enrollment between 1996 and 2006. They were all menstruating, had no history of cancer, and no prophylactic mastectomy or bilateral oophorectomy. Blood samples, taken twice during the menstrual cycle on day 5-10 and again on day 18-23 (5-10 days before the predicted onset of the next menstrual period), were used for genetic and hormonal analyses. Women were asked to call back with the date of the first day of their next menstrual period. Plasma and blood cells were separated and frozen at minus 70 degrees Celsius at the laboratory of the Department of Oncology, Lund. Body measurements were taken at both visits when blood samples were collected. Each woman also filled out a questionnaire including questions on lifestyle and reproductive factors such as use of oral contraceptives. Written informed consent was obtained from all women. IGF-1 levels were analyzed in 258 women and 267 consented for genetic analyses. *BRCA1/2* mutation testing was not performed as a part of this study but data on mutation status was obtained from the Lund Oncogenetic Clinic.

In addition, 56 women (40 *BRCA1* and 16 *BRCA2* mutation carriers from the South Swedish Health Care Region born between 1950 and 1988), were included irrespective of cancer status and whether they had undergone prophylactic mastectomies or oophorectomies. No hormone analyses were performed on these blood samples but the women had consented to genetic analyses. Questionnaire data was available for some, but not all of these women.

The local ethics committee at Lund University approved the study.

Genotyping of IGF1 htSNPs

Genomic DNA Purification Kit (Promega, Madison, WI, USA) was used to extract DNA from 300 μ L of peripheral blood. All SNPs were genotyped at Region Skåne Competence Centre (RSKC Malmö), Malmö University Hospital, Malmö, Sweden. The SEQUENOM massARRAY® designer software was used for multiplex SNP analysis design. The analyses were performed on a matrix-assisted laser desorption/ionization time-of-flight mass spectrometry on a Sequenom MassARRAY® platform (Sequenom, San Diego, CA, USA) using iPLEX reagents according to the manufacturers' protocol. Three SNPs did not work well on iPLEX. The SNPs (rs855211, rs6220, and rs2033178) were genotyped using a Taqman SNP allelic discrimination assay in 384-well format on ABI PRISM 7900 Sequence Detection System (Applied Biosystems, Foster City, CA, USA). For quality control 40 samples were run in duplicate. The concordance was 100% for all SNPs. The call rates were 90% for rs855211 and over 97% for the remaining SNPs.

Diplotype construction

HtSNPs can be used to construct haplotypes and since we have two copies of *IGF1*, we constructed paired haplotypes called diplotypes. By crosstabulation of each htSNP in one block against the other SNPs in the same block, all possible linkage combinations were identified. Sometimes several combinations were equally likely, see figure 1. For statistical analyses, diplotypes with less than 10 carriers were grouped into a composite category of "rare diplotypes". In block 1 the "rare variant" diplotype consisted of 14 women with six or seven different diplotypes (six genotypes, one with two possible combinations). Block 2 "rare diplotypes" included 22 women with six or seven different diplotypes and block 3 included 29 women with six different diplotypes.

Hormone analysis

IGF-1 was analyzed in EDTA-plasma taken during cycle day 5-10 and day 18-23 from 258 women of the original study population. The analyses were performed at the Endocrinological Laboratory of Karolinska Hospital, Stockholm, Sweden, using a radioimmunoassay (RIA) method as previously described by Bang *et al.* [19]. IGF-1 was separated from IGFBPs by acid ethanol extraction followed by cryo-precipitation. Recombinant human IGF-1 (Kabi-Pharmacia, Stockholm, Sweden) was used as standard. The international IGF-1 standard was delivered by the National Institute of Biological Standards and Control, London, Great Britain. The intra assay variation was 4% and the inter assay variation was 11%. The limit of detection was 6 µg/L. IGF-1 levels were missing during cycle day 5-10 for one woman. For several of the OC users, the samples obtained during cycle day 5-10 were collected during their pill-free week.

Follow-up

Women were considered at risk for breast cancer from age 20 and were followed until the development of a first breast cancer according to the Regional Cancer Registry, until the date of a self-reported prophylactic mastectomy or oophorectomy, or until May 31 2009, whichever came first. The women in the study who were considered to have a high-risk of developing breast cancer were offered clinical follow-up including annual mammograms, ultrasounds or magnetic resonance imaging and physical examination of the breasts. The report rate of the Swedish cancer registries is close to 100 percent.

Data analyses

All statistical analyses were performed using the statistical software PASW 17.0 (formerly known as SPSS). Women who were currently breast feeding (n=4) or using hormonal contraceptives other than combined OCs (n=19), or both (n=1), were excluded from analyses of IGF-1 levels. Two women did not report the onset of their next menstrual period. Non-standardized and standardized means and 95% CIs were calculated for OC users and non-users combined as well as stratified according to current OC use. Non-standardized means for IGF-1 levels according to diplotypes were obtained using one-sample t-tests. For standardized levels, multivariate linear regression analyses were performed for each single SNP variant with the ancestral variant as reference and the hetero- and homozygous variant alleles coded as dummy variables in the analyses. Each diplotype was entered in separate multivariate analyses. IGF-1 levels were standardized at age 29 years, weight 67 kgs, day 7 or 7 days until the next period (i.e. day 21 in most women). When analyses of OC users and non-users were combined, the IGF-1 levels were standardized to non-use. The diplotype distribution among women from different *BRCA*-family categories was studied using Chi-square tests. Breast cancer free survival rates (from 20 years of age) in relation to different genotypes were assessed using Kaplan-Meier log-rank test and Cox regression models. Due to the large number of individual SNPs tested, we considered a *P*-value of <0.001 as statistically significant when analyzing IGF-1 levels in relation to individual SNPs. Nominal *P*-values are presented, All *P*-values are two-sided.

Results

Table 1 shows the characteristics of all 325 women and of the subgroup of 231 women included in the IGF-1 analyses. Genotype frequencies for each SNP, block compositions and frequencies of most likely diplotypes are shown in figure 1. Based on cross-tabulations, we found that the combination of rs2033178 CC, rs7136446 TC and rs6220 TC in block 3, most likely generates a mix of the two possible diplotypes CCC/CTT and CCT/CTC and we therefore present the genotypes as CC/CT/TC.

IGF-1 levels and IGF1 SNPs

Standardized levels of IGF-1 during cycle days 5-10 and 18-23 were calculated for all 231 women as well as for the 143 non-users and the 88 current OC users separately (table 2). IGF-1 levels did not differ significantly according to individual htSNPs or the three additional SNPs among all women, or when stratifying according to current OC status. No trend was observed for increasing or decreasing IGF-1 levels for any of the SNP variants.

IGF-1 levels and IGF1 diplotypes

We examined IGF-1 levels in relation to diplotypes in all women, current OC users and non-users during cycle days 5-10 and 18-23, figures 2a and 2b. Only the composite group of rare variant diplotypes in block 1 had lower levels during cycle day 5-10 compared with the women with other diplotypes in all women ($P=0.006$) and non-users ($P=0.03$), adjusted for age, weight, menstrual cycle day, and OC use. No association between rare variant diplotype and IGF-1 levels was seen in current OC users.

Diplotype distribution according to BRCA family status

The diplotype frequencies were examined according to *BRCA* family status for all women with available diplotypes belonging to either a family with a confirmed *BRCA1* or a *BRCA2* mutation, or to a family where no mutation has been identified ("BRCA_X"). Women from untested families were excluded. No significant differences in diplotype distribution was observed. Including only the first assigned woman from each family did not change the result.

Diplotype distribution according to BRCA mutation status

BRCA1 mutation carriers more often had a rare diplotype in block 1, OR 4.1 (95% CI 1.4 - 12.2; $P=0.007$) compared with other women. *BRCA1* mutation carriers had also more often a rare diplotype in block 2 compared with other women OR 2.8 (95% CI 1.1 - 6.8; $P=0.021$). Moreover, *BRCA1* mutation carriers had higher odds of having one of the variants of a htSNPs in block 1, rs2162679 GG, OR 11.2 (95% CI 1.2 - 109.8, $P=0.009$) than other women. Figures 3a, 3b, 3c.

IGF1 CA-repeats in relation to diplotypes

The *IGF1* CA-repeat allele was successfully genotyped for 260 women and diplotypes in block 1 were available for 232 of these. Women with a rare diplotype in block 1 were more likely than the other women to lack the common *IGF1* 19 CA-repeat allele OR 33.3 (95% CI 6.6-166.7; $P<0.001$). IGF-1 measurements were only available for two women with rare diplotypes in the group of current OC users.

The rs5742612 SNP was successfully genotyped for 259 women. In line with a previous study in Singapore Chinese women [11], we also found LD between the two polymorphisms. The 21 CA-repeat was found in 9% of the women with the rs5742612 TT genotype (n=235), 52%

of the women with the TC genotype (n=23) and the only woman with the CC genotype was homozygous for the 21 CA-repeat allele, (100%). This SNP was not associated with IGF-1 or breast cancer.

Diploypes in relation to breast cancer

After inclusion of the 56 additional *BRCA* mutation carriers, there were 23 women diagnosed with breast cancer. Having a rare variant diploype in block 1 was significantly associated with a higher risk of developing breast cancer (Log Rank $P < 0.001$), figure 4a. The number of women in this group was very small (n=14) and the rare variant diploype group consisted of several different diploypes. Four women with rare diploypes in block 1 had been diagnosed with breast cancer, and they had three different diploypes. No other diploype was associated with breast cancer risk.

Since women with rare diploypes in block 1 were more likely to have been diagnosed with breast cancer and also more likely to be *BRCA1* mutation carriers, the hazard analysis was also restricted to *BRCA1* mutation carriers, figure 4b. Having a rare diploype in block 1 was not only associated with a higher risk of developing breast cancer but was also associated with earlier diagnosis of a first breast cancer among *BRCA1* mutation carriers (Log Rank $P = 0.031$).

Discussion

Minor allele frequencies of the nine htSNPs were essentially in line with the findings of a previous study of a healthy Swedish male population [12]. Only the composite group of rare diplotypes in block 1 was associated with IGF-1 levels. Stratification on OC use did not indicate effect modification of OC use on IGF-1 levels for any of the diplotypes. The odds of having a rare diplotype in block 1 was higher in *BRCA1* mutation carriers and in women lacking the common *IGF1* 19 CA-repeat allele. The rare diplotype group of block 1 was associated with an increased hazard of a first breast cancer and with a younger age at diagnosis in *BRCA1* mutation carriers.

SNPs in relation to IGF-1 levels and cancer risk and risk factors

The SNPs in the present study were chosen based on previous studies by Johansson *et al.* who studied *IGF1* htSNPs in relation to prostate cancer and IGF-1 levels in Swedish populations. They found that the TCC haplotype of block 3 was associated with an increased risk for prostate cancer [12]. They also reported that the TCC haplotype, and two of the three htSNPs tagging this haplotype, were associated with increased levels of IGF-1 in men [20]. In the later study they did not study block 1 or 2 in relation to IGF-1 levels.

In the present study, none of the three SNPs included in the TCC haplotype was associated with IGF-1 levels. The group of women with the diplotype potentially containing the TCC haplotype had non-significantly higher IGF-1 levels when all women were included in the analysis, as well as when the analysis was restricted to women not currently using OCs. Major differences in age and in gender as well as potential differences in lifestyle and diet/nutrition status may account for the differing results between this study and Johansson's *et al.*'s study. A large study assessing some of the htSNPs included in the present study (rs5742612, rs1019731, rs7956547, and rs2033178) as well as a number of other SNPs in the

IGF1 gene found associations between diplotypes and circulating IGF-1 levels [21].

However, the study did not take into account age, BMI, OC use, parity, or cycle day of blood draw[22]. We measured IGF-1 levels in two blood samples from each woman and estimated values standardized for age, weight, and menstrual cycle day. We also stratified according to OC status. The stratifications resulted in smaller groups than would otherwise have been the case. A larger study population would be beneficial given that the current study already includes the majority of the healthy women from high-risk breast cancer families in the South Swedish health care region, a larger study population would have to include women from other areas of Sweden or other countries. IGF-1 levels measured in several blood samples during a menstrual cycle for each woman could have improved the estimates. Nutrition status influences circulating IGF-1 levels [23], but we lacked the information required to be able to adjust for this variable in the current study. However, fasting status at the time of blood draw did not affect the results (data not shown).

Rs1019731 has been associated with increased risk for ovarian cancer [24]. Another large study of IGF-1 polymorphisms, IGF-1 levels and breast cancer risk reported no association between this SNP and circulating IGF-1 levels or breast cancer risk [25]. We did not study ovarian cancer in the present study.

In one study of IGF-1 levels and breast density, the three htSNPs in block 3 (rs2033178, rs7136446 and rs6220) were shown to be associated with elevated IGF-1 levels, but the relationship with breast density was indecisive [26]. Another study of *IGF1* SNPs and breast density among postmenopausal women included a haplotype of rs6220 and rs2162679 that was associated with breast density but the association did not remain significant after stratification by use of hormone replacement therapy [27]. Tamimi *et al.* found *IGF1* SNPs

associated with mammographic density but only one of the SNPs in the current study was included (rs1019731) [28]. We did not study breast density in the present study.

Rs 2033178, rs1019731, rs2162679 and rs35765 (among others) were studied in relation to age at menarche in Caucasian women. These SNPs were not associated with age at menarche while others were [29].

Diplotypes

We studied the htSNPs combined as diplotypes since we believe that diplotypes better represent the genetic composition of each woman. The use of diplotypes results in smaller subgroups, since women without complete SNP genotyping are excluded and more combinations are possible than for individual SNPs. In the analyses of diplotypes in relation to IGF-1 levels, diplotypes with less than 10 carriers were grouped into the composite category of "rare diplotypes". Johansson *et al.* excluded such rare haplotypes from the analyses [12, 21], and we were therefore unable to compare our results with their data. We found that the rare block 1 diplotypes were associated with absence of the *IGF1* 19 CA-repeat allele, *BRCA1* mutation status and with an increased risk of developing early-onset breast cancer. These rare diplotypes could hypothetically be due to genotyping errors, but we do not consider such an explanation likely since the validation samples showed 100 percent concordance. Rather, these rare diplotypes may indicate genetic conditions in the *IGF1* gene that are evolutionary unfavorable, thus explaining why they are not very common. They may also tag for abnormalities in the *IGF1* gene that may contribute to the risk of developing breast cancer since four of the nine women included in the group of rare diplotypes of block 1 were diagnosed with breast cancer. Since our women were selected from families with high penetrance of breast cancer, it is likely that other low penetrance polymorphisms, associated

with breast cancer, segregate at a higher frequency in this population than in the general population [30]. However, as stated above, we were unable to compare our data to Johansson *et al.* [12, 21].

Absence of the *IGF1* 19 CA-repeat allele has been associated with a four-fold increased risk of developing breast cancer among women with a family history of breast cancer and even higher risk if the woman had ever used OCs [8]. The majority (>90%) of the women included in the present study had used or were current users of OCs. Therefore, we believe that the higher proportion of breast cancer cases among the women with rare diplotypes in block 1 may partly be driven by the lack of the *IGF1* 19 CA-repeat allele. A previous study based on results from a subset of the current cohort's population showed that women lacking the *IGF1* 19 CA-repeat allele had higher IGF-1 levels during OC use[7]. Since the block 1 rare variant group was associated with absence of 19 CA-repeat allele, and use of OCs among women with absence of the 19 CA-repeat allele is associated with increased breast cancer risk, the higher frequency of breast cancer in the women with rare block 1 diplotypes was not surprising. However, since IGF-1 measurements were only available for two women with rare diplotypes in the group of current OC users, we could not investigate the relationship between IGF-1 levels and block 1 rare variant diplotypes in OC users. We do not know whether the increased risk for earlier onset breast cancer was driven by the absence of the common 19 CA-repeat allele or by the rare variant diplotype. However, these rare *IGF1* genotype variants were more common in *BRCA1* carriers and confirmed an increased risk for very early-onset breast cancer among *BRCA1* mutation carriers. Since almost all women had used OCs at some point, we hypothesized that rare *IGF1* genotypes in combination with OCs are especially detrimental in high-risk women.

One study of *IGF1* htSNPs among *BRCA1* and *BRCA2* mutation carriers did not see any association between diplotypes and breast cancer [31]. However, Neuhausen *et al.*'s. study differs from the present study in a number of ways. Their htSNPs were not selected specifically for a Caucasian population even though they studied a Caucasian cohort and D'Alosio *et al.* has shown the importance of ethnic specific tag SNPs in *IGF1* [21]. Aside from SNP rs6220, Neuhausen *et al.* investigated different htSNPs. Moreover, they constructed haplotypes instead of diplotypes, which makes less biological sense, since *IGF1* is expressed on two alleles. In our study, rare diplotypes in block 1 and 2 were more common in *BRCA1* mutation carriers than in other women. We hypothesize that the rare diplotype variants may confer some form of survival advantage during fetal life among *BRCA1* mutation carriers, maybe through crosstalk between the genes[15].

In conclusion, neither the individual SNPs, nor the htSNPs combined as diplotypes were associated with IGF-1 levels in this study. These findings do not exclude the possibility that these SNPs or diplotypes maybe useful markers for breast cancer risk. A high proportion of women diagnosed with breast cancer had rare diplotype variants in block 1. Having a rare variant was also associated with two other genotypes associated with an increased risk of developing breast cancer, absence of the *IGF1* 19 CA-repeat allele and *BRCA1* mutation, and predicted early-onset breast cancer beyond the effect of a *BRCA1* mutation. These rare diplotypes may identify women with particularly high risk for early-onset breast cancer and data on rare variant diplotypes should be included in forthcoming studies. However, our study warrants confirmation in independent population.

Acknowledgments

We thank our research nurses Kerstin Nilsson, Monica Pehrsson, Anita Schmidt-Casslén for assistance with body measurements and blood drawing and Johanna Wagenius, Johanna Frenander, Helen Sundberg, Malin Sternby, and Susanna Holmquist for their assistance with recruitment. We also thank Dr. Eric T. Dryver for proofreading the manuscript.

This study was supported by grants from Vetenskapsrådet (the Swedish Research Council, K2001-27GX-14120-01A and K2008-68X20802-01-3), the Medical Faculty, Lund University, the Mrs. Berta Kamprad's Foundation, the Gunnar Nilsson Foundation, the Crafoord Foundation, the South Swedish Health Care Region (Region Skåne), Lund Hospital Fund and the Swedish Cancer Society.

References

1. Dickson RB, Russo J (2000) Biochemical Control of Breast Development. 2nd ed. Philadelphia, Williams and Wilkins
2. Stewart CE, Rotwein P (1996) Growth, differentiation, and survival: multiple physiological functions for insulin-like growth factors. *Physiol Rev* 76(4): 1005-26
3. LeRoith D, Roberts CT, Jr. (2003) The insulin-like growth factor system and cancer. *Cancer Lett* 195(2): 127-37 DOI S0304383503001599 [pii]
4. Toniolo P, Bruning PF, Akhmedkhanov A, et al. (2000) Serum insulin-like growth factor-I and breast cancer. *Int J Cancer* 88(5): 828-32
5. Hankinson SE, Willett WC, Colditz GA, et al. (1998) Circulating concentrations of insulin-like growth factor-I and risk of breast cancer. *Lancet* 351(9113): 1393-6
6. Harrela M, Koistinen H, Kaprio J, et al. (1996) Genetic and environmental components of interindividual variation in circulating levels of IGF-I, IGF-II, IGFBP-1, and IGFBP-3. *J Clin Invest* 98(11): 2612-5
7. Jernström H, Sandberg T, Bålgeman E, Borg Å, Olsson H (2005) Insulin-like growth factor-1 (IGF1) genotype predicts breast volume after pregnancy and hormonal contraception and is associated with circulating IGF-1 levels: implications for risk of early-onset breast cancer in young women from hereditary breast cancer families. *Br J Cancer* 92(5): 857-66
8. Cleveland RJ, Gammon MD, Edmiston SN, et al. (2006) IGF1 CA repeat polymorphisms, lifestyle factors and breast cancer risk in the Long Island Breast Cancer Study Project. *Carcinogenesis* 27(4): 758-65
9. Al-Zahrani A, Sandhu MS, Luben RN, et al. (2006) IGF1 and IGFBP3 tagging polymorphisms are associated with circulating levels of IGF1, IGFBP3 and risk of breast cancer. *Hum Mol Genet* 15(1): 1-10 DOI ddi398 [pii]
10.1093/hmg/ddi398
10. Palles C, Johnson N, Coupland B, et al. (2008) Identification of genetic variants that influence circulating IGF1 levels: a targeted search strategy. *Hum Mol Genet* 17(10): 1457-64 DOI ddn034 [pii]
10.1093/hmg/ddn034
11. Wong HL, Delellis K, Probst-Hensch N, et al. (2005) A new single nucleotide polymorphism in the insulin-like growth factor I regulatory region associates with colorectal cancer risk in singapore chinese. *Cancer Epidemiol Biomarkers Prev* 14(1): 144-51 DOI 14/1/144 [pii]
12. Johansson M, McKay JD, Stattin P, et al. (2007) Comprehensive evaluation of genetic variation in the IGF1 gene and risk of prostate cancer. *Int J Cancer* 120(3): 539-42 DOI 10.1002/ijc.22344
13. Shukla V, Coumoul X, Cao L, et al. (2006) Absence of the full-length breast cancer-associated gene-1 leads to increased expression of insulin-like growth factor signaling axis members. *Cancer Res* 66(14): 7151-7 DOI 66/14/7151 [pii]
10.1158/0008-5472.CAN-05-4570
14. Abramovitch S, Werner H (2003) Functional and physical interactions between BRCA1 and p53 in transcriptional regulation of the IGF-IR gene. *Horm Metab Res* 35(11-12): 758-62 DOI 10.1055/s-2004-814154
15. Maor S, Yosepovich A, Papa MZ, et al. (2007) Elevated insulin-like growth factor-I receptor (IGF-IR) levels in primary breast tumors associated with BRCA1 mutations. *Cancer Lett* 257(2): 236-43 DOI S0304-3835(07)00343-6 [pii]
10.1016/j.canlet.2007.07.019

16. Jernström H, Johannsson O, Borg Å, Ivarsson H, Olsson H (1998) BRCA1-positive patients are small for gestational age compared with their unaffected relatives. *Eur J Cancer* 34(3): 368-71
17. Henningson M, Bageman E, Sandberg T, Borg A, Olsson H, Jernstrom H (2007) Absence of the common IGF1 19 CA-repeat allele is more common among BRCA1 mutation carriers than among non-carriers from BRCA1 families. *Fam Cancer* 6(4): 445-52 DOI 10.1007/s10689-007-9141-0
18. Ford D, Easton DF, Stratton M, et al. (1998) Genetic heterogeneity and penetrance analysis of the BRCA1 and BRCA2 genes in breast cancer families. The Breast Cancer Linkage Consortium. *Am J Hum Genet* 62(3): 676-89
19. Bang P, Eriksson U, Sara V, Wivall IL, Hall K (1991) Comparison of acid ethanol extraction and acid gel filtration prior to IGF-I and IGF-II radioimmunoassays: improvement of determinations in acid ethanol extracts by the use of truncated IGF-I as radioligand. *Acta Endocrinol (Copenh)* 124(6): 620-9
20. Johansson M, McKay JD, Wiklund F, et al. (2007) Implications for prostate cancer of insulin-like growth factor-I (IGF-I) genetic variation and circulating IGF-I levels. *J Clin Endocrinol Metab* 92(12): 4820-6 DOI jc.2007-0887 [pii] 10.1210/jc.2007-0887
21. D'Aloisio AA, Schroeder JC, North KE, et al. (2009) IGF-I and IGFBP-3 polymorphisms in relation to circulating levels among African American and Caucasian women. *Cancer Epidemiol Biomarkers Prev* 18(3): 954-66 DOI 1055-9965.EPI-08-0856 [pii] 10.1158/1055-9965.EPI-08-0856
22. Jernstrom H (2009) IGF1 and IGFBP3 polymorphisms and plasma levels in women. *Cancer Epidemiol Biomarkers Prev* 18(10): 2794; author reply 5 DOI 18/10/2794 [pii] 10.1158/1055-9965.EPI-09-0324
23. Norat T, Dossus L, Rinaldi S, et al. (2007) Diet, serum insulin-like growth factor-I and IGF-binding protein-3 in European women. *Eur J Clin Nutr* 61(1): 91-8 DOI 1602494 [pii] 10.1038/sj.ejcn.1602494
24. Terry KL, Tworoger SS, Gates MA, Cramer DW, Hankinson SE (2009) Common genetic variation in IGF1, IGFBP1 and IGFBP3 and ovarian cancer risk. *Carcinogenesis* 30(12): 2042-6 DOI bgp257 [pii] 10.1093/carcin/bgp257
25. Patel AV, Cheng I, Canzian F, et al. (2008) IGF-1, IGFBP-1, and IGFBP-3 polymorphisms predict circulating IGF levels but not breast cancer risk: findings from the Breast and Prostate Cancer Cohort Consortium (BPC3). *PLoS One* 3(7): e2578 DOI 10.1371/journal.pone.0002578
26. Verheus M, McKay JD, Kaaks R, et al. (2008) Common genetic variation in the IGF-1 gene, serum IGF-I levels and breast density. *Breast Cancer Res Treat* 112(1): 109-22 DOI 10.1007/s10549-007-9827-x
27. Biong M, Gram IT, Brill I, et al. Genotypes and haplotypes in the insulin-like growth factors, their receptors and binding proteins in relation to plasma metabolic levels and mammographic density. *BMC Med Genomics* 3: 9 DOI 1755-8794-3-9 [pii] 10.1186/1755-8794-3-9
28. Tamimi RM, Cox DG, Kraft P, et al. (2007) Common genetic variation in IGF1, IGFBP-1, and IGFBP-3 in relation to mammographic density: a cross-sectional study. *Breast Cancer Res* 9(1): R18 DOI bcr1655 [pii] 10.1186/bcr1655

29. Zhao J, Xiong DH, Guo Y, Yang TL, Recker RR, Deng HW (2007) Polymorphism in the insulin-like growth factor 1 gene is associated with age at menarche in caucasian females. Hum Reprod 22(6): 1789-94 DOI dem052 [pii]
10.1093/humrep/dem052
30. Easton DF, Ford D, Bishop DT (1995) Breast and ovarian cancer incidence in BRCA1-mutation carriers. Breast Cancer Linkage Consortium. Am J Hum Genet 56(1): 265-71
31. Neuhausen SL, Brummel S, Ding YC, et al. (2009) Genetic variation in insulin-like growth factor signaling genes and breast cancer risk among BRCA1 and BRCA2 carriers. Breast Cancer Res 11(5): R76 DOI bcr2414 [pii]
10.1186/bcr2414

Figure legends

Figure 1. The SNP distribution in the *IGF1* gene and allele and diplotype frequencies. Genotypes and minor allele frequencies (MAF) are presented for each htSNP. Possible combinations of the htSNPs are listed together with the number of carriers and the diplotype(s) that can result from each block combination. A diplotype seen in less than 10 women was grouped into a composite group of rare diplotypes.

Figure 2 Non-standardized and standardized IGF-1 levels, days 5-10 (a) and days 18-23 (b), according to diplotypes in all blocks, in all women (n=231), in non-users of OC (n=143), and in current OC users (n=88). The y-axis cuts the x-axis at the mean level of each of the three groups, the broken line represents the mean of the 231 women. The shaded span around the y-axis represents the standard error of the mean (SEM). Diamonds represents mean levels for each genotype group and horizontal lines represents SEM. Computed values standardized at age 29 years, 67 ksg and cycle day 7 (a) or 21 (b) are indicated by smaller dots.

Figure 3 *IGF1* diplotype distribution for block 1 (a), block 2 (b) and block 3 (c) in relation to *BRCA* mutation status (*BRCA1* mutation carriers (*BRCA1+*) *BRCA1* non-carriers (*BRCA-*), *BRCA2* mutation carriers (*BRCA2+*) *BRCA2* non carriers (*BRCA2-*), and women where no *BRCA1/2* mutation had been found in the family (*BRCAX*). The number of women included in each *BRCA* group is indicated under the x-axis. Each diplotype group is represented as their percentage of contribution to the bar for women in the respective *BRCA* group.

Figure 4 Influence of rare *IGF1* diplotypes in block 1 on cumulative hazard of a first breast cancer among a) all women and (Log rank $P < 0.001$) b) *BRCA1* mutation carriers (Log Rank $P = 0.031$). The number of women at each follow-up time is indicated below the graphs.

Table 1. Characteristics for all women and for the women included in the analyses of IGF-1 levels

	All women		Included in IGF-1 analyses	
	Median (IQR) ¹ , % or n	n	Median (IQR) ¹ , % or n	n
Medians (IQR)				
Age (years)	30 (25-36)	325	29 (24-35)	231
Year of birth	1969 (1964-1975)	291	1969 (1964-1976)	231
Birthweight (g)	3410 (3070-3720)	291	3405 (3070-3700)	224
Weight (kg)	64 (58-74)	269	64 (58-73)	231
Height (cm)	168 (164-172)	269	168 (164-172)	231
BMI (kg/m ²)	22.8 (20.0-25.6)	269	22.6 (20.8-25.1) (25.2/233)	231
Waist-to-hip ratio	0.76 (0.73-0.80)	269	0.76 (0.73-0.80)	231
Age at menarche (years)	13 (12-14)	312	13 (12-14)	230
Percentages (%)				
Parous	54.4	305	47.2	231
Ever use of oral contraception	92.2	246	91.3	231
Current use of combined oral contraception	34.2	269	37.7	231 (233)
Current use of progestin-only pills	4.5	269	0	231
Current use of other hormonal contraception or IUD*	3.3	269	0	231
Ever use of hormone replacement therapy	5.4	268	0	231
Ever smoker	41.8	268	43.1	232
Current smoker	22.8	269	23.3	232
Number of women (n)				
BRCA1 family	134	325	76	233
Mutation carrier	70		23	
Negative	49		44	
Untested	15		9	
BRCA2 family	34	325	14	233
Mutation carrier	23		7	
Negative	7		5	
Untested	4		2	
BRCAX family	110	325	102	233
Untested family	45	325	41	233

* Intra Uterine Device

Table 2. Standardized IGF-1 levels in relation to *IGF1* SNP genotypes

Menstrual cycle days 5-10										Menstrual cycle days 18-23											
		All women			No OC use			Current OC					All women			No OC use			Current OC		
SNP	Genotype	N	IGF-1 (ug/L)	P-value	N	IGF-1 (ug/L)	P-value	N	IGF-1 (ug/L)	P-value	N	IGF-1 (ug/L)	P-value	N	IGF-1 (ug/L)	P-value	N	IGF-1 (ug/L)	P-value		
Block 1																					
rs855211	GG	149	212		93	209		56	215		149	214		93	211		56	194			
	GA	53	224	0,211	36	233	0,035	17	209	0,718	53	231	0,049	36	241	0,008	17	184	0,334		
	AA	5	158	0,048	5	162	0,079	0			5	196	0,447	5	201	0,69	0				
rs35765	CC	188	212		114	209		74	212		188	216		114	215		74	183			
	CA	43	209	0,754	30	215	0,608	13	195	0,348	43	219	0,761	30	227	0,313	13	170	0,33		
	AA	0			0			0			0			0			0				
rs2162679	AA	174	213		106	209		68	215		174	217		106	215		68	186			
	AG	53	210	0,732	35	220	0,356	18	192	0,157	53	216	0,851	35	227	0,281	18	164	0,05		
	GG	3	157	0,109	3	160	0,15	0			3	217	0,988	3	221	0,846	0				
Block 2																					
rs1019731	CC	168	209		104	210		64	202		168	214		104	215		64	175			
	CA	53	216	0,47	35	205	0,622	18	231	0,083	53	225	0,199	35	221	0,62	18	196	0,064		
	AA	9	229	0,322	5	251	0,124	4	193	0,781	9	233	0,28	5	245	0,271	4	176	0,979		
rs7956547	AA	127	211		79	211		48	205		127	220		79	221		48	181			
	AG	77	216	0,529	48	212	0,974	29	224	0,183	77	214	0,462	48	214	0,496	29	181	0,95		
	GG	26	201	0,456	17	203	0,603	9	193	0,599	26	212	0,503	17	212	0,558	9	175	0,701		
rs5742632	AA	129	211		82	212		47	206		129	220		82	222		47	182			
	AG	77	212	0,896	47	210	0,857	30	214	0,548	77	212	0,272	47	211	0,324	30	177	0,593		
	GG	24	209	0,902	15	208	0,816	9	209	0,883	24	216	0,735	15	216	0,749	9	180	0,883		
Block 3																					
rs2033178	CC	197	211		124	208		73	215		197	216		124	215		73	182			
	CT	30	211	0,995	20	227	0,189	10	182	0,123	30	222	0,531	20	233	0,2	10	169	0,366		
	TT	0			0			0			0			0			0				
rs7136446	TT	85	215		52	216		33	214		85	221		52	226		33	181			
	TC	102	213	0,816	65	209	0,507	37	216	0,897	102	216	0,517	65	213	0,228	37	186	0,611		
	CC	38	197	0,129	23	198	0,228	15	195	0,32	38	210	0,32	23	213	0,369	15	172	0,484		
rs6220	TT	113	210		71	209		42	210		113	219		71	219		42	185			
	TC	88	218	0,393	56	218	0,344	32	215	0,739	88	217	0,862	56	219	0,988	32	180	0,621		
	CC	24	202	0,517	14	202	0,713	10	203	0,736	24	214	0,705	14	219	0,959	10	172	0,395		
Block 4																					
rs5742612	TT	210	211		129	211		81	209		210	216		129	217		81	182			
	CT	19	214	0,854	14	214	0,827	5	212	0,914	19	219	0,802	14	225	0,63	5	170	0,533		
	CC	1	222	0,853	1	235	0,68	0			1	322	0,046	1	328	0,057	0				
rs35765817	CC	201	211		126	210		75	210		201	214		126	216		75	179			
	CT	28	213	0,841	16	213	0,844	12	202	0,677	28	229	0,18	16	227	0,447	12	190	0,399		
	TT	2	240	0,486	2	241	0,464	0			2	274	0,113	2	274	0,162	0				
rs35455143	CC	208	211		130	210		78	210		208	215		130	216		78	179			
	CT	21	211	0,983	12	213	0,883	9	201	0,688	21	232	0,151	12	232	0,355	9	193	0,346		
	TT	2	240	0,49	2	241	0,465	0			2	274	0,113	2	274	0,162	0				

Figure 1.

Figure 2a. IGF-1 levels during menstrual cycle days 5-10

Figure 2b. IGF-1 levels during menstrual cycle days 18-23

Figure 3. IGF1 diplotype distribution in relation to BRCA status

3b.

3c.

a) All women

b) BRCA1 mutation carriers

Rare IGF1 diplotype in block 1

Yes, N=14	13	2	0	0	N=7	6	1	0	0
No, N=275	231	117	13	0	N=54	45	30	4	0