



# LUND UNIVERSITY

## Vitronectin in host pathogen interactions and antimicrobial therapeutic applications

Singh, Birendra; Su, Yu-Ching; Riesbeck, Kristian

*Published in:*

Central European Journal of Biology

*DOI:*

[10.2478/s11535-011-0077-x](https://doi.org/10.2478/s11535-011-0077-x)

2011

[Link to publication](#)

*Citation for published version (APA):*

Singh, B., Su, Y.-C., & Riesbeck, K. (2011). Vitronectin in host pathogen interactions and antimicrobial therapeutic applications. *Central European Journal of Biology*, 6(6), 973-980. <https://doi.org/10.2478/s11535-011-0077-x>

*Total number of authors:*

3

### General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117  
221 00 Lund  
+46 46-222 00 00

**Review:**

**Vitronectin in host pathogen interactions and antimicrobial therapeutic applications**

Birendra Singh, Yu-Ching Su and Kristian Riesbeck

*Medical Microbiology, Department of Laboratory Medicine Malmö, Lund University, Skåne University Hospital, SE-205 02 Malmö, Sweden*

**Corresponding author:**

Kristian Riesbeck  
Medical Microbiology, Department of Laboratory Medicine Malmö  
Lund University, Skåne University Hospital  
SE-205 02 Malmö, Sweden.  
Tel.: +46 40 331340; fax: +46 40 336234.  
E-mail address: kristian.riesbeck@med.lu.se

**Running title:** Vitronectin and bacterial pathogens

**Key words:** bacteria, complement system, extracellular matrix, innate immunity, integrins, vitronectin

## Footnotes

- (1) Potential conflicts of interest: non reported.
  
- (2) Financial support: This work was supported by grants from the Alfred Österlund, the Anna and Edwin Berger, the Marianne and Marcus Wallenberg, the Greta and Johan Kock Foundations, the Swedish Medical Research Council, the Cancer Foundation at the University Hospital in Malmö, and Skåne county council´s research and development foundation.
  
- (3) Reprints or correspondence: Dr. Kristian Riesbeck, Medical Microbiology, Department of Laboratory Medicine Malmö, Lund University, Skåne University Hospital, SE-205 02 Malmö, Sweden (kristian.riesbeck@med.lu.se).

## **Summary**

Vitronectin (Vn) is a multifunctional glycoprotein profusely present in serum and bound to epithelial cell surfaces. It plays an important role in cell migration, tissue repair and regulation of membrane attack complex (MAC) formation. In the last decade the role of Vn has been extensively investigated in eukaryotic signalling and cell migration leading to the possibility of developing novel anticancer drugs. In parallel, several studies have suggested that pathogens utilize Vn in invasion of the host. Here we review the properties of Vn and its role in host-pathogen interactions that might be a future target for therapeutical intervention.

## 1. Introduction

Vitronectin (Vn) is a multifunctional glycoprotein that ensures optimum homeostasis in the body by regulating angiogenesis, cell migration and tissue repair [1]. However, it also plays an important role in tumor growth and metastasis [2], atherosclerotic disease [3] and viral infections [4]. Vn is known as an effective regulator of the complement system that limits the self-reactivity of the membrane attack complex (MAC) in the host. There are also other complement regulators, such as the classical complement pathway regulator C4b binding protein (C4BP) that binds to C4b followed by degradation of C4b with the help of factor I, and finally degrades C3 convertase. Additionally, factor H (FH) inhibits the activation of the alternative complement activation pathway by binding to factor C3b and inactivating it in the presence of factor I [65]. Activation of the complement system by the classical, alternative, or lectin-mediated pathways leads to a common lytic pathway after C5b activation. Subsequently, a C5b-C7 complex is formed and immobilized on the bacterial surface by C8. The C5b-C8 complex is recognized by C9 and results in C9 polymerization to form a cytolytic pore. Vn inhibits the association of C5b-C7, and surface-bound Vn may bind the C5b-7 complex that consequently does not associate with C8 for the initiation of the MAC. In addition, Vn bound to the bacterial surface may also remain active to interact with C9 and inhibit C9 polymerization.

The major complement regulators described above are frequently recruited by bacterial pathogens at their surfaces in order to circumvent the host innate immune response [5]. Since the first discovery of an interaction between Vn and pathogens by Chhatwal and co-workers in 1987, the role of Vn in microbial pathogenesis has been widely appreciated and investigated [6]. Of note, a large body of studies has suggested the pivotal role of heparin binding domains (HBDs) of Vn in the interaction with pathogens. Three distinguished HBDs have been defined on the Vn

molecule, residing at the amino acids 82-137 (HBD-1), 175-219 (HBD-2), and finally 348-361 (HBD-3). The most C-terminal one (HBD-3) has been characterized as the major binding site recognised by pathogens [7] as illustrated in Figure 1.

Vitronectin initiates cellular signaling by interacting with integrins ( $\alpha 3\beta 1$ ,  $\alpha v\beta 1$ ,  $\alpha v\beta 3$ ,  $\alpha v\beta 5$ , and  $\alpha IIb\beta 3$ ) at their N-terminal region containing the amino acids RGD. Binding of fibronectin and Vn induces integrin clustering and recruitment of host proteins involved in the signalling cascade, resulting in actin rearrangement and internalization of the Vn-integrin complex. Pathogens therefore bind Vn and manipulate the interaction for adherence on epithelial cells as well as for internalization into host cells [8, 9]. The role of Vn in bacterial pathogenesis has been comprehensively described in our recent publication [7]. In the present review, we highlight the molecular aspects of Vn in host-pathogen interactions and further conclude its therapeutic application as an antimicrobial agent.

## **2. Molecular forms of vitronectin**

Vn exists in two isoforms with molecular weights of 75 kDa and 65 kDa. The circulating form of Vn is usually present as a monomer (native Vn) that represents approximately 93% of the total Vn content in human plasma. The remaining 2 to 7% is activated Vn and undergoes a unique conformational change resulting in a multimeric form [10]. It was initially believed that native Vn does not have a heparin-binding capacity due to the cryptic HBD in native Vn [11]. This is in bright contrast to other ECM proteins, *e.g.*, fibronectin that has a heparin-binding capacity even at their native conformations. However, Zhuang and co-workers (1997) discovered that monomeric Vn can also bind heparin *in vitro*, while increased binding was observed with the activated form. Moreover, the heparin-binding property of multimeric Vn involved ionic

interactions [12]. The most commonly protocol used in the purification of Vn from serum is based upon denaturing conditions in the presence of urea. Exposure to urea and subsequent renaturation cause multimeric association of activated Vn, resulting in multivalent of heparin-binding sites [13].

Native (monomeric) Vn is involved in the regulation of plasminogen activation by interacting with plasminogen activator inhibitor-1 (PAI-1) and urokinase plasminogen activator (uPA) / uPA receptor (uPAR). This interaction leads to Vn multimerization that in turn increases Vn efficiency to bind with the surface receptor of epithelial cells and promote other eukaryotic functions [14, 15]. Of note, only multimeric Vn can bind to the epithelial cell surfaces and induce cell signalling [10, 13], and in some cases the glycoprotein is also endocytosed intracellularly. PAI-1 and heparin both promote multimerization of the Vn molecule, but their mechanism of interaction and impact on the production of multimeric Vn is different [16]. Conformational transitions of the Vn molecules as described above are solely concluded from *in vitro* studies. In contrast, the generation of multimeric Vn from monomers is only partially known *in vivo*. It thus remains to investigate whether activated Vn occurs under continuous exposure to its ligands, such as PAI-1 and glycosaminoglycans, or whether the *in vitro* induced multimerization of Vn resembles the *in vivo* conditions.

In the complement system, Vn regulates the self-reactivity of the MAC by inhibiting the C5b-7 complex assembly and C9 polymerization. It is, however, an enigma whether the regulation of the MAC is conducted by a specific conformation of Vn. In some experimental models it has been shown that injection of [<sup>32</sup>P]-labelled monomeric Vn into mice leads to incorporation of radioactive Vn into a high molecular weight soluble (S) C5b-9 complex, whereas injection of urea-activated [<sup>32</sup>P]-labelled polymeric Vn is rapidly cleared from serum

[17]. In general it is thus believed that monomeric Vn performs its MAC inhibitory role by making a soluble Vn/ C5b-7/ C5b-8/C5b9 complex. These complexes are lytically inactive, unable to be inserted into the cell membrane, and finally cleared from the circulation (Sodetz & Plumb, 1998). The role of polymeric Vn in the regulation of MAC is yet to be defined. On the other hand, it has been observed that only polymeric (activated) Vn is importantly associated with bacterial pathogenesis during adhesion to host epithelial cells. Recently, it was observed that adhesion and internalization of *S. pneumoniae* to lung epithelial cells is several-fold increased in the presence of polymeric Vn compared to native Vn [8]. More recently, similar observations have been noticed with *Neisseria meningitidis*; only activated Vn enhanced bacterial adhesion and internalization of HBMEC cells [9]. These studies suggested that the polymeric Vn has a suitable conformation that makes a bridge between epithelial cells and the pathogens. Thus, the polymeric form of Vn that consists of the minor fraction of total Vn (2 to 7%) might be a limiting factor for many pathogens to acquire Vn-mediated serum resistance or adherence. Hence, accurate estimation of the availability of Vn to pathogens during infection can only be achieved by using *in vivo* models.

### **3. The PAI-1/Vn complex and bacterial pathogenesis**

Vn is unique among other extracellular matrix (ECM) proteins, considering its role as an effective regulator of the fibrinolytic system. This is achieved by binding of Vn to plasminogen activator inhibitor (PAI-1). The PAI-1/Vn complex inhibits the plasminogen activity by inactivating tissue plasminogen activator (tPA) and urokinase plasminogen activator (uPA) [18]. Vn increases the half life of PAI-1 by 2- to 4-fold, thus prolongs inhibitory activity of PAI-1 on the activation of plasminogen [19]. It has been shown in the mouse model that PAI-1 expression



is necessary to protect the host from Gram-negative pathogens by recruiting neutrophils to the alveolar compartment. PAI-1 knockout animals are prone to infection by *Klebsiella pneumoniae*, and restoring the expression of PAI-1 in these knockout models generates resistance against *K. pneumoniae* infection [20]. Similarly, the role of PAI-1 in bacterial pathogenesis is also supported by the increased level of the plasminogen inhibitor during sepsis and other severe pathological conditions [21-23]. Some bacterial pathogens such as *Pseudomonas aeruginosa*, *Staphylococcus aureus*, and *Bacillus subtilis* cause degradation of PAI-1 during invasion [24-26]. Bacterial proteases also degrade Vn during invasion [27, 28], but there is no information available on degradation of the PAI-1/Vn complex. Recently it was shown that proteins of the omptins family (bacterial surface proteases) including plasminogen activator (Pla) of *Yersinia pestis*, PgtE of *Salmonella typhimurium*, and Kop of *K. pneumoniae* can degrade the PAI-1/Vn complex *in vitro* [19]. It has been hypothesized that abolished regulation of the tPA/uPA activity is attributed to the pathogen-dependent degradation of PAI-1/Vn complex and eventually lead to the uncontrolled fibrinolysis. Thus, a high production of plasmin will accelerate tissue damage that would be beneficial to the pathogens during invasion (Fig. 1).

#### **4. Interactions between bacterial pathogens and vitronectin**

The well studied complement regulators (C4BP, FH and Vn) are frequently recruited by bacterial pathogens to circumvent the host innate immune response[29]. During the bacterial infection only one or all complement pathways will be activated, depending on the surface molecule of the infecting pathogen. In contrast to the FH and C4BP, Vn inhibits the lytic (terminal) pathway that is ultimately common amongst all the complement pathways. Therefore, recruitment of Vn at the

bacterial surface might thus be a more effective strategy and important compared to the other two complement regulators.

There are several examples of successful utilization of vitronectin by various pathogens. The MAC complex is, however, more effective in killing of Gram-negative bacteria. This is in bright contrast to Gram-positive pathogens that are shielded against complement due to a thick peptidoglycan layer. Both groups of pathogens have, however, the capacity to bind Vn and utilize the molecule for inhibition of the MAC and adhesion to host tissues. The Gram-negative *H. influenzae*, for example, recruits Vn by the surface exposed adhesin protein E (PE), and *Haemophilus* surface fibrils (Hsf) [30-32], whereas *H. ducreyi* binds to Vn by the protein DsrA [33]. On the other hand, *Moraxella catarrhalis* interacts with Vn by ubiquitous surface protein (Usp) A2 [7, 34, 35]. *N. meningitidis* [9], and *N. gonorrhoeae* [36] also acquire serum resistance by attracting Vn. Several other Gram negative bacterial pathogens are known to bind Vn, but the importance of these interactions in serum resistance have not been explored in detail [35].

Similarly, Gram-positive pathogens, *e.g.*, *S. pneumoniae*, *S. pyogenes*, *S. bovis*, *S. suis*, *S. dysgalactiae*, *Staphylococcus epidermidis*, *S. aureus* and *Enterococcus faecalis* bind Vn and utilize the protein for adhesion to host epithelial surfaces. Details of these pathogens and their interactions with Vn have recently been reviewed [7]. The inhibitory capacity of Vn regarding the MAC and Gram-positive pathogens have not been reported until now. However, the role of Vn-dependent bacterial adhesion and internalization into host cells has been thoroughly explored. It is well known that several invasive pathogens utilize integrin-mediated internalization by using host phagocytic mechanisms [37]. Vn binds to epithelial cell surface integrin receptors ( $\alpha\beta3$ ,  $\alpha\beta1$ ,  $\alpha3\beta$ ,  $1\alpha\text{IIb}\beta3$  and  $\alpha\beta5$ ) via RGD-binding motifs, where vitronectin and integrin receptors interacts to induce signalling by multiple pathways [35]. In

conclusion, during adhesion bacteria will only successfully cross-link Vn that has the ability to bind epithelial surface receptors. Similarly, host epithelial surface-bound Vn will only be recognised by bacteria if the pathogen binding sites on the Vn molecule are available for interaction.

## **5. Vitronectin and future therapeutic applications**

The biological functions of Vn are associated with its conformation either in an unfolded (activated) or folded (native) form. However, activated Vn is highly associated with bacterial pathogenesis during infection as well as in tumor metastasis, which is mainly attributed to its frequent distribution in ECM or on epithelial cell surfaces, a preferable platform for both bacterial colonization and cancer cell invasion. Microbes bind Vn and utilize the molecule to establish infection such as to subvert host complement killing, adherence to host epithelial cells followed by internalization through Vn-integrin signaling [7]. Most of pathogenic bacteria bind to activate Vn, particularly either at the N-terminal polyanionic region (between amino acids 43-68) [9], the central domain containing hemopexin like domains (comprising residues 131-323) [38], or at the basic carboxy terminal HBD-3 [7]. The bacterial-Vn binding usually does not interfere with the Vn domain involved in the inhibition of C5b-9 complex (internal region between amino acid 51-310), thus allowing the complement regulator to remain active in inhibiting MAC formation while binding on the bacterial surface [30, 39, 40]. HBD-3 was proposed to be involved in the inhibition of the MAC [41]. However, the inhibitory domain was later relocated as described above [39, 40].

Interestingly, antimicrobial peptides (AMP) including LL37 and  $\alpha$ -defensin share similar heparin binding motifs as seen in HBD-3 [42, 43]. These AMPs lose their bactericidal properties

when interacting with glycosaminoglycans. Several cationic peptides representing the heparin binding domain of fibronectin, various laminin isoforms, von Willebrand factor, Protein C inhibitor as well as C3 show antimicrobial effects [44, 45]. HBD-3 of Vn which consists of Cardin motifs (XBBXBX, where X represents hydrophobic or non-polar amino acids and B represents basic amino acids) [46, 47] was bactericidal towards *Enterococcus faecalis* at concentrations between 0.6 to 3  $\mu\text{M}$ , a killing effect which is comparable to the classical LL37-defensin. At low salt conditions in a radial diffusion assay, HBD-3 (100 $\mu\text{M}$ ) showed a more potent antibacterial activity to *Escherichia coli*, *Candida albicans*, *P. aeruginosa* and *Proteus mirabilis* as compared to the defensin LL37. However, the actual mode of action of antibacterial activity of the HBD-3 remains unknown, instead more work is needed to expand the understanding of the utilization of Vn as an AMP. To date, reports concerning the exploration of Vn peptides as antimicrobial agents are still limited.

Bacterial colonization on commonly used biomaterial implants and medical devices causes serious complications. For example, implant-associated infections have caused a high mortality among patients with heart implants, and serious disabilities in patients with orthopaedic devices [48, 49]. Relative immune system fitness of the implant recipient and virulence of microorganisms such as the capacity to form biofilm are the main factors contributing to infections on implant surfaces. When a biomaterial is implanted and exposed to blood, a reservoir of host proteins including fibrinogen, Vn, fibronectin, albumin and immunoglobulins usually tends to adsorb to its surface. Moreover, Vn and fibronectin adsorbed on foreign surfaces unfold and activate glycoproteins, and potentially increases interactions with their ligands and bacterial specific receptors [50, 51]. Thus, the thin layer of host proteins would favour the interaction of microbes with the proteinous film, and the microbes would consequently use

specific receptors through hydrophobic interactions or Van der Waals and electrostatic charges, facilitating the formation of biofilm. In the presence of cerebrospinal fluid (CSF), Vn was found to mediate adherence of coagulase-negative staphylococci on the polyvinylchloride (PVC), a material used in prosthetic devices [52]. The biofilm on biomaterial surfaces is usually resistant to antibiotic killing and infection can often only be eradicated by the removal of the infected biomaterial. Therefore, several strategies have been developed to control the biofilm related to implant-associated infections, such as i) by using non-protein absorptive surfaces[53, 54], ii) pre-colonization of surfaces with non-pathogenic bacteria for nich masking [55], iii) coating of antibiotics [53, 56, 57], and iv) surfaces containing biocidal substances [58, 59]. Providing the advantages of the stability *in vivo* of AMP and lack of resistance development, pre-coating biomaterial surfaces with antimicrobial agents including HBD-3 of Vn would be an effective effort to prevent biofilm formation, and would in the long run reduce survival of microorganisms. Several antimicrobial coating strategies have recently been reviewed in detail [60]. In addition, it is possible to reduce bacterial adherence to biomaterial surfaces by targeting activated Vn with specific antibodies to the biomaterial surface after immediate exposure to body fluids containing Vn [61, 62]. A Vn-binding *S. epidermidis* showed reduced binding to PVC of prosthetic devices when the PVC priorly exposed to CSF was incubated with anti-Vn antibodies [52, 63].

In contrast to AMP derived from Vn, the full length Vn molecule has not been shown to exert any bactericidal effect, instead Vn exhibits a role as a pro-bacterial pathogenesis molecule. However, Vn is a potent inhibitor of complement activation that is prone to occur at the biomaterial surface in implant devices that are in contact with blood and aqueous humor [64, 65]. Complement activation in the vicinity of an implant results in unspecific inflammation and

attachment of thrombocytes, bacteria and fungi. Intriguingly, the complement inhibitory role of Vn at the biomaterial surface is faded by the presence of Vn-binding microorganisms [66, 67]. Binding of the Vn-binding *Staphylococcus hemolyticus* to surfaces precoated with the glycoprotein results in hindrance of Vn from inhibiting complement activation. Conversely, interaction of non-Vn binding strains of *S. epidermidis* with surface-coated Vn does not affect the complement-inhibitory role of Vn.

## 6. Conclusion

Vn functions as one of the major host factors that regulate multiple biological processes. The regulatory mechanisms by Vn in cellular migration have been targeted for the development of anticancer therapeutics. In addition, this glycoprotein has also been recognized as a biomarker molecule in pathophysiological processes. The adhesive and MAC inhibitory properties of Vn is utilized by pathogens for adherence to host cells and protection against the host immune system. Moreover, some bacterial pathogens have also manipulated the host signalling pathway (Vn-integrin mediated) for internalization and invasion. In a nutshell, the Vn-host interaction could be targeted for the development of novel antimicrobial agents. In addition, heparin-binding Vn peptides can be used as antibacterial agents on implant devices or use in antiseptic formulations. The implementation of Vn as an antimicrobial therapeutic agent will thus shed light on alternative drugs to overcome the issue of antibiotic resistance. Such development requires careful consideration and configuration based on targeted anatomical sites, taking into account the multi-essential role of Vn for host physiological needs.

## 7. References:

- [1] Preissner, K.T. and Seiffert, D., Role of vitronectin and its receptors in haemostasis and vascular remodeling, *Thromb. Res.*, 1998, 89, 1-21
- [2] Yeh, W.L., Lu, D.Y., Liou, H.C. and Fu, W.M., A forward loop between glioma and microglia: Glioma-derived extracellular matrix-activated microglia secrete IL-18 to enhance the migration of glioma cells, *J. Cell Physiol.*, 2011, DOI: 10.1002/jcp.22746.
- [3] Ekmekci, O.B. and Ekmekci, H., Vitronectin in atherosclerotic disease, *Clin. Chim. Acta.*, 2006, 368, 77-83
- [4] Xiao, J., Natarajan, K., Rajala, M.S., Astley, R.A., Ramadan, R.T. and Chodosh, J., Vitronectin: a possible determinant of adenovirus type 19 tropism for human corneal epithelium, *Am. J. Ophthalmol.*, 2005, 140, 363-369
- [5] Zipfel, P.F. and Skerka, C., Complement regulators and inhibitory proteins, *Nat. Rev. Immunol.*, 2009, 9, 729-740
- [6] Chhatwal, G.S., Preissner, K.T., Muller-Berghaus, G. and Blobel, H., Specific binding of the human S protein (vitronectin) to streptococci, *Staphylococcus aureus*, and *Escherichia coli*, *Infect. Immun.*, 1987, 55, 1878-1883
- [7] Singh, B., Su, Y.C. and Riesbeck, K., Vitronectin in bacterial pathogenesis: a host protein used in complement escape and cellular invasion, *Mol. Microbiol.*, 2010, 78, 545-560
- [8] Bergmann, S., Lang, A., Rohde, M., Agarwal, V., Rennemeier, C., Grashoff, C., et al., Integrin-linked kinase is required for vitronectin-mediated internalization of *Streptococcus pneumoniae* by host cells, *J. Cell Sci.*, 2009, 122, 256-267
- [9] Sa, E.C.C., Griffiths, N.J. and Virji, M., *Neisseria meningitidis* Opc invasin binds to the sulphated tyrosines of activated vitronectin to attach to and invade human brain endothelial cells, *PLoS Pathog.*, 2010, 6, e1000911
- [10] Preissner, K.T., Structure and biological role of vitronectin, *Annu. Rev. Cell. Biol.*, 1991, 7, 275-310
- [11] Peterson, C.B., Binding sites on native and multimeric vitronectin exhibit similar affinity for heparin the influence of self-association and multivalence on ligand binding, *Trends Cardiovasc. Med.*, 1998, 8, 124-131
- [12] Zhuang, P., Chen, A.I. and Peterson, C.B., Native and multimeric vitronectin exhibit similar affinity for heparin. Differences in heparin binding properties induced upon denaturation are due to self-association into a multivalent form, *J. Biol. Chem.*, 1997, 272, 6858-6867
- [13] Volker, W., Hess, S., Vischer, P. and Preissner, K.T., Binding and processing of multimeric vitronectin by vascular endothelial cells, *J. Histochem. Cytochem.*, 1993, 41, 1823-1832
- [14] Blouse, G.E., Dupont, D.M., Schar, C.R., Jensen, J.K., Minor, K.H., Anagli, J.Y., et al., Interactions of plasminogen activator inhibitor-1 with vitronectin involve an extensive binding surface and induce mutual conformational rearrangements, *Biochemistry*, 2009, 48, 1723-1735
- [15] Minor, K.H. and Peterson, C.B., Plasminogen activator inhibitor type 1 promotes the self-association of vitronectin into complexes exhibiting altered incorporation into the extracellular matrix, *J. Biol. Chem.*, 2002, 277, 10337-10345
- [16] Underwood, P.A., Kirkpatrick, A. and Mitchell, S.M., New insights into heparin binding to vitronectin: studies with monoclonal antibodies, *Biochem. J.*, 2002, 365, 57-67

- [17] Peake, P.W., Greenstein, J.D., Pussell, B.A. and Charlesworth, J.A., The behaviour of human vitronectin in vivo: effects of complement activation, conformation and phosphorylation, *Clin. Exp. Immunol.*, 1996, 106, 416-422
- [18] Mondino, A. and Blasi, F., uPA and uPAR in fibrinolysis, immunity and pathology, *Trends Immunol.*, 2004, 25, 450-455
- [19] Haiko, J., Laakkonen, L., Juuti, K., Kalkkinen, N. and Korhonen, T.K., The omptins of *Yersinia pestis* and *Salmonella enterica* cleave the reactive center loop of plasminogen activator inhibitor 1, *J. Bacteriol.*, 2010, 192, 4553-4561
- [20] Renckens, R., Roelofs, J.J., Bonta, P.I., Florquin, S., de Vries, C.J., Levi, M., et al., Plasminogen activator inhibitor type 1 is protective during severe Gram-negative pneumonia, *Blood*, 2007, 109, 1593-1601
- [21] Lijnen, H.R., Pleiotropic functions of plasminogen activator inhibitor-1, *J. Thromb. Haemost.*, 2005, 3, 35-45
- [22] Keates, A.C., Tummala, S., Peek, R.M., Jr., Csizmadia, E., Kunzli, B., Becker, K., et al., *Helicobacter pylori* infection stimulates plasminogen activator inhibitor 1 production by gastric epithelial cells, *Infect. Immun.*, 2008, 76, 3992-3999
- [23] Chung, M.C., Jorgensen, S.C., Popova, T.G., Tonry, J.H., Bailey, C.L. and Popov, S.G., Activation of plasminogen activator inhibitor implicates protease InhA in the acute-phase response to *Bacillus anthracis* infection, *J. Med. Microbiol.*, 2009, 58, 737-744
- [24] Urano, T., Ihara, H., Umemura, K., Suzuki, Y., Oike, M., Akita, S., et al., The profibrinolytic enzyme subtilisin NAT purified from *Bacillus subtilis* Cleaves and inactivates plasminogen activator inhibitor type 1, *J. Biol. Chem.*, 2001, 276, 24690-24696
- [25] Beaufort, N., Seweryn, P., de Bentzmann, S., Tang, A., Kellermann, J., Grebenchtchikov, N., et al., Activation of human pro-urokinase by unrelated proteases secreted by *Pseudomonas aeruginosa*, *Biochem. J.*, 2010, 428, 473-482
- [26] Beaufort, N., Wojciechowski, P., Sommerhoff, C.P., Szmyd, G., Dubin, G., Eick, S., et al., The human fibrinolytic system is a target for the staphylococcal metalloprotease aureolysin, *Biochem. J.*, 2008, 410, 157-165
- [27] Kapur, V., Topouzis, S., Majesky, M.W., Li, L.L., Hamrick, M.R., Hamill, R.J., et al., A conserved *Streptococcus pyogenes* extracellular cysteine protease cleaves human fibronectin and degrades vitronectin, *Microb. Pathog.*, 1993, 15, 327-346
- [28] Janoir, C., Pechine, S., Grosdidier, C. and Collignon, A., Cwp84, a surface-associated protein of *Clostridium difficile*, is a cysteine protease with degrading activity on extracellular matrix proteins, *J. Bacteriol.*, 2007, 189, 7174-7180
- [29] Blom, A.M., Hallstrom, T. and Riesbeck, K., Complement evasion strategies of pathogens-acquisition of inhibitors and beyond, *Mol. Immunol.*, 2009, 46, 2808-2817
- [30] Hallstrom, T., Blom, A.M., Zipfel, P.F. and Riesbeck, K., Nontypeable *Haemophilus influenzae* protein E binds vitronectin and is important for serum resistance, *J. Immunol.*, 2009, 183, 2593-2601
- [31] Hallstrom, T., Trajkovska, E., Forsgren, A. and Riesbeck, K., *Haemophilus influenzae* surface fibrils contribute to serum resistance by interacting with vitronectin, *J. Immunol.*, 2006, 177, 430-436
- [32] Singh, B., Jalalvand, F., Morgelin, M., Zipfel, P., Blom, A.M. and Riesbeck, K., *Haemophilus influenzae* protein E recognizes the C-terminal domain of vitronectin and modulates the membrane attack complex, *Mol. Microbiol.*, 2011, 81, 80-98



- [33] Leduc, I., Olsen, B. and Elkins, C., Localization of the domains of the *Haemophilus ducreyi* trimeric autotransporter DsrA involved in serum resistance and binding to the extracellular matrix proteins fibronectin and vitronectin, *Infect. Immun.*, 2009, 77, 657-666
- [34] Attia, A.S., Ram, S., Rice, P.A. and Hansen, E.J., Binding of vitronectin by the *Moraxella catarrhalis* UspA2 protein interferes with late stages of the complement cascade, *Infect. Immun.*, 2006, 74, 1597-1611
- [35] Singh, B., Blom, A.M., Unal, C., Nilson, B., Morgelin, M. and Riesbeck, K., Vitronectin binds to the head region of *Moraxella catarrhalis* ubiquitous surface protein A2 and confers complement-inhibitory activity, *Mol. Microbiol.*, 2010, 75, 1426-1444
- [36] Dehio, M., Gomez-Duarte, O.G., Dehio, C. and Meyer, T.F., Vitronectin-dependent invasion of epithelial cells by *Neisseria gonorrhoeae* involves alpha(v) integrin receptors, *FEBS Lett.*, 1998, 424, 84-88
- [37] Isberg, R.R. and Tran Van Nhieu, G., Binding and internalization of microorganisms by integrin receptors, *Trends Microbiol.*, 1994, 2, 10-14
- [38] Liang, O.D., Rosenblatt, S., Chhatwal, G.S. and Preissner, K.T., Identification of novel heparin-binding domains of vitronectin, *FEBS Lett.*, 1997, 407, 169-172
- [39] Milis, L., Morris, C.A., Sheehan, M.C., Charlesworth, J.A. and Pussell, B.A., Vitronectin-mediated inhibition of complement: evidence for different binding sites for C5b-7 and C9, *Clin. Exp. Immunol.*, 1993, 92, 114-119
- [40] Sheehan, M., Morris, C.A., Pussell, B.A. and Charlesworth, J.A., Complement inhibition by human vitronectin involves non-heparin binding domains, *Clin. Exp. Immunol.*, 1995, 101, 136-141
- [41] Tschopp, J., Masson, D., Schafer, S., Peitsch, M. and Preissner, K.T., The heparin binding domain of S-protein/vitronectin binds to complement components C7, C8, and C9 and perforin from cytolytic T-cells and inhibits their lytic activities, *Biochemistry*, 1988, 27, 4103-4109
- [42] Schmidtchen, A., Frick, I.M., Andersson, E., Tapper, H. and Bjorck, L., Proteinases of common pathogenic bacteria degrade and inactivate the antibacterial peptide LL-37, *Mol. Microbiol.*, 2002, 46, 157-168
- [43] Schmidtchen, A., Frick, I.M. and Bjorck, L., Dermatan sulphate is released by proteinases of common pathogenic bacteria and inactivates antibacterial alpha-defensin, *Mol. Microbiol.*, 2001, 39, 708-713
- [44] Andersson, E., Rydengard, V., Sonesson, A., Morgelin, M., Bjorck, L. and Schmidtchen, A., Antimicrobial activities of heparin-binding peptides, *Eur. J. Biochem.*, 2004, 271, 1219-1226
- [45] Malmsten, M., Davoudi, M. and Schmidtchen, A., Bacterial killing by heparin-binding peptides from PRELP and thrombospondin, *Matrix Biol.*, 2006, 25, 294-300
- [46] Cardin, A.D. and Weintraub, H.J., Molecular modeling of protein-glycosaminoglycan interactions, *Arteriosclerosis*, 1989, 9, 21-32
- [47] Fromm, J.R., Hileman, R.E., Caldwell, E.E., Weiler, J.M. and Linhardt, R.J., Pattern and spacing of basic amino acids in heparin binding sites, *Arch. Biochem. Biophys.*, 1997, 343, 92-100
- [48] Vila, J., Soriano, A. and Mensa, J., [Molecular basis of microbial adherence to prosthetic materials. Role of biofilms in prosthesis-associated infection], *Enferm. Infecc. Microbiol. Clin.*, 2008, 26, 48-54; quiz 55

- [49] Darouiche, R.O., Treatment of infections associated with surgical implants, *N. Engl. J. Med.*, 2004, 350, 1422-1429
- [50] Narasimhan, C. and Lai, C.S., Conformational changes of plasma fibronectin detected upon adsorption to solid substrates: a spin-label study, *Biochemistry*, 1989, 28, 5041-5046
- [51] Wolff, C. and Lai, C.S., Fluorescence energy transfer detects changes in fibronectin structure upon surface binding, *Arch. Biochem. Biophys.*, 1989, 268, 536-545
- [52] Lundberg, F., Schliamser, S. and Ljungh, A., Vitronectin may mediate staphylococcal adhesion to polymer surfaces in perfusing human cerebrospinal fluid, *J. Med. Microbiol.*, 1997, 46, 285-296
- [53] Cagavi, F., Akalan, N., Celik, H., Gur, D. and Guciz, B., Effect of hydrophilic coating on microorganism colonization in silicone tubing, *Acta Neurochir. (Wien)*, 2004, 146, 603-610; discussion 609-610
- [54] Glinel, K., Jonas, A.M., Jouenne, T., Leprince, J., Galas, L. and Huck, W.T., Antibacterial and antifouling polymer brushes incorporating antimicrobial peptide, *Bioconjug. Chem.*, 2009, 20, 71-77
- [55] de Carvalho, C.C., Biofilms: recent developments on an old battle, *Recent Pat. Biotechnol.*, 2007, 1, 49-57
- [56] Jose, B., Antoci, V., Jr., Zeiger, A.R., Wickstrom, E. and Hickok, N.J., Vancomycin covalently bonded to titanium beads kills *Staphylococcus aureus*, *Chem. Biol.*, 2005, 12, 1041-1048
- [57] Kilic, D., Agalar, C., Ozturk, E., Denkbaz, E.B., Cime, A. and Agalar, F., Antimicrobial activity of cefazolin-impregnated mesh grafts, *ANZ. J. Surg.*, 2007, 77, 256-260
- [58] Leung, D., Spratt, D.A., Pratten, J., Gulabivala, K., Mordan, N.J. and Young, A.M., Chlorhexidine-releasing methacrylate dental composite materials, *Biomaterials*, 2005, 26, 7145-7153
- [59] Ravikumar, T., Murata, H., Koepsel, R.R. and Russell, A.J., Surface-active antifungal polyquaternary amine, *Biomacromolecules*, 2006, 7, 2762-2769
- [60] Costa, F., Carvalho, I.F., Montelaro, R.C., Gomes, P. and Martins, M.C., Covalent immobilization of antimicrobial peptides (AMPs) onto biomaterial surfaces, *Acta Biomater.*, 2011, 7, 1431-1440
- [61] Aaboe, M., Offersen, B.V., Christensen, A. and Andreasen, P.A., Vitronectin in human breast carcinomas, *Biochim. Biophys. Acta*, 2003, 1638, 72-82
- [62] Bloemendal, H.J., de Boer, H.C., Koop, E.A., van Dongen, A.J., Goldschmeding, R., Landman, W.J., et al., Activated vitronectin as a target for anticancer therapy with human antibodies, *Cancer Immunol. Immunother.*, 2004, 53, 799-808
- [63] Li, D.Q., Lundberg, F. and Ljungh, A., Characterization of vitronectin-binding proteins of *Staphylococcus epidermidis*, *Curr. Microbiol.*, 2001, 42, 361-367
- [64] Nydegger, U.E., Rieben, R. and Aeschbacher, B., Biocompatibility of apheresis harness, *Transfus. Sci.*, 1990, 11, 43-54
- [65] Salama, A., Hugo, F., Heinrich, D., Hoge, R., Muller, R., Kiefel, V., et al., Deposition of terminal C5b-9 complement complexes on erythrocytes and leukocytes during cardiopulmonary bypass, *N. Engl. J. Med.*, 1988, 318, 408-414
- [66] Lundberg, F., Lea, T. and Ljungh, A., Vitronectin-binding staphylococci enhance surface-associated complement activation, *Infect. Immun.*, 1997, 65, 897-902

- [67] Lundberg, F., Li, D.Q., Falkenback, D., Lea, T., Siesjo, P., Soderstrom, S., et al., Presence of vitronectin and activated complement factor C9 on ventriculoperitoneal shunts and temporary ventricular drainage catheters, *J. Neurosurg.*, 1999, 90, 101-108
- [68] Liang, O.D., Preissner, K.T. and Chhatwal, G.S., The hemopexin-type repeats of human vitronectin are recognized by *Streptococcus pyogenes*, *Biochem. Biophys. Res. Commun.*, 1997, 234, 445-449

## Legend

**Figure 1:** Vn has a multidomain structure, wherein the N-terminal somatomedin-B (SMB) domain (amino acids 1-43) is involved in functional regulation by binding to the plasminogen activator inhibitor-1 (PAI-1). This interaction controls the activity of the plasminogen system in order to limit the overall damage of tissues. Bacterial pathogens degrade the PAI-1/Vn complex in order to accelerate plasmin activity and thus enhance the tissue damage. Downstream of the SMB domain is a Arg-Gly-Asp (RGD) sequence that recognises integrins and is involved in signalling mediated by the integrin-linked kinase (ILK). Bacterial pathogens can be internalized by host cells using this mechanism. Furthermore, Vn has four putative hemopexin like domains (green colour). There are three distinguished heparin binding domains (HBDs) spanning amino acids 82-137, 175-219 and, 348-361 on the Vn molecule. *S. pneumoniae* binds to the region HBD-3 (unpublished data). This interaction helps in bacterial adhesion and internalization mediated by Vn-integrin signalling. *M. catarrhalis* is also known to bind at the region 312-396 and utilizes Vn at its surface in order to inhibit the MAC. In addition, *H. influenzae* interacts with HBD-3 at the amino acid region 353-363. This interaction leads to MAC inhibition and extended survival of *H. influenzae* in serum. *N. meningitide* utilizes Vn recruitment for serum resistance as well as internalization. This species is known to bind at the HBD-3 region, but also at the N-terminal domain 48-68, in particular with sulphated residues of Y56 and Y59. Moreover, *S.*

*pyogenes* has multiple binding sites on Vn (the hemopexin domains) in addition to the C-terminal HBD [38, 68].

**Figure 1:**

