



LUND UNIVERSITY

Abnormal atrial activation is common in patients with arrhythmogenic right ventricular cardiomyopathy.

Platonov, Pyotr; Christensen, Alex H; Holmqvist, Fredrik; Carlson, Jonas; Haunsø, Stig; Svendsen, Jesper H

Published in:
Journal of Electrocardiology

DOI:
[10.1016/j.jelectrocard.2010.08.008](https://doi.org/10.1016/j.jelectrocard.2010.08.008)

2011

[Link to publication](#)

Citation for published version (APA):

Platonov, P., Christensen, A. H., Holmqvist, F., Carlson, J., Haunsø, S., & Svendsen, J. H. (2011). Abnormal atrial activation is common in patients with arrhythmogenic right ventricular cardiomyopathy. *Journal of Electrocardiology*, 44, 237-241. <https://doi.org/10.1016/j.jelectrocard.2010.08.008>

Total number of authors:
6

General rights

Unless other specific re-use rights are stated the following general rights apply:
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00



LUND UNIVERSITY
Faculty of Medicine

LUP

Lund University Publications

Institutional Repository of Lund University

This is an author produced version of a paper published in Journal of Electrocardiology. This paper has been peer-reviewed but does not include the final publisher proof-corrections or journal pagination.

Citation for the published paper:
Pyotr Platonov, Alex H Christensen,
Fredrik Holmqvist, Jonas Carlson, Stig Haunsø,
Jesper H Svendsen

"Abnormal atrial activation is common in patients
with arrhythmogenic right ventricular
cardiomyopathy."

Journal of Electrocardiology
2010 Nov 19

<http://dx.doi.org/10.1016/j.jelectrocard.2010.08.008>

Access to the published version may require journal
subscription.

Published with permission from: Elsevier

Abnormal atrial activation is common in patients with arrhythmogenic right ventricular cardiomyopathy

Short title: *"Atrial abnormality in ARVC"*

Pyotr G. Platonov MD, PhD, ^{*} Alex H. Christensen MD, [‡] Fredrik Holmqvist MD, PhD, ^{*} Jonas
Carlson PhD, ^{*} Stig Haunso MD, PhD, ^{††} and Jesper H. Svendsen, MD, PhD. ^{††}

^{}Department of Cardiology and Center for Integrative Electrophysiology at Lund University,
Lund, Sweden*

*[‡]Department of Cardiology, Rigshospitalet, Copenhagen University Hospital, The Danish
National Research Foundation Center for Cardiac Arrhythmia*

*[†]Department of Medicine and Surgery, Faculty of Health Sciences, University of Copenhagen,
Denmark.*

Conflicts of interests: None

Correspondence Address:

Pyotr G Platonov MD, PhD

Dept of Cardiology

Lund University Hospital

SE-221 85 LUND

SWEDEN

Phone: +46 46 17 35 18

Fax: +46 46 15 78 57

E-mail: pyotr.platonov@med.lu.se

ABSTRACT

Introduction: Structural right atrial abnormalities have been described in patients with arrhythmogenic right ventricular cardiomyopathy (ARVC). However, little is known about ECG signs of atrial involvement in ARVC as no systematic studies have been conducted.

Methods: P-wave triggered signal-averaged orthogonal ECG from 40 ARVC patients (46 ± 15 years, 16 females) was compared with recordings from age- and gender-matched healthy control subjects for assessment of P-wave duration and morphology. P-wave morphology was classified in regard to the P-wave polarity in leads X, Y and Z.

Results: P wave duration was longer in patients (135 ± 18 vs. 124 ± 12 ms, $p=0,003$). Two typical P wave morphologies were identified in the controls: positive in X and Y and negative (45%) or biphasic (55%) in Z. In ARVC pts, typical P waves were seen in only 60% while 15 pts (37%) had atypical P waves positive in all three leads ($p<0,0001$). The presence of atypical P waves in ARVC group was not associated with the presence of either structural or functional right ventricular abnormality.

Conclusions: Patients with ARVC commonly demonstrate deteriorated atrial activation expressed either as prolonged P-wave duration or abnormal P wave morphology. The P wave abnormalities were not secondary to right ventricular dilatation. These findings show that atrial involvement is common in ARVC and may represent yet another manifestation of the disease to be considered for inclusion in ARVC diagnostic work-up.

Keywords: P-wave morphology, ARVC, signal-averaged ECG

INTRODUCTION

Arrhythmogenic right ventricular cardiomyopathy (ARVC) is one of the most common causes of sudden cardiac death in young.¹ The reported ARVC prevalence varies from 1:1000 to 1:10000^{2, 3} and is believed to be underdiagnosed due to subtle and variable manifestations of the disease. During early stages of the disease life-threatening cardiac arrhythmias may occur and sudden cardiac death may be the first symptom of the disease.⁴ ARVC commonly affects the heart on the ventricular level and current diagnostics is based mostly on the signs of electrical, structural or functional right and/or left ventricular dysfunction.⁵

Despite the desmosomal proteins that are commonly involved in the pathogenesis of the disease⁶ are present both in ventricles and atria, there has not been any study that systematically evaluates possible signs of atrial involvement in ARVC. Studies in animal models of the disease consistently report the presence of fibro-fatty replacements in atrial myocardium in affected animals,^{7, 8} while human data on this topic are limited to a handful of case reports.⁹⁻¹¹

The aim of the present study was to assess the presence of atrial abnormalities using signal-averaged P-wave (PSA-ECG) analysis of P-wave duration and morphology in patients with ARVC.

METHODS

Patient population

In total, 40 consecutive, unrelated patients meeting 1994 Task Force diagnostic criteria for ARVC⁵ were enrolled to the study at Rigshospitalet, Copenhagen, Denmark (n=25) and Lund University Hospital, Sweden (n=15). The diagnostic work-up of the patients was conducted in accordance with local guidelines that include initial non-invasive evaluation with resting 12-

lead ECG, 24-48 hr Holter ECG, signal-averaged ECG for assessment of ventricular late potentials, transthoracic echocardiography (n=40) and/or cardiac MRI (n=32). MRI was not routinely performed for all subjects if patients met diagnostic criteria based on other diagnostic modalities. If initial work-up was suggestive of ARVC but score did not reach diagnostic threshold, right ventricular angiography was performed (n=24). All patients included in the study had a transthoracic echocardiography performed within the last three years. Forty ethnically-, age- and gender-matched, healthy subjects were selected for a 1:1 case-control comparison from a database of healthy volunteers recruited in Southern Sweden.¹²

The study was approved by the local ethics committees and complied with the Declaration of Helsinki. All subjects gave informed consent to participation.

Data acquisition and analysis

Standard 12-lead ECG data, of at least 10 seconds duration were recorded using clinical equipment (a minimal sampling frequency of 500 Hz and a sampling resolution of 5 μ V was required).¹³ To enable the analysis of orthogonal P wave morphology, orthogonal-lead ECG data were derived from the 12-lead ECG using the inverse Dower transform.^{14, 15} Unfiltered, signal-averaged P waves were analyzed to determine P wave morphology.¹³ Following high-pass (0.5 Hz) and bandstop (50 Hz) filtering the QRS complexes were automatically identified and grouped according to similarity (a cross-correlation coefficient, $\rho > 0.9$). P waves were extracted using 250 ms wide signal windows preceding each QRS complex. The signal windows were then shifted in time to estimate the maximal correlation in each lead. P waves with a cross-correlation coefficient of $\rho > 0.9$ (analyzed separately in all leads) were grouped together and averaged. The method used is described in detail elsewhere.^{13, 14} P

wave duration was defined by manual setting of the earliest onset and latest end in any of the three orthogonal leads. All analyses were carried out in a blinded fashion.

The P wave morphology was subsequently automatically classified into one of three predefined classes described earlier^{13, 16, 17} (Type 1: positive Leads X and Y and negative Lead Z; Type 2: positive Leads X and Y and biphasic Lead Z (-/+); and Type 3: positive Lead X and biphasic signals in Leads Y (+/-) and Z (-/+)). Different P wave morphology types are illustrated in Figure 1.

Statistics

Data are expressed as the mean \pm standard deviation. The Mann-Whitney U or χ^2 tests were used where appropriate for statistical testing. All tests were two-sided and $P < 0.05$ was considered statistically significant. All statistical analyses were performed using PASW Statistics, version 18.0 (SPSS Inc., Chicago, IL, USA).

RESULTS

Patient population

A total of 40 ARVC patients were included in the study (age 46 ± 15 , range 15-76 years, 16 females). Of those, 13 fulfilled minor and 16 major Task Force criteria in regard to the presence of either structural or functional right ventricular abnormality on imaging (echo, cardiac MRI, or right ventricular angiography). Mean age at diagnosis or symptom onset was 37 ± 19 years. Left ventricular ejection fraction (LVEF) was normal in 38 patients. None of them had any right or left atrial dilatation of clinical significance defined as right atrial short-axis diameter > 45 mm in apical four-chamber view and left atrial diameter > 45 mm in parasternal view. None of the patients had moderate or severe tricuspid regurgitation. The

remaining two patients had markedly reduced right ventricular contractility, LVEF (25% and 30% respectively) and dilatation of all cardiac chambers. In the whole ARVC group, median left atrial dimension in the parasternal view was 39 mm (range 26-44 mm) compared with 35 mm in the control group (range 32-43 mm), not showing any significant difference between the groups.

Ventricular depolarization abnormality defined as the presence of either epsilon-wave (major), isolated QRS prolongation ($>110\text{ms}$) in the right precordial leads (major), or positive late potentials (minor) was identified in 70% of patients. Majority of patients were not on antiarrhythmic medication at the time of ECG registration, while six were treated with beta-blockers and five were taking either sotalol or amiodarone. Characteristics of patients and control subjects are presented in Table 1.

PSA-ECG analysis

There was no significant difference between the groups in regard to heart rate or duration of PQ-interval (Table 2). ARVC patients had longer P-waves than control subjects (135 ± 18 vs 124 ± 12 ms, $p=0.003$). The differences remained significant after correction for the use of class III antiarrhythmic drugs. In ARVC group, the presence of major structural/functional right ventricular abnormality was associated with longer P wave duration (140 ± 15 vs 126 ± 20 , $p=0.049$ in comparison with patients without detectable RV abnormality). None of other Task Force diagnostic criteria was associated with P wave prolongation (Table 3). There was no association between the P wave duration and the left atrial diameter.

P wave morphology was significantly different in the ARVC group when compared with healthy subjects (Table 2). In particular, 38% of ARVC patients had atypical P waves not meeting predefined morphology criteria with activation vector directed posteriorly in the

sagittal plane (Atypical or Type 4) that appeared as positive P waves in lead Z. One ARVC patient had a P wave that met criteria for advanced interatrial block with retrograde activation of the left atrium,^{18, 19} i.e. biphasic P wave morphology in inferior leads and P wave duration >120 ms (Type 3). None of healthy subjects had either Type 3 or Type 4 P waves. Type 4 P wave morphology was not associated with ventricular depolarization or repolarization abnormalities, occurrence of ventricular arrhythmias and was not linked to the presence of structural or functional right ventricular abnormalities.

DISCUSSION

Our study showed that ARVC patients have not only prolonged interatrial conduction expressed as a longer P wave duration but also significant abnormalities in P-wave morphology. These findings consistently support the notion of atrial involvement in the ARVC disease process. To the best of our knowledge, our study is the first that in a systematic manner compared P wave duration and morphology in patients with ARVC and healthy subjects.

Atrial involvement in ARVC in animal models and humans

Atrial involvement in the ARVC disease process has been neglected until recently, when several animal models of the disease reported the loss of atrial myocardium and bilateral focal fibro-fatty atrial lesions in up to one third of affected boxer dogs⁷ and cats.⁸ Even higher number of affected animals were present with atrial enlargement due to the dilatation of the right ventricle and tricuspid regurgitation.

In humans, information on atrial involvement in ARVC is scarce. Few case reports specifically mention atrial dilatation associated with moderate/severe tricuspid regurgitation in patients

with ARVC^{9, 20} and association of ARVC with sick sinus syndrome.⁹⁻¹¹ In a single case report, right atrial biopsies verified the presence of fibrotic replacements in atrial myocardium in a patient with ARVC, sick sinus syndrome and atrio-ventricular conduction disturbance.¹⁰ In a more recent report, ARVC was diagnosed in a patient without ventricular arrhythmias but severe dilatation of right atrium, sick sinus syndrome and extended atrial scarring assessed using voltage mapping of the right atrium during electrophysiological study.¹¹

In the light of these observations, the finding of the prolonged P wave duration in ARVC patients in the present study further indicates that atrial involvement is commonly present in ARVC. Interestingly, none of ARVC patients in our study had either sick sinus syndrome or any significant right atrial dilatation by echocardiography. In agreement with earlier observations,²¹ left atrial diameter was not different between ARVC patients and control subjects.

We observed a borderline significant P wave prolongation in patients with advanced structural/functional right ventricular abnormality (major criterion) compared with structurally/functionally unaffected patients. This may be a consequence of the primary fibro-fatty replacement of atrial myocardium or an effect of 'pure' hemodynamic atrial overload due to reduced contractility of the right ventricle and/or tricuspid regurgitation. However, the lack of clinically significant atrial enlargement in the vast majority of study subjects makes the hemodynamic overload an unlikely cause of the observed P wave prolongation in ARVC group.

P wave morphology

In our earlier studies that used morphology analysis of unfiltered signal-averaged P waves, we demonstrated that P waves with Type 1 and Type 2 morphologies, i.e. either positive or

biphasic in orthogonal lead Z and positive in leads X and Y, are the most common morphological types that are equally common in healthy population. While Type 1 is more common in adolescents and young adults,^{12, 22} Type 2 is the predominant morphology in elderly,¹² and commonly observed in patients with paroxysmal atrial fibrillation.^{16, 23} Type 3 (biphasic P wave in inferior leads) corresponding to the advanced interatrial block with retrograde left atrial activation has not been observed in our cohort of healthy controls, but was present in one ARVC patient who did not have any structural right ventricular abnormality by echo or MRI.

P-waves with activation vector directed posteriorly (Type 4) were present in 15 ARVC patients (38%) and none of control subjects. While this morphology has not been specifically reported in orthogonal leads earlier, several reports indicate that P waves that are negative in lead V₁ and positive in inferior and lateral leads are commonly observed in atrial rhythms originating from right atrial appendage.^{24, 25} It is therefore tempting to suggest that shift of the origin of sinus rhythm toward anterior part or the right atrium or right atrial appendage in ARVC patients may explain the unusually high prevalence of this P wave morphology. Variability of the location of sinus rhythm origin including its extreme anterior locations has been reported earlier by Boineau et al.²⁶

STUDY LIMITATIONS

Our study is a cross-sectional by design and it was therefore not possible to assess whether signs of deteriorated atrial conduction occur early or late in the course of the disease. However, our findings may serve as a background for future studies aimed to answer this question. There was no correlation between P wave duration and either age at symptom onset or duration of symptomatic phase of the disease. Secondly, echocardiography has

been performed as a part of clinical routine and therefore detailed description of right atrial dimension was not available. However, apart from the two patients with bilateral atrial enlargement presented above, ARVC subjects had no signs of clinically significant atrial enlargement and were reported as having 'normal size' atria. Recently proposed revised Task Force ARVC diagnostic criteria²⁷ introduced quantification of RV structural abnormalities which we have not been able to reassess retrospectively in all our patients. However, as recently shown,²⁸ the new criteria are more likely to increase the number of possible and borderline ARVC cases rather than to revise diagnosis in patients who were diagnosed with ARVC earlier.

CONCLUSIONS

Patients with ARVC commonly demonstrate deteriorated atrial activation expressed either as prolonged P wave duration or abnormal P wave morphology. These findings show that atrial involvement in ARVC is common and may represent yet another manifestation of the disease to be considered for inclusion in ARVC diagnostic work-up.

ACKNOWLEDGEMENTS

The present study was supported by Governmental funding of Clinical research within the Swedish National Healthcare System, Crafoord Foundation, donation funds at Lund University Hospital and Maggie Stievens Foundation. AHC has been a recipient of a research fellowship from the Danish Cardiovascular Research School and The Danish National Research Foundation Centre for Cardiac Arrhythmia. In addition, the study was supported from The John and Birthe Meyer Foundation, The Research Foundation of the Heart Centre

Rigshospitalet, and Lægernes Forsikringsforening af 1891. There are no conflicts of interest to report.

REFERENCES

1. Thiene G, Nava A, Corrado D, Rossi L, Pennelli N. Right ventricular cardiomyopathy and sudden death in young people. *N Engl J Med*. 1988;318(3):129-133.
2. Nava A, Bauce B, Basso C, Muriago M, Rampazzo A, Villanova C, Daliento L, Buja G, Corrado D, Danieli GA, Thiene G. Clinical profile and long-term follow-up of 37 families with arrhythmogenic right ventricular cardiomyopathy. *J Am Coll Cardiol*. 2000;36(7):2226-2233.
3. Peters S, Trummel M, Meyners W. Prevalence of right ventricular dysplasia-cardiomyopathy in a non-referral hospital. *Int J Cardiol*. 2004;97(3):499-501.
4. Hulot JS, Jouven X, Empana JP, Frank R, Fontaine G. Natural history and risk stratification of arrhythmogenic right ventricular dysplasia/cardiomyopathy. *Circulation*. 2004;110(14):1879-1884.
5. McKenna WJ, Thiene G, Nava A, Fontaliran F, Blomstrom-Lundqvist C, Fontaine G, Camerini F. Diagnosis of arrhythmogenic right ventricular dysplasia/cardiomyopathy. Task Force of the Working Group Myocardial and Pericardial Disease of the European Society of Cardiology and of the Scientific Council on Cardiomyopathies of the International Society and Federation of Cardiology. *Br Heart J*. 1994;71(3):215-218.
6. Basso C, Corrado D, Marcus FI, Nava A, Thiene G. Arrhythmogenic right ventricular cardiomyopathy. *Lancet*. 2009;373(9671):1289-1300.
7. Basso C, Fox PR, Meurs KM, Towbin JA, Spier AW, Calabrese F, Maron BJ, Thiene G. Arrhythmogenic right ventricular cardiomyopathy causing sudden cardiac death in boxer dogs: a new animal model of human disease. *Circulation*. 2004;109(9):1180-1185.
8. Fox PR, Maron BJ, Basso C, Liu SK, Thiene G. Spontaneously occurring arrhythmogenic right ventricular cardiomyopathy in the domestic cat: A new animal model similar to the human disease. *Circulation*. 2000;102(15):1863-1870.
9. Kazmierczak J, Kornacewicz-Jach Z, Wojtarowicz A. Atrial epicardial pacing with long stimulus to P wave interval in a patient with arrhythmogenic right ventricular dysplasia complicated by right atrial thrombosis. *Pacing Clin Electrophysiol*. 1999;22(7):1111-1113.
10. Nogami A, Adachi S, Nitta J, Taniguchi K, Marumo F, Aonuma K, Iesaka Y, Hiroe M. Arrhythmogenic right ventricular dysplasia with sick sinus syndrome and atrioventricular conduction disturbance. *Jpn Heart J*. 1990;31(3):417-423.
11. Takemura N, Kono K, Tadokoro K, Shinbo G, Ito I, Abe C, Matsuhashi N, Iemura T, Nishikimi T, Horinaka S, Matsuoka H. Right atrial abnormalities in a patient with arrhythmogenic right ventricular cardiomyopathy without ventricular tachycardia. *J Cardiol*. 2008;51(3):205-209.
12. Havmoller R, Carlson J, Holmqvist F, Herreros A, Meurling CJ, Olsson B, Platonov P. Age-related changes in P wave morphology in healthy subjects. *BMC Cardiovasc Disord*. 2007;7:22.
13. Holmqvist F, Platonov PG, Havmoller R, Carlson J. Signal-averaged P wave analysis for delineation of interatrial conduction - further validation of the method. *BMC Cardiovasc Disord*. 2007;7:29.
14. Carlson J, Havmoller R, Herreros A, Platonov P, Johansson R, Olsson B. Can orthogonal lead indicators of propensity to atrial fibrillation be accurately assessed from the 12-lead ECG? *Europace*. 2005;7 Suppl 2:39-48.

15. Edenbrandt L, Pahlm O. Vectorcardiogram synthesized from a 12-lead ECG: superiority of the inverse Dower matrix. *J Electrocardiol.* 1988;21(4):361-367.
16. Holmqvist F, Platonov PG, Carlson J, Zareba W, Moss AJ. Altered interatrial conduction detected in MADIT II patients bound to develop atrial fibrillation. *Ann Noninvasive Electrocardiol.* 2009;14(3):268-275.
17. Holmqvist F, Platonov PG, Carlson J, Havmoller R, Waktare JE, McKenna WJ, Olsson SB, Meurling CJ. Variable interatrial conduction illustrated in a hypertrophic cardiomyopathy population. *Ann Noninvasive Electrocardiol.* 2007;12(3):227-236.
18. Cosio FG, Martin-Penato A, Pastor A, Nunez A, Montero MA, Cantale CP, Schames S. Atrial activation mapping in sinus rhythm in the clinical electrophysiology laboratory: observations during Bachmann's bundle block. *J Cardiovasc Electrophysiol.* 2004;15(5):524-531.
19. Bayes de Luna A, Cladellas M, Oter R, Torner P, Guindo J, Marti V, Rivera I, Iturralde P. Interatrial conduction block and retrograde activation of the left atrium and paroxysmal supraventricular tachyarrhythmia. *Eur Heart J.* 1988;9(10):1112-1118.
20. Galloway JM, Koepke L, Ott P, Marcus FI. Arrhythmogenic right ventricular dysplasia in an American Indian woman. *Pacing Clin Electrophysiol.* 1999;22(7):1093-1096.
21. Kjaergaard J, Hastrup Svendsen J, Sogaard P, Chen X, Bay Nielsen H, Kober L, Kjaer A, Hassager C. Advanced quantitative echocardiography in arrhythmogenic right ventricular cardiomyopathy. *J Am Soc Echocardiogr.* 2007;20(1):27-35.
22. Dilaveris P, Raftopoulos L, Giannopoulos G, Katinakis S, Maragiannis D, Roussos D, Gatzoulis K, Michaelides A, Stefanadis C. Prevalence of interatrial block in healthy school-aged children: definition by P-wave duration or morphological analysis. *Ann Noninvasive Electrocardiol.* 15(1):17-25.
23. Platonov PG, Carlson J, Ingemansson MP, Roijer A, Hansson A, Chireikin LV, Olsson SB. Detection of inter-atrial conduction defects with unfiltered signal-averaged P-wave ECG in patients with lone atrial fibrillation. *Europace.* 2000;2(1):32-41.
24. Zhang T, Li XB, Wang YL, Yin JX, Zhang P, Zhang HC, Xu Y, Guo JH. Focal atrial tachycardia arising from the right atrial appendage: electrophysiologic and electrocardiographic characteristics and catheter ablation. *Int J Clin Pract.* 2009;63(3):417-424.
25. Kistler PM, Roberts-Thomson KC, Haqqani HM, Fynn SP, Singarayar S, Vohra JK, Morton JB, Sparks PB, Kalman JM. P-wave morphology in focal atrial tachycardia: development of an algorithm to predict the anatomic site of origin. *J Am Coll Cardiol.* 2006;48(5):1010-1017.
26. Boineau JP, Canavan TE, Schuessler RB, Cain ME, Corr PB, Cox JL. Demonstration of a widely distributed atrial pacemaker complex in the human heart. *Circulation.* 1988;77(6):1221-1237.
27. Marcus FI, McKenna WJ, Sherrill D, Basso C, Bauce B, Bluemke DA, Calkins H, Corrado D, Cox MG, Daubert JP, Fontaine G, Gear K, Hauer R, Nava A, Picard MH, Protonotarios N, Saffitz JE, Sanborn DM, Steinberg JS, Tandri H, Thiene G, Towbin JA, Tsatsopoulou A, Wichter T, Zareba W. Diagnosis of arrhythmogenic right ventricular cardiomyopathy/dysplasia: proposed modification of the task force criteria. *Circulation.* 2010;121(13):1533-1541.
28. Cox MG, van der Smagt JJ, Noorman M, Wiesfeld AC, Volders PG, van Langen IM, Atsma DE, Dooijes D, Houweling AC, Loh P, Jordaens L, Arens Y, Cramer MJ, Doevendans PA, van Tintelen JP, Wilde AA, Hauer RN. Arrhythmogenic right ventricular dysplasia/cardiomyopathy diagnostic task force criteria: impact of new task force criteria. *Circ Arrhythm Electrophysiol.* 2010;3(2):126-133.

Figure




Figure 1. P wave morphologies in X, Y, Z orthogonal leads and corresponding leads V₁, I and aVF observed in ARVC patients and healthy subjects. Healthy controls had only Type 1 and Type 2 morphologies while 1 ARVC patient had Type 3 and 15 patients had Type 4 P waves.

Table 1: Subject characteristics

		ARVC	Controls
		n=40	n=40
Age at symptom onset	[years]	37 ±16	NA
Male gender*	[n]	24	22
Age at ECG registration*	[years]	46±15	45±15
LA dimension*	[mm]	37±5	36±3
Antiarrhythmic drugs			
None	[n]	29	40
Class II	[n]	6	0
Class III	[n]	5	0
<i>ARVC Task Force Criteria (1994)⁵</i>			
Family history	[none/minor/major]	24/14/2	0
Depolarization abnormality	[none/minor/major]	12/17/11	0
Repolarization abnormality	[none/minor]	20/20	0
Arrhythmia (VT/PVC)	[none/minor]	5/35	0
RV functional/ anatomical abnormality	[none/minor/major]	11/13/16	0
Histology	[normal/major/not done]	8/14/18	N/A

LA= left atrium; N/A= data not available; PVC=premature ventricular contractions;
 VT=ventricular tachycardia

*- non-significant difference between the groups

Table 2: Results of the P-wave signal-averaged ECG analysis

		ARVC n=40	Controls n=40	P-value
Heart rate	[bpm]	64±14	69±11	ns
PQ interval	[ms]	176±44	163±20	ns
P-wave duration	[ms]	135±18	125±12	0.007
P-wave morphology	[n]			
Type 1		10	18	0.000
Type 2		14	22	
Type 3		1	0	
Type 4		15	0	

ns= not significant

Table 3: P wave duration in relation to the presence of ARVC Task Force diagnostic criteria⁵, ms

	None	Minor	Major	P-value
Family history	136±13	135±26	140±15	n.s.
Depolarization abnormality	128±21	138±17	138±16	n.s.
Repolarization abnormality	130±18	140±19	N/A	0.06
Arrhythmia (VT/PVC)	132±11	136±19	N/A	n.s.
RV functional/anatomical abnormality	126±20	136±18	140±15	0.049*

N/A= not applicable; PVC= premature ventricular contractions; RV= right ventricular; VT= ventricular tachycardia

*- p-value concerns comparison between the patients with major structural/functional right ventricular abnormality and those without any detectable abnormality.