



LUND UNIVERSITY

Reaction Invariants and Their Importance in the Analysis of Eigenvectors, State Observability and Controllability of the Continuous Stirred Tank Reactor

Fjeld, Magne; Asbjørnsen, O. A.; Åström, Karl Johan

Published in:
Chemical Engineering Science

1974

Document Version:
Publisher's PDF, also known as Version of record

[Link to publication](#)

Citation for published version (APA):
Fjeld, M., Asbjørnsen, O. A., & Åström, K. J. (1974). Reaction Invariants and Their Importance in the Analysis of Eigenvectors, State Observability and Controllability of the Continuous Stirred Tank Reactor. *Chemical Engineering Science*, 29, 1917-1926.

Total number of authors:
3

General rights

Unless other specific re-use rights are stated the following general rights apply:
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

Reprinted from

Chemical Engineering Science, 1974, Vol. 29, pp. 1917-1926. Pergamon Press. Printed in Great Britain

REACTION INVARIANTS AND THEIR IMPORTANCE
IN THE ANALYSIS OF EIGENVECTORS, STATE
OBSERVABILITY AND CONTROLLABILITY OF THE
CONTINUOUS STIRRED TANK REACTOR

M. FJELD

The University of Nairobi, Department of Electrical Engineering, P.O. Box 30197, Nairobi, Kenya

O. A. ASBJØRNSEN

Chemical Engineering Laboratory at the University of Trondheim, The Norwegian Institute of
Technology, N-7034, Trondheim, Norway

and

K. J. ÅSTRÖM

The Technical University of Lund, Division of Automatic Control, Fack 725, Lund 7, Sweden



PERGAMON PRESS
OXFORD · NEW YORK

7419

REACTION INVARIANTS AND THEIR IMPORTANCE IN THE ANALYSIS OF EIGENVECTORS, STATE OBSERVABILITY AND CONTROLLABILITY OF THE CONTINUOUS STIRRED TANK REACTOR

M. FJELD

The University of Nairobi, Department of Electrical Engineering, P.O. Box 30197, Nairobi, Kenya

O. A. ASBJØRNSEN

Chemical Engineering Laboratory at the University of Trondheim, The Norwegian Institute of Technology, N-7034, Trondheim, Norway

and

K. J. ÅSTRÖM

The Technical University of Lund, Division of Automatic Control, Fack 725, Lund 7, Sweden

(Received 10 May, 1973; accepted 21 February, 1974)

Abstract—Two different projection operators are found, projecting the state space of a continuous stirred tank reactor (C.S.T.R.) dynamic model on the space of asymptotically invariant states (or on the space of true invariant states for a batch reactor). The first projection gives an explicit expression for the invariants, but it requires a certain partitioning of the stoichiometric matrix. The other projection avoids this difficulty, but gives a description of the invariants by a set of equations which do not have full rank. Concerning the observability of the dynamical modes, it turns out that the invariants put certain restrictions on the observation matrix for the system to be observable. The corresponding conditions for the system to be controllable by feed composition control are briefly outlined. Certain properties of the solution of the linearized system are also analyzed.

1. INTRODUCTION

The study of chemical reactor dynamics becomes more and more important in the attempts to improve the production, the selectivity and the dynamic control of the reactors. The basic thermodynamic variables like composition and enthalpy are important state variables in the analysis of these phenomena.

A subset of reaction invariant variables may be found by a simple partitioning of the reactor equations. These variables are linear combinations of the basic concentrations and the temperature, even if the differential state equations are strongly nonlinear due to the reaction rates. The motion of these variables is linear and constrained to a subset orthogonal to the remaining state space. For a C.S.T.R., these variables may be called asymptotic invariants[1]. In a batch reactor, they are true invariants[2]. Hence, the dimensionality of such a state space description can be reduced, as has been realized for a long time for closed systems. Recently, conditions for such a reduction to be valid in

open, continuous flow reactors have been analyzed by Asbjørnsen and Fjeld[3], Fjeld[1] and Asbjørnsen[4]. For global stability analysis, all the states are necessary, unless special conditions are met. However, for stability analysis in the small, only a reduced number of state variables needs to be considered. In this paper, a closer study of the projection of the state space on the set of asymptotically invariants (eigenmodes) is made.

The concepts of observability and controllability are both very important for the control of dynamical systems. State observability is important also in parameter estimation schemes. It turns out that invariant states of chemical reactors are of interest in the analysis of state observability of such processes, and this is the topic studied in the last part of the paper.

2. INVARIANTS AND EIGENVECTORS

Consider a nonlinear system described by the differential equation

$$\dot{\mathbf{x}}(t) = \mathbf{f}(\mathbf{x}(t), \mathbf{u}(t), t) \quad (1)$$

where for $t \geq 0$, $\mathbf{x} \in X \subset E_n$, $\mathbf{u} \in U \subset E_r$ and $\mathbf{x}(0) = \mathbf{x}_0 \in X$. Furthermore, $\mathbf{u}(\cdot) \in \mathcal{F}$, \mathcal{F} is a given class of integrable functions. The solution of Eq. (1) is assumed to exist for all $t > 0$ and all $\mathbf{x}_0 \in X$. The notion of invariance is introduced as follows.

Definition 2.1

A true invariant of the solution Eq. (1) is a function $h(\mathbf{x})$ of the state vector $\mathbf{x}(t)$ such that $\dot{h} = dh/dt = \langle \text{grad } h, \dot{\mathbf{x}} \rangle = \langle \text{grad } h, \mathbf{f} \rangle = 0$.

If $\langle \text{grad } h, \mathbf{f} \rangle$ is neither a constant nor equal to zero, but its limit as $t \rightarrow \infty$ is zero, then h is an asymptotic invariant of the solution of Eq. (1).

If \mathbf{x} is the state vector of a chemical reactor and $h(t)$ does not explicitly depend on any reaction parameter, such as stoichiometric coefficients, kinetic parameters or heats of reactions, then an invariant, true or asymptotic, is called a true or asymptotic reaction invariant.

Linear systems

For linear autonomous systems, i.e. $\dot{\mathbf{x}} = A\mathbf{x}$, it is easily seen that the projection on the nullspace of A is an invariant. Similarly it is found that the projection on the space spanned by the eigenvectors corresponding to stable eigenvalues is an asymptotic invariant.

Nonlinear systems

In attempting to extend this idea to the nonlinear case the deviation $\Delta\mathbf{x} = \mathbf{x} - \mathbf{x}_s$ is introduced; here \mathbf{x}_s is a singular point at which $\mathbf{f}(\mathbf{x}_s, \mathbf{u}_s) = \mathbf{0}$. Then

$$\Delta\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}_s + \Delta\mathbf{x}) \triangleq \mathbf{h}(\Delta\mathbf{x}) \triangleq \mathbf{F}(\Delta\mathbf{x}) \Delta\mathbf{x} \quad (2)$$

where the matrix \mathbf{F} has elements $F_{ij} = h_i(\Delta\mathbf{x})/\Delta x_j$. These are assumed to be uniformly bounded for finite Δx_j as the other $\Delta x_i \rightarrow 0$. For any fixed $\Delta\mathbf{x}$, the eigenvalue problem for \mathbf{F} is

$$[\lambda\mathbf{I} - \mathbf{F}(\Delta\mathbf{x})]\mathbf{e} = \mathbf{0}. \quad (3)$$

At $\mathbf{x} = \mathbf{x}_s$, i.e. as $\Delta\mathbf{x} \rightarrow \mathbf{0}$, \mathbf{F} becomes the Jacobian of the functions \mathbf{f} with respect to the variables \mathbf{x} evaluated at the critical point. For the problem (3) the eigenvectors and eigenvalues will depend on $\Delta\mathbf{x}$ unless Eq. (1) is linear. However certain nonlinear systems, including chemical reactors in general, may have a subset of state space where linear motions take place. Certain eigenvalues and the corresponding eigenvectors will then be independent of $\Delta\mathbf{x}$. For an absolute invariant to be present there must be a zero eigenvalue. An asymptotic reaction invariant implies that an eigenmode becomes constant as $t \rightarrow \infty$ independently of the reaction.

3. STIRRED TANK REACTOR EQUATIONS

The dynamic mass and enthalpy balances for a stirred tank reactor has been studied in details by Asbjørnsen and Fjeld[3] and Asbjørnsen[4, 10] under isothermal and adiabatic operation, and under operation with heat exchange.

Provided all physical properties as well as the heat of reactions are constant, the enthalpy and species balances are combined in one single vector differential equation[3, 4]

$$\dot{\mathbf{x}} = \frac{1}{\tau}(\mathbf{x}_f - \mathbf{x}) - A\mathbf{r}(\mathbf{x}) \quad (4)$$

where A for isothermal operation is the transpose of the stoichiometric matrix. For truly adiabatic operation, the adiabatic temperature rises for each reaction is added as an extra row, as a consequence of the inclusion of the temperature in state vector \mathbf{x} . This vector then comprises the temperature and the molar concentrations of the n participants of n_r truly independent chemical reactions[3, 4].

The characteristic feature of the differential equation describing the C.S.T.R. is that the nonlinear term at the right hand side of Eq. (4) maps as follows

$$A\mathbf{r}: E_n \xrightarrow{r} E_{n_r} \xrightarrow{A} E_n. \quad (5)$$

This particular property of the stirred tank reactor equations is the basis for the analysis presented in this paper.

Provided the reactions are truly independent, the matrix A will have full rank n_r , and a square submatrix A_1 , with rank n_r , can always be found.

In this case, Asbjørnsen and Fjeld[3] showed by a partitioning procedure that a reaction invariant $\mathbf{z}(t)$ is obtained

$$\mathbf{z} = \mathbf{x}_2 - A_2 A_1^{-1} \mathbf{x}_1 \quad (6)$$

where A_2 contains the remaining $n - n_r$ rows of A and $\dim(\mathbf{z}) = n - n_r$.

This variable is described by the differential equation for a stirred tank without any reactions

$$\dot{\mathbf{z}} = \frac{1}{\tau}(\mathbf{z}_f - \mathbf{z}). \quad (7)$$

The solution for the reaction invariant is simply

$$\mathbf{z}(t) = \exp\left(-\int_0^t d\nu/\tau\right)\mathbf{z}(0) + \int_0^t \exp\left(-\int_\theta^t d\nu/\tau\right)\mathbf{z}_f(\theta) d\theta \quad (8)$$

which allows the parameter τ to be dependent of time[11].

Provided \mathbf{z}_f is constant, then $\mathbf{z}(t)$ is an asymptotic

reaction invariant, since:

$$\lim_{t \rightarrow \infty} [z(t)] = z_f \tag{9}$$

If $1/\tau = 0$, i.e. for a batch reactor, z is a true invariant $z(0)$. This leads to the following theorem [3, 4]:

Theorem 1. For a C.S.T.R. with n_r reactions between n species and with a stoichiometric matrix of full rank n_r , an asymptotic invariant exists, which is

$$z(t) = x_2(t) - A_2 A_1^{-1} x_1(t) \tag{10}$$

provided the feed is such that

$$z_f = x_{2f} - A_2 A_1^{-1} x_{1f} \tag{11}$$

is constant. For a batch reactor, $z(t) = z(0)$ is a true invariant.

Note that all concentrations and the temperature have to be positive

$$x_i \geq 0, x_{ij} \geq 0, i = 1, 2, \dots, n. \tag{12}$$

This constrains the states to a certain subset of E_n in the positive orthant E_n^+ . This means that the state space is not a linear space, and therefore the term "subspace" should not be used for certain subsets of E_n .

The deviation from a steady state x_s is introduced as $\Delta x = x - x_s$. A new set of state variables may be chosen, as Δx_1 and Δz . For a given $x_f =$ constant, the n -dimensional non-linear reactor equation then becomes

$$\begin{bmatrix} \Delta \dot{x}_1 \\ \Delta \dot{z} \end{bmatrix} = -\frac{1}{\tau} \begin{bmatrix} \Delta x_1 \\ \Delta z \end{bmatrix} - \begin{bmatrix} A_1 \\ 0 \end{bmatrix} \Delta r \tag{13}$$

where

$$\Delta r \triangleq r(x) - r(x_s). \tag{14}$$

Furthermore

$$\Delta \tilde{x} \triangleq \begin{bmatrix} \Delta x_1 \\ \Delta z \end{bmatrix} \text{ and } \Delta x = \begin{bmatrix} -I & 0 \\ A_2 A_1^{-1} & I \end{bmatrix} \Delta \tilde{x}. \tag{15}$$

4. INVARIANT SUBSETS AND STATE VARIABLES

4.1 The partitioning procedure

It is now seen that the subset on which $\Delta z = 0$ defines a n_r -dimensional subset R_{n_r} of the state space $X \subset E_n^+$ which is invariant [5] under the non-linear transformation $F(\Delta \tilde{x})$, since Eqs. (18)–(19) may be written as

$$\Delta \tilde{x} = \begin{bmatrix} -\frac{1}{\tau} I - A_1 R(\Delta \tilde{x}) & -A_1 R_2(\Delta \tilde{x}) \\ 0 & -\frac{1}{\tau} I \end{bmatrix} \Delta \tilde{x} = F(\Delta \tilde{x}) \Delta \tilde{x} \tag{16}$$

where

$$\Delta r \triangleq [R_1(\Delta \tilde{x}) \mid R_2(\Delta \tilde{x})] \Delta x \tag{17}$$

$$\dim(R_1) = n_r \times n_r, \quad \dim(R_2) = n_r \times (n - n_r).$$

The transformation projecting $\Delta x(t)$ on the space of asymptotical invariants $\Delta z(t)$ is given by Eq. (14) or

$$\Delta z(t) = [-A_2 A_1^{-1} \mid I] \Delta x(t). \tag{18}$$

Therefore, the invariant subset R_{n_r} is described through all those Δx satisfying the constraints (12) and the $n - n_r$ equations

$$V \Delta x \triangleq [-A_2 A_1^{-1} \mid I] \Delta x = 0 \tag{19}$$

from which the simple relation follows:

$$R_{n_r} = \mathcal{N}(V)^+ \tag{20}$$

The $^+$ reminds about the restrictions corresponding to Eq. (12). It is observed that the rows of V are linearly independent, and span a space R_{n-n_r} orthogonal to R_{n_r} .

The relation above may be expressed by the projection operator P_1 . When $X = R_{n_r} \oplus R_{n-n_r}$ [care should be taken in the use the direct sum symbol \oplus since the spaces are not linear because of Eq. (12)], Δx may be expressed as

$$\Delta x = P_1 \Delta x + (I - P_1) \Delta x \tag{21}$$

where

$$P_1 \Delta x \in R_{n-n_r} \text{ and } (I - P_1) \Delta x \in R_{n_r}$$

P_1 is a projection operator on R_{n-n_r} along R_{n_r} if and only if $P_1 = P_1^2$ when $R_{n-n_r} = \mathcal{R}(P_1)$ and $R_{n_r} = \mathcal{N}(P_1)$ [5]. It is seen that the conditions are satisfied with

$$P_1 = \begin{bmatrix} 0 & 0 \\ -A_2 A_1^{-1} & I \end{bmatrix}. \tag{22}$$

Then

$$P_1 \Delta x = \begin{pmatrix} 0 \\ \Delta z \end{pmatrix} \tag{23}$$

$$(I - P_1) \Delta x = \begin{pmatrix} \Delta x_1 \\ \Delta x_2 - \Delta z \end{pmatrix} \tag{24}$$

Note that neither $P_1 \Delta x$ nor $(I - P_1) \Delta x$ are state vectors, but Δz and Δx_1 are. The asymptotic invariants are explicitly given by the last $n - n_r$ components of $P_1 \Delta x$ as expressed by Eq. (2)

4.2 Finding invariants through direct use of projections

The problem of finding the reaction invariants has so far been solved through a particular choice of state variables. To get further insight into the problem another coordinate free solution is presented. The analysis will also result in a different

representation of the asymptotic invariants. This approach is motivated by the corresponding description of a batch reaction, i.e. $1/\tau = q(t)/V = 0$:

$$\dot{x}(t) = -Ar[x(t)] \quad (25)$$

From matrix algebra, it is known that if A is a linear matrix transformation from E_n to E_n , then

$$E_n = \mathcal{R}(A) \oplus \mathcal{N}(A^T) \quad (26)$$

and

$$\mathcal{N}(A^T) = [\mathcal{R}(A)]^\perp \quad (27)$$

Let $w \in \mathcal{N}(A^T)$, i.e. w satisfies $A^T w = 0$, any vector x in E_n given by the solution of (25) can be expressed as

$$x = w + s = w + Av \quad (28)$$

where $s \in \mathcal{R}(A)$. The latter may simply be taken as $s = Av$, where v is a n_r -dimensional vector, in general with nonzero entries. The projection operator for the projection from E_n to $\mathcal{N}(A^T)$ is

$$P_2 = I - AA^\dagger \quad (29)$$

Where A^\dagger is a pseudo-inverse of A . Since A has full rank, A^\dagger is simply:

$$A^\dagger = (A^T A)^{-1} A^T \quad (30)$$

It is observed that P_2 satisfies all the conditions for P_2 to be a projection operator. Applying P_2 to Eq. (25),

$$P_2 \dot{x}(t) = \dot{w}(t) = 0 \quad (31)$$

i.e.

$$w(t) = \text{constant} = w_0. \quad (32)$$

Further, the projection onto $\mathcal{R}(A) = \mathcal{N}(A^T)^\perp$ is AA^\dagger , i.e.

$$(I - P_2)\dot{x}(t) = -Ar(w_0 + Av) \quad (33)$$

that is

$$A\dot{v}(t) = -Ar(w_0 + Av). \quad (34)$$

Projecting again $A\dot{v}$ on R_{n_r} by multiplying Eq. (34) with A^\dagger from the left (this projects $A\dot{v}$ onto $\mathcal{R}(A^T) = \mathcal{N}(A)^\perp$, i.e. R_{n_r})

$$A^\dagger A\dot{v}(t) = \dot{v}(t) = -r[w_0 + Av(t)]. \quad (35)$$

This result shows that the variable v is equivalent to the 'extent of reaction' so frequently referred to in the chemical engineering literature. To conclude, an arbitrary $x(t)$ satisfying Eq. (25) can be expressed as

$$x(t) = P_2 x(t) + (I - P_2)x(t) = w_0 + Av(t) \quad (36)$$

where P_2 is an orthogonal projection operator. Note that $\text{rank}(P_2) = n - n_r$.

Theorem 2. For a stirred tank batch reactor with n_r reactions between n species and with a stoichiometric matrix of rank n_r , the function

$$w(t) = [I - A(A^T A)^{-1} A^T]x(t) \quad (37)$$

is a true reaction invariant.

Furthermore, the function

$$v(t) = (A^T A)^{-1} A^T x(t) \quad (38)$$

is the extent of reaction in the batch reactor.

The projection operator P_2 is applied to the C.S.T.R. equation, Eq. (4), which becomes

$$P_2 \dot{x} = \frac{1}{\tau} (P_2 x_f - P_2 x) - P_2 Ar. \quad (39)$$

Since the properties of the pseudo-inverse A^\dagger is such that $A^\dagger A = I$, the following equation is obtained

$$P_2 \dot{x} = \frac{1}{\tau} (P_2 x_f - P_2 x). \quad (40)$$

Defining as before $w = P_2 x$ and $w_f = P_2 x_f$, it follows that

$$\dot{w} = \frac{1}{\tau} (w_f - w). \quad (41)$$

Furthermore

$$(I - P_2)\dot{x} = AA^\dagger \dot{x} = \frac{1}{\tau} AA^\dagger (x_f - x) - Ar. \quad (42)$$

Since Eq. (28) implies that $v = A^\dagger x$ and $v_f = A^\dagger x_f$, it follows that

$$A\dot{v} = \frac{1}{\tau} A(v_f - v) - Ar. \quad (43)$$

With $Av \in \mathcal{R}(A)$, a projection of this vector on to $\mathcal{N}(A^T)$ as in Eq. (35), gives

$$\dot{v} = \frac{1}{\tau} (v_f - v) - r. \quad (44)$$

Observe that $w(0) \in \mathcal{N}(A^T)$ and $w_f \in \mathcal{N}(A^T)$ implies that $w(t)$ is invariant in this respect under the solution of Eq. (41) so $w(t) \in \mathcal{N}(A^T)$. The solution is equivalent to Eq. (8), but with z replaced by w

$$w(t) = \exp\left(-\int_0^t dv/\tau\right)w(0) + \int_0^t \exp\left(-\int_\theta^t dv/\tau\right)w_f(\theta) d\theta. \quad (45)$$

Finally, $x(t)$ is given by

$$x(t) = w(t) + Av(t). \quad (46)$$

If the feed is such that $w_f = P_2 x_f = \text{constant}$, then Eq. (41) implies that

$$\lim_{t \rightarrow \infty} w(t) = w_f. \quad (47)$$

In analogy to Eqs. (11), (12) and (13), this leads to the following theorem:

Theorem 3. For a C.S.T.R. with n_r reactions between n species, and with a stoichiometric matrix of full rank n_r , an asymptotic reaction invariant exists, which is

$$\mathbf{w}(t) = [I - A(A^T A)^{-1} A^T] \mathbf{x}(t) \quad (48)$$

provided the feed is such that

$$\mathbf{w}_f = [I - A(A^T A)^{-1} A^T] \mathbf{x}_f \quad (49)$$

is constant.

Remark. Theorem 3 includes Theorem 1 as a special case, which is demonstrated by the introduction of the partitioned forms of A and \mathbf{x} . Then

$$[I - A(A^T A)^{-1} A^T] \mathbf{x} = \begin{bmatrix} \mathbf{x}_1 - A_1(A_1^T A_1 + A_2^T A_2)^{-1} (A_1^T \mathbf{x}_1 + A_2^T \mathbf{x}_2) \\ -A_2(A_1^T A_1 + A_2^T A_2)^{-1} (A_1^T \mathbf{x}_1 + A_2^T \mathbf{x}_2) + \mathbf{x}_2 \end{bmatrix} \quad (50)$$

and it follows from Theorem 3 that for large t , the two vectors

$$\mathbf{x}_2 - A_2(A_1^T A_1 + A_2^T A_2)^{-1} [A_1^T \mathbf{x}_1 + A_2^T \mathbf{x}_2] \rightarrow \text{constant} \quad (51)$$

$$\mathbf{x}_1(A_1(A_1^T A_1 + A_2^T A_2)^{-1} [A_1^T \mathbf{x}_1 + A_2^T \mathbf{x}_2]) \rightarrow \text{constant}$$

are asymptotic reaction invariants. Eliminating the expression within the square brackets, gives

$$\mathbf{x}_2 - A_2 A_1^{-1} \mathbf{x}_1 \rightarrow \text{constant}$$

which is equivalent to Theorem 1.

5. EXAMPLE

Some of the relationships in the previous sections is illustrated by a simple example, given by the stoichiometric matrix

$$A^T = \begin{bmatrix} -1 & 2 & 0 & 0 \\ 0 & -2 & 2 & 1 \end{bmatrix} \quad (52)$$

From Eq. (19), the subset $R_{n_r} = R_2$ is obtained through

$$V \Delta \mathbf{x} = \begin{bmatrix} 2 & 1 & 1 & 0 \\ 2 & 1 & 0 & 1 \end{bmatrix} \Delta \mathbf{x} = \mathbf{0} \quad (53)$$

P_1 is given by

$$P_1 = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 2 & 1 & 1 & 0 \\ 2 & 1 & 0 & 1 \end{bmatrix} \quad (54)$$

From Eq. (29), P_2 is given by

$$P_2 = \frac{1}{29} \begin{bmatrix} 20 & 10 & 8 & 4 \\ 10 & 5 & 4 & 2 \\ 8 & 4 & 9 & -10 \\ 4 & 2 & -10 & 24 \end{bmatrix} \quad (55)$$

which has a rank equal two. $P_2 \Delta \mathbf{x} = \mathbf{0}$ can be reduced to Eq. (53), but the form $P_2 \Delta \mathbf{x}$ is not dependent on the condition that A_1 is nonsingular: Any rearrangement of the columns of A^T does not change the result.

6. LINEARIZED SOLUTION

The use of Eq. (53) leads to a particular simple form of the autonomous ($\Delta \mathbf{x}_f = \mathbf{0}$) solution of a C.S.T.R. The linearized description can be found from Eq. (44) where $\Delta \mathbf{x}_f = \mathbf{0}$ and hence $\Delta \mathbf{v}_f = \mathbf{0}$

$$\Delta \dot{\mathbf{v}} = -\frac{1}{\tau} \Delta \mathbf{v} - \Delta \mathbf{r} \quad (56)$$

where $\Delta \mathbf{r}$ is replaced by $(\partial \mathbf{r} / \partial \mathbf{v}) \Delta \mathbf{v}$

$$\left. \frac{\partial \Delta \mathbf{r}}{\partial \Delta \mathbf{v}} \right|_{\substack{\mathbf{w}=\mathbf{w}_s \\ \mathbf{v}=\mathbf{v}_s}} = \left. \frac{\partial \mathbf{r}}{\partial \mathbf{x}} \frac{\partial \mathbf{x}}{\partial \mathbf{v}} \right|_{\substack{\mathbf{x}=\mathbf{x}_s \\ \mathbf{v}=\mathbf{v}_s}} = \mathbf{R} \mathbf{A} \quad (57)$$

At steady state, one has from Eqs. (28), (41) and (44)

$$\begin{aligned} \mathbf{x}_s &= \mathbf{w}_s + \mathbf{A} \mathbf{v}_s \\ \mathbf{w}_s &= \mathbf{w}_f \end{aligned} \quad (58)$$

$$\mathbf{v}_s = \mathbf{v}_f - \tau \mathbf{r}(\mathbf{x}_s).$$

This leads to a linearized approximation to the n_r -dimensional differential equation for the vector $\Delta \mathbf{v}$:

$$\Delta \dot{\mathbf{v}} = -\frac{1}{\tau} \Delta \mathbf{v} - \mathbf{R} \mathbf{A} \Delta \mathbf{v}. \quad (59)$$

The solution for $\Delta \mathbf{x} = \boldsymbol{\xi}(t)$ is

$$\boldsymbol{\xi}(t) = \Delta \mathbf{w}(t) + \mathbf{A} \Delta \mathbf{v}(t) \quad (60)$$

that is:

$$\begin{aligned} \boldsymbol{\xi}(t) &= \exp(-t/\tau) \Delta \mathbf{w}(0) \\ &+ \exp\left[\left(-\frac{1}{\tau} I - \mathbf{R} \mathbf{A}\right)t\right] \mathbf{A} \Delta \mathbf{v}(0) \end{aligned} \quad (61)$$

7. EIGENVECTORS

7.1 The nonlinear system

Considering the non-linear system, the deviations in the reaction rate from its steady state value $\mathbf{r}_s = \mathbf{r}(\mathbf{x}_s)$ are defined by

$$\Delta \mathbf{r}(\mathbf{x}_s + \Delta \mathbf{x}) \triangleq \mathbf{R}(\mathbf{x}) \Delta \mathbf{x}. \quad (62)$$

Then the autonomous balance equations become

$$\Delta \dot{\mathbf{x}}(t) = -\frac{1}{\tau} \Delta \mathbf{x}(t) - \mathbf{A} \mathbf{R}(\mathbf{x}) \Delta \mathbf{x}(t) \quad (63)$$

applying Eq. (3), the right column eigenvectors \mathbf{e}_j satisfy (for given $\Delta\mathbf{x}$) the equation

$$\left[\left(\lambda_j + \frac{1}{\tau} \right) I + AR(\mathbf{x}) \right] \mathbf{e}_j = \mathbf{0} \quad (64)$$

Similarly, the left row eigenvectors \mathbf{v}_j satisfy the equation:

$$\mathbf{v}_j \left[\left(\lambda_j + \frac{1}{\tau} \right) I + AR(\mathbf{x}) \right] = \mathbf{0}. \quad (65)$$

For $\lambda_j = -1/\tau$, Eq. (65) may be written as a matrix equation:

$$VAR(\mathbf{x}) = [0] \quad (66)$$

where V is the matrix of the $n - n_r$ left row eigenvectors belonging to the $n - n_r$ eigenvalues $-1/\tau$. Equation (66) is satisfied by

$$V = [-A_2 A_1^{-1} \quad I_{n-n_r}] \quad (67)$$

independently of the matrix $R(\mathbf{x})$. Here I_{n-n_r} is the identity matrix of dimension $[n - n_r] \times [n - n_r]$. Hence, it follows

Theorem 4. The $n - n_r$ left eigenvectors of the non-linear C.S.T.R. equation belonging to the $n - n_r$ 'uncoupled' eigenvalues $-1/\tau$ are linearly independent and equal to the rows of the matrix V in Eq. (67). 'Uncoupled' eigenvalues meaning that the algebraic multiplicity is equal to the geometrical multiplicity, since the system matrix is diagonalizable in the $n - n_r$ eigenvalues $-1/\tau$, due to the properties of the reaction invariants.

The right column eigenvectors \mathbf{e}_i corresponding to the $n - n_r$ left row eigenvectors above, can be found from the orthogonality condition

$$\mathbf{v}_i \mathbf{e}_i = \delta_{ii} \quad (68)$$

7.2 The linearized system

The variable \mathbf{v} in the partitioning $\mathbf{x} = \mathbf{w} + A\mathbf{v}$, was shown by Eq. (41) to be described by a set of n_r non-linear equations which were given a linearised approximation in Eq. (59). This equation has n_r eigenvalues, which are assumed to be distinct. Correspondingly, there are n_r eigenvectors of this equation.

The basic balance equations, Eq. (4), in the variable \mathbf{x} may be linearised in a similar way [3, 4] by defining as in Eq. (57)

$$\left. \frac{\partial \Delta \mathbf{r}}{\partial \Delta \mathbf{x}} \right|_{\mathbf{x}=\mathbf{x}_s} = R(\mathbf{x}_s). \quad (69)$$

Then the linearised approximation to Eq. (4) may be written for an autonomous system ($\Delta \mathbf{x}_f = \mathbf{0}$ [3, 4],

$$\Delta \dot{\mathbf{x}} = -\frac{1}{\tau} \Delta \mathbf{x} - AR \Delta \mathbf{x}. \quad (70)$$

It may be of interest to see how the eigenvectors of Eq. (59) (the $\Delta \mathbf{v}$ -system) are related to the eigenvectors of Eq. (70) (the $\Delta \mathbf{x}$ -system). The right eigenvectors of the $\Delta \mathbf{x}$ -system are given by

$$\left[\left(\lambda_i + \frac{1}{\tau} \right) I_n + AR \right] \mathbf{e}_i = \mathbf{0} \quad (71)$$

where $\dim(\mathbf{e}_i) = n$. Similarly, the right eigenvectors of the $\Delta \mathbf{v}$ -system are given by

$$\left[\left(\lambda_i + \frac{1}{\tau} \right) I_{n_r} + RA \right] \tilde{\mathbf{e}}_i = \mathbf{0}, \lambda_i \neq -\frac{1}{\tau} \quad (72)$$

where $\dim(\tilde{\mathbf{e}}_i) = n_r$. Multiplying Eq. (71) from the left with R , it follows that

$$\left[\left(\lambda_i + \frac{1}{\tau} \right) I_{n_r} + RA \right] (R\mathbf{e}_i) = \mathbf{0}, \lambda_i \neq -\frac{1}{\tau}. \quad (73)$$

Hence, if \mathbf{e}_i is a right eigenvector of the $\Delta \mathbf{x}$ -system, \mathbf{e}_i belonging to an eigenvalue $\lambda_i \neq -1/\tau$, then the corresponding right eigenvector of the $\Delta \mathbf{v}$ -system, $\tilde{\mathbf{e}}_i$ belonging to the same eigenvalue, is given by

$$\tilde{\mathbf{e}}_i = R\mathbf{e}_i, i = 1, 2, \dots, n_r. \quad (74)$$

The left row eigenvectors in the $\Delta \mathbf{x}$ -system satisfy the equation:

$$\mathbf{v}_i \left[\left(\lambda_i + \frac{1}{\tau} \right) I_n + AR \right] = \mathbf{0}. \quad (75)$$

Similarly, the left row eigenvectors of the $\Delta \mathbf{v}$ -system satisfy the equation

$$\tilde{\mathbf{v}}_i \left[\left(\lambda_i + \frac{1}{\tau} \right) I_{n_r} + RA \right] = \mathbf{0}, \lambda_i \neq -\frac{1}{\tau}. \quad (76)$$

Multiplying Eq. (76) with R from the right, it follows

$$(\tilde{\mathbf{v}}_i R) \left[\left(\lambda_i + \frac{1}{\tau} \right) I_{n_r} + AR \right] = \mathbf{0}, \lambda_i \neq -\frac{1}{\tau}. \quad (77)$$

In analogy to Eq. (73), it is clear that if $\tilde{\mathbf{v}}_i$ is a left row eigenvector of the $\Delta \mathbf{v}$ -system, $\tilde{\mathbf{v}}_i$ belonging to an eigenvalue $\lambda_i \neq -1/\tau$, then the corresponding left row eigenvector of the $\Delta \mathbf{x}$ -system, \mathbf{v}_i belonging to the same eigenvalue, is given by

$$\tilde{\mathbf{v}}_i R = \mathbf{v}_i, i = 1, 2, \dots, n_r. \quad (78)$$

Note that $\mathbf{e}_i \notin \mathcal{N}(R)$ ($\Rightarrow R\mathbf{e}_i \neq \mathbf{0}$, $\tilde{\mathbf{v}}_i R \neq \mathbf{0}$) is implied by $\lambda_i \neq -1/\tau$. Therefore, only n_r eigenvalues come into this discussion. Furthermore, neither $R\mathbf{e}_i$ nor $\tilde{\mathbf{v}}_i R$ are zero, when $\text{rank}(R) = n_r$.

This is true, however, only when the Jacobian of the reaction rate expressions indicates true independence between the expressions.

8. INVARIANTS AND OBSERVABILITY

Consider the linearized system given by Eq. (63),

$$\dot{\xi}(t) = \left[-\frac{1}{\tau} I - AR(x_s) \right] \xi(t) = F\xi(t). \quad (79)$$

Lemma 1. Given (63), let y_j be an observation of the form $y_j = v_j x$, i.e.

$$y_j = v_j(x_s + \xi) = v_j x_s + \Delta y_j \quad (80)$$

i.e.

$$\Delta y_j = v_j \xi \quad (81)$$

where v_j is a left row eigenvector of F . The rank of the observability matrix O [6] is equal to one, and the system Eq. (63) is not *locally observable* at the point x_s [6].

Proof. To prove this, we do not need to consider the term $v_j x_s$ in y_j , since this is a constant for each chosen x_s . The observability matrix is then

$$O = [v_j^T \quad F^T v_j^T \quad \dots \quad (F^T)^{n-1} v_j^T]. \quad (82)$$

Since v_j is a left eigenvector,

$$v_j F = \lambda_j v_j$$

or

$$F^T v_j^T = \lambda_j v_j^T$$

that is,

$$(F^T)^k v_j^T = \lambda_j^k v_j^T. \quad (83)$$

This means that all v_j is contained in an F^T -invariant subset of E_n . For the observability matrix we have

$$O = [v_j^T \quad \lambda_j v_j^T \quad \dots \quad \lambda_j^{n-1} v_j^T]$$

which obviously has rank equal to one, and the system is not locally observable [6].

Remark. In particular, if $y = Vx$, then y is an asymptotic reaction invariant vector of the nonlinear system, and the system is neither locally nor globally observable.

Theorem 6. For the nonlinear system Eq. (4), given an observation in the neighbourhood of a steady state x_s , such that

$$y = d^T x = d^T [x_s + \xi(t)] \quad (84)$$

i.e.

$$y = d^T \xi \quad (85)$$

where $\xi(t)$ is given by the solution of Eq. (79). If d^T is contained in an F^T -invariant subspace of dimension less than n , then the system is not locally observable at x_s .

Proof. The proof follows easily from Lemma 1. $y = d^T \xi$ yields an observability matrix of rank less than n : since v_j is a basis, there exists numbers β_j , not all equal to zero, $j = 1, 2, \dots, q$, such that

$$d^T = \sum_{j=1}^q \beta_j v_j \quad (86)$$

where $q < n$. Therefore the rank of

$O =$

$$\left[\sum_{j=1}^q \beta_j v_j^T \quad F^T \sum_{j=1}^q \beta_j v_j^T \quad \dots \quad (F^T)^{n-1} \sum_{j=1}^q \beta_j v_j^T \right] \quad (87)$$

obviously has rank $q < n$ at most.

9. THE MINIMAL POLYNOMIAL OF F AND THE SOLUTION OF THE LINEARIZED SYSTEM

From matrix algebra it is known, if a matrix F is semisimple [7], the minimal polynomial of F is

$$r(\lambda) = \Pi'(\lambda - \lambda_i) \quad (88)$$

where Π' denotes the product of all distinct factors. The direct solution of Eq. (79)

$$\xi(t) = e^{Ft} \xi(0) = e^{-(1/\tau + AR)t} \xi(0) \quad (89)$$

may be expressed as

$$\xi(t) = \sum_{i=0}^{n-1} p_i F^i \xi(0) \quad (90)$$

by means of the theorem of Cayley-Hamilton. Assuming that F is semi-simple, F is diagonalizable with eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_{n_r}$ plus $(n - n_r)$ eigenvalues equal to $-1/\tau$. The minimal polynomial is hence at most of order $n_r + 1$. For a matrix function $f(F)$ we obtain

$$f(F) = \sum_{i=1}^q f(\lambda_i) H_i \quad (91)$$

where $q \leq n_r + 1$ and

$$H_i = \prod_{\substack{j=1 \\ j \neq i}}^q \frac{F - \lambda_j I}{\lambda_i - \lambda_j} \quad (92)$$

(Sylvester's Theorem). For the matrix exponential we then obtain

$$e^{Ft} = \sum_{i=1}^q e^{\lambda_i t} H_i \quad (93)$$

or more specifically

$$\xi(t) = [e^{\lambda_1 t} H_1 + e^{\lambda_2 t} H_2 + \dots + e^{\lambda_{n_r} t} H_{n_r} + e^{(-1/\tau)t} H_{n_r+1}] \xi(0). \quad (94)$$

10. CONTROLLABILITY

10.1 Feed flow rate control, constant feed composition

When flow rate control is applied, the mean residence time will change, and Eqs. (7) or (40) may be written

$$\dot{z}(t) = \frac{q(t)}{V} [z_f(t) - z(t)]. \quad (95)$$

It is a well-known feature of a stirred tank without any chemical reactions, that the thermodynamic states are insensitive to flow rate changes when the balance equation takes the form of Eq. (95)[11]. This is so, if the system operates at a steady state in z , as then $z(t) = z_s = z_f$ and $\dot{z} = 0$. Consequently, the subset $R_{n_r}^+$ is uncontrollable for all t .

If, however, $z(0) \neq z_f$, then a certain transient control is possible, via $z(0)$ initially, but gradually $R_{n_r}^+$ becomes uncontrollable as t approaches infinity.

10.2 Feed composition control, constant flow rate


From Eqs. (7) or (40)

$$\dot{z}(t) = \frac{1}{\tau} [z_f(t) - z(t)] \quad (96)$$

where the parameter τ is constant, it is seen that the state z is controllable only via the input variable z_f . Consequently, if the feed is such that $z_f = \mathbf{0}$, i.e. $x_f(t) \in \mathcal{N}(V)^+$, then $R_{n_r}^+$ is an uncontrollable subset for all t .

11. THE STOICHIOMETRIX DOES NOT HAVE FULL RANK

A general model for a set of chemical reactions involves parallel and consecutive mechanisms. Parallel reactions are in principle indicating different possible reaction paths to form the same products from the same original reactants. The stoichiometry along such parallel paths are linear combinations of each other, provided the reactants at the start and the final products at the end of a path are the same for the parallel paths. The rank of the stoichiometric matrix will in this case be less than the number of reactions. For example, the steam reforming of methane may be considered in the first approximation as two apparent parallel reactions.




as far as methane, water vapour and hydrogen is concerned. However, they have linearly independent stoichiometry due to the formation of carbon monoxide and dioxide. If however, the water shift reaction takes place at the same time:



The stoichiometric conditions for this reaction is

just the difference between the other two. This is due to the fact that the shift reaction acts as an intermediate, consecutive reaction in two parallel paths both using methane and water to produce hydrogen and carbon dioxide. The introduction of the shift reaction does not alter the rank of the stoichiometric matrix from 2, as it was before.

Another example on reaction models with several parallel paths, is the hydrogenation of fatty oils from soybeans. The stoichiometry along these paths is shown to be linear combinations of each other. The mechanism is best illustrated by a graph:




as shown by Hertzberg and Asbjørnsen[9]. Here A is linolenic, B linoleic, C *trans*-oleic, D *cis*-oleic and E stearic acid.

As it is stoichiometric there are seven reactions suggested, five components, while the stoichiometric matrix in the equation:

$$Nm = \begin{bmatrix} 1 & -1 & 0 & 0 & 0 \\ 0 & 1 & -1 & 0 & 0 \\ 0 & 1 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 & -1 \\ 0 & 0 & 1 & -1 & 0 \\ 0 & 0 & 1 & 0 & -1 \\ 0 & 0 & 0 & 1 & -1 \end{bmatrix} \begin{bmatrix} A \\ B \\ C \\ D \\ E \end{bmatrix} \rightarrow \mathbf{0} \quad (100)$$

is seen to have the rank four. Let the rows in N be \mathbf{a}_i and the columns \mathbf{b}_i . By developing the echelon matrix, the following relationships appear: $\mathbf{a}_5 = \mathbf{a}_3 - \mathbf{a}_2$, $\mathbf{a}_6 = \mathbf{a}_4 - \mathbf{a}_2$, $\mathbf{a}_7 = \mathbf{a}_4 - \mathbf{a}_3$ and $\mathbf{b}_1 = (\mathbf{b}_2 + \mathbf{b}_3 + \mathbf{b}_4 + \mathbf{b}_5)$. The graph contains three basic elements with parallel paths of linearly dependent stoichiometry. Those are



As shown by Asbjørnsen[10], an asymptotic reaction invariant may be found also in this case, by the procedure of partitioning the material balance equations, as well as the linearly dependent reactions:

$$\begin{aligned} \dot{x}_1 &= \frac{1}{\tau} (x_{1f} - x_1) - [A_{11} \quad A_{12}] \begin{bmatrix} r_1 \\ r_2 \end{bmatrix} \\ \dot{x}_2 &= \frac{1}{\tau} (x_{2f} - x_2) - [A_{21} \quad A_{22}] \begin{bmatrix} r_1 \\ r_2 \end{bmatrix} \end{aligned} \quad A = N^T \quad (102)$$

Here, A_{11} is a square $p \times p$ matrix of rank p , which is also the rank of A . Since all rows in the matrix $[A_{21} \ A_{22}]$ are linear combinations of the rows in the matrix $[A_{11} \ A_{12}]$, it follows that

$$A_{21}A_{11}^{-1}A_{12} - A_{22} = 0. \quad (103)$$

Hence, the asymptotic reaction invariant z is as before in Eq. (6)

$$z = x_2 - A_{21}A_{11}^{-1}x_1. \quad (104)$$

The projection matrix P_1 is then:

$$P_1 = \left[\begin{array}{cccc|ccc} 0 & & & & & & 0 \\ \hline -A_{21}A_{11}^{-1} & & & & & & I \end{array} \right]. \quad (105)$$

For the hydrogenation model, let the matrix A be partitioned:

$$A = \left[\begin{array}{cccc|cccc} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 & -1 & 0 & 0 & 1 \\ 0 & 0 & 0 & -1 & 0 & -1 & -1 & -1 \end{array} \right] \\ = \left[\begin{array}{ccc|ccc} A_{21} & & & A_{22} & & \\ \hline A_{11} & & & A_{12} & & \end{array} \right]. \quad (106)$$

Then the reaction invariant z is seen to be a scalar and expressed through

$$z = x_1 - \begin{bmatrix} -1 & -1 & -1 & -1 \end{bmatrix} \begin{bmatrix} x_2 \\ x_3 \\ x_4 \\ x_5 \end{bmatrix} \\ = x_1 + x_2 + x_3 + x_4 + x_5 \quad (107)$$

This equation expresses z as the sum of all the mole numbers, which naturally has to be reaction invariant in the elementary first order reactions suggested. The projection operator P_1 is:

$$P_1 = \left[\begin{array}{cccc|ccc} 0 & & & & & & \\ \hline 1 & 1 & 1 & 1 & 1 & 1 & 1 \end{array} \right]. \quad (108)$$

For the projection operator P_2 , the procedures in the previous Section 4.2 may still be applied, with some minor modifications in the computation of the pseudoinverse. Of course, the matrix A_1^{-1} in Sections 3-4.1 does not exist now, since $\text{rank}(A_1) < n$, with $n < n_r$.

Suppose that A has rank $p < n$. Suppose that we know a real $n \times p$ matrix B and a real $p \times n_c$ matrix C , both of rank p , having the property

$$A = BC. \quad (109)$$

Then the pseudoinverse of A may be expressed as [5],

$$A^\dagger = C^T(CC^T)^{-1}(B^TB)^{-1}B^T \quad (110)$$

The pseudoinverse A^\dagger is unique.

Considering now $x(t)$ as given by Eq. (25), the projection from E_n onto $\mathcal{N}(A^T) [= \mathcal{R}(A)^\perp]$ is

$$P_2\dot{x}(t) = (I - AA^\dagger)\dot{x}(t) = -Ar + AA^\dagger Ar. \quad (111)$$

A general property of the pseudo-inverse is that [5]:

$$AA^\dagger A = A \quad (112)$$

such that for a batch reaction, we still have

$$\dot{w}(t) \triangleq P_2\dot{x}(t) = 0 \quad (113)$$

where $w \in \mathcal{N}(A^T)$.

The orthogonal projection on $\mathcal{R}(A)$ is still

$$(I - P_2)\dot{x}(t) = A\dot{v}(t) = AA^\dagger\dot{x}(t) = -AA^\dagger Ar \quad (114)$$

i.e.

$$A\dot{v}(t) = -Ar. \quad (115)$$

Projecting again on $\mathcal{R}(A^T) [= \mathcal{N}(A)^\perp]$,

$$\dot{v} = -r(w + Av). \quad (116)$$

This procedure may be illustrated by the hydrogenation model, as introduced above. Let the matrix B be the first 5×4 part of matrix A . Then the rank of B is four (p), and the matrix C follows such that $A = BC$:

$$B = \begin{bmatrix} 1 & 0 & 0 & 0 \\ -1 & 1 & 1 & 1 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{bmatrix} \\ C = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & -1 & -1 & 0 \\ 0 & 0 & 1 & 0 & 1 & 0 & -1 \\ 0 & 0 & 0 & 1 & 0 & 1 & -1 \end{bmatrix} \quad (117)$$

The pseudo-inverse of A is computed from Eq. (30), and found to be:

$$A^\dagger = \frac{1}{20} \begin{bmatrix} 16 & -4 & -4 & -4 & -4 \\ 4 & 4 & -6 & -1 & -1 \\ 4 & 4 & -1 & -6 & -1 \\ 4 & 4 & -1 & -1 & -6 \\ 0 & 0 & 5 & -5 & 0 \\ 0 & 0 & 5 & 0 & -5 \\ 0 & 0 & 0 & 5 & -5 \end{bmatrix} \quad (118)$$

Furthermore:

$$AA^{\dagger} = \frac{1}{5} \begin{bmatrix} 4 & -1 & -1 & -1 & -1 \\ -1 & 4 & -1 & -1 & -1 \\ -1 & -1 & 4 & -1 & -1 \\ -1 & -1 & -1 & 4 & -1 \\ -1 & -1 & -1 & -1 & 4 \end{bmatrix} \quad (119)$$

giving the projection operator P_2

$$P_2 = \frac{1}{5} \begin{bmatrix} 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 \end{bmatrix} \quad (120)$$

which has rank one. As shown before, the rank of P_2 corresponds to the dimension of the reaction invariant z , which in this case is one. Hence, it is shown by this simple illustration how the concepts of reaction rate invariants apply equally well to systems where the stoichiometric matrix does not have full rank. If too many parallel reactions occur

with linearly related stoichiometry, a reaction invariant subset may still be found by the two projection methods suggested.

REFERENCES

- [1] Fjeld M., Doctoral Thesis, The Norwegian Institute of Technology, Division of Automatic Control, 1972.
- [2] Aris R., *Arc. Rational mech. Anal.* 1965 **19** 81.
- [3] Asbjørnsen O. A. and Fjeld M., *Chem. Engng Sci.* 1970 **25** 1627.
- [4] Asbjørnsen O. A., *Chem. Engng Sci.* 1972, **27**, 709.
- [5] Zadeh L. A. and Desoer C. A., *Linear System Theory: The State Space Approach*. McGraw-Hill, New York 1963.
- [6] Lee E. B. and Markus L., *Foundations of Optimal Control Theory*. Wiley, New York 1967.
- [7] Pease III M. C., *Methods of Matrix Algebra*. Academic Press, New York 1965.
- [8] Aris R., *Introduction to the Analysis of Chemical Reactors*. Prentice Hall, Englewood Cliffs, N.J. 1965.
- [9] Hertzberg T. and Asbjørnsen O. A., Paper presented at *Chisa IV*, Prague 1972.
- [10] Asbjørnsen O. A., *Kemian Teollisuus* 1972 **29** 633.
- [11] Asbjørnsen O. A., *Process Dynamics*. Åbo Akademi, Åbo 1974.